

Original Article
Emergency &
Critical Care Medicine



Effects of Emergency Care-related Health Policies during the COVID-19 Pandemic in Korea: a Quasi-Experimental Study

Yun-Suk Pak ,¹ Young Sun Ro ,^{1,2} Se-Hyung Kim ,¹ So-Hyun Han ,¹ Sung-keun Ko ,¹ Taehui Kim ,¹ Young Ho Kwak ,² Tag Heo ,³ and Sungwoo Moon ^{1,4}

OPEN ACCESS

Received: Dec 24, 2020

Accepted: Apr 6, 2021

Address for Correspondence:

Young Sun Ro, MD, DrPH

National Emergency Medical Center, National Medical Center and Department of Emergency Medicine, Seoul National University Hospital, 101 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea.

E-mail: ro.youngsun@gmail.com

Tag Heo, MD, PhD

Department of Emergency Medicine, Chonnam National University Hospital, 42 Jebong-ro, Dong-gu, Gwangju 61469, Republic of Korea.

E-mail: docheo@hanmail.net

© 2021 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Yun-Suk Pak

<https://orcid.org/0000-0002-9617-1757>

Young Sun Ro

<https://orcid.org/0000-0003-3634-9573>

Se-Hyung Kim

<https://orcid.org/0000-0002-8891-3634>

So-Hyun Han

<https://orcid.org/0000-0001-7113-3482>

¹National Emergency Medical Center, National Medical Center, Seoul, Korea

²Department of Emergency Medicine, Seoul National University Hospital, Seoul, Korea

³Department of Emergency Medicine, Chonnam National University Hospital, Gwangju, Korea

⁴Department of Emergency Medicine, Korea University Ansan Hospital, Ansan, Korea

ABSTRACT



Background: The purpose of this study was to review the nationwide emergency care-related health policies during the coronavirus disease 2019 (COVID-19) pandemic disaster in Korea and to analyze the effects of the policies on the safety of patients who visit emergency departments (EDs) during this period.

Methods: This study is a quasi-experiment study. The study population was patients who visited all 402 EDs in Korea between December 31, 2019 and May 13, 2020, using the National Emergency Department Information System (NEDIS) database. The study period was classified into 5 phases according to the level of national crisis warning of infectious disease and the implementation of emergency care-related health policies, and all study phases were 27 days. The primary outcome was in-hospital mortality, and the secondary outcome was length of stay (LOS) in the ED during the COVID-19 outbreak.

Results: The number of ED visits during the study period was 2,636,341, and the in-hospital mortality rate was 1.4%. The number of ED visits decreased from 803,160 in phase 1 to 496,619 in phase 5 during the study period. For in-hospital mortality, the adjusted odds ratio (OR) (95% confidence interval) was 0.77 (0.74–0.79) in phase 5 compared to phase 3. Additionally, by subgroup, the ORs were 0.69 (0.57–0.83) for the patients with acute myocardial infarction and 0.76 (0.67–0.87) for severe trauma in phase 5 compared to phase 3. The ED LOS increased while the number of ED visits decreased as the COVID-19 pandemic progressed, and the ED LOS declined after policy implementation (beta coefficient: -5.3 [-6.5 to -4.2] minutes in phase 5 compared to phase 3).

Conclusion: Implementing appropriate emergency care policies in the COVID-19 pandemic would have contributed to improving the safety of all emergency patients and reducing in-hospital mortality by preventing excessive deaths.

Keywords: COVID-19; Emergency Care Policies; Quasi-experimental Study

Sung-keun Ko <https://orcid.org/0000-0002-0549-2388>Taehui Kim <https://orcid.org/0000-0001-6897-3712>Young Ho Kwak <https://orcid.org/0000-0003-2062-7575>Tag Heo <https://orcid.org/0000-0002-2333-2169>Sungwoo Moon <https://orcid.org/0000-0001-9950-3449>**Disclosure**

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Moon S, Ro YS, Pak YS, Han S, Heo T, Kwak YH. Data curation: Pak YS, Han S, Ko SK, Kim T. Formal analysis: Pak YS, Kim SH. Investigation: Pak YS, Kim SH, Ko SK, Kim T. Methodology: Ro YS, Moon S. Software: Pak YS, Kim SH, Ko SK, Kim T. Supervision: Moon S, Heo T, Kwak YH. Validation: Han S, Ro YS. Visualization: Pak YS. Writing - original draft: Pak YS, Ro YS. Writing - review & editing: Pak YS, Ro YS, Kim SH, Han S, Ko SK, Kim T, Kwak YH, Heo T, Moon S.

INTRODUCTION

Since the first case of novel coronavirus disease 2019 (COVID-19) was reported in China on December 28, 2019, it has spread rapidly around the world.^{1,2} According to World Health Organization (WHO) report, by June 29, a total of 10,021,401 people were infected, and 499,913 people had died.³ Global public health is facing a crisis according to WHO definitions,⁴ and 160,000 new confirmed cases are reported worldwide a day. To mitigate the public health crisis caused by infectious diseases, WHO promotes cross-sectoral linkages and cross-governmental and cross-social integration through the Emergency and Disaster Risk Management for Health guidelines.⁵

The government response system for the public health crisis is different between countries. There are organizations in charge of crisis management for infectious diseases in each country, and several countries have implemented policies, including social or physical distancing, traffic restrictions, closure of schools and workplaces, or lockdown to prevent person-to-person transmission of COVID-19 and to reduce the burden of pandemic disasters.^{6,7} Regarding emergency care, most symptomatic and asymptomatic infected patients can primarily use emergency medical services (EMS). There is a need to isolate infected patients and appropriate emergency care in a timely manner, and it is also important to protect other noninfected emergency patients from infection and to provide adequate treatment according to acuity in disaster situations.⁸⁻¹¹ Therefore, the importance of the EMS system has become more prominent in pandemic disaster situations to protect the safety of society.¹² Several countries have published emergency care-related health policies, such as designating centers for exclusive COVID-19 treatment.¹³ In Korea, the Ministry of Health and Welfare (MoHW) and National Emergency Medical Center (NEMC) implemented emergency care-related policies to respond to the COVID-19 pandemic.

The purpose of this study was to review nationwide emergency care-related health policies during the COVID-19 pandemic disaster in Korea and to analyze the effects of national emergency care-related policies on the safety of patients who visit emergency departments (EDs) during this period.

METHODS**Study design and data source**

This study is a quasi-experimental study that consists of a review of emergency care-related health policies (natural experiments) and a before-and-after design to evaluate the effects of emergency care-related health policies on clinical outcomes for emergency patients during the COVID-19 pandemic.

The study period was classified into 5 phases according to the level of national crisis warning of infectious disease and the implementation of emergency care-related health policies: warning level 1 (Attention) and 2 (Caution) period (study phase 1: December 31, 2019 to January 26, 2020, for 27 days), warning level 3 (Alert) period (study phase 2: January 27 to February 22, 2020, for 27 days), warning level 4 (Serious) and preintervention (study phase 3: February 23 to March 20, 2020, for 27 days), warning level 4 (Serious) and run-in-intervention (study phase 4: March 21 to April 16, 2020, for 27 days), and warning level 4 (Serious) and postintervention (study phase 5: April 17 to May 13, 2020, for 27 days). National

crisis warning levels and related health policies during the COVID-19 pandemic are described in **Supplementary Data 1**.

The National Emergency Department Information System (NEDIS) database was established to measure emergency care qualities and to provide evidence for developing national and federal emergency care policy in 2003. The NEDIS data collected from all 402 nationwide EDs in real time included demographics (such as gender, age, and insurance), symptoms (chief complaints and onset), prehospital (EMS use and treatment and means of transportation), and ED hospital (level of consciousness at presentation, emergency operative procedures, critical care requirement, disposition, hospital stay after admission and final clinical outcomes) information. For data quality management, the NEDIS data should be approved annually by Statistics Korea. NEMC, an administrative agency under the MoHW, is designated as an organization for managing the NEDIS data in accordance with Article 25 of the Emergency Medical Services Act.

Study population

The study population was defined as patients who visited all 402 emergency medical institutions in Korea between December 31, 2019, and May 13, 2020. Patients who visited the ED for nonmedical purposes, such as issuing a medical certificate, were excluded. Cases who were classified as dead on arrival were also excluded.

Study setting

Korea has a tax-based public EMS system operated by the National Fire Agency. Qualified emergency medical technicians provide necessary prehospital emergency care to patients. There were standard operation protocols for 4 major emergency conditions: out-of-hospital cardiac arrest, severe trauma, acute myocardial infarction (AMI), and acute stroke. During the COVID-19 outbreak, the critical protocol for patients with fever ($\geq 37.5^{\circ}\text{C}$) conditions was added.

The MoHW in Korea designed 3 levels of ED according to emergency medical resources and capabilities (including facilities, equipment, and medical staffs) and operated 38 Level 1 EDs, 125 Level 2 EDs, and 239 Level 3 EDs (a total of 402 EDs) in June 2020. A Level 1 ED is designed to accommodate and to provide definite care to serious and severe emergency patients and to prepare and respond to disasters and mass casualty incidents. Level 1 EDs should designate special areas for isolation rooms, areas for severe and acute patients, and areas for pediatric patients. Before this pandemic, there were no standards of isolation beds for the level 2 and level 3 EDs.

The Central Accident Countermeasures Headquarters under the Ministry of Public Administration and Security is in charge of national disaster response in accordance with Article 14 of the Framework Act on the Management of Disasters and Safety. In the case of infectious disease, the Korea Disease Control and Prevention Agency responds to a national crisis warning in accordance with Article 38(2) of the Framework Act on the Management of Disasters and Safety.

National crisis warning level and emergency care-related health policies

On December 31, 2019, a total of 27 cases of unexplained pneumonia were reported in Wuhan, China, and the Korean government started the infection monitoring and surveillance response system (Crisis warning level 1 [Attention], study phase 1). The first COVID-19 case in Korea was confirmed on January 20, 2020 (Crisis warning level 2 [Caution], study

phase 1). With the occurrence of 4 additional confirmed cases on January 27, 2020, the crisis warning level for infectious diseases was raised (Crisis warning level 3 [Alert], study phase 2). The quarantine of Korean immigrants began, and 288 medical centers operated separate COVID-19 screening stations (46 healthcare centers and 242 hospitals). Since some community infections had been identified, the level of national crisis warning had also been raised to the highest level to prevent the spread of COVID-19 infection nationwide on February 23, 2020 (Crisis warning level 4 [Serious] and national emergency care-related policies: preintervention, study phase 3).

Regarding emergency care, there were several shut-down events of the EDs because COVID-19 confirmed cases were detected among the patients admitted to the emergency room with the spread of the COVID-19 pandemic. It became necessary to designate EDs that could manage and treat suspected COVID-19 patients since the emergency care of patients with fever was delayed and the safety of patients was threatened. The MoHW and NEMC established the criteria for the designation of the Severe Emergency Medical Care Centers (SEMC) to treat appropriately suspected and confirmed COVID-19 patients with severe symptoms, to protect other uninfected emergency patients from infection and to provide appropriate care based on their severity; 5 or more isolation rooms with/without negative pressure, separate air-conditioning facilities, isolated areas for febrile patients, and designated triage areas in front of the emergency room. Of these level 1 and 2 EDs, 57 EDs were designated as the SEMC sequentially from Mar 21 to Apr 16, 2020 (Crisis warning level 4 [Serious] and the national emergency care-related policies: run-in-intervention, study phase 4). A total of 57 SEMC were operated since April 17, 2020. The number of isolation units increased from 1,558 to 1,700 after designation of the SEMC, and 49.2% (837 among 1,700) of isolation units belonged to the SEMC. In addition, the information on resources, including available isolation units and beds for suspected patients in the SEMC, was shared in real time through the web-based national emergency care information system (crisis warning level 4 [Serious] and the national emergency care-related policies: postintervention, study phase 5). Emergency care-related health policies during the COVID-19 pandemic are described in **Table 1** and **Supplementary Data 2**.

Outcome variables and statistics

The primary outcome was in-hospital mortality, and the secondary outcome was length of stay (LOS) in the ED during the COVID-19 outbreak. In-hospital mortality was defined as death at the time of discharge from the ED or hospital and did not include hopeless discharge.

Study subgroups were defined as emergency patients with clinically important diagnoses at discharge from the ED and/or hospital among those who visited the ED within 7 days from the onset of the symptoms and who were aged 15 years old or older; AMI included I21.0–I21.9; acute ischemic stroke included I63.0–I63.9; acute hemorrhagic stroke included I60.0–I62.9; severe trauma included any S or T code and ICD-derived injury severity score less than 0.9.¹⁴

The characteristics and outcomes of the study population and subgroups were compared by study phase. The descriptive analysis for the study subject was compared using univariate analysis. The χ^2 test for categorical variables and the Wilcoxon signed-rank test method for nonnormally distributed continuous variables were used.

Multivariable logistic regression analysis and general linear models were conducted to evaluate the effects of the study phase on study outcomes. Adjusted odds ratios (aORs) with

Table 1. The implementation of the policy by crisis alert level

National crisis warning	National public health policies	Emergency care-related policies
Level 1 (December 31, 2019–January 19, 2020)	Initiation of Korean government infection monitoring and surveillance response (December 31)	
Level 2 (January 20–January 26, 2020)	The first confirmed COVID-19 case in Korea was detected (January 20)	
Level 3 (January 27–February 22, 2020)	Operation of the Central Disease Control Headquarters (January 27) Provision of special immigration procedures and quarantine of Korean immigrants (January 28) Operation of the separate COVID-19 screening stations (46 health centers and 242 medical institutions) (January 28) Expansion of the indications for the COVID-19 screening test (February 6) Initiation of “Self-Quarantine Safety Protection App” or “Self-Diagnosis App” to monitor for quarantine (February 6)	Survey and support for operation of the separate COVID-19 screening stations (January 28) Monitoring and sharing information on ED shutdowns (January 29) Establishment of prehospital transport protocols related to COVID-19 (February 3) System development for detecting COVID-19 related patient information (February 5) Implementation of administrative measures to support operating the separate COVID-19 screening station (February 19) Preparation of installation of a 100-bed mobile hospital in Daegu (February 21, operation on March 4)
Level 4 (February 23, 2020–end of the year)	Installation of the Central Disaster and Safety Countermeasure Headquarters headed by the Prime Minister (February 23) Implementation of a policy to increase the negative pressure in isolation units and isolation units in hospitals (February 23) Designation of 43 dedicated COVID-19 hospitals (February 27) Designation of 127 public relief hospitals (February 27) Initiation of the social distancing program (February 29) Designation of the living treatment centers for patients with mild symptoms in each province (March 1) Designation of 67 infectious disease hospitals (March 3) Establishment of the information system on operational isolated units (March 6) Declaration of COVID-19 pandemic (March 12) Establishment of control system for the long-term care facilities (March 20) Strengthened social distancing program (March 22) Introduction of 14-day self-isolation quarantine for all travelers entering Korea (April 1)	Preparation for criteria for the designation of the Severe Emergency Medical Care centers (such as 5 or more isolation beds, individual air-conditioning facilities, isolation device installation, and designated triage areas in front of the emergency room) (March 2) On-site inspection for tentative EDs (March 3) Designation of 57 Severe Emergency Medical Care centers (March 13) Real-time sharing of the information resources of the Severe Emergency Medical Care center (March 17)

COVID-19 = coronavirus disease 2019, ED = emergency department.

95% confidence intervals (CIs) were calculated for in-hospital mortality, and beta coefficients and 95% CIs were calculated for ED LOS.

Regarding the regional variation of COVID-19 outbreak, the subgroup analyses were conducted for 1) Daegu Metropolitan City and Gyeongsangbuk-do (hereafter, Daegu and Gyeongbuk), where cluster infection occurred among a religious group called “Shincheonji” at the time of the initial COVID-19 outbreak, and 2) Other 15 provinces (**Supplementary Tables 1 and 2**). All statistical analyses were performed using SAS software ver. 9.4 (SAS Institute Inc., Cary, NC, USA).

Ethics statement

The present study protocol was reviewed and approved by the Institutional Review Board of National Medical Center (approval No. NMC-2007-026). Informed consent was waived because of the retrospective nature of the study.

RESULTS

Demographics of emergency patients during the COVID-19 pandemic

The number of ED visits during the study period was 2,636,341. Compared to phase 1, the number of ED visits decreased rapidly according to the COVID-19 outbreak: 803,160 (30.5%) in phase 1, 551,903 (20.9%) in phase 2, 377,212 (14.3%) in phase 3, 407,447 (15.5%) in phase 4, and 496,619 (18.8%) in phase 5. The median and interquartile range of ED LOS were 92 (43–162) minutes in phase 1, 95 (43–174) minutes in phase 2, 94 (37–185) minutes in phase 3, 95 (38–188) minutes in phase 4, and 93 (36–183) minutes in phase 5, respectively (*P*-for-trend 0.04). In-hospital mortality rates were 1.4% during the study period, 1.1% in phase 1, 1.4% in phase 2, 1.9% in phase 3, 1.7% in phase 4, and 1.5% in phase 5 (*P*-for-trend < 0.01) (Table 2). The trends of daily ED visits and COVID-19 confirmed patients during the study period are shown in Fig. 1.

The proportion of emergency patients with clinically important diagnoses among ED visits increased significantly (all *P*-for-trends < 0.01). For AMI, the in-hospital mortality rate was 10.0% during the study period (10.1% in phase 1 and 8.3% in phase 5); for acute ischemic stroke, the in-hospital mortality rate was 5.3% (5.3% in phase 1 and 5.0% in phase 5); for acute hemorrhagic stroke, the in-hospital mortality rate was 14.2% (13.3% in phase 1 and 14.9% in phase 5); and for severe trauma, the in-hospital mortality rate was 2.1% (2.1% in phase 1 and 1.9% in phase 5) (all *P*-for-trend < 0.01) (Table 3).

Table 2. Characteristics of the study population according to national crisis warning level and intervention

Variables	Total	Phase 1	Phase 2	Phase 3	Phase 4	Phase 5	<i>P</i> value
Crisis warning level		1 & 2	3	4	4	4	-
Intervention		-	-	Before	Run-in	After	-
Total visits	2,636,341	803,160	551,903	377,212	407,447	496,619	< 0.001
Age, yr							
0–17	374,847 (14.2)	148,744 (18.5)	77,327 (14.0)	40,403 (10.7)	47,126 (11.6)	61,247 (12.3)	< 0.001
18–64	1,619,528 (61.4)	479,554 (59.7)	339,843 (61.6)	240,789 (63.8)	253,287 (62.2)	306,055 (61.6)	< 0.001
65–120	641,927 (24.3)	174,852 (21.8)	134,723 (24.4)	96,014 (25.5)	107,027 (26.3)	129,311 (26.0)	< 0.001
Gender, female	1,286,477 (48.8)	408,335 (50.8)	268,936 (48.7)	178,716 (47.4)	192,685 (47.3)	237,805 (47.9)	< 0.001
Insurance, medicaid	167,370 (6.3)	44,002 (5.5)	35,457 (6.4)	26,115 (6.9)	28,160 (6.9)	33,636 (6.8)	< 0.001
Reason for visits, injury	667,511 (25.3)	161,259 (20.1)	130,586 (23.7)	99,870 (26.5)	123,171 (30.2)	152,625 (30.7)	< 0.001
EMS use	468,894 (17.8)	111,590 (13.9)	99,858 (18.1)	77,977 (20.7)	85,099 (20.9)	94,370 (19.0)	< 0.001
Transferred in	202,481 (7.7)	52,927 (6.6)	46,165 (8.4)	31,680 (8.4)	33,425 (8.2)	38,284 (7.7)	< 0.001
Initial BT, ≥ 37.5°C	290,807 (11.0)	117,187 (14.6)	58,609 (10.6)	35,034 (9.3)	37,006 (9.1)	42,971 (8.7)	< 0.001
Initial triage, KTAS 1 & 2	140,499 (5.3)	34,369 (4.3)	29,429 (5.3)	23,877 (6.3)	25,118 (6.2)	27,706 (5.6)	< 0.001
Triage							
Appropriate	2,438,388 (92.5)	740,668 (92.2)	506,702 (91.8)	348,641 (92.4)	379,214 (93.1)	463,163 (93.3)	< 0.001
Undertriage	70,869 (2.7)	18,414 (2.3)	15,238 (2.8)	10,917 (2.9)	12,131 (3.0)	14,169 (2.9)	< 0.001
Overtriage	34,173 (1.3)	11,294 (1.4)	7,344 (1.3)	4,624 (1.2)	5,030 (1.2)	5,881 (1.2)	< 0.001
ED LOS, min, mean (STD)	163.0 (275.2)	149.0 (256.7)	159.6 (268.0)	174.3 (298.1)	176.8 (291.3)	169.2 (279.2)	< 0.001
ED LOS, min, median (IQR)	93 (40–175)	92 (43–162)	95 (43–174)	94 (37–185)	95 (38–188)	93 (36–183)	< 0.001
ED Disposition							
Discharge	2,055,396 (78.0)	660,693 (82.3)	428,238 (77.6)	282,132 (74.8)	304,020 (74.6)	380,313 (76.6)	< 0.001
Transfer out	45,243 (1.7)	11,913 (1.5)	10,107 (1.8)	7,201 (1.9)	7,580 (1.9)	8,442 (1.7)	< 0.001
Admission	504,765 (19.1)	124,632 (15.5)	107,515 (19.5)	80,978 (21.5)	89,796 (22.0)	101,844 (20.5)	< 0.001
Death	12,375 (0.5)	2,567 (0.3)	2,401 (0.4)	2,483 (0.7)	2,478 (0.6)	2,446 (0.5)	< 0.001
Unknown	18,562 (0.7)	3,355 (0.4)	3,642 (0.7)	4,418 (1.2)	3,573 (0.9)	3,574 (0.7)	< 0.001
ICU Admission	77,874 (3.0)	17,142 (2.1)	16,346 (3.0)	13,846 (3.7)	14,918 (3.7)	15,622 (3.1)	< 0.001
Operation	30,423 (1.2)	6,552 (0.8)	6,161 (1.1)	5,507 (1.5)	5,976 (1.5)	6,227 (1.3)	< 0.001
In-hospital mortality, total	37,982 (1.4)	8,496 (1.1)	7,845 (1.4)	7,170 (1.9)	7,128 (1.7)	7,343 (1.5)	< 0.001
ED	12,375 (0.5)	2,567 (0.3)	2,401 (0.4)	2,483 (0.7)	2,478 (0.6)	2,446 (0.5)	< 0.001
Ward	25,607 (1.0)	5,929 (0.7)	5,444 (1.0)	4,687 (1.2)	4,650 (1.1)	4,897 (1.0)	< 0.001

Data are presented as number (%).

EMS = emergency medical services, BT = body temperature, KTAS = Korean triage and acuity scale, ED = emergency department, LOS = length of stay, STD = standard deviation, IQR = interquartile range, ICU = intensive care unit.




Fig. 1. Daily number of ED visits and confirmed COVID-19 patients during the study period. ED = emergency department, COVID-19 = coronavirus disease 2019.

Table 3. Characteristics of patients with clinically important diagnoses according to the study phase

Variables	Total	Phase 1	Phase 2	Phase 3	Phase 4	Phase 5	P value
Crisis warning level		1 & 2	3	4	4	4	-
Intervention		-	-	Before	Run-in	After	-
Total visits	2,636,341	803,160	551,903	377,212	407,447	496,619	-
Acute myocardial infarction							
Total	13,778 (0.5)	3,131 (0.4)	3,011 (0.5)	2,257 (0.6)	2,533 (0.6)	2,846 (0.6)	< 0.001
In-hospital mortality	1,371 (10.0)	317 (10.1)	299 (9.9)	262 (11.6)	256 (10.1)	237 (8.3)	< 0.001
ICU admission	7,878 (57.2)	1,725 (55.1)	1,684 (55.9)	1,342 (59.5)	1,528 (60.3)	1,599 (56.2)	< 0.001
ED LOS, min, mean (STD)	318.6 (444.0)	312.9 (436.2)	291.8 (444.2)	308.2 (428.1)	327.0 (434.9)	353.9 (469.9)	< 0.001
ED LOS, min, median (IQR)	161 (71-356)	166 (75-343)	153 (69-316)	149 (66-349)	162.5 (67-378)	172 (75-421)	< 0.001
Ischemic stroke							
Total	27,169 (1.0)	6,114 (0.8)	5,578 (1.0)	4,749 (1.3)	5,071 (1.2)	5,657 (1.1)	< 0.001
In-hospital mortality	1,428 (5.3)	324 (5.3)	277 (5.0)	271 (5.7)	273 (5.4)	283 (5.0)	< 0.001
ICU admission	7,250 (26.7)	1,572 (25.7)	1,460 (26.2)	1,294 (27.2)	1,426 (28.1)	1,498 (26.5)	< 0.001
ED LOS, min, mean (STD)	306.7 (387.1)	290.5 (384.5)	273.8 (337.7)	309.5 (411.7)	325.6 (396.9)	337.2 (401.8)	< 0.001
ED LOS, min, median (IQR)	196 (124-328)	192 (124-310)	189 (120-300)	189.5 (121-316)	203 (125-356)	209 (130-382)	< 0.001
Hemorrhagic stroke							
Total	11,460 (0.4)	2,505 (0.3)	2,437 (0.4)	2,046 (0.5)	2,213 (0.5)	2,259 (0.5)	< 0.001
In-hospital mortality	1,633 (14.2)	333 (13.3)	338 (13.9)	292 (14.3)	333 (15.0)	337 (14.9)	< 0.001
ICU admission	7,554 (65.9)	1,604 (64.0)	1,616 (66.3)	1,394 (68.1)	1,482 (66.7)	1,458 (54.5)	< 0.001
ED LOS, min, mean (STD)	259.3 (398.3)	231.8 (381.2)	235.9 (352.0)	261.7 (401.4)	283.1 (415.0)	289.9 (439.1)	< 0.001
ED LOS, min, median (IQR)	143 (92-249)	138 (88-232)	135 (90-232)	142 (93-245)	146 (94-267)	155 (95-278)	< 0.001
Severe trauma, ICISS < 0.90							
Total	111,345 (4.2)	25,641 (3.2)	22,729 (4.1)	17,851 (4.7)	20,929 (5.1)	24,195 (4.9)	< 0.001
In-hospital mortality	2,341 (2.1)	527 (2.1)	488 (2.1)	444 (2.5)	430 (2.1)	452 (1.9)	< 0.001
ICU admission	10,804 (9.7)	2,236 (8.7)	2,204 (9.7)	1,935 (10.8)	2,181 (10.4)	2,248 (9.3)	< 0.001
ED LOS, min, mean (STD)	199.3 (281.1)	190.6 (273.4)	188.0 (276.1)	208.7 (299.2)	210.9 (288.1)	201.9 (273.3)	< 0.001
ED LOS, min, median (IQR)	117 (63-224)	114 (61-214)	114 (62-213)	121 (65-229)	121 (64-236)	117 (63-231)	< 0.001

Data are presented as number (%).

ICU = intensive care unit, ED = emergency department, LOS = length of stay, STD = standard deviation, IQR = interquartile range.

Effects of emergency care interventions on safeties for emergency patients

For in-hospital mortality among the study population, compared to the study phase 3, the adjusted ORs (95% CIs) were 0.63 (0.61-0.65) for phase 1, 0.77 (0.74-0.79) for phase 2, 0.90 (0.87-0.93) for phase 4, and 0.77 (0.74-0.79) for phase 5. By subgroup for region, for the patients in Daegu and Gyeongbuk, the adjusted ORs (95% CIs) were 0.78 (0.71-0.87) for phase 4 and 0.57 (0.51-0.64) for phase 5 compared to phase 3, and for other 15 provinces, the

Table 4. Adjusted odds ratios and 95% CIs for in-hospital mortality for the study population

Variables	Phase 1	Phase 2	Phase 3	Phase 4	Phase 5
Crisis warning level	1 & 2	3	4	4	4
Intervention	-	-	Before	Run-in	After
Total	0.63 (0.61–0.65)	0.77 (0.74–0.79)	1.00	0.90 (0.87–0.93)	0.77 (0.74–0.79)
Region					
Daegu and Gyeongbuk	0.49 (0.44–0.54)	0.66 (0.60–0.73)	1.00	0.78 (0.71–0.87)	0.57 (0.51–0.64)
Other 15 provinces	0.66 (0.64–0.69)	0.79 (0.76–0.82)	1.00	0.93 (0.90–0.97)	0.81 (0.78–0.84)
Clinically important diagnosis					
Acute myocardial infarction	0.82 (0.69–0.98)	0.82 (0.69–0.98)	1.00	0.86 (0.71–1.03)	0.69 (0.57–0.83)
Ischemic stroke	0.90 (0.76–1.07)	0.85 (0.71–1.01)	1.00	0.93 (0.71–1.11)	0.86 (0.72–1.02)
Hemorrhagic stroke	0.92 (0.78–1.09)	0.97 (0.82–1.15)	1.00	1.07 (0.90–1.27)	1.07 (0.91–1.27)
Severe trauma	0.85 (0.74–0.96)	0.88 (0.77–1.00)	1.00	0.83 (0.73–0.95)	0.76 (0.67–0.87)

Data are presented as adjusted odds ratios (95% CI).

Odds ratios were adjusted for age and gender.

CI = confidence interval.

Table 5. Multivariable regression analysis for length of stay in emergency departments (minutes) for the study population

Variables	Phase 1	Phase 2	Phase 3	Phase 4	Phase 5
Crisis warning level	1 & 2	3	4	4	4
Intervention	-	-	Before	Run-in	After
Total	-18.1 (-19.2 to -17.1)	-12.3 (-13.5 to -11.2)	Ref.	1.7 (0.52 to 2.9)	-5.3 (-6.5 to -4.2)
Region					
Daegu and Gyeongbuk	-36.4 (-42.8 to -30.0)	-17.1 (-23.9 to -10.3)	Ref.	6.2 (-1.1 to 13.5)	2.0 (-5.0 to 9.0)
Other 15 provinces	-14.6 (-15.8 to -13.5)	-11.6 (-12.9 to -10.3)	Ref.	1.9 (0.6 to 3.2)	-4.6 (-5.9 to -3.3)
Clinically important diagnosis					
Acute myocardial infarction	2.0 (-21.9 to 25.9)	-17.5 (-41.7 to 6.6)	Ref.	19.2 (-5.9 to 44.2)	45.8 (21.3 to 70.2)
Ischemic stroke	-19.9 (-34.5 to -5.2)	-36.2 (-51.1 to -21.2)	Ref.	15.6 (0.3 to 30.9)	27.3 (12.4 to 42.2)
Hemorrhagic stroke	-30.8 (-53.9 to -7.5)	-26.6 (-50.0 to -3.2)	Ref.	20.8 (-3.1 to 44.7)	28.5 (4.7 to 52.3)
Severe trauma	-18.0 (-23.3 to -12.7)	-20.4 (-25.8 to -14.9)	Ref.	3.1 (-2.4 to 8.7)	-5.7 (-11.0 to -0.3)

Data are presented as β coefficients (95% CI).

Beta coefficients were adjusted for age and gender.

CI = confidence interval.

adjusted ORs (95% CIs) were 0.93 (0.90–0.97) for phase 4 and 0.81 (0.78–0.84) for phase 5 compared to phase 3. By subgroup for diagnosis, for the patients with AMI, the adjusted ORs (95% CIs) were 0.69 (0.57–0.83) for phase 5 compared to phase 3, and for severe trauma, the adjusted ORs (95% CIs) were 0.83 (0.73–0.95) and 0.76 (0.67–0.87) for phase 4 and phase 5, respectively, compared to phase 3. For acute ischemic stroke and hemorrhagic stroke, the effect measures were not significant during any of the study phases (Table 4).

For ED LOS among the study population, compared to the study phase 3, the adjusted coefficients (95% CIs) were -18.1 (-19.2 to -17.1) minutes for phase 1, -12.3 (-13.5 to -11.2) minutes for phase 2, 1.7 (0.52 to 2.9) minutes for phase 4, and -5.3 (-6.5 to -4.2) minutes for phase 5. By subgroup analysis, for severe trauma, the ED LOS decreased by -5.7 (-11.0 to -0.3) minutes in phase 5 compared to phase 3. Compared to phase 3, the ED LOS in phase 5 was increased by 45.8 (21.3 to 70.2) minutes for the patients with AMI, 27.3 (12.4 to 42.2) minutes for acute ischemic stroke, and 28.5 (4.7 to 52.3) minutes for acute hemorrhagic stroke (Table 5).

DISCUSSION

Using a nationwide emergency patient database, we investigated the effects of emergency care-related health policies on the safety and clinical outcomes of emergency patients during the COVID-19 pandemic. In Korea, emergency care-related health policies aimed to treat

suspected and confirmed COVID-19 patients with severe symptoms appropriately, to protect other uninfected emergency patients from infection and to provide appropriate care based on their severity in disaster situations. The number of ED visits during the study period decreased from 803,160 in phase 1 to 496,619 in phase 5. The in-hospital mortality rate increased from 1.1% in phase 1 to 1.9% in phase 3, as the COVID-19 pandemic progressed, and decreased to 1.5% in phase 5 after policy implementation (adjusted OR [95% CI], 0.77 (0.74–0.79) in phase 5 compared to phase 3). The ED LOS increased while the number of ED visits decreased as the COVID-19 pandemic progressed, and the ED LOS declined after policy implementation (beta coefficient: –5.3 minutes in phase 5 compared to phase 3). Implementing appropriate emergency care policies in the COVID-19 pandemic would have contributed to improving the safety of all emergency patients and reducing in-hospital mortality by preventing excessive deaths for not only for infected patients but also for uninfected patients.

During the COVID-19 pandemic, the number of emergency visits decreased, but the proportion of emergency patients with clinically important diagnoses among ED visits increased significantly. In previous studies, the number of medical visits decreased, but mortality rates in patients with specific diseases, including cancer, increased during the COVID-19 pandemic.^{15,16} These phenomena were explained by several hypotheses; because of the risk and fear of transmission of COVID-19, the patients with acute emergency symptoms would be hesitant to visit the ED, even if the patient needed immediate emergency care, which may have increased the time delay before medical contact and delayed treatment.¹⁷ In addition, the lack of medical resources in the ED to treat emergency patients other than COVID-19 would be one of the possibilities. Because the treatment of COVID-19 patients requires more medical resources, in a pandemic disaster situation, there may be insufficient space and resources to provide adequate care to other emergency patients, which can increase in-hospital mortality and threaten safety.¹⁸ For this reason, public health policies must be implemented to ensure adequate emergency medical use for severe emergency disease patients, even in the COVID-19 pandemic situation.

In this study, implementation of emergency care-related health policies increased the number of isolation units and ICU beds in SEMC and enabled the web-based real-time sharing of resource information. The in-hospital mortality rates for all emergency patients decreased after policy implementation (phase 5 compared to phase 3), for not only Daegu and Gyeongbuk but also other 15 provinces. However, the effects of health policies varied among subgroups of emergency patients with clinically important diagnoses. Despite the designation of the SEMC, ED LOS increased in patients with AMI and stroke, probably due to the rapid increase in the number of COVID-19 confirmed patients with severe illness leading to a shortage of ICU beds. The mortality rates of emergency patients with clinically important diagnoses increased during the COVID-19 pandemic¹⁹⁻²² and the length of hospital stay for patients with the diseases increased²³⁻²⁶ in several countries with no relevant health policies. The in-hospital mortality rates of AMI and severe trauma, which increased during the early phase of COVID-19 outbreak, could be reduced by implementing the emergency care-related health policies.

During the COVID-19 pandemic, the mortality rates of confirmed cases were different for each country, from over 16% in France to less than 0.3% in Singapore.^{27,28} Such broad variation implies that there are factors, such as government response, other than patient characteristics that determine COVID-19 mortality.²⁹ Effective government policies responded

to the COVID-19 pandemic proactively to ensure a sufficient supply of personal protective equipment, provision of good public health services in treating infected patients, and quick implementation of screening programs.^{30,31} In addition, emergency care-related health policies could address not only reduced accessibility to healthcare services but also a shortage of human and economic resources that strain the healthcare system for uninfected emergency patients. Korea was the second-largest country with confirmed COVID-19 patients, which was due to the rapid increase in COVID-19 patients in March 2020, but it is evaluated as one of the countries that is currently successfully carrying out COVID-19 prevention with a strong preemptive response. Since the ED is a place where many patients with various diseases are concentrated, it is necessary to establish a system to respond properly to disaster situations. The emergency care-related health policies, including the designation of 57 SEMC, expansion of isolation units, and real-time sharing of resource information of SEMC on the NEMC portal (portal.nemc.or.kr), would have allowed timely provision of appropriate care and quick determination of ED disposition after acute care for not only for infected patients but also for uninfected patients. As the COVID-19 pandemic is currently underway, it is necessary to analyze further the impact of emergency care-related health policies in the COVID-19 pandemic on the safety of all emergency patients after the end of the pandemic. Our findings have important implications for implementing emergency care policies in the potential risk of chronic COVID-19 pandemics or other communicable disease outbreaks.

The limitations of this study are as follows. First, this is a quasi-experimental study and does not use a randomization controlled trial. Types of selection bias that can occur in quasi-experimental studies include maturation bias and instrumentation bias. These bias effects may have different outcomes by country and circumstance. Second, study subgroups were defined by the Korean Standard Classification of Diseases-7 code. Misclassification bias in subgroups was possibly present in studies. Therefore, it is necessary to conduct detailed studies on changes before and after the COVID-19 outbreak through future research. As the COVID-19 pandemic is currently underway, it is necessary to analyze further the impact of emergency care-related health policies on the safety of patients after the end of the communicable disease pandemic.

During the COVID-19 pandemic, implementing emergency care-related health policies would contribute to improving the safety of all emergency patients and reducing in-hospital mortality by preventing excessive deaths in critically ill patients visiting EDs. Implementing a multidisciplinary approach and evidence-based policy in a disaster situation will be the driving force to protect the safety and lives of the people despite the ongoing threat.

SUPPLEMENTARY MATERIALS

Supplementary Data 1

National crisis warning and health policies during the COVID-19 pandemic

[Click here to view](#)

Supplementary Data 2

National experiments: Emergency care-related health policies during the COVID-19 pandemic

[Click here to view](#)

Supplementary Table 1

Characteristics of the study population in Daegu and Gyeongbuk according to national crisis warning level and intervention

[Click here to view](#)

Supplementary Table 2

Characteristics of the study population in other 15 provinces according to national crisis warning level and intervention

[Click here to view](#)

REFERENCES

1. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061-9.
[PUBMED](#) | [CROSSREF](#)
2. Stawicki SP, Jeanmonod R, Miller AC, Paladino L, Gaieski DF, Yaffee AQ, et al. The 2019–2020 novel coronavirus (severe acute respiratory syndrome coronavirus 2) pandemic: a joint American College of Academic International Medicine-World Academic Council of Emergency Medicine Multidisciplinary COVID-19 Working Group consensus paper. *J Glob Infect Dis* 2020;12(2):47-93.
[PUBMED](#) | [CROSSREF](#)
3. World Health Organization. Coronavirus disease (COVID-19) pandemic. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Updated 2020. Accessed July 10, 2020.
4. World Health Organization. International health regulations (second edition). <http://www.who.int/ihr/>. Updated 2020. Accessed July 10, 2020.
5. World Health Organization. Health emergency and disaster risk management framework. <https://www.who.int/hac/techguidance/preparedness/health-emergency-and-disaster-risk-management-framework-eng.pdf>. Updated 2019. Accessed July 10, 2020.
6. Hale T, Petherick A, Phillips T, Webster S. *Variation in Government Responses to COVID-19. Blavatnik School of Government Working Paper. No. 31*. Oxford, England: University of Oxford; 2020.
7. Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan, China. *JAMA* 2020;323(19):1915-23.
[PUBMED](#) | [CROSSREF](#)
8. Carlsson AC, Wessman T, Larsson A, Leijonberg G, Tofik R, Ärnlov J, et al. Endostatin predicts mortality in patients with acute dyspnea - a cohort study of patients seeking care in emergency departments. *Clin Biochem* 2020;75:35-9.
[PUBMED](#) | [CROSSREF](#)
9. Rahmatinejad Z, Tohidinezhad F, Reihani H, Rahmatinejad F, Pourmand A, Abu-Hanna A, et al. Prognostic utilization of models based on the APACHE II, APACHE IV, and SAPS II scores for predicting in-hospital mortality in emergency department. *Am J Emerg Med* 2020;38(9):1841-6.
[PUBMED](#) | [CROSSREF](#)
10. Murao S, Yamakawa K, Kabata D, Kinoshita T, Umemura Y, Shintani A, et al. Effect of earlier door-to-CT and door-to-bleeding control in severe blunt trauma: a retrospective cohort study. *Res Sq*. Forthcoming 2020. DOI: 10.21203/rs.3.rs-23452/v1.
[CROSSREF](#)
11. World Health Organization. The regulation and management of international emergency medical teams. <https://www.ifrc.org/Global/Publications/IDRL/newsletters/December%202015/EMT%20Report%20HR.pdf>. Updated 2017. Accessed July 10, 2020.
12. Cabañas JG, Williams JG, Gallagher JM, Brice JH. COVID-19 pandemic: the role of EMS physicians in a community response effort. *Prehosp Emerg Care* 2021;25(1):8-15.
[PUBMED](#) | [CROSSREF](#)
13. Boyarsky BJ, Po-Yu Chiang T, Werbel WA, Durand CM, Avery RK, Getsin SN, et al. Early impact of COVID-19 on transplant center practices and policies in the United States. *Am J Transplant* 2020;20(7):1809-18.
[PUBMED](#) | [CROSSREF](#)

14. Osler T, Rutledge R, Deis J, Bedrick E. ICISS: an international classification of disease-9 based injury severity score. *J Trauma* 1996;41(3):380-6.
[PUBMED](#) | [CROSSREF](#)
15. Maringe C, Spicer J, Morris M, Purushotham A, Nolte E, Sullivan R, et al. The impact of the COVID-19 pandemic on cancer deaths due to delays in diagnosis in England, UK: a national, population-based, modelling study. *Lancet Oncol* 2020;21(8):1023-34.
[PUBMED](#) | [CROSSREF](#)
16. Jeffery MM, D'Onofrio G, Paek H, Platts-Mills TF, Soares WE 3rd, Hoppe JA, et al. Trends in emergency department visits and hospital admissions in health care systems in 5 states in the first months of the COVID-19 pandemic in the US. *JAMA Intern Med* 2020;180(10):1328-33.
[PUBMED](#) | [CROSSREF](#)
17. Won LE, Hawkins JE, Langness S, Murrell KL, Iris P, Sammann A. Where are all the patients? Addressing Covid-19 fear to encourage sick patients to seek emergency care. *NEJM Catal*. Forthcoming 2020.
18. Birkmeyer JD, Barnato A, Birkmeyer N, Bessler R, Skinner J. The impact of the COVID-19 pandemic on hospital admissions in the United States. *Health Aff (Millwood)* 2020;39(11):2010-7.
[PUBMED](#) | [CROSSREF](#)
19. Egol KA, Konda SR, Bird ML, Dedhia N, Landes EK, Ranson RA, et al. Increased mortality and major complications in hip fracture care during the COVID-19 pandemic: a New York City perspective. *J Orthop Trauma* 2020;34(8):395-402.
[PUBMED](#) | [CROSSREF](#)
20. De Rosa S, Spaccarotella C, Basso C, Calabrò MP, Curcio A, Filardi PP, et al. Reduction of hospitalizations for myocardial infarction in Italy in the COVID-19 era. *Eur Heart J* 2020;41(22):2083-8.
[PUBMED](#)
21. Wu J, Mamas M, Rashid M, Weston C, Hains J, Luescher T, et al. Patient response, treatments and mortality for acute myocardial infarction during the COVID-19 pandemic. *Eur Heart J Qual Care Clin Outcomes*. Forthcoming 2020. DOI: 10.1093/ehjqcc/qcaa062.
[CROSSREF](#)
22. Meza HT, Lambea Gil Á, Saldaña AS, Martínez-Zabaleta M, Juez PR, Martínez EL, et al. Impact of COVID-19 outbreak on ischemic stroke admissions and in-hospital mortality in North-West Spain. *Int J Stroke* 2020;15(7):755-62.
[PUBMED](#) | [CROSSREF](#)
23. Aldujeli A, Hamadeh A, Briedis K, Tecson KM, Rutland J, Krivickas Z, et al. Delays in presentation in patients with acute myocardial infarction during the COVID-19 pandemic. *Cardiol Rev* 2020;11(6):386-91.
[PUBMED](#) | [CROSSREF](#)
24. de Havenon A, Ney JP, Callaghan B, Delic A, Hohmann S, Shippey E, et al. Impact of COVID-19 on outcomes in ischemic stroke patients in the United States. *J Stroke Cerebrovasc Dis* 2021;30(2):105535.
[PUBMED](#) | [CROSSREF](#)
25. Benussi A, Premi E, Pilotto A, Libri I, Pezzini A, Paolillo C, et al. Effects of COVID-19 outbreak on stroke admissions in Brescia, Lombardy, Italy. *Eur J Neurol* 2021;28(1):e4-5.
[PUBMED](#) | [CROSSREF](#)
26. Devarakonda AK, Wehrle CJ, Chibane FL, Drevets PD, Fox ED, Lawson AG. The effects of the COVID-19 pandemic on trauma presentations in a level one trauma center. *Am Surg*. Forthcoming 2020. DOI: 10.1177/0003134820973715.
[CROSSREF](#)
27. Fouillet A, Pontais I, Caserio-Schönemann C. Excess all-cause mortality during the first wave of the COVID-19 epidemic in France, March to May 2020. *Euro Surveill* 2020;25(34):2001485.
28. Fouda A, Mahmoudi N, Moy N, Paolucci F. The COVID-19 pandemic in Greece, Iceland, New Zealand, and Singapore: health policies and lessons learned. *Health Policy Technol* 2020;9(4):510-24.
[PUBMED](#) | [CROSSREF](#)
29. Liang LL, Tseng CH, Ho HJ, Wu CY. Covid-19 mortality is negatively associated with test number and government effectiveness. *Sci Rep* 2020;10(1):1.
[PUBMED](#) | [CROSSREF](#)
30. Flaxman S, Mishra S, Gandy A, Unwin HJ, Mellan TA, Coupland H, et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020;584(7820):257-61.
[PUBMED](#) | [CROSSREF](#)
31. Tian H, Liu Y, Li Y, Wu CH, Chen B, Kraemer MU, et al. An investigation of transmission control measures during the first 50 days of the COVID-19 epidemic in China. *Science* 2020;368(6491):638-42.
[PUBMED](#) | [CROSSREF](#)