

Internet-based therapy versus face-to-face therapy for alcohol use disorder, a randomized controlled non-inferiority trial

Magnus Johansson¹ , Kristina Sinadinovic^{2,3} , Mikael Gajecki^{2,3} , Philip Lindner^{2,3} , Anne H. Berman^{2,3,4} , Ulric Hermansson^{2,3}  & Sven Andréasson¹

Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden,¹ Centre for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden,² Stockholm Health Care Services, Stockholm County Council, Stockholm, Sweden³ and Department of Psychology, Uppsala University, Uppsala, Sweden⁴

ABSTRACT

Background and aims Most people with alcohol use disorder (AUD) are never treated. Internet-based interventions are effective in reducing alcohol consumption and could help to overcome some of the barriers to people not seeking or receiving treatment. The aim of the current study was to compare internet-delivered and face-to-face treatment among adult users with AUD. **Design** Randomized controlled non-inferiority trial with a parallel design, comparing internet-delivered cognitive-behavioural therapy (ICBT) ($n = 150$) with face-to-face CBT ($n = 151$), at 3- and 6-month follow-ups. **Setting** A specialized clinic for people with AUD in Stockholm, Sweden. Participants were recruited between 8 December 2015 and 5 January 2018. **Participants** A total of 301 patients [mean age 50 years, standard deviation (SD) = 12.3] with AUD, of whom 115 (38%) were female and 186 (62%) were male. **Intervention and comparator** Participants were randomized in blocks of 20 at a ratio of 1 : 1 to five modules of therapist-guided ICBT or to five modules of face-to-face CBT, delivered over a 3-month period. The same treatment material and the same therapists were used in both groups. **Measurements** The primary outcome was standard drinks of alcohol consumed during the previous week at 6-month follow-up, analysed according to intention-to-treat. The pre-specified non-inferiority limit was five standard drinks of alcohol and $d = 0.32$ for secondary outcomes. **Results** The difference in alcohol consumption between the internet and the face-to-face group was non-inferior in the intention-to-treat analysis of data from the 6-month follow-up [internet = 12.33 and face-to-face = 11.43, difference = 0.89, 95% confidence interval (CI) = -1.10 to 2.88]. The secondary outcome, Alcohol Use Disorder Identification Test score, failed to show non-inferiority of internet compared with face-to-face in the intention-to-treat analysis at 6-month follow-up (internet = 12.26 and face-to-face = 11.57, $d = 0.11$, 95% CI = -0.11 to 0.34). **Conclusions** Internet-delivered treatment was non-inferior to face-to-face treatment in reducing alcohol consumption among help-seeking patients with alcohol use disorder but failed to show non-inferiority on some secondary outcomes.

Keywords Alcohol, alcohol use disorder, cognitive behaviour therapy, internet intervention, non-inferiority, randomized controlled trial, treatment.

Correspondence to: Magnus Johansson, Department of Global Public Health, Karolinska Institutet, Riddargatan 1 – mottagningen för alkohol och hälsa, Riddargatan 1. SE-114 35 Stockholm 171 77, Sweden. E-mail: magnus.johansson.1@ki.se

Submitted 11 March 2020; initial review completed 8 June 2020; final version accepted 11 September 2020

INTRODUCTION

Background

Alcohol use is a leading risk factor for premature death and disability-adjusted life-years. Among the population aged 15–49 years, alcohol use was the leading risk factor in 2016, accounting for nearly 10% of global deaths. A

dose-response relationship exists between alcohol use and many major diseases [1]. Alcohol use disorder (AUD) is characterized by impaired control and continued heavy use, despite these known negative consequences [2]. The prevalence of AUD is estimated to be 4.9% world-wide and 6.1% in western Europe [3]. Effective treatment options are available, both psychological and

pharmacological, that can help an individual reduce their alcohol consumption [4,5]. However, of all mental disorders, alcohol and substance use disorders have the widest treatment gap between the number of individuals affected and the number in treatment, with only approximately 7% estimated to receive treatment [6]. Shame, stigma and poor access to attractive treatment options are common reasons why people with AUDs do not receive treatment [7].

Digital interventions for problematic alcohol use have been developed for access via computers and the internet during the last 30 years. Meta-analyses have shown that internet interventions yield significantly better effects in terms of reduced alcohol consumption compared to control groups [8,9]. The majority of studies on digital interventions for alcohol have been conducted on brief single-session interventions aimed at students or similar risk populations [10]. Extended digital interventions, aimed at people with AUDs, are intended to be used over a number of weeks and are often based on treatment principles such as cognitive-behavioural therapy (CBT) and/or motivational interviewing (MI) [11]. An individual patient meta-analysis has reported a significant effect of guided internet interventions for alcohol, with a reduction of 10 weekly standard units [95% confidence intervals (CI) = 5.4–14.6, $P = 0.001$] compared to controls [9].

Although several studies have shown that digital interventions are effective in reducing alcohol use, it is not clear how effective they are compared to face-to-face interventions. A recent Cochrane Review of internet interventions found no evidence of a difference in effect on alcohol consumption between internet and face-to-face intervention arms in five trials of brief interventions targeting college students' alcohol use [8]. Research on other psychiatric and somatic disorders has suggested that internet interventions that are therapist-guided have effects comparable to those of face-to-face interventions [12]. Previous studies conducted in clinical settings have tested digital interventions for alcohol as an alternative to treatment as usual (TAU) [13–15] or as an addition to TAU [15–17]. A recent meta-analysis found that when digital CBT for alcohol was tested as an addition to TAU, compared with TAU only, the effect size from seven studies was positive and significant. It also found that the pooled effect size from two studies comparing digital CBT for alcohol use to TAU did not suggest superior efficacy [18].

In order to investigate whether a new treatment modality such as internet-delivered treatment is actually no less effective than an established treatment, a non-inferiority or equivalence design is needed. Such a design can also help to avoid ethical problems with using placebo or inactive control groups in the evaluation of a new treatment for a serious condition such as AUD, where there is already evidence of efficacy for an existing standard treatment. The

use of a non-inferiority design can be problematic, however, if the new treatment does not offer other advantages compared to an already established treatment [19]. Possible advantages of internet-delivered treatment for alcohol use disorders include overcoming problems with cost, accessibility and stigma in traditional face-to-face treatment [20]. One previous non-inferiority study of an internet intervention for alcohol has been conducted in primary care, with the proportion of hazardous drinkers in each group as the primary outcome. The study failed to show non-inferiority for facilitated access to an internet self-help programme compared to a face-to-face brief intervention at 3- and 12-month follow-ups, when a biased outcome measure was removed from the analysis [21].

The purpose of this randomized controlled non-inferiority trial was to investigate the effects of internet-delivered CBT, compared to face-to-face CBT, among adult users with AUD at a specialized clinic for AUDs. The main hypothesis of the trial was that internet-delivered CBT for reducing alcohol use would not be less effective in reducing alcohol consumption, in terms of standard units consumed per week, compared to face-to-face CBT. The margin for the difference between treatments, within which internet-delivered therapy would be considered non-inferior, was set to five standard units of alcohol consumed per week.

METHODS

Study design

The present study was a two-armed randomized controlled non-inferiority trial, with a parallel design. Participants were randomly assigned in blocks of 20 at a ratio of 1 : 1 to internet-delivered CBT or to face-to-face CBT.

Recruitment

Participants were recruited through the open access website of Riddargatan 1, an outpatient clinic within the Stockholm Centre for Dependence Disorders, specializing in treating AUDs. Visitors to the website were informed about the study on the front page as well as on the page with contact information for the clinic. Website visitors were invited to participate in a study on both internet- and clinic-based services, where they would be randomized and the number of visits to the clinic would vary, depending on study group. By clicking on a link, they accessed more information about the study, provided their informed consent and completed screening with the Alcohol Use Disorders Identification Test (AUDIT) [22], self-reported alcohol dependence criteria [International Classification of Disease 10 (ICD-10)] [23] and demographic questions. After screening, registrants under the age of 18 years or those having a low risk for AUD (fewer than three ICD-10

criteria or a score lower than 15 points on the AUDIT) were informed that they did not qualify for the study and were referred to the clinic's ordinary services for further help. Eligible registrants were instructed to create a personal account, with a unique user-name and password, and then asked to complete baseline measures.

Participants who completed the baseline measures were contacted via telephone by a research nurse, who registered them as patients at the clinic, instructed them on how to provide a blood sample for biological markers and booked an appointment for assessment with a physician at the clinic. At that visit, the physician established whether the patient had an AUD, and furthermore did not meet any of the following exclusion criteria: risk for severe withdrawal symptoms, risk for suicide, requiring medication for AUD, having a mental illness requiring separate care, not residing in Stockholm County, not undergoing other treatment for AUD or lacking sufficient understanding of the Swedish language. Participants had no other contact with the researchers or clinic staff before being randomized.

Randomization and masking

After the visit, all participants who were not excluded were given one part of a sealed opaque double envelope, prepared and shuffled in bunches of 20 by the first author. The envelope included information on the plan for the assigned treatment. The other part of the envelope was opened by one of the therapists, who then contacted the participant to start treatment via internet or face-to-face. The participants did not receive information on which of the groups was of interest for the research and which was considered a control for study purposes. All research and clinical staff were blinded to allocation until after the participant was included and given their sealed envelope.

Interventions

Internet-delivered treatment

The internet CBT (ICBT) was delivered on-line at the same website that was used for recruitment and assessment in the study, programmed in the open-source platform Drupal (drupal.org) by the first author. The CBT programme was based on self-help material previously used in studies on the internet and in specialist care [24–27]. The content and exercises in the programme were based on motivational interviewing, relapse prevention and behavioural self-control. The programme included user instructions, videos with example situations and short interviews with experts. The length of the programme was equivalent to 50 pages of printed text. The programme was divided into five main modules: (1) motivation, (2) drinking-goal and self-control, (3) behavioural analysis of drinking and

risk-situations, (4) general problem-solving and (5) preventing relapse, three optional problem-solving modules (handling feelings, drink-refusal skills and handling cravings) and 11 fact sheets (blood alcohol level, anxiety, depression, anger, stress, handling thoughts, relaxation, sleep, leisure activities and communication). The first module included brief feedback on the assessment. Users were also encouraged to register their alcohol consumption and craving estimates, with space to provide details on the situation where they drank or experienced craving. All the therapists were CBT-trained psychologists or psychotherapists. Participants were recommended to spend between 1 and 2 weeks on each module and between 10 and 12 weeks for the whole treatment. In the internet-based treatment the modules were released one-by-one by the therapist and all the communication between the therapist and the patient was conducted on-line asynchronously, without any visits at the clinic during the treatment. Automated e-mails were sent to the patient at the release of every new module and with every new message from the therapist. Patients who did not start the internet programme or stopped using it were sent reminders via text messages to use the intervention. Telephone numbers were provided on the website for technical assistance and to the clinic in case of emergency.

Face-to-face treatment

The same programme as in the ICBT was provided as a paper printout for face-to-face patients. Each of the main modules and the optional additional modules and content were discussed in five 30–45-minute pre-booked sessions at the clinic. If the patient missed a session, it was re-scheduled by telephone. The participants could not contact their therapist via telephone, but they could contact the clinic to re-book their session or in case of emergency. The same eight therapists who delivered the internet CBT provided the face-to-face CBT. The instructions to the therapist were to keep the delivery of the CBT modules as similar as possible in both interventions. Sessions were recorded on audio and all therapists received regular supervision from the first author.

Follow-up

At 3 and 6 months after randomization, participants were e-mailed a link to the follow-up questionnaires. If they spontaneously logged-on to the programme at 3 or 6 months after randomization, they were also re-directed to the follow-up questionnaires. The same questionnaires as at baseline were used at follow-up, adjusted for the time since the last assessment (3 months). Participants who did not respond to this initial request received up to five automated e-mail reminders, followed by a manual e-mail reminder and a mobile text message from the first author.

Once a participant completed the follow-up measures, no further reminders were sent.

Measures

Primary outcome

The primary outcome was the difference between groups in alcohol consumption, measured by the total number of standard drinks consumed during the previous week at 6-month follow-up. The number of standard drinks consumed on each of the 7 days was self-reported using the time-line follow-back (TLFB) method [28]. According to the Swedish definition, one standard drink contains 12 g of pure alcohol. The TLFB has also been shown to be a valid and reliable procedure to document recent drinking when administered via the internet [29] and in a 7-day version [30].

Secondary outcomes

Secondary outcomes were alcohol consumption at 3-month follow-up, measured by the total number of standard drinks consumed during the previous week. Additional outcomes, calculated for the preceding week based on the TLFB score, included the number of non-drinking days, the number of binge-drinking days (defined as days with three or more drinks for women and four or drinks for men), the average number of drinks on drinking days and low-risk score, as well as the 'zone' categories described below, and use-categories. Consumption (according to Swedish guidelines) was no more than 14 drinks per week and no binge drinking for men or, for women, no more than nine drinks per week and no binge drinking. Problematic alcohol use was assessed with the AUDIT total score, as well as the 'zone' categories described below, and use-categories. AUDIT is a 10-item instrument that has been validated in Swedish and via computer [31,32]. For descriptive purposes at baseline, as well as for assessment of non-improved and deteriorated participants, the AUDIT total score was categorized into four risk-level zones as follows: (I) 0–8p, (II) 9–15p, (III) 16–19p and (IV) 20–40p. AUD was assessed by the self-reported number of AUD criteria fulfilled according to the Diagnostic and Statistical Manual version 5 (DSM-5) [2] during the past year. Health-related quality of life was assessed with the EuroQol-5 dimension (EQ-5D-5L). Index scores were calculated with Crosswalk value sets, using the United Kingdom as a reference [33,34]. Readiness to change was measured with two visual analogue scale (VAS) entries, where users responded on a scale of 0–10 to the statements: 'I am not ready to reduce/quit my drinking' (0) and 'I am very much ready to reduce/quit my drinking' [35]. Symptoms of depression were measured by the Montgomery Asberg Depression Rating Scale-Self Rated (MADRS-S) [36,37]. Symptoms of anxiety were measured

by the Generalized Anxiety Disorder assessment–7 items (GAD-7) [38,39]. Use of other support for reducing or eliminating alcohol use was assessed by four questions covering with whom and in what context participants had spoken to someone about their alcohol problems, which medication they had used or which other internet resources they had used regarding alcohol.

The primary and secondary outcomes were assessed on-line at baseline and at 3 and 6 months post-randomization. At 3- and 6-month follow-ups the time-frame for AUDIT and DSM-5 was adjusted from 12 to 3 months.

Additional measures

All participants received e-mail invitations 3 and 6 weeks after beginning treatment to answer the Working Alliance Inventory (WAI) and Session Rating Scale (SRS) [40]. Questions concerning satisfaction with treatment and perceived knowledge-gains from treatment were completed at 6 months. Blood samples were collected at baseline and 6-month follow-up and were analysed for levels of carbohydrate-deficient transferrin (CDT), which rises following recent heavy alcohol use.

Non-inferiority limit

The non-inferiority limit should be based on the effect of the reference treatment [19]. It is important to note that the basis for the limit is not the overall effect of interventions aimed at reducing alcohol use or effects of previous internet interventions, which have often been self-help interventions studied in public health or prevention settings [8]. The margin was calculated with the fixed-margin method [41] and a preserved effect of 50% of the lower margin of the 95% CI, as suggested by guidelines [42]. The non-inferiority limit in the current study was based on the effect of CBT for substance use disorder compared to no intervention [$g = 0.85$ (0.69–1.01)], shown in meta-analysis [4]. The non-inferiority limit was set to five standard drinks for primary outcome and $d = 0.32$ for secondary outcomes.

Sample size

Based on preliminary data from a previous study, the power calculation included in the original protocol was performed with the G*Power program, showing that the required number of study participants needed in the statistical analysis was 176 per group. An updated power calculation for non-inferiority using sealedenvelope.com was performed in the autumn of 2017 based on a standard deviation (SD) of 13.5, observed at baseline for recruited participants to this study. With no difference between the face-to-face and internet-based CBT, 115 patients per group would be

required for 80% certainty that the lower limit of a 95% two-sided confidence interval would be above the non-inferiority limit of five drinks at 6-month follow-up.

Analytical plan and statistical procedure

The analysis of the primary outcome was performed using a linear model, with alcohol consumption at 6 months as the dependent variable and treatment group as the factor variable, adjusting for baseline consumption. The same statistical model was used for analyses of secondary outcomes, adjusting for the baseline value of each respective outcome. Outcomes that, according to visual inspection of data and residuals, were not normally distributed were analysed using generalized linear models with either a negative binomial or a Poisson family function, depending on the goodness-of-fit statistics. Between-group effect sizes (Cohen's *d*) were calculated by dividing differences in estimated mean values (at 3- and 6-month follow-ups) by the pooled SD. Additional analyses used *t*-tests and χ^2 -tests. All analyses used two-sided tests and a significance level of $P < 0.05$. The primary outcome analysis was based on the intention-to-treat (ITT). Under the assumption that data were missing at random, missing data were handled with multiple imputation with 100 different imputed data sets that were combined using the pooled estimate. The imputations were performed using predictive mean matching and fully conditional specification (10 maximum iterations), with the constraint that baseline variables were predictors-only. The study featured no design aspects that risks introducing non-random missingness, including separating outcome reporting from treatment delivery. In addition, anecdotal experience from reaching participants who had initially not provided follow-up data suggested no bias in any direction. Multiple imputation was chosen for the primary analysis over baseline carried forward imputation, as the latter type of simple imputation has been shown to give biased estimates of treatment effects [43] and decreases power to detect between-group differences [44], which could unduly favour a hypothesis of non-inferiority. Two sensitivity analyses were performed; one according to per-protocol (PP), only including those who completed the 3- or 6-month follow-ups, respectively, and who accessed the allocated treatment but not any other treatment; and the second according to the assumption of missing not at random (MNAR), where missing values at the 3- or 6-month follow-ups were replaced by baseline values. Data imputation and all tests were carried out using SPSS version 25. Little's overall test of randomness indicated that missing data on the primary outcome occurred completely at random [$\chi^2 = 1.163$, degrees of freedom (d.f.) = 7, $P = 0.992$]. Imputation was used for 38% of the data in the ITT and the MNAR analysis.

RESULTS

Participants

During the 2-year recruitment period, 735 individuals were screened for participation in the study. A total of 301 participants were randomly allocated to the two study arms (see flow-chart in Fig. 1). This was lower than the initially intended sample size, but in line with the updated sample calculation made after recruitment start. During the last year prior to treatment, six (2%) randomized participants had talked to someone in specialized care and 78 (26%) to any professional about their alcohol use. Full demographic and clinical variables at baseline are presented in Table 1.

Loss to follow-up

Attrition was 33% ($n = 98$) at the 3-month follow-up and 43% ($n = 129$) at the 6-month follow-up. There was no difference in attrition between the internet and the face-to-face group, neither at 3 months ($\chi^2 = 2.062$, $P = 0.151$) nor at 6 months ($\chi^2 = 0.004$, $P = 0.947$). The number of participants completely missing from both follow-ups was 25% in both groups ($n = 38$ versus $n = 37$; $\chi^2 = 0.004$, $P = 0.947$). A difference in baseline GAD-7 scores occurred between participants who were lost and those who completed the 6-month follow-up (7.31 versus 6.14, $t = 2.116$, $P = 0.035$). No differences in any other baseline variables were found between those lost to follow-up at 3 or 6 months and those who completed both follow-ups. No significant interactions between group assignment and being lost at 3- or 6-month follow-ups were found on any baseline variables, indicating no overall confounding effects of missing data.

Treatment usage

The face-to-face group completed more modules than the internet-based group (4.19 versus 3.74, $t = 2.49$, $P = 0.013$). In the internet-based treatment, the mean number of comments per module sent by patients was 1.6 (SD = 1.1) and by therapists 4.1 (SD = 2.5). Treatment was never accessed by seven participants in the internet group and 12 in the face-to-face group. Other treatment than that according to protocol was given to 13 participants: in the internet group, three participants received additional treatment visits at the clinic and two received pharmacological treatment; in the face-to-face group, eight participants received pharmacological treatment. In a few cases the last treatment session was after the post-treatment follow-up, which was conducted 3 months after randomization for all participants.




Figure 1 Flow-chart. ITT = intention-to-treat; PP = per-protocol; MNAR = missing not at random. [Colour figure can be viewed at wileyonlinelibrary.com]

Primary outcome analysis

Controlling for baseline consumption, the difference between internet and face-to-face treatment was non-inferior according to the pre-specified limit of five standard drinks of alcohol the previous week at 6-month follow-up. In the PP sensitivity analysis, only including those who completed 6-month follow-up, who received the allocated treatment but not any other treatment, and in the separate MNAR sensitivity analysis, where missing values were replaced by baseline alcohol consumption, the results also showed non-inferiority of internet compared to face-to-face treatment at 6-month follow-up (see Fig. 2 and Table 2 for detailed results).

Secondary outcome analyses

At 6 months, internet treatment was inferior to face-to-face treatment in all secondary outcomes, with the exception of the AUDIT score and the number of binge drinking days in the ITT analysis, as well as all outcomes in the MNAR analysis. In the PP analysis, the AUDIT score, the number of fulfilled AUD criteria and the biomarker of alcohol consumption CDT did not show non-inferiority for internet treatment. At 3 months, several of the outcomes did not show non-inferiority for internet treatment

compared to face-to-face treatment (see Table 3 for estimated means, effect sizes and CI).

Clinical outcomes

Low-risk alcohol consumption according to Swedish guidelines was reported by 35 of 86 (41%) of follow-up participants in the internet group and 32 of 86 (37%) participants in the face-to-face group at 6 months, with no significant differences found between the groups ($\chi^2 = 0.22$, $P = 0.64$).

Deterioration and non-response

Among follow-up participants at 6 months, one participant in the internet group and two in the face-to-face group deteriorated to a more severe category of alcohol use according to AUDIT, while 14 participants (16%) in the internet group and seven in the face-to-face group (8%) had not changed and remained within the highest AUDIT category. The difference in deterioration and non-response between the groups was not statistically significant ($\chi^2 = 1.26$, $P = 0.26$).

Adverse events

There were no reported adverse events due to the intervention reported by participants. Two participants reported

Table 1 Demographic and baseline values.

Sex	Face-to-face		Internet			
	n	%	n	%		
Women	60	39.7	55	36.7		
Men	91	60.3	94	62.7		
Civil state						
Married	63	41.7	66	44.0		
Cohabiting	38	25.2	40	26.7		
Single	30	19.9	27	18.0		
Divorced	16	10.6	11	7.3		
Widow	4	2.6	5	3.3		
Country of birth						
Sweden	143	94.7	138	92.0		
Other Nordic country	7	4.6	4	2.7		
North America	0	0.0	1	0.7		
Rest of Europe	1	0.7	6	4.0		
Education						
University/college	104	68.9	100	66.7		
Upper secondary school. Training school or equivalent	41	27.2	41	27.3		
Primary school/folk school	6	4.0	7	4.7		
Source of income						
Employment	124	82.1	124	82.7		
Pension	13	8.6	13	8.7		
Study allowance	6	4.0	4	2.7		
Other	8	5.4	7	4.7		
Residence						
Condominium	71	47.0	64	42.7		
Rental apartment	43	28.5	37	24.7		
Villa or townhouse	31	20.5	44	29.3		
Other	6	4.0	4	2.7		
Living arrangements						
With partner and children	60	39.7	51	34.0		
With partner only	43	28.5	50	33.3		
With children only	9	6.0	8	5.3		
Alone	22	14.6	18	12.0		
Other/varied	17	11.2	22	14.7		
AUDIT risk-level						
Zone II (9–15)	9	6	9	6		
Zone III (16–19)	55	36	40	27		
Zone IV (20–40)	87	58	101	67		
ICD alcohol dependence	24	84	24	84		
	n	Mean	SD	n	Mean	SD
Readiness to reduce alcohol consumption (scale 0–10)	151	8.9	1.7	150	9.1	1.7
Drinks/last week	151	22.9	13.3	150	24.9	13.7
Sober days (last week)	151	2.9	2.1	150	2.7	1.8
Binge drinking days (last week)	151	3.3	2.1	150	3.5	1.8
AUDIT-score (0–40)	151	21.0	4.8	150	21.9	4.8
DSM-5 AUD	151	6.3	2.1	150	6.6	2.1
GAD-7	151	6.4	4.5	150	6.9	5.0
MADRS-S	151	13.5	8.3	150	15.0	8.6
EQ-5D-5L	151	0.831	0.131	150	0.832	0.148
CDT	134	2.0	1.6	128	2.0	1.2

AUDIT = Alcohol Use Disorders Identification Test; DSM = Diagnostic and Statistical Manual; GAD = Generalized Anxiety Disorder; CDT = carbohydrate-deficient transferrin; ICD = International Classification of Disease; MADRS-S = Montgomery Asberg Depression Rating Scale–Self Rated; EQ-5D-5L = EuroQol-5 dimension.




Figure 2 Mean difference in estimated standard drinks of alcohol previous week at 6 months follow up with 95% confidence interval. PP = per-protocol; MNAR = missing not at random

negative consequences from being registered as patients at a clinic for AUD. One was refused additional life insurance and one was refused an *in-vitro* fertilization procedure.

Working alliance during treatment

Participants who completed the alliance rating during treatment gave a significantly higher rating of the working alliance in the face-to-face group compared to the internet group, according to both WAI and SRS (see Tables 3 and 4 for details).

Satisfaction with treatment

A higher proportion of participants in the internet group indicated that they missed other forms of contact, and a lower proportion viewed the received treatment as effective compared to the face-to-face group. The internet group also viewed the treatment as less personal (see Tables 3 and 4 for details).

DISCUSSION

Principal results

This randomized controlled non-inferiority trial investigated the effects of internet-delivered CBT compared to face-to-face CBT among participants with AUD who sought help in a specialized clinic. As hypothesized, the internet-delivered treatment was non-inferior to

face-to-face treatment in the primary ITT analysis of alcohol consumption 6 months after recruitment, as well as in the two sensitivity analyses. At 6 months, non-inferiority was also shown in most secondary outcomes. Nonetheless, non-inferiority could not be shown at 6 months in the secondary outcomes measured by the total AUDIT score and the number of binge drinking days in the ITT analysis, or in the PP analysis, for the total AUDIT score, AUD criteria and CDT biomarker levels. At the 3-month follow-up, non-inferiority could not be shown in the PP analysis for number of days sober and EQ-5D-5L in any analysis, or in drinks per week and binge drinking days. Overall, there were small differences in outcomes between the treatment groups.

The fact that the findings from some of the secondary outcomes did not show non-inferiority, especially in the PP analysis of outcomes immediately after treatment, could indicate that there may have been some additional benefits from having met a therapist face-to-face. Treatment use was slightly higher among participants in the face-to-face group and they were more satisfied with the treatment compared to the internet group. Many internet patients said they missed other forms of therapist contact. Receiving treatment through internet interventions has both advantages and disadvantages, and might not be for everyone. The special method of communicating asynchronously can give both participants and therapists more time to think and reflect about what they want to express, but can make it harder to understand details [45]. Communicating asynchronously via the internet can negatively affect the working alliance between the patient and the therapist, and this might explain some of the differences between the groups observed in the current study.

Limitations

This study had several limitations. Similar to previous studies on the internet-based treatment [46], attrition at follow-up was relatively high, although lower than in previous studies on internet-based alcohol interventions. This

Table 2 Estimated mean standard drinks of alcohol consumed the previous week adjusted for standard drinks at baseline.

	Internet	Face-to-face	Diff.	95% CI	
ITT	12.33	11.43	0.89	-1.10	2.88 ^a
PP	13.02	11.42	1.6	-1.04	4.24 ^a
MNAR	17.67	16.32	1.35	-1.06	3.76 ^a

CI = confidence interval; ITT = intention-to-treat; PP = per protocol; MNAR = missing not at random. ^aNon-inferior according to the 5 standard drink limit.

Table 3 Estimated means adjusted for baseline and between group effect sizes with 95% CI for secondary outcomes at three- and six-months follow-up.

6 months							
ITT	Internet		Face-to-face		<i>d</i>	95% CI	
	Mean	SD	Mean	SD			
AUDIT	12.26	6.25	11.57	5.70	0.11	−0.11	0.34
DSM-5 AUD ^a	3.83	2.80	3.59	2.76	0.09	−0.14	0.31 ^c
Binge days ^a	1.13	1.33	0.99	1.14	0.11	−0.11	0.34
Sober days	3.87	1.98	3.93	2.01	−0.03	−0.26	0.20 ^c
GAD-7 ^a	3.75	4.07	3.59	4.07	0.04	−0.19	0.27 ^c
MADRS-S ^a	7.79	6.92	7.22	6.92	0.08	−0.14	0.31 ^c
EQ-5D-5L	0.842	0.142	0.852	0.137	−0.07	−0.29	0.16 ^c
CDT	2.01	1.72	1.85	1.72	0.09	−0.13	0.32 ^c
PP	Mean	SD	Mean	SD	<i>d</i>	95% CI	
AUDIT	12.22	6.65	10.75	6.32	0.23	0.00	0.45
DSM-5 AUD ^b	3.75	2.77	3.28	2.87	0.17	−0.06	0.39
Binge days ^a	1.20	1.56	1.05	1.56	0.10	−0.13	0.32 ^c
Sober days	3.76	2.00	3.88	2.13	−0.06	−0.28	0.17 ^c
GAD-7 ^a	2.09	2.80	2.32	2.82	−0.08	−0.31	0.14 ^c
MADRS-S ^a	5.26	7.57	4.66	6.11	0.09	−0.14	0.31 ^c
EQ-5D-5L	0.870	0.110	0.880	0.140	−0.08	−0.31	0.15 ^c
CDT	1.83	1.06	1.62	1.06	0.20	−0.03	0.42
MNAR	Mean	SD	Mean	SD	<i>d</i>	95% CI	
AUDIT	16.35	7.70	15.70	7.55	0.08	−0.14	0.31 ^c
DSM-5 AUD ^b	4.72	2.90	4.59	2.90	0.04	−0.18	0.27 ^c
Binge days ^a	1.95	1.95	1.79	1.89	0.08	−0.14	0.31 ^c
Sober days	3.39	1.92	3.39	2.06	0.00	−0.23	0.23 ^c
GAD-7 ^a	4.62	5.05	4.39	4.85	0.05	−0.18	0.27 ^c
MADRS-S ^a	10.52	9.00	9.73	9.00	0.09	−0.14	0.31 ^c
EQ-5D-5L	0.842	0.137	0.854	0.137	−0.08	−0.31	0.14 ^c
CDT	2.01	1.42	1.91	1.42	0.07	−0.16	0.30 ^c
3 months							
Drinks/week	Internet		Face-to-face		Diff	95% CI	
ITT	Mean	SD	Mean	SD			
ITT	12.00		9.74		2.26	−0.19	4.71 ^c
PP	13.17		9.47		3.7	0.656	6.74
MNAR	16.54		14.82		1.72	−0.89	4.33 ^c
ITT	Mean	SD	Mean	SD	<i>d</i>	95% CI	
AUDIT	11.82	6.16	11.56	6.16	0.04	−0.18	0.27 ^c
DSM-5 AUD ^b	4.47	2.86	4.30	2.85	0.06	−0.17	0.29 ^c
Binge days ^a	1.01	1.43	0.91	1.40	0.07	−0.16	0.30 ^c
Sober days	4.16	1.89	4.57	1.96	−0.21	−0.44	0.01
GAD-7 ^a	3.12	3.76	2.80	3.16	0.09	−0.13	0.32 ^c
MADRS-S ^a	7.61	7.17	6.87	6.14	0.11	−0.12	0.34
EQ-5D-5L	0.856	0.145	0.876	0.117	−0.15	−0.37	0.08
PP	Mean	SD	Mean	SD	<i>d</i>	95% CI	
AUDIT	11.74	6.32	11.29	6.32	0.07	−0.16	0.30 ^c
DSM-5 AUD ^b	4.28	2.87	4.23	2.87	0.02	−0.21	0.24 ^c
Binge days ^a	1.10	1.55	0.86	1.56	0.15	−0.07	0.38
Sober days	3.98	3.96	4.45	2.13	−0.15	−0.37	0.08
GAD-7 ^a	3.31	3.79	2.77	2.82	0.16	−0.07	0.39
MADRS-S ^a	7.43	7.57	6.81	6.11	0.09	−0.14	0.32 ^c

(Continues)

Table 3. (Continued)

3 months							
Drinks/week	Internet		Face-to-face		Diff	95% CI	
	Mean	SD	Mean	SD		<i>d</i>	95% CI
EQ-5D-5L	0.850	0.140	0.880	0.140	-0.21	-0.44	0.01
MNAR							
AUDIT	14.67	7.66	15.20	7.77	-0.07	-0.29	0.16 ^b
DSM-5 AUD ^a	4.79	2.84	4.89	2.84	-0.04	-0.26	0.19 ^b
Binge days ^c	1.62	2.10	1.53	2.07	0.04	-0.18	0.27 ^b
Sober days	3.62	2.02	3.94	2.22	-0.15	-0.38	0.08
GAD-7 ^c	4.15	4.88	3.81	4.68	0.07	-0.16	0.30 ^b
MADRS-S ^c	9.34	8.20	9.03	8.50	0.04	-0.19	0.26 ^b
EQ-5D-5L	0.856	0.140	0.876	0.140	-0.14	-0.37	0.09

^aPoisson; ^bnon-inferior according to the 0.32 limit; ^cneg bin. AUDIT = Alcohol Use Disorders Identification Test; DSM = Diagnostic and Statistical Manual; GAD = Generalized Anxiety Disorder; CDT = carbohydrate-deficient transferrin; ICD = International Classification of Disease; MADRS-S = Montgomery Asberg Depression Rating Scale-Self Rated; EQ-5D-5L = EuroQol-5 dimension; MNAR = missing not at random; SD = standard deviation; CI = confidence interval; ITT = intention to treat.

might be a consequence of allowing users a fast and accessible way of signing up for participation. The current trial only measured outcomes at three time-points. More measuring time-points during the intervention could allow more flexibility in the modelling approach and also in examination of moderators and mediators [47]. The non-inferiority limit of five standard drinks might be considered too lenient; for example, if the limit is compared to the effects of brief internet-based or face-to-face interventions aimed at reducing alcohol use in community and primary care settings [8,48]. An additional limitation is that the non-inferiority limit, and consequently the

results of this study, mainly have relevance in CBT treatment of AUDs in specialized clinical settings. The participants in the current study were mainly well-educated, employed full time and had stable living arrangements. All participants had also shown an interest in internet-based services, as they were recruited from the web and informed that parts of the study were conducted via the internet. The results cannot be generalized to all people with AUD, and especially not to patients with more severe or complex problems. Finally, as always, causal interpretations of PP analyses should be made with caution, as the reasons for attrition are largely unknown.

Table 4 Working alliance and satisfaction with treatment in each group.

	Internet			Face-to-face					
	<i>n</i>	<i>Mean</i>	<i>SD</i>	<i>n</i>	<i>Mean</i>	<i>SD</i>	<i>t</i>	<i>P</i>	
WAI total score	79	5.23	1.11	63	5.71	0.91	2.82	0.005	
WAI_bond	79	5.17	1.41	63	5.57	1.09	1.82	0.07	
WAI_goal	79	5.56	0.97	63	6.02	0.83	3.02	0.003	
WAI_task	79	4.95	1.23	63	5.54	1.05	3.05	0.003	
SRS total score	79	30.21	8.69	63	34.87	5.23	3.75	> 0.001	
SRS relationship	79	8.32	2.22	63	8.92	2.18	1.60	0.11	
SRS goals and topics	79	7.97	2.17	63	8.96	1.41	3.14	0.002	
SRS approach or method	79	7.31	2.92	63	8.62	1.53	3.24	0.001	
SRS overall	79	6.62	2.83	63	8.37	1.65	4.37	> 0.001	
Experienced the treatment received as									
Pleasant (0–5)	73	4.03	1.04	70	4.24	1.07	1.22	0.22	
Personal (0–5)	73	2.37	1.56	70	3.84	1.38	5.99	> 0.001	
Safe (0–5)	73	3.90	1.45	70	4.14	1.49	0.97	0.33	
				Count	%	Count	%	χ^2	
Missed other form of contact with their therapist				44	60	8	11	38.16	> 0.001
Would recommend the programme to others				57	78	61	87	2.67	0.26
Considered the programme an effective method for changing their drinking				39	53	50	71	7.49	0.02

WAI = Working Alliance Inventory; SRS = Session Rating Scale.

Strengths

This is the first study, to our knowledge, showing that internet-based treatment is non-inferior to face-to-face treatment for AUD in reducing alcohol consumption as well as symptoms of AUD, anxiety and depression. A major strength was that it was conducted in a dependency clinic with regular patients, lending the study high ecological validity. Treatments were well established in both modes of delivery, with several previous studies conducted in the current setting. Another strength was that the same therapists and the same treatment material was used in both groups. A majority of the participants reduced their drinking and many lowered their drinking to a low-risk level of use at follow-ups.

Comparison with prior work

The findings are in line with results found in other forms of internet-based psychological treatment, such as depression and anxiety [49], and are also in line with previous studies comparing face-to-face and internet alcohol interventions [8]. The level of dependence was similar to a recent Swedish clinical trial in specialized and primary care for AUD, but lower than in regular specialized care for AUD [26]. Participants had lower AUD severity than those who usually receive treatment, but they are representative of the majority of individuals with AUD [50], a population that can be reached via internet interventions.

Future research

There is a need for more studies that compare internet interventions with other treatments for AUD, including studies with long-term follow-up. The difficulties, risks and advantages of delivering alcohol treatment via the internet should be studied more carefully. Internet interventions seem to be effective in changing alcohol consumption, but more studies are needed to understand how they can be used and combined with other interventions to improve treatment for people with AUD. In sparsely populated areas, where people often need to travel a long distance to visit a clinic in person, internet interventions might be the only available alternative for receiving treatment for AUD.

CONCLUSION

In this study we found that internet-delivered treatment was not inferior to face-to-face treatment in reducing alcohol consumption among help-seeking patients with alcohol use disorder. Differences in secondary outcomes indicate that internet-delivered treatment might be inferior to face-to-face treatment on some alcohol-related outcomes.

Clinical trial registration details

The original study protocol including the non-inferiority margins was registered with and approved by the Stockholm Regional Ethical Review Board (ref 2014/1758–31/2, 5 November 2014). The trial was also registered at Clinicaltrials.gov (NCT02671019).

Declaration of interests

M.J. configured the internet platform and co-authored the programme used in the study. The programme is licensed under creative commons and the internet platform Drupal is open-source. A.H.B is co-owner of a company, TeleCoach AB, aiming to disseminate digital interventions; the company is not currently active. M.J., S.A. and U.H. are employed at the clinic Riddargatan 1, M.J., P.L. and M.G. are employed at the Addiction eClinic, both part of the Stockholm Center of Dependency disorders that offers face-to-face and internet-based treatments for alcohol use disorders. All other authors declare no competing interests.

Acknowledgements

This study was supported by grants to K.S. and U.H. provided by Region Stockholm (ALF project), numbers 20130688 and 20150322, and to M.J. by the Doctoral School in Health Care Sciences. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript. The authors wish to thank the therapists: Karin Romberg, Ola Siljeholm, Sara Wallhed Finn and Olle Wiklund from Riddargatan 1, as well as Rosette Lind and Miriam Jacobsson from the eSupport unit of Stockholm Center for Dependency Disorders. We also wish to thank the physicians Sven Wåhlin and Karin Hyland, the research nurse Victoria Andersson and the rest of the staff at Riddargatan 1.

Author Contributions

Magnus Johansson: Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; project administration; resources; software; validation; visualization. **Kristina Sinadinovic:** Conceptualization; funding acquisition; methodology; project administration; supervision. **Mikael Gajeci:** Investigation; methodology; resources. **Philip Lindner:** Formal analysis; investigation; visualization. **Anne H. Berman:** Conceptualization; funding acquisition; methodology; software; supervision. **Ulric Hermansson:** Conceptualization; funding acquisition; methodology; project administration; resources; supervision. **Sven Andreasson:** Conceptualization; funding acquisition; methodology; project administration; resources; supervision.

References

- GBD 2016 Alcohol Collaborators, Griswold M. G., Fullman N., Hawley C., Arian N., Zimsen S. R. *et al.* Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the global burden of disease study 2016. *Lancet* 2018; **392**: 1015–35.
- American Psychiatric Association *Diagnostic and Statistical Manual of Mental Disorders (DSM-5[®])*. Washington, DC: American Psychiatric Association Publishing; 2013.
- Gowing L. R., Ali R. L., Allsop S., Marsden J., Turf E. E., West R. *et al.* Global statistics on addictive behaviours: 2014 status report. *Addiction* 2015; **110**: 904–19.
- Magill M., Ray L. A. Cognitive-behavioral treatment with adult alcohol and illicit drug users: a meta-analysis of randomized controlled trials. *J Stud Alcohol Drugs* 2009; **70**: 516–27.
- Maisel N. C., Blodgett J. C., Wilbourne P. L., Humphreys K., Finney J. W. Meta-analysis of naltrexone and acamprostate for treating alcohol use disorders: when are these medications most helpful? *Addiction* 2013; **108**: 275–93.
- Degenhardt L., Glantz M., Evans-Lacko S., Sadikova E., Sampson N., Thornicroft G. *et al.* Estimating treatment coverage for people with substance use disorders: an analysis of data from the world mental health surveys. *World Psychiatry* 2017; **16**: 299–307.
- Probst C., Mantney J., Martinez A., Rehm J. Alcohol use disorder severity and reported reasons not to seek treatment: a cross-sectional study in European primary care practices. *Subst Abuse Treat Prev Policy BioMed Central* 2015; **10**: 32.
- Kaner E. F. S., Beyer F. R., Garnett C., Crane D., Brown J., Muirhead C. *et al.* Personalised digital interventions for reducing hazardous and harmful alcohol consumption in community-dwelling populations. *Cochrane Database of Systematic Reviews* 2017. <https://doi.org/10.1002/14651858.cd011479.pub2>.
- Riper H., Hoogendoorn A., Cuijpers P., Karyotaki E., Boumparis N., Mira A. *et al.* Effectiveness and treatment moderators of internet interventions for adult problem drinking: an individual patient data meta-analysis of 19 randomised controlled trials. *PLOS Med* 2018; **15**: e1002714.
- White A., Kavanagh D., Stallman H., Klein B., Kay-Lambkin E., Proudfoot J. *et al.* Online alcohol interventions: a systematic review. *J Med Internet Res* 2010; **12**: e62.
- Riper H., Spek V., Boon B., Conijn B., Kramer J., Martin-Abello K. *et al.* Effectiveness of E-self-help interventions for curbing adult problem drinking: a meta-analysis. *J Med Internet Res* 2011; **13**: e42.
- Carlbring P., Andersson G., Cuijpers P., Riper H., Hedman-Lagerlöf E. Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: an updated systematic review and meta-analysis. *Cogn Behav Ther* 2018; **47**: 1–18.
- Kay-Lambkin F. J., Baker A. L., Lewin T. J., Carr V. J., Kay-Lambkin F. J., Baker A. L. Computer-based psychological treatment for comorbid depression and problematic alcohol and/or cannabis use: a randomized controlled trial of clinical efficacy. *Addiction* 2009; **104**: 378–88.
- Kay-Lambkin F. J., Baker A. L., Kelly B., Lewin T. J. Clinician-assisted computerised versus therapist-delivered treatment for depressive and addictive disorders: a randomised controlled trial. *Med J Aust* 2011; **195**: S44–S50.
- Kiluk B. D., Devore K. A., Buck M. B., Nich C., Frankforter T. L., LaPaglia D. M. *et al.* Randomized trial of computerized cognitive behavioral therapy for alcohol use disorders: efficacy as a virtual stand-alone and treatment add-on compared with standard outpatient treatment. *Alcohol Clin Exp Res* 2016; **40**: 1991–2000.
- Acosta M. C., Possemato K., Maisto S. A., Marsch L. A., Barrie K., Lantinga L. *et al.* Web-delivered CBT reduces heavy drinking in OEF-OIF veterans in primary care with symptomatic substance use and PTSD. *Behav Ther* 2017; **48**: 262–76.
- Campbell A. N. C., Nunes E. V., Matthews A. G., Stitzer M., Miele G. M., Polsky D. *et al.* Internet-delivered treatment for substance abuse: a multisite randomized controlled trial. *Am J Psychiatry* 2014; **171**: 683–90.
- Magill M., Ray L., Kiluk B., Hoadley A., Bernstein M., Scott Tonigan J. *et al.* A meta-analysis of cognitive-behavioral therapy for alcohol or other drug use disorders: treatment efficacy by contrast condition. *J Consult Clin Psychol* 2019; **87**: 1093–105.
- Piaggio G., Elbourne D. R., Pocock S. J., Evans S. J. W., Altman D. G. Reporting of noninferiority and equivalence randomized trials: extension of the CONSORT 2010 statement. *JAMA* 2012; **308**: 2594–604.
- Andersson G. The promise and pitfalls of the internet for cognitive behavioral therapy. *BMC Med BioMed Central* 2010; **8**: 82.
- Wallace P., Struzzo P., Della Vedova R., Scafuri F., Tersar C., Lygidakis C. *et al.* Randomised controlled non-inferiority trial of primary care-based facilitated access to an alcohol reduction website. *BMJ Open* 2017; **7**: e014576.
- Saunders J. B., Aasland O. G., Babor T. F., De La Fuente J. R., Grant M. Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction* 1993; **88**: 791–804.
- World Health Organization *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. Geneva: World Health Organization; 1992.
- Johansson M., Sinadinovic K., Hammarberg A., Sundstrom C., Hermansson U., Andreasson S. *et al.* Web-based self-help for problematic alcohol use: a large naturalistic study. *Int J Behav Med* 2017; **24**: 749–59. <https://doi.org/10.1007/s12529-016-9618-z>.
- Sundström C., Gajecki M., Johansson M., Blankers M., Sinadinovic K., Stenlund-Gens E. *et al.* Guided and unguided internet-based treatment for problematic alcohol use - a randomized controlled pilot trial. *PLOS ONE* 2016; **11**: e0157817. <https://doi.org/10.1371/journal.pone.0157817>.
- Wallhed Finn S., Hammarberg A., Andreasson S. Treatment for alcohol dependence in primary care compared to outpatient specialist treatment—a randomized controlled trial. *Alcohol Alcohol* 2018; **53**: 376–85.
- Sinadinovic K., Wennberg P., Johansson M., Berman A. H. Targeting individuals with problematic alcohol use via web-based cognitive-behavioral self-help modules, personalized screening feedback or assessment only: a randomized controlled trial. *Eur Addict Res* 2014; **20**: 305–18.
- Sobell L. C., Sobell M. B. Timeline follow-back. In: Litten R. Z., Allen J. P., editors. *Measuring Alcohol Consumption*. Totowa, NJ: Humana Press; 1992, pp. 41–72.
- Rueger S. Y., Trela C. J., Palmeri M., King A. C. Self-administered web-based timeline followback procedure for drinking and smoking behaviors in young adults. *J Stud Alcohol Drugs* 2012; **73**: 829–33.
- Thomas B. A., McCambridge J. Comparative psychometric study of a range of hazardous drinking measures administered

- online in a youth population. *Drug Alcohol Depend* 2008; **96**: 121–7.
31. Bergman H., Kallmen H., Källmén H., Kallmen H. Alcohol use among Swedes and a psychometric evaluation of the Alcohol Use Disorders Identification Test. *Alcohol Alcohol* 2002; **37**: 245–51.
 32. Sinadinovic K., Wennberg P., Berman A. H. Population screening of risky alcohol and drug use via internet and interactive voice response (IVR): a feasibility and psychometric study in a random sample. *Drug Alcohol Depend* 2011; **114**: 55–60.
 33. Van Hout B., Janssen M. E., Feng Y. S., Kohlmann T., Busschbach J., Golicki D. *et al.* Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health* 2012; **15**: 708–15.
 34. Herdman M., Gudex C., Lloyd A., Janssen M., Kind P., Parkin D. *et al.* Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011; **20**: 1727–36.
 35. Bertholet N., Cheng D. M., Palfai T. P., Samet J. H., Saitz R. Does readiness to change predict subsequent alcohol consumption in medical inpatients with unhealthy alcohol use? *Addict Behav* 2009; **34**: 636–40.
 36. Svanborg P., Åsberg M. A comparison between the Beck depression inventory (BDI) and the self-rating version of the Montgomery Åsberg depression rating scale (MADRS). *J Affect Disord* 2001; **64**: 203–16.
 37. Holländare E., Andersson G., Engström I. A comparison of psychometric properties between internet and paper versions of two depression instruments (BDI-II and MADRS-S) administered to clinic patients. *J Med Internet Res* 2010; **12**: e49.
 38. Spitzer R. L., Kroenke K., Williams J. B. W., Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006; **166**: 1092–7.
 39. Dear B. F., Titov N., Sunderland M., McMillan D., Anderson T., Lorian C. *et al.* Psychometric comparison of the generalized anxiety disorder Scale-7 and the Penn State worry questionnaire for measuring response during treatment of generalised anxiety disorder. *Cogn Behav Ther TF* 2011; **40**: 216–27.
 40. Campbell A., Hemsley S. Outcome rating scale and session rating scale in psychological practice: clinical utility of ultra-brief measures. *Clin Psychol* 2009; **13**: 1–9.
 41. Angeli F., Verdecchia P., Vaudo G., Masnaghetti S., Reboldi G. Optimal use of the non-inferiority trial design. *Pharmaceut Med* 2020; **34**: 159–65.
 42. Food and Drug Administration. Non-inferiority clinical trials to establish effectiveness: guidance for industry [internet]. US Department of Health and Human Services. 2016. Available at: <https://www.fda.gov/media/78504/download> (accessed 10 July 2020).
 43. Rioux C., Little T. D. Missing data treatments in intervention studies: what was, what is, and what should be. *Int J Behav Dev* 2019. <https://doi.org/10.1177/0165025419880609>.
 44. Wiens B. L., Zhao W. The role of intention to treat in analysis of noninferiority studies. *Clin Trials* 2007; **4**: 286–91.
 45. Ekström V., Johansson M. Sort of a nice distance: a qualitative study of the experiences of therapists working with internet-based treatment of problematic substance use. *Addict Sci Clin Pract* 2019; **14**: 44.
 46. Eysenbach G. The law of attrition. *J Med Internet Res* 2005; **7**: e11.
 47. Hesser H. Modeling individual differences in randomized experiments using growth models: recommendations for design, statistical analysis and reporting of results of internet interventions. *Internet Interv* 2015; **2**: 110–20.
 48. Kaner E. F., Beyer F. R., Muirhead C., Campbell E., Pienaar E. D., Bertholet N. *et al.* *Cochrane Database of Systematic Reviews*. Hoboken, NJ: John Wiley & Sons, Ltd; 2018.
 49. Cuijpers P., Donker T., Van Straten A., Li J., Andersson G. Is guided self-help as effective as face-to-face psychotherapy for depression and anxiety disorders? A systematic review and meta-analysis of comparative outcome studies. *Psychol Med* 2010; **40**: 1943–57.
 50. Andreasson S., Danielsson A.-K. K., Hallgren M., Andréasson S., Danielsson A.-K. K., Hallgren M. *et al.* Severity of alcohol dependence in the Swedish adult population: association with consumption and social factors. *Alcohol* 2013; **47**: 21–5.