

Thiophene-2-carbaldehyde 2,4-dinitrophenylhydrazone

Zhi-gang Yin,^{a*} Heng-yu Qian,^a He-ping Li,^b Jie Hu^a and Chun-xia Zhang^a

^aKey Laboratory of Surface and Interface Science of Henan, School of Materials and Chemical Engineering, Zhengzhou University of Light Industry, Zhengzhou 450002, People's Republic of China, and ^bDepartment of Materials and Chemical Engineering, Guiling University of Technology, People's Republic of China
Correspondence e-mail: yinck@263.net


Received 13 November 2008; accepted 18 November 2008

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.045; wR factor = 0.087; data-to-parameter ratio = 12.4.

In the approximately planar molecule of the title compound, $\text{C}_{11}\text{H}_8\text{N}_4\text{O}_4\text{S}$, the dihedral angle between the thiophene and benzene rings is $5.73(10)^\circ$. In the crystal structure, bifurcated inter/intramolecular $\text{N}-\text{H}\cdots(\text{O},\text{O})$ hydrogen bonds are present. The intermolecular links lead to inversion dimers containing an $R_2^2(12)$ graph-set motif.

Related literature

For general background, see: Okabe *et al.* (1993). For graph-set notation, see: Etter *et al.* (1990); Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{11}\text{H}_8\text{N}_4\text{O}_4\text{S}$	$c = 25.708(8)\text{ \AA}$
$M_r = 292.27$	$\beta = 92.71(2)^\circ$
Monoclinic, $P2_1/c$	$V = 1197.7(7)\text{ \AA}^3$
$a = 4.8994(17)\text{ \AA}$	$Z = 4$
$b = 9.520(3)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.29\text{ mm}^{-1}$
 $T = 291(2)\text{ K}$

$0.30 \times 0.26 \times 0.24\text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.915$, $T_{\max} = 0.929$

11010 measured reflections
2285 independent reflections
1718 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.087$
 $S = 1.02$
2285 reflections
184 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H1 \cdots O1 ⁱ	0.87 (2)	2.57 (2)	3.338 (2)	148.2 (19)
N3—H1 \cdots O1	0.87 (2)	2.01 (2)	2.630 (2)	127.4 (19)

Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL* and *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXTL*.

The authors express their deep appreciation to the Startup Fund for PhDs of Natural Scientific Research of Zhengzhou University of Light Industry (grant No. 2005001) and the Startup Fund for Masters of Natural Scientific Research of Zhengzhou University of Light Industry (grant No. 000455).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2406).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2000). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst. B* **46**, 256–262.
- Okabe, N., Nakamura, T. & Fukuda, H. (1993). *Acta Cryst. C* **49**, 1678–1680.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o2421 [doi:10.1107/S1600536808038397]

Thiophene-2-carbaldehyde 2,4-dinitrophenylhydrazone

Z. Yin, H. Qian, H. Li, J. Hu and C. Zhang

Comment

2,4-Dinitrophenylhydrazine is a reagent which is widely used for condensation with aldehydes and ketones. Several phenylhydrazone derivatives have been shown to be potentially DNA-damaging and are mutagenic agents(Okabe *et al.* 1993). As part of our work, we have synthesized the title compound (I) and reported its crystal structure.

The title molecule is roughly planar. Indeed, the benzene and the thiophene rings are only slightly twisted, making a dihedral angle of 5.73 (10) $^{\circ}$ (Fig. 1). The two nitro groups, O1/N1/O2 and O3/N2/O4 are coplanar with the benzene ring to which they are attached.

Intermolecular N-H \cdots O hydrogen bonds link the molecule two by two around inversion center, building then a R₂²(12) graph set motif (Etter *et al.*, 1990; Bernstein *et al.*, 1995) if the intramolecular N-H \cdots O hydrogen bonds are ignored (Table 1, Fig.2).

Experimental

2,4-dinitrophenylhydrazine(1 mmol, 0.198 g) was dissolved in anhydrous methanol, H₂SO₄ (98% 0.5 ml) was added to this, the mixture was stirred for several minutes at 351 K, thiophene-2-carbaldehyde (1 mmol 0.112 g) in methanol (8 ml) was added dropwise and the mixture was stirred at refluxing temperature for 2 h. The product was isolated and recrystallized in methanol, purple single crystals of (I) was obtained after 5 d.

Refinement

All H atoms were placed in calculated positions and treated as riding with C—H=0.93 \AA (aromatic) and N—H=0.86 \AA with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C or N})$.

Figures




Fig. 1. Molecular view of the title compound with the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

supplementary materials




Fig. 2. View showing the N-H \cdots O hydrogen bonds building a $R_2^2(12)$ graph set motif. Hydrogen bonds are shown as dashed lines.

Thiophene-2-carbaldehyde 2,4-dinitrophenylhydrazone

Crystal data

C ₁₁ H ₈ N ₄ O ₄ S	$F_{000} = 600$
$M_r = 292.27$	$D_x = 1.621 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 4.8994 (17) \text{ \AA}$	Cell parameters from 5391 reflections
$b = 9.520 (3) \text{ \AA}$	$\theta = 2.3\text{--}27.4^\circ$
$c = 25.708 (8) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$\beta = 92.71 (2)^\circ$	$T = 291 (2) \text{ K}$
$V = 1197.7 (7) \text{ \AA}^3$	Block, purple
$Z = 4$	$0.30 \times 0.26 \times 0.24 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer	2285 independent reflections
Radiation source: sealed tube	1718 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.047$
$T = 291(2) \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$h = -6 \rightarrow 5$
$T_{\text{min}} = 0.915$, $T_{\text{max}} = 0.929$	$k = -11 \rightarrow 11$
11010 measured reflections	$l = -31 \rightarrow 29$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.087$	$w = 1/[\sigma^2(F_o^2) + (0.04P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2285 reflections	$\Delta\rho_{\text{max}} = 0.16 \text{ e \AA}^{-3}$

184 parameters

 $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$ Primary atom site location: structure-invariant direct
methods

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1154 (4)	0.2360 (2)	0.04553 (7)	0.0342 (4)
C2	-0.0759 (4)	0.1287 (2)	0.04720 (7)	0.0350 (4)
H2	-0.2010	0.1150	0.0193	0.042*
C3	-0.0801 (4)	0.04351 (18)	0.08984 (7)	0.0315 (4)
C4	0.1065 (4)	0.0673 (2)	0.13207 (7)	0.0383 (5)
H4	0.1048	0.0087	0.1610	0.046*
C5	0.2882 (4)	0.1738 (2)	0.13141 (7)	0.0350 (4)
H5	0.4096	0.1870	0.1599	0.042*
C6	0.2980 (4)	0.26512 (18)	0.08856 (7)	0.0291 (4)
C7	0.8165 (4)	0.49823 (19)	0.13079 (7)	0.0389 (5)
H7	0.7974	0.5614	0.1033	0.047*
C8	1.0163 (4)	0.52284 (19)	0.17283 (7)	0.0341 (4)
C9	1.1994 (5)	0.6353 (2)	0.17481 (8)	0.0418 (5)
H9	1.2066	0.7050	0.1496	0.050*
C10	1.3738 (5)	0.6289 (2)	0.22076 (9)	0.0465 (5)
H10	1.5107	0.6939	0.2290	0.056*
C11	1.3173 (5)	0.5174 (2)	0.25090 (8)	0.0522 (6)
H11	1.4113	0.4975	0.2823	0.063*
N1	0.0973 (4)	0.31477 (18)	0.00000 (6)	0.0394 (4)
N2	-0.2778 (4)	-0.06932 (17)	0.09065 (7)	0.0414 (4)
N3	0.4846 (4)	0.37052 (17)	0.08923 (6)	0.0378 (4)
H1	0.483 (5)	0.426 (2)	0.0624 (9)	0.045*
N4	0.6631 (3)	0.38663 (15)	0.13180 (6)	0.0321 (4)
O1	0.2484 (4)	0.41150 (16)	-0.00353 (6)	0.0545 (4)
O2	-0.0594 (3)	0.29004 (14)	-0.03604 (5)	0.0461 (4)
O3	-0.4272 (3)	-0.08987 (16)	0.05310 (6)	0.0526 (4)
O4	-0.2847 (3)	-0.13710 (15)	0.12986 (6)	0.0513 (4)
S1	1.05706 (12)	0.41736 (6)	0.22559 (2)	0.04762 (18)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0302 (10)	0.0438 (10)	0.0277 (10)	-0.0016 (8)	-0.0096 (8)	-0.0034 (7)
C2	0.0330 (10)	0.0410 (10)	0.0303 (10)	-0.0007 (8)	-0.0046 (8)	-0.0039 (7)
C3	0.0326 (10)	0.0290 (8)	0.0325 (9)	-0.0001 (7)	-0.0024 (8)	-0.0025 (7)
C4	0.0414 (12)	0.0417 (10)	0.0314 (10)	0.0038 (8)	-0.0038 (9)	0.0075 (8)
C5	0.0259 (10)	0.0463 (10)	0.0323 (10)	0.0090 (8)	-0.0035 (8)	-0.0005 (8)
C6	0.0270 (9)	0.0339 (8)	0.0262 (9)	0.0002 (7)	-0.0004 (7)	-0.0062 (7)
C7	0.0510 (12)	0.0394 (10)	0.0257 (9)	-0.0026 (9)	-0.0057 (8)	-0.0013 (8)
C8	0.0316 (10)	0.0353 (9)	0.0349 (9)	-0.0023 (8)	-0.0025 (8)	-0.0079 (7)
C9	0.0473 (13)	0.0361 (9)	0.0412 (11)	-0.0042 (9)	-0.0073 (9)	-0.0059 (8)
C10	0.0387 (12)	0.0416 (10)	0.0581 (14)	-0.0045 (9)	-0.0105 (10)	-0.0220 (10)
C11	0.0629 (15)	0.0558 (13)	0.0357 (11)	0.0052 (11)	-0.0222 (11)	-0.0131 (9)
N1	0.0383 (10)	0.0452 (9)	0.0335 (9)	-0.0104 (8)	-0.0120 (8)	-0.0013 (7)
N2	0.0360 (10)	0.0421 (9)	0.0453 (10)	-0.0026 (7)	-0.0074 (8)	0.0047 (8)
N3	0.0387 (10)	0.0443 (9)	0.0290 (9)	-0.0077 (7)	-0.0126 (7)	-0.0001 (7)
N4	0.0345 (8)	0.0344 (8)	0.0266 (8)	-0.0003 (6)	-0.0087 (7)	-0.0040 (6)
O1	0.0659 (11)	0.0525 (9)	0.0435 (9)	-0.0225 (8)	-0.0163 (8)	0.0158 (7)
O2	0.0471 (9)	0.0505 (8)	0.0383 (8)	-0.0317 (7)	-0.0222 (7)	0.0182 (6)
O3	0.0488 (10)	0.0534 (9)	0.0537 (9)	-0.0260 (7)	-0.0185 (8)	-0.0002 (7)
O4	0.0566 (10)	0.0451 (8)	0.0519 (9)	-0.0169 (7)	-0.0008 (8)	0.0131 (7)
S1	0.0480 (3)	0.0553 (3)	0.0385 (3)	-0.0071 (2)	-0.0093 (2)	0.0040 (2)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.388 (3)	C8—C9	1.396 (3)
C1—N1	1.389 (2)	C8—S1	1.692 (2)
C1—C6	1.417 (2)	C9—C10	1.426 (3)
C2—C3	1.365 (3)	C9—H9	0.9300
C2—H2	0.9300	C10—C11	1.351 (3)
C3—C4	1.404 (2)	C10—H10	0.9300
C3—N2	1.447 (2)	C11—S1	1.696 (2)
C4—C5	1.350 (3)	C11—H11	0.9300
C4—H4	0.9300	N1—O1	1.188 (2)
C5—C6	1.406 (3)	N1—O2	1.1983 (19)
C5—H5	0.9300	N2—O4	1.199 (2)
C6—N3	1.357 (2)	N2—O3	1.199 (2)
C7—N4	1.302 (2)	N3—N4	1.377 (2)
C7—C8	1.442 (2)	N3—H1	0.87 (2)
C7—H7	0.9300		
C2—C1—N1	114.01 (15)	C9—C8—S1	112.01 (14)
C2—C1—C6	121.47 (17)	C7—C8—S1	123.66 (14)
N1—C1—C6	124.40 (17)	C8—C9—C10	110.86 (19)
C3—C2—C1	119.87 (17)	C8—C9—H9	124.6
C3—C2—H2	120.1	C10—C9—H9	124.6
C1—C2—H2	120.1	C11—C10—C9	112.19 (18)

C2—C3—C4	119.41 (17)	C11—C10—H10	123.9
C2—C3—N2	119.22 (15)	C9—C10—H10	123.9
C4—C3—N2	121.36 (16)	C10—C11—S1	113.08 (15)
C5—C4—C3	121.14 (18)	C10—C11—H11	123.5
C5—C4—H4	119.4	S1—C11—H11	123.5
C3—C4—H4	119.4	O1—N1—O2	118.16 (16)
C4—C5—C6	121.43 (17)	O1—N1—C1	117.89 (14)
C4—C5—H5	119.3	O2—N1—C1	123.94 (16)
C6—C5—H5	119.3	O4—N2—O3	123.26 (17)
N3—C6—C5	119.70 (15)	O4—N2—C3	117.21 (15)
N3—C6—C1	123.69 (17)	O3—N2—C3	119.53 (16)
C5—C6—C1	116.55 (17)	C6—N3—N4	119.65 (16)
N4—C7—C8	119.31 (16)	C6—N3—H1	117.6 (14)
N4—C7—H7	120.3	N4—N3—H1	122.8 (14)
C8—C7—H7	120.3	C7—N4—N3	114.87 (15)
C9—C8—C7	124.33 (18)	C8—S1—C11	91.85 (11)
N1—C1—C2—C3	180.00 (18)	C8—C9—C10—C11	-0.6 (3)
C6—C1—C2—C3	-3.8 (3)	C9—C10—C11—S1	0.1 (3)
C1—C2—C3—C4	1.3 (3)	C2—C1—N1—O1	177.51 (19)
C1—C2—C3—N2	-178.99 (18)	C6—C1—N1—O1	1.5 (3)
C2—C3—C4—C5	0.6 (3)	C2—C1—N1—O2	-3.3 (3)
N2—C3—C4—C5	-179.16 (18)	C6—C1—N1—O2	-179.4 (2)
C3—C4—C5—C6	0.1 (3)	C2—C3—N2—O4	-176.10 (19)
C4—C5—C6—N3	-179.66 (19)	C4—C3—N2—O4	3.6 (3)
C4—C5—C6—C1	-2.5 (3)	C2—C3—N2—O3	3.1 (3)
C2—C1—C6—N3	-178.60 (18)	C4—C3—N2—O3	-177.21 (19)
N1—C1—C6—N3	-2.8 (3)	C5—C6—N3—N4	0.2 (3)
C2—C1—C6—C5	4.3 (3)	C1—C6—N3—N4	-176.76 (17)
N1—C1—C6—C5	-179.88 (19)	C8—C7—N4—N3	-178.20 (17)
N4—C7—C8—C9	177.13 (19)	C6—N3—N4—C7	-173.88 (18)
N4—C7—C8—S1	-2.5 (3)	C9—C8—S1—C11	-0.66 (18)
C7—C8—C9—C10	-178.8 (2)	C7—C8—S1—C11	179.01 (19)
S1—C8—C9—C10	0.8 (2)	C10—C11—S1—C8	0.3 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H1···O1 ⁱ	0.87 (2)	2.57 (2)	3.338 (2)	148.2 (19)
N3—H1···O1	0.87 (2)	2.01 (2)	2.630 (2)	127.4 (19)

Symmetry codes: (i) $-x+1, -y+1, -z$.

supplementary materials

Fig. 1




Fig. 2

