

Research Article

Determination of Aflatoxin B₁ in Feedstuffs without Clean-Up Step by High-Performance Liquid Chromatography

Sasiprapa Choochuay,^{1,2} Jutamas Phakam,^{1,2} Prakorn Jala,³
Thanapoom Maneeboon,⁴ and Natthasit Tansakul ^{1,2}

¹Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, Kasetsart University, 10900, Bangkok, Thailand

²Center for Advanced Studies for Agriculture and Food, KU Institute for Advanced Studies, Kasetsart University (CASAF, NRU-KU), 10900, Bangkok, Thailand

³Faculty of Veterinary Medicine, Kamphaengsaen Campus, 73140, Nakhon Pathom, Thailand

⁴Kasetsart University Research and Development Institute, 10900, Bangkok, Thailand

Correspondence should be addressed to Natthasit Tansakul; natthasitt@yahoo.com

Received 16 March 2018; Accepted 10 May 2018; Published 24 June 2018

Academic Editor: Mohamed Abdel-Rehim

Copyright © 2018 Sasiprapa Choochuay et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A reliable and rapid method has been developed for the determination of aflatoxin B₁ (AFB₁) in four kinds of feedstuffs comprising broken rice, peanuts, corn, and fishmeal. A sample preparation was carried out based on the QuEChERS method with the exclusion of the clean-up step. In this study, AFB₁ was extracted using acetonitrile/methanol (40/60 v/v), followed by partitioning with sodium chloride and magnesium sulfate. High-performance liquid chromatography with precolumn derivatization and fluorescence detection was performed. The coefficients of determination were greater than 0.9800. Throughout the developed method, the recovery of all feedstuffs achieved a range of 82.50-109.85% with relative standard deviation lower than 11% for all analytes at a concentration of 20-100 ng/g. The limit of detection (LOD) ranged from 0.2 to 1.2 ng/g and limit of quantitation (LOQ) ranged from 0.3 to 1.5 ng/g. The validated method was successfully applied to a total of 120 samples. The occurrence of AFB₁ contamination was found at the following concentrations: in broken rice (0.44-2.33ng/g), peanut (3.97-106.26ng/g), corn (0.88-50.29 ng/g), and fishmeal (1.06-10.35 ng/g). These results indicate that the proposed method may be useful for regularly monitoring AFB₁ contamination in feedstuffs.

1. Introduction

Aflatoxins (AFs) are secondary metabolites of fungi (e.g., *Aspergillus flavus* and *A. parasiticus*). AFs occur naturally and can be found in common food and feedstuffs such as rice, peanuts, corn, fishmeal, and soybean meal [1-3]. The four major analogues— aflatoxins B₁ (AFB₁), B₂ (AFB₂), G₁ (AFG₁), and G₂ (AFG₂)—are the most important members because they all pose a potential risk to human and animal health if food and feedstuffs have been contaminated. In particular, the toxicity of AFB₁ can range from levels that may cause immune system suppression to the induction of teratogenic, mutagenic, and carcinogenic activities [4, 5], which is collectively classified as a carcinogen (Group 1) by

the International Agency for Research on Cancer (IARC) [6]. In addition, feedstuffs comprise the first link of the food chain; therefore, there is a risk of AFB₁ carryover from feedstuffs into animal tissues and/or biological fluids such as meat, milk, and eggs, which may eventually be hazardous for human consumption [7-9]. As a result of such adverse health effects of the toxin, it is necessary to have a sensitive, reliable, and accurate method for monitoring AFB₁ level in feedstuffs.

High-performance liquid chromatography with fluorescence detection (HPLC-FLD) is considered as the most reliable instrument for the quantification of AFs due to its accuracy and high sensitivity. However, it requires clean-up steps involving immune affinity columns (IAC) to remove interferences and preconcentration of AFB₁. IAC is also

quite expensive, difficult to use, and time-consuming [10]. In this regard, a QuEChERS (Quick, Easy, Cheap, Effective, Rugged, and Safe) method, which was originally developed for pesticides and veterinary drug residue analysis, has been applied for the measurement of mycotoxins in a variety of matrices [10–12]. Commonly, it involves two steps: the first step is the extraction step based on a partitioning phase, and the second step is the dispersive-solid phase extraction step (d-SPE) using a combination of magnesium sulfate with different sorbents. The advantages of this method are that it is simple and reduces time consumption [13, 14].

In order to reduce sample handling and increase sample output efficiency, the aim of this study is focused on the optimization of sample preparation using QuEChERS-based extraction and optimizing the chromatographic conditions of an HPLC-FLD method with pre-column derivatization for quantification of AFB₁ in the feedstuffs as well as a cross-sectional investigation of the contamination of AFB₁ content in the feedstuffs using a QuEChERS-based method.

2. Materials and Methods

2.1. Chemicals and Reagents. AFB₁ standard was purchased from Sigma-Aldrich (Madrid, Spain). Stock solution of AFB₁ was prepared in methanol at a concentration of 10 mg/mL. From the stock standard solution, working solution was prepared by diluting AFB₁ in methanol at a concentration of 1 mg/mL and stored at 4°C. Before use, the solution was brought to laboratory room temperature (25°C) and thoroughly shaken. Acetonitrile and methanol were of HPLC grade and supplied by ACILabsan (Thailand). Sodium chloride was obtained from Ajax Finechem Pty. Ltd. (New Zealand) and anhydrous magnesium sulfate was from Applicati Chem Panreac ITW Companies (Germany). Trifluoroacetic acid was purchased from ACROS Organics (Belgium) and water was purified (18 M Ω) on a Milli-Q Plus apparatus.

2.2. Optimization of the HPLC-FLD Analytical Method. To fulfill the purposes, several conditions, including a mobile phase, extraction solvents combined with the QuEChERS salt, volume of extraction, and precolumn derivatization solvents, were optimized to ensure an efficient determination of AFB₁. Different combinations of the mobile phase solution, including methanol/water (50/50 v/v), acetonitrile/methanol/water (20/20/60 v/v/v), acetonitrile/methanol/water (16/22/62 v/v/v), and acetonitrile/methanol/water (15/15/70 v/v/v), were evaluated in order to optimize the retention time of the AFB₁ and to enhance the sensitivity. In addition, the different mixtures of extract solvents that consisted of acetonitrile (100%), acetonitrile/methanol (40/60 v/v), and acetonitrile/water (50/50 v/v) were investigated. Rather than optimizing the ratio of extraction solvents alone, this study simultaneously evaluated the addition of different drying agents or salt; these included sodium chloride/magnesium sulfate (1/4 w/w) or sodium acetate/magnesium sulfate (1/4 w/w). The effect of extraction volume of the satisfied extraction solvent was tested by two different volumes of 10 and 20 mL. Moreover, the critical points of reconstitution of the

precolumn derivatization with suitable solvent between 10% acetonitrile and the selected mobile phase were compared.

2.3. Optimization of Sample Preparation. The optimization of sample preparation was carried out using broken rice as a representative matrix. Briefly, the effects of different extraction solvents (acetonitrile 100%, acetonitrile/methanol 40/60 v/v, and acetonitrile/water 50/50 v/v) and salts (sodium chloride and sodium acetate) on the signal response of AFB₁ were compared. After milling and homogenization, a 10 g sample was weighed into the extraction tube. Then, the sample was spiked by adding AFB₁ standard solution and left at laboratory room temperature (25°C) for approximately 30 min to allow the solvent to evaporate. After leaving the samples to undergo a process of equilibration, the extraction solvents were added and shaken at 3000 g for 3 min to ensure that the solvents had mixed thoroughly with the entire sample. Thereafter, 1 g of the selected salt (sodium chloride) with 4 g of anhydrous magnesium sulfate was added to the mixture and the shaking procedure was repeated in order to induce phase separation and AFB₁ partitioning. Subsequently, an aliquot of the supernatant layer (1 mL) was evaporated until dry under nitrogen gas. Following that, the precolumn derivatization of AFB₁ was done in terms of how to enhance its sensitivity. The residue was reconstituted in 900 μ l of 10% acetonitrile or in the mobile phase that had been optimized in this study. After that, 100 μ l of trifluoroacetic acid was added, followed by an incubation period at 50°C for 15 min. The derivatized solution was then centrifuged at 1000 g for 5 min before HPLC-FLD analysis took place.

The influence of the volume of extraction solvent on the signal response of AFB₁ was optimized (in this case, a mixture of acetonitrile/methanol 40/60 v/v was used as the extraction solvent). The volumes of extraction solvent compared were 10 mL and 20 mL. Subsequent extraction procedures were done as described above.

2.4. Validation of Methods

2.4.1. Final HPLC Analysis Used in Validation Experiments. The apparatus used for the analysis was a reverse phase HPLC (Shimadzu, Kyoto, Japan) equipped with a fluorescence detector. The analytical column used was a Symmetry[®] C-18 3.9x150 mm with 5 μ m particle size from Waters (Massachusetts) and the guard column was Silfilter STD C-18 3.0x10 mm. The column was maintained at 40°C. Analysis was run at a flow rate of 1 ml/min by an isocratic mobile phase using a mixture of acetonitrile/methanol/water (15/15/70 v/v/v). The total run time was 20 min. An aliquot of a 10 μ l sample extract was injected into the chromatographic system and detection was carried out by a fluorescence detector (excitation and emission wavelengths were 360 and 440 nm, respectively). Chromatograms were displayed with Class VP LC software.

2.4.2. Final Sample Preparation Method Used in Validation Experiments. 10 g of the sample and 20 mL of acetonitrile/methanol (40/60 v/v) were added to an extraction tube. Extraction was achieved through shaking the sample at

3000 g for 3 min. Thereafter, phase partition was induced using 1 g of sodium chloride and 4 g of anhydrous magnesium sulfate with shaking at 3000 g for 3 min. Subsequently, 1 ml of the organic layer was evaporated until dry under nitrogen gas. After evaporation, a precolumn derivatization of AFB₁ was prepared by reconstituting the residue in 900 μ l of 10% acetonitrile solution and adding 100 μ l of trifluoroacetic acid followed by incubation at 50 °C for 15 min. Finally, the derivatized solution was centrifuged at 1000 g for 5 min and an aliquot was transferred to the vials for HPLC-FLD analysis which was described previously.

The method was validated according to SANCO/12571/2013 which demonstrates the conformity of the analytical performances with criteria established in regulation (EC) no. 178/2010 [15]. The guidelines recommend the validation procedure for linearity, specificity, limit of detection (LOD), limit of quantitation (LOQ), accuracy, and precision. This study used the maximum permitted levels (MPLs) for the legislated mycotoxin in animal feed under the Animal Feed Quality Control Act B.E. 2525 of Thailand at 100 ng/g to consider as a maximum fortified concentration for the samples.

Linearity was tested by external standardization using matrix calibration curves constructed from AFB₁ standard solutions at 6 different concentrations within the range of 5–100 ng/g (5, 10, 20, 40, 60, and 100 ng/g). Analytical curves were established by plotting the peak areas which were used as the analytical signal response (y) versus the concentration of AFB₁ (x).

The specificity of the method was evaluated by comparing the retention times in the blank sample matrices and the samples were spiked with 100 ng/g of AFB₁ to ensure there was no interference in the retention time of the target analyte.

The sensitivity of the method was considered according to the LOD and LOQ. The LOD was calculated as the lowest concentration of the AFB₁ giving a signal response 3 times greater than the average of the baseline noise obtained from 10 independent blank samples of each matrix (S/N 3:1) and the LOQ was defined as an AFB₁ signal response 10 times greater than the average of the baseline noise obtained from 10 independent blank samples of each matrix (S/N 10:1).

The accuracy was tested through recovery studies by spiked AFB₁ standard solution at 3 different concentration levels (equivalent to 20, 40, and 100 ng/g) into blank sample matrices. Six replicates of each concentration were prepared for each matrix. The level of precision, which is expressed as relative standard deviations (RSDs), was estimated by performing intraday repeatability, expressed as RSD_r, and interday reproducibility, expressed as RSD_R, by spiked blank sample with AFB₁ standard solution at 3 concentration levels (20, 40, and 100 ng/g). Again, 6 replicates of each concentration were prepared for each matrix and determined within 1 day and in 3 consecutive days.

2.5. Limited Survey of AFB₁ Contamination in Selected Feedstuffs. A total of 120 samples, including 30 samples of each kind of feedstuffs (broken rice, peanuts, corn, and fishmeal), were randomly collected from swine farms in the western region of Thailand. All samples were sealed in plastic bags and stored at -20 °C until analysis.

3. Results and Discussion

3.1. Optimization of HPLC-FLD Conditions. To optimize the chromatographic separation, the study was carried out using four different solutions of mobile phase solvent as described above. The results showed that using methanol/water (50/50 v/v), acetonitrile/methanol/water (20/20/60 v/v/v), and acetonitrile/methanol/water (16/22/62 v/v/v) obtained poor signal responses and the AFB₁ was not separated from the unretained compound. The best mobile phase which gave the highest signal response was obtained and that offered adequate chromatographic separation with acetonitrile/methanol/water (15/15/70 v/v/v). This might be due to the fact that a high polarity of the mobile phase combination was able to exclude nonpolar substances from the C-18 analytical column [16]. However, broadened peak may occur when the mobile phase is composed of water over 70% [10].

HPLC-FLD has been the most popular analytical instrument for AFB₁ analysis. Nevertheless, fluorescent detection sensitivity of AFB₁ requires a conversion to more highly fluorescent derivatives (AFB_{2a}). Several derivatization methods are available, including precolumn treatment with trifluoroacetic acid or postcolumn derivatization with pyridinium hydrobromide [2, 17]. With an existence instrument, the method used the precolumn derivatization throughout the quantitative determinations. Generally, the residues were reconstituted using a mobile phase to reduce the solvent effects. However, in this study, the chromatographic response signal of the AFB₁ derivative was unsatisfied when dissolved with mobile phase (acetonitrile/methanol/water 15/15/70 v/v/v). This might be due to the fact that methanol affects the trifluoroacetic acid derivatization and AFB_{2a} could potentially be degraded by methanol, which would then result in a poor signal response [18, 19].

3.2. Optimization of Sample Preparation. In quantifying the substance, the sample preparation has a crucial impact on the accuracy and precision of the results, especially when the sample was contaminated at very low level. Usually, the sample was extracted and partitioned using acidified acetonitrile and salt as a drying agent followed by the purification of extractant with d-SPE [11, 12]. To reduce sample handling, in this study, a simple extraction QuEChERS-based procedure without d-SPE clean-up step was applied.

Following the current protocol, the optimization method was carried out using 10 g of broken rice as a representative. The conventional extraction procedures of AFB₁ from different matrices used a mixture of water and organic solvents such as acetonitrile, methanol, or acetone [20]. Therefore, different mixtures of extract solvents, which were acetonitrile (100%), acetonitrile/methanol (40/60 v/v), and methanol/water (50/50 v/v), were evaluated. An important characteristic of the QuEChERS procedure is the addition of salts, such as sodium chloride, sodium acetate, or sodium sulfate, which the typical QuEChERS method uses to separate interfering compounds from matrix and increase the solubility of the analyte into the organic phase [10, 14, 21]. Therefore, all of the extraction solvents were tested by adding either sodium chloride/magnesium sulfate (1/4 w/w)

TABLE 1: The results of validation method with mean value*(accuracy and precision), n=6.

Matrices	Spiked level (ng/g)	%Recovery*			Intra-day* (%RSD _r)			Inter- day* (%RSD _R)	LOD (ng/g)	LOQ (ng/g)
		Day 1	Day 2	Day 3	Day 1	Day 2	Day 3			
Broken rice	20	86.83	91.83	96.83	3.52	5.48	3.16	5.44	0.2	0.3
	40	82.50	89.83	90.83	3.73	4.30	4.25	4.12		
	100	84.00	93.66	92.50	3.61	3.35	2.34	5.77		
Peanut	20	106.57	109.37	109.37	3.34	5.89	1.98	1.42	0.6	0.9
	40	106.65	104.13	109.41	1.02	2.45	5.42	2.47		
	100	101.64	93.89	107.10	4.06	1.83	3.97	6.58		
Corn	20	90.10	90.98	91.72	6.77	5.63	4.89	0.89	0.6	0.8
	40	91.84	95.06	84.95	5.05	5.27	1.69	5.70		
	100	86.36	85.64	85.43	3.28	4.20	6.33	0.57		
Fish meal	20	109.85	108.88	102.31	5.02	2.96	6.18	6.33	1.2	1.5
	40	100.35	89.13	97.92	4.26	2.99	11.0	6.16		
	100	86.63	86.70	99.49	2.76	2.34	2.38	8.29		

or sodium acetate/magnesium sulfate (1/4 w/w). The results clearly showed that the highest efficiency to extract AFB₁ was achieved when using acetonitrile/methanol (40/60 v/v) combined with sodium chloride/magnesium sulfate (1/4 w/w). Generally, the initial QuEChERS method was developed for the determination of pesticide residue in fruits and vegetables which contain a high concentration of water (>80%), while feedstuffs have lower percentages of water composition than in those matrices. An earlier report suggested that the sample should be soaked in water prior to extraction [22]. However, the extraction procedure in this study was done without this soaking step.

Moreover, the present results showed that using acetonitrile (100%) and a mixture of acetonitrile/water (50/50 v/v) as an extraction solvent resulted in a low recovery of determination which is in accordance with Capriotti [23]. This might be due to the fact that AFB₁ is normally soluble in moderately polar organic solvents, e.g., methanol or chloroform, and scarcely soluble in water. Consequently, AFB₁ could not be completely extracted and remained in matrix [24]. Although acetonitrile/water (50/50 v/v) offered high signal response, its broadened tailing chromatogram overlapped with interference peaks. Moreover, the presence of a large amount of water in the solvent was adsorbed by matrices and tended to aggregate in the tube. Generally, the drying agents are anhydrous inorganic salts that acquire water for hydration. Therefore, the crystal form with large clumps of drying agent commonly occurred when exposed to water. Then, the results created a high viscosity of the extractant, reduced volume of supernatant layer, and extended duration for the evaporation step and a low signal response as well as percentage of recovery [10, 20].

The volume of extraction solvent (10 mL and 20 mL) was optimized. The results clearly showed that using 20 mL of extraction volume gave more satisfying data than using 10 mL. This is in agreement with an earlier study that used an adequate volume for extraction that increased the extraction efficiency by decreasing the matrix effect and enhanced the

release of the AFB₁ from the matrix resulting in high recoveries [10]. Therefore, 20 mL of acetonitrile/methanol (40/60 v/v) combined with sodium chloride (1g) and magnesium sulfate (4g) without any clean-up step was chosen as an extraction solution for AFB₁ analysis.

3.3. Method Performance. Performance characteristics of the optimized method were established by a validation procedure with spiked feeding stuff samples to test linearity, specificity, accuracy, precision, LOD, and LOQ. As a result, the analytical curves showed good linearity within the working range (5-100 ng/g), with coefficients (R^2) of determination higher than 0.9800. The discrimination between AFB₁ and interference compounds was defined as specificity. The absence of any signal response close to the retention time of AFB₁ in all matrices indicated that there were no matrix interferences in spite of the high complexity of the matrices. The obtained results have no coeluting peaks close to the retention time of the AFB₁ (Figure 1), demonstrating that it was a specification method.

The sensitivity of the method was considered according to the LOD and LOQ (Table 1). As presented, the calculated values in all matrices were found to have the LOD between 0.2 and 1.2 ng/g. The LOQ was in the range of 0.3-1.5 ng/g. Both the LOD and LOQ values were satisfactory for the AFB₁ detection in the feedstuffs, being far below the MPLs legislated values.

In order to check the accuracy of the method, recovery experiments were tested with a spiked AFB₁ standard solution at 3 different concentration levels (20, 40, and 100 ng/g). Average recoveries for all feedstuffs were in the range of 82.50-109.85% (Table 1) which demonstrated the conformity of the guidelines for the recovery of analysis range of 70-110%, according to SANCO/12571/2013. RSDs were calculated under RSD_r and RSD_R conditions to evaluate the precision of the method. In the present study, the RSD_r and RSD_R of most samples were 1.02-6.77% and 0.57-6.58%, respectively, except for the fishmeal matrix which obtained




FIGURE 1: Chromatograms of AFB₁ standard solution containing 100 ng/g and sample matrices (broken rice, peanut, corn, and fishmeal) were spiked with AFB₁ standard solution at 100 ng/g.

RSD_r at 11% on the third day and RSD_R was found to be between 6.16 and 8.29% (Table 1). However, the performed method was in the acceptable range of RSDs of less than 20% with the performance criteria requirement of the EC [15]. Additionally, the obtained results comply well with the Thai regulations for the detection of AFB₁ in feedstuff.

3.4. Occurrence of AFB₁ from Collected Samples. The developed method was applied for the analysis of a total of 120 feedstuff samples including 30 samples of each type of feedstuffs of broken rice, peanuts, corn, and fishmeal which were randomly collected from swine farms in the western region of Thailand where there is a high density of live-stock available. Several types of AFs analogues are naturally produced in food and feed. Obviously, AFB₁ contamination

in animal feed is the most predominant compared to other metabolites. Currently, of four major AFs analogues, AFB₁ is the only metabolite with maximum permitted levels (MPLs) for animal feed set under Directive 2002/32. A guidance value of the AFs contamination by the European Food Safety Authority (EFSA) in animal feed is limited between 5 ng/g for compound feed of dairy and young animals and 20 ng/g for all other feed materials. In Thailand, the safe limits of AFB₁ at 30-100 ng/g in animal feed (depending on the type of feed and animal species) have been regulated under the Animal Feed Quality Control Act B.E. 2525 of Thailand.

With regard to the data (Table 2), a low incidence (23.33%) with AFB₁ trace level contamination in broken rice samples was determined by the validated method. It is possible that AFB₁ is present mainly at the fungal invading

TABLE 2: Occurrence and levels of AFB₁ contamination in feedstuffs.

Sample matrices (n=120)	Number of samples and detected level (ng/g)						Concentration range (ng/g) (Mean±SD)
	n.d.	<4	4-20	21-50	51-100	>100	
Broken rice (30)	23	7	-	-	-	-	0.44-2.33 (0.27±0.61)
Peanut (30)	18	1	-	4	6	1	3.97-106.26 (23.29±33.76)
Corn (30)	7	10	7	5	1	-	0.88-50.29 (10.67±11.72)
Fishmeal (30)	24	4	2	-	-	-	1.06-10.35 (0.82±2.20)

area and remains on the outer layer of the grain which was removed by the degree of milling. As a result, AFB₁ in broken rice is usually found at trace level [25–27].

However, postharvest factors, for instance, storage under inappropriate conditions and being exposed to rain, may lead to increased AFB₁ contamination levels [28]. Likewise, Anjum [1] reported that AFB₁ contamination in broken rice is affected by rain. In addition, Thieu [25] has reported that broken rice could be contaminated with AFB₁ during harvesting and via a dehumidification process through sun-drying technique.

A high incidence of AFB₁ contamination in peanuts (40%) with a wide range from 3.97 to 106.27 ng/g was found in this study. Among those, 11 samples were contaminated with an AFB₁ level that exceeded the maximum levels set in Directive 2003/100/EC, amended Directive 2002/32 (20 ng/g), and 7 samples were contaminated with AFB₁ level higher than the permissible limits as established by the Animal Feed Quality Control Act B.E. 2525 of Thailand (50 ng/g). In contrast, our current study showed that the contamination level was higher than that found in peanuts from Thailand in a study conducted by Thanida [29]. Moreover, 93-99% of the 200 peanut samples collected from 2012 to 2015 were found to have lower than 4 ng/g (unpublished in-house data). This may be due to underestimation detected by the ELISA method used in previous survey.

Corn (maize) is one of the major ingredients used in animal feed which susceptible to infection by mycotoxin-producing fungi and, consequently, has the potential to become contaminated with mycotoxins [30]. The current results showed that the incidence of AFB₁ remained in 23 out of 30 corn samples (76.67%) with a wide range of concentrations (0.88-50.29 ng/g). Similarly, previous reports have shown that corn can become contaminated with AFB₁ in various ranges. Recently, AFs in corn ingredient were found with a high prevalence [31]. Thanida [29] reported AFB₁ contamination in corn samples from Thailand found ranging from 0.275 to 40 ng/g. As compared to the in-house data, 15% of the total 400 corn feed samples collected during 2012 to 2015 were contaminated with AFB₁ ranging from 51 to 100 ng/g while 7% of the samples contained AFB₁ over 100 ng/g (unpublished data). Our current study presented the notion that the AFB₁ contents were within the safe limits under the Animal Feed Quality Control Act B.E. 2525 of Thailand (for feedstuff set as 100 ng/g) and the FDA guidance (100-300 ng/g).

Typically, AFB₁ is not found in fishmeal because its matrix is unsuitable for the production of mycotoxin. It has

been presented that AFB₁ was not detected in any fishmeal samples [32]. However, fishmeal contaminated with AFB₁ has been reported. Anjum [1] found AFB₁ 24 ng/g from Pakistan during rainy season which was in accordance with the results of Mayahi [33] who found AFB₁ contamination at an average of 15 ng/g. As detected in the present study, 20% of the sample was found to be contaminated with AFB₁ in the range of 1.06-10.35 ng/g. This coincides with earlier results reported by Thanida [29] that AFB₁ contamination in fishmeal was found in the range of 1.67-11.90 ng/g. This indicates that fishmeal as a feedstuff presents no risk of AFB₁ contamination to animal health as the detected AFB₁ was not above the MPLs.

4. Conclusion

A modified method based on the QuEChERS procedure by a single extraction step without employing a clean-up step was developed for AFB₁ determination in feedstuffs. Good analytical results were obtained, including good linearity, specificity, accuracy, precision (repeatability and reproducibility), and analytical limits (LOD and LOQ). Therefore, it can be suggested as an alternative to expensive and time-consuming methods by using immune affinity columns or two steps of liquid/solid extraction procedure. Finally, an efficient and sensitive method was developed here that was applied to four different kinds of matrices which were detectable at low level. Considering that, the usefulness of the method could be applied as a regular monitoring method of AFB₁ in broken rice, peanuts, corn, fishmeal, and their related matrices.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Acknowledgments

This work was partially supported by the Kasetsart University Research and Development Institute (KURDI) and Center for Advanced Studies for Agriculture and Food, Institute for Advanced Studies (CASAF), Kasetsart University, under the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission, Ministry of Education, Thailand.

The authors gratefully acknowledge the staffs of the diagnostic and laboratory service units (Mycotoxin Unit) in Kamphaengsaen Campus for their technical and equipment support.

References

- [1] M. A. Anjum, S. H. Khan, A. W. Sahota, and R. Sardar, "Assessment of aflatoxin B1 in commercial poultry feed and feed ingredients," *Journal of Animal and Plant Sciences*, vol. 22, no. 2, pp. 268–272, 2012.
- [2] N. Tansakul, S. Limsuwan, J. Böhm, M. Hollmann, and E. Razzazi-Fazeli, "Aflatoxins in selected Thai commodities," *Food Additives & Contaminants: Part B. Surveillance*, vol. 6, no. 4, pp. 254–259, 2013.
- [3] M. Oplawska-Stachowiak, N. Sajic, Y. Xu et al., "Fast and sensitive aflatoxin B1 and total aflatoxins ELISAs for analysis of peanuts, maize and feed ingredients," *Food Control*, vol. 63, pp. 239–245, 2016.
- [4] N. Arroyo-Manzanares, L. Gámiz-Gracia, A. M. Gámiz-Gracia, J. J. Soto-Chinchilla, A. M. García-Campaña, and L. E. García-Ayuso, "On-line preconcentration for the determination of aflatoxins in rice samples by micellar electrokinetic capillary chromatography with laser-induced fluorescence detection," *Electrophoresis*, vol. 31, no. 13, pp. 2180–2185, 2010.
- [5] W. S. Khayoon, B. Saad, T. P. Lee, and B. Salleh, "High performance liquid chromatographic determination of aflatoxins in chilli, peanut and rice using silica based monolithic column," *Food Chemistry*, vol. 133, no. 2, pp. 489–496, 2012.
- [6] International Agency for Research on Cancer, "Some traditional herbal medicines, some mycotoxins, naphthalene and styrene," *IARC Monograph on the evaluation of carcinogenic risk to human*, vol. 82, no. 2-3, pp. 171–300, 2002.
- [7] S. A. Aly and W. Anwer, "Effect of naturally contaminated feed with aflatoxins on performance of laying hens and the carryover of aflatoxin B1 residues in table eggs," *Pakistan Journal of Nutrition*, vol. 8, no. 2, pp. 181–186, 2009.
- [8] S. M. Herzallah, "Determination of aflatoxins in eggs, milk, meat and meat products using HPLC fluorescent and UV detectors," *Food Chemistry*, vol. 114, no. 3, pp. 1141–1146, 2009.
- [9] R. W. Han, N. Zheng, J. Q. Wang, Y. P. Zhen, X. M. Xu, and S. L. Li, "Survey of aflatoxin in dairy cow feed and raw milk in China," *Food Control*, vol. 34, no. 1, pp. 35–39, 2013.
- [10] A. Y. Sirhan, G. H. Tan, A. Al-Shunnaq, L. Abdulra'uf, and R. C. S. Wong, "QuEChERS-HPLC method for aflatoxin detection of domestic and imported food in Jordan," *Journal of Liquid Chromatography & Related Technologies*, vol. 37, no. 3, pp. 321–342, 2014.
- [11] N. Arroyo-Manzanares, A. M. García-Campaña, and L. Gámiz-Gracia, "Multiclass mycotoxin analysis in *Silybum marianum* by ultra high performance liquid chromatography-tandem mass spectrometry using a procedure based on QuEChERS and dispersive liquid-liquid microextraction," *Journal of Chromatography A*, vol. 1282, pp. 11–19, 2013.
- [12] U. Koesukwiwat, K. Sanguankaew, and N. Leepipatpiboon, "Evaluation of a modified QuEChERS method for analysis of mycotoxins in rice," *Food Chemistry*, vol. 153, pp. 44–51, 2014.
- [13] N. Arroyo-Manzanares, J. F. Huertas-Pérez, A. M. García-Campaña, and L. Gámiz-Gracia, "Simple methodology for the determination of mycotoxins in pseudocereals, spelt and rice," *Food Control*, vol. 36, no. 1, pp. 94–101, 2014.
- [14] M. Á. González-Curbelo, B. Socas-Rodríguez, A. V. Herrera-Herrera, J. González-Sálamo, J. Hernández-Borges, and M. Á. Rodríguez-Delgado, "Evolution and applications of the QuEChERS method," *TrAC - Trends in Analytical Chemistry*, vol. 71, pp. 169–185, 2015.
- [15] EC, "Commission regulation (EU) No 178/2010 of 2 March 2010 amending regulation (EC) No 401/2006 of 23 February 2006 laying down the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs," *Official Journal of the European Union*, 2010.
- [16] A. Cismileanu, G. Voicu, V. Ciuca, and M. Ionescu, "Determination of aflatoxin B1 in cereal-based feed by a high-performance chromatographic method," *Lucrari Stiintific Medicinaverinara*, vol. 41, pp. 565–569, 2008.
- [17] M. Muscarella, M. Iammarino, D. Nardiello et al., "Validation of a confirmatory analytical method for the determination of aflatoxins B1, B2, G1 and G2 in foods and feed materials by HPLC with on-line photochemical derivatization and fluorescence detection," *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, vol. 26, no. 10, pp. 1402–1410, 2009.
- [18] W. T. Kok, "Derivatization reactions for the determination of aflatoxins by liquid chromatography with fluorescence detection," *Journal of Chromatography B: Biomedical Sciences and Applications*, vol. 659, no. 1-2, pp. 127–137, 1994.
- [19] S. M. Pearson, A. A. G. Candlish, K. E. Aidoo, and J. E. Smith, "Determination of aflatoxin levels in pistachio and cashew nuts using immunoaffinity column clean-up with HPLC and fluorescence detection," *Biotechnology Techniques*, vol. 13, no. 2, pp. 97–99, 1999.
- [20] E. Reiter, J. Zentek, and E. Razzazi, "Review on sample preparation strategies and methods used for the analysis of aflatoxins in food and feed," *Molecular Nutrition & Food Research*, vol. 53, no. 4, pp. 508–524, 2009.
- [21] E. A. Orlando and A. V. C. Simionato, "Extraction of tetracycline antibiotic residues from fish file: Comparison and optimization of different procedures using liquid chromatography with fluorescence detection," *Journal of Chromatography A*, vol. 1307, pp. 111–118, 2013.
- [22] S. C. Cunha and J. O. Fernandes, "Development and validation of a method based on a QuEChERS procedure and heart-cutting GC-MS for determination of five mycotoxins in cereal products," *Journal of Separation Science*, vol. 33, no. 4-5, pp. 600–609, 2010.
- [23] A. L. Capriotti, C. Cavaliere, S. Piovesana, R. Samperi, and A. Laganà, "Multiclass screening method based on solvent extraction and liquid chromatography-tandem mass spectrometry for the determination of antimicrobials and mycotoxins in egg," *Journal of Chromatography A*, vol. 1268, pp. 84–90, 2012.
- [24] W. S. Khayoon, B. Saad, C. B. Yan et al., "Determination of aflatoxins in animal feeds by HPLC with multifunctional column clean-up," *Food Chemistry*, vol. 118, no. 3, pp. 882–886, 2010.
- [25] N. Q. Thieu, B. Ogle, and H. Pettersson, "Screening of Aflatoxins and Zearalenone in feedstuffs and complete feeds for pigs in Southern Vietnam," *Tropical Animal Health and Production*, vol. 40, no. 1, pp. 77–83, 2008.
- [26] S. Z. Iqbal, M. R. Asi, A. Arino, N. Akram, and M. Zuber, "Aflatoxin contamination in different fractions of rice from Pakistan and estimation of dietary intakes," *Mycotoxin Research*, vol. 28, no. 3, pp. 175–180, 2012.

- [27] S. Choi, H. Jun, J. Bang et al., "Behaviour of *Aspergillus flavus* and *Fusarium graminearum* on rice as affected by degree of milling, temperature, and relative humidity during storage," *Food Microbiology*, vol. 46, pp. 307–313, 2015.
- [28] K. R. N. Reddy, C. S. Reddy, and K. Muralidharan, "Detection of *Aspergillus* spp. and aflatoxin B1 in rice in India," *Food Microbiology*, vol. 26, no. 1, pp. 27–31, 2009.
- [29] W. Thanita, C. Surapol, K. Saowanit, and W. Sutha, "Aflatoxin contamination in feedstuffs and complete feed for layer, broiler and swine in Songkhla Province," *Thaksin*, vol. 11, no. 1, pp. 8–18, 2008.
- [30] F. Wu and G. P. Munkvold, "Mycotoxins in ethanol co-products: Modeling economic impacts on the livestock industry and management strategies," *Journal of Agricultural and Food Chemistry*, vol. 56, no. 11, pp. 3900–3911, 2008.
- [31] F. Granados-Chinchilla, A. Molina, G. Chavarría, M. Alfaro-Cascante, D. Bogantes-Ledezma, and A. Murillo-Williams, "Aflatoxins occurrence through the food chain in Costa Rica: Applying the One Health approach to mycotoxin surveillance," *Food Control*, vol. 82, pp. 217–226, 2017.
- [32] B. M. Kokic, I. S. Cabarkapa, I. D. Levicet, and etal., "Screening of mycotoxins in animal feed from the region of Vojodina," *Proc Nat SciMaticaSrpska Novi Sad*, vol. 117, pp. 87–96, 2009.
- [33] M. Mayahi, R. J. Mohammad, and S. Negin, "Isolation of *Aspergillus* spp. and determination of aflatoxin level in fish meal, maize and soya meal," *ShahidChamran University Journal of Science*, vol. 17, pp. 95–105, 2007.