

Do Major League Baseball Team Physicians Harvest the Semitendinosus From the Drive Leg or Landing Leg When Performing Ulnar Collateral Ligament Reconstruction on Elite Baseball Pitchers?

Brandon J. Erickson,^{*†} MD, Peter N. Chalmers,[‡] MD, Jeffrey R. Dugas,[§] MD, Bernard R. Bach Jr,[†] MD, Gregory P. Nicholson,[†] MD, Nikhil N. Verma,[†] MD, Christopher S. Ahmad,^{||} MD, and Anthony A. Romeo,[†] MD

Investigation performed at Midwest Orthopaedics at Rush University Medical Center, Chicago, Illinois, USA

Background: Hamstring autograft is a common graft choice when performing ulnar collateral ligament reconstruction (UCLR).

Purpose/Hypothesis: The purpose of this study was to survey Major League Baseball (MLB) team physicians and determine whether these physicians harvest the hamstring (semitendinosus or gracilis) from the drive leg (ipsilateral to surgical site) or landing leg (contralateral to surgical site) when performing UCLR on elite-level pitchers. The hypothesis was that the majority of surgeons harvest the hamstring from the drive leg when performing a UCLR.

Study Design: Descriptive epidemiology study.

Methods: Overall, 52 MLB team orthopaedic surgeons were sent the 5-question online survey. The survey assessed surgeon UCLR volume, surgical technique, which leg the hamstring graft was harvested from, the reasoning for choosing that particular leg, and whether the surgeon would change their practice if evidence showed the hamstring from one of the legs was more important than the other. The survey was sent out 5 separate times to maximize the response rate.

Results: Forty (77%) MLB team physicians completed the survey. The largest number of surgeons ($n = 16$; 40%) performed between 5 and 14 UCLRs annually, while 6 (15%) performed more than 50 UCLRs annually. Most surgeons ($n = 23$; 57.5%) used the docking technique. Significantly more surgeons harvested the hamstring from the landing leg ($n = 29$; 72.5%) compared with the drive leg ($n = 11$; 27.5%) ($P = .007$). More surgeons cited the reason for their choice of leg as a belief that the hamstring they harvested plays less of a role in the ability of a pitcher to generate a forceful pitch ($n = 25$; 62.5%) than for logistical reasons in the operating room ($n = 15$; 37.5%); this difference was not statistically significant. Significantly more surgeons would change their practice ($n = 35$; 87.5%) if evidence showed the hamstrings from a specific (drive or landing) leg to be more active in the throwing motion compared with those who would not ($P < .001$).

Conclusion: When performing a UCLR using hamstring autograft, the majority of MLB team physicians harvest the hamstring from the landing leg as opposed to the drive leg, as they believe the hamstring from the landing leg plays less of a role in generating a forceful pitch than that from the drive leg.

Keywords: ulnar collateral ligament reconstruction; UCLR; Major League Baseball; MLB; pitcher; hamstring; Tommy John

Injuries to the ulnar collateral ligament (UCL) have become commonplace among elite-level and adolescent baseball pitchers.^{3,6,11,12,15,26,34} While some pitchers are fortunate enough to return to sport (RTS) after a period of rest and rehabilitation, most pitchers who wish to RTS at a high

level after sustaining such an injury require a UCL reconstruction (UCLR).^{24,35} Studies have described many UCLR techniques that afford pitchers the ability to RTS at rates greater than 85%, although these studies have not found a specific technique to be superior to the rest.^{3,5,10,12,16,36,44} While UCL repair with an internal brace has received recent attention with promising short-term results, the current standard treatment for baseball pitchers who wish to RTS is a UCLR.^{3,7-9,26}

The Orthopaedic Journal of Sports Medicine, 5(7), 2325967117713987
DOI: 10.1177/2325967117713987
© The Author(s) 2017

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<http://creativecommons.org/licenses/by-nc-nd/3.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For reprints and permission queries, please visit SAGE's website at <http://www.sagepub.com/journalsPermissions.nav>.

When performing a UCLR, several graft options for reconstruction include palmaris longus autograft, hamstring autograft, allograft, and others.^{3,10,38} Studies have shown reliable RTS rates from each graft choice, with no superiority of one choice over another, although no randomized comparative trials exist.^{3,13} When a patient does not have a palmaris longus tendon in either arm, the palmaris tendon appears small in size or damaged (torn, ossified, etc), or the patient is undergoing a revision UCLR, many surgeons choose to harvest a hamstring tendon (gracilis or semitendinosus) from either the ipsilateral leg or contralateral leg, similar to an anterior cruciate ligament reconstruction (ACLR).^{1,28} The ipsilateral leg is commonly known as the *drive or lag leg* while the contralateral leg is known as the *landing or lead leg*.

Hamstring tendon harvest is a relatively benign procedure with low reported graft site complication rates in the ACLR literature, including injury to the infrapatellar branch of the saphenous nerve and wound dehiscence.^{29,32,37} However, one complication that can have a long-lasting effect on patients is persistent hamstring weakness on the side of the harvest. While studies have shown varying amounts of hamstring regrowth after gracilis and semitendinosus harvest, some studies have shown persistent weakness in knee flexion at final follow-up.^{2,27,40,42} Long-term follow-up has demonstrated essentially permanent atrophy of those muscles attached to the harvested tendons.⁴¹ It is unclear whether this residual weakness has an effect on pitching mechanics in high-level baseball pitchers.

Therefore, the purpose of this study was to survey Major League Baseball (MLB) team physicians and determine if these physicians harvest the hamstring from the drive leg or landing leg when performing a UCLR on elite-level pitchers. We hypothesized that the majority of MLB team physicians harvest the hamstring tendon from the drive leg when performing a UCLR on elite-level pitchers using a hamstring autograft.

METHODS

We created a 5-question survey (Table 1) on the website SurveyMonkey (<http://www.surveymonkey.com>) and sent

TABLE 1
Five-Question Survey Administered to Major League Baseball Team Orthopaedic Surgeons^a

1. How many ulnar collateral ligament reconstructions (UCLRs) do you perform in a year?
 - a. <5
 - b. 5-14
 - c. 15-29
 - d. 30-50
 - e. >50
2. What is your preferred surgical technique when performing a UCLR?
 - a. ASMI
 - b. DANE-TJ
 - c. Docking
 - d. Double docking
 - e. Modified Jobe
 - f. Other (please specify)
3. When performing a UCLR on a pitcher using a hamstring autograft, which leg do you routinely harvest the graft from?
 - a. Drive leg (also known as the trail, push-off, or back leg)
 - b. Landing leg (also known as the stride, lead, or front leg)
4. What is your reasoning for your answer to the previous question (question #3)?
 - a. Logistical reasons for set-up, prepping, etc. in the operating room
 - b. I believe that the hamstring I use plays less of a role in the ability of a pitcher to generate a forceful pitch than the opposite leg, and can be sacrificed with less of a detriment to the pitch.
 - c. Other (please specify)
5. If evidence showed the hamstrings from one specific (drive or landing) leg to be more active in the throwing motion, would you change your practice to avoid harvesting the more active hamstrings?
 - a. Yes
 - b. No

^aASMI, American Sports Medicine Institute; DANE-TJ, Dave Altchek-Neal ElAttrache Tommy John.

the survey to 52 MLB team orthopaedic surgeons. Surgeons were identified by team websites, Major League Baseball Team Physicians Association website (<http://mlbtpa.org/index.php/team-physicians>), and press releases. Only

*Address correspondence to Brandon J. Erickson, MD, Midwest Orthopaedics at Rush, Rush University Medical Center, 1611 West Harrison Street, Suite 300, Chicago, IL 60612, USA (email: brandon.j.erickson@gmail.com).

†Midwest Orthopaedics at Rush, Rush University Medical Center, Chicago, Illinois, USA.

‡Department of Orthopaedic Surgery, University of Utah, Salt Lake City, Utah, USA.

§American Sports Medicine Institute, Birmingham, Alabama, USA.

||Department of Orthopaedic Surgery, Columbia University, New York, New York, USA.

One or more of the authors has declared the following potential conflicts of interest or source of funding: J.R.D. receives royalties from Arthrex and Oakstone Publishing, is a paid consultant for Arthrex and Theralase, and has stock/stock options in Theralase and Topical Gear. B.R.B. is a team physician for the Chicago White Sox; receives research support from Arthrex, ConMed Linvatec, DJ Orthopaedics, Ossur, Smith & Nephew, and Tornier; and receives publishing royalties from SLACK Inc. G.P.N. is a physician for the Chicago White Sox, receives royalties from Innomed and SLACK Inc, is a paid consultant for Tornier, receives research support from Tornier, and has stock/stock options in Zimmer. N.N.V. is a team physician for the Chicago White Sox; receives research support from Arthrex, Arthrosurface, DJ Orthopaedics, Ossur, Smith & Nephew, Athletico, ConMed Linvatec, Miomed, and Mitek; has stock/stock options in Cymedica, Minivasive, and Omeros; is a paid consultant for Minivasive, Orthospace, and Smith & Nephew; and receives royalties from SLACK Inc, Smith & Nephew, and Vindico Medical-Orthopedics Hyperguide. C.S.A. is the head team physician for the New York Yankees, receives royalties from Arthrex, is a paid consultant or Athrex, has stock/stock options in At Peak, receives publishing royalties from Lead Player, and receives research support from Major League Baseball and Stryker. A.A.R. is a team physician for the Chicago White Sox; receives royalties from Arthrex and SLACK Inc; is a paid consultant for Arthrex; is a paid presenter or speaker for Arthrex; and receives research support from DJO Surgical, Ossur, and Smith & Nephew.

Ethical approval was not sought for the present study.




Figure 1. Responses to “How many ulnar collateral ligament reconstructions (UCLRs) do you perform in a year?”

orthopaedic surgeons with a valid email address were included in this study. The team surgeons were determined by internet web searches. In 2016, the survey was sent out to the surgeons first on October 19 (round 1), and a reminder email was sent on November 3 (round 2), November 10 (round 3), December 7 (round 4), and December 21 (round 5) to those who had not yet completed the survey. The responses were kept confidential, and the data were organized via the SurveyMonkey web tool. We were blinded as to the surgeons’ names. The team physicians were instructed to respond to all relevant questions, and the survey was arranged such that the participant could not complete the survey without answering all the questions. Institutional review board approval was not necessary for this survey study.

Statistics

Descriptive statistics were calculated. Chi-square and Fisher exact tests were used as appropriate, depending on cell populations to compare the distribution of responses to a null hypothesis of a random distribution. In addition, the distribution of responses of those surgeons who preferred the landing leg were compared with those who preferred the drive leg using chi-square and Fisher exact tests as appropriate, depending on cell populations.

RESULTS

A total of 40 (77%) MLB team physicians responded to the survey. The majority of surgeons (n = 16; 40%) self-reported the performance of between 5 and 14 UCLRs annually, while 6 (15%) performed >50 UCLRs annually (Figure 1). Most surgeons (n = 23; 57.5%) performed UCLR using the docking technique, with 3 using the modified docking/docking plus technique (Figure 2).

Significantly more surgeons harvested the hamstring from the landing leg (n = 29; 72.5%) than the drive leg (n = 11; 27.5%) (P = .007). Overall, most surgeons cited the




Figure 2. Responses to “What is your preferred surgical technique when performing a UCLR?” ASMI: American Sports Medicine Institute; DANE-TJ: Dave Altchek–Neal ElAttrache Tommy John; UCLR, ulnar collateral ligament reconstruction.




Figure 3. Comparison of responses to why surgeons choose to harvest the leg from the landing leg versus the drive leg based on which leg they harvest their graft from.

reason for their choice of leg as a belief that the hamstring they harvested plays less of a role in the ability of a pitcher to generate a forceful pitch than the opposite leg (n = 25; 62.5%) rather than for logistical reasons in the operating room (n = 15; 37.5%); however, this difference was not statistically significant (P = .155). When responses were separated out based on which leg the surgeon harvested the hamstring from, more surgeons who harvested the hamstring from the landing leg did so because of a belief that the landing leg plays less of a role in generating a forceful pitch (21/29, 72%) while surgeons who harvested the hamstring from the drive leg did so more commonly for logistical reasons such as operating simultaneously on the arm and leg (7/11, 64%), although this difference was not statistically significant (P = .065) (Figure 3). There were no significant differences in technique between those surgeons who preferred the drive leg (8/11, 73% docking and modifications and 3/11, 27% Jobe and modifications) over the landing leg (21/29, 72% docking and modifications and 8/29, 28% Jobe and modifications) (P > .999); nor were there any significant differences in case volume between those surgeons

who preferred the drive leg (4/11, 36% <15 UCLRs per year and 7/11, 64% ≥15 UCLRs per year) over the landing leg (17/29, 59% <15 UCLRs per year and 12/29, 41% ≥15 UCLRs per year) ($P = .293$).

Significantly more surgeons would change their practice ($n = 35$; 87.5%) than would not ($n = 5$; 12.5%) to avoid harvesting the more active hamstring if evidence showed the hamstrings from a specific leg (drive or landing) to be more active in the throwing motion ($P < .001$). Although not statistically significant, those surgeons who preferred the drive leg were no more likely to change their practice if evidence showed that hamstrings from a specific leg were more active in the throwing motion (10/11, 91%) compared with surgeons who preferred the landing leg (24/29, 83%) ($P > .999$).

DISCUSSION

It has been well documented that the number of UCLRs performed in both elite and adolescent baseball pitchers is rising.^{3,14,15,26} As the number of UCLRs continues to increase, it is important to determine how to optimize this procedure to afford pitchers the best chance at returning to sport at the same or higher level as before surgery. Because no clinical data exist on the importance of the hamstring tendons of the drive leg and landing leg in the baseball pitch, the expert opinion of MLB team physicians who perform UCLR on the most elite baseball pitchers and are most familiar with high-level pitching mechanics may provide the highest level of evidence available. Our hypothesis was incorrect, as the majority of MLB team physicians (72.5%) harvested the hamstring tendon from the landing leg when performing a UCLR using hamstring autograft as opposed to the drive leg.

Hamstring autograft has recently gained popularity as a viable graft choice when performing a UCLR in adolescent as well as elite-level athletes.^{10,34} The gracilis, as well as the semitendinosus, have both been used as graft options for UCLR, although typically only 1 tendon is harvested when performing a UCLR as opposed to an ACLR, in which both tendons are commonly harvested.³² Multiple studies have reported encouraging results when performing a UCLR with hamstring autograft.^{3,34} Erickson et al¹³ performed a study in which the authors compared palmaris longus autograft, hamstring autograft, and allograft as graft choices in UCLR to determine if differences in outcomes (RTS rate, clinical outcomes scores, and reoperation rates) existed between grafts. These authors found no significant difference in clinical outcome scores (Kerlan-Jobe Orthopaedic Clinic, Timmerman-Andrews), RTS rates, reoperations, or complications between graft choices. However, it is currently unclear which leg the hamstring graft should be harvested from in a baseball pitcher to minimize any effect the graft harvest may have on the pitching motion.

Harvesting the hamstring tendons leads to minimal acute graft site morbidity, such as leg soreness and the possibility of numbness in the distribution of the infrapatellar branch of the saphenous nerve.³² However, a more

significant issue with harvesting the hamstrings is persistent knee flexion weakness as well as weakness in internal rotation.^{27,39,43} Konrath et al²⁷ reviewed 20 patients who underwent an ACLR using hamstring autograft and found that knee flexion, extension, and internal rotation strength were significantly lower in the leg where the hamstring had been harvested compared with the contralateral limb (all P s <.05). The authors also noted that only 35% of patients showed regeneration of both the gracilis and semitendinosus on magnetic resonance imaging. Similarly, Tashiro et al⁴³ found significant knee flexion weakness after hamstring harvest for ACLR. Hence, the residual weakness present in the leg from which the graft was harvested may affect the pitcher's throwing motion, as motion analysis studies have shown significant lower-extremity muscle activity during the pitching motion.⁴ Therefore, mitigating this effect by harvesting the hamstring graft from the leg that contributes less to the overall pitch is imperative.

While a significant amount of research has focused on upper-extremity mechanics as it relates to the overhand pitch, much less attention has been paid to the lower extremity.¹⁷⁻²³ However, the limited data that are available have shown significant muscle activation in the lower extremities during the baseball pitch.^{4,25,31,45} These studies have found that, in pitching, ground reaction and braking forces are significant; and significant torques pass through the hip and knee that are created and balanced by the hamstrings, suggesting that these muscles are important for pitch velocity, command, and control.^{4,25,31} Furthermore, Montgomery and Knudson³³ found that pitch speed could be increased by increasing the pitcher's stride length, something that can be directly related to hamstring strength.

While the focus of the pitch is often the shoulder and elbow, a significant amount of the force necessary for velocity creation is generated in the legs and trunk and is then transferred up the body into the shoulder, elbow, and finally the ball.^{4,25,30,31} The lower extremities are also important for balance and braking and may therefore be important for command and control. Therefore, loss of hip extensor and knee flexor power may not be trivial for an elite pitcher. As a result, appreciating which leg is more important to the overhand pitch and therefore which leg is optimal for hamstring autograft for a UCLR may lead to better postoperative outcomes. No current literature exists regarding this dilemma.

The results of the current study showed that significantly more MLB team orthopaedic surgeons harvest the hamstring for a UCLR from the landing leg. Furthermore, of the team orthopaedic surgeons who harvested the hamstring from the landing leg, the majority (72.4%) of them did so because they believe the landing leg is less important to the overhand pitch than the drive leg. Conversely, of the team orthopaedic surgeons who harvested the hamstring from the drive leg, the majority (63.6%) did so for logistical purposes in the operating room. Such logistical reasons can include operating simultaneously on the leg and elbow, inability to prep and drape contralateral extremities due to the size of the operating room, and others. Hence, only 10% of the surgeons surveyed believe the hamstrings from

the landing leg are more important in generating a forceful baseball pitch (meaning only 4 surgeons responded that they would harvest the hamstring from the drive leg because they believed the hamstrings from the landing leg are more important in the baseball pitch). This is an important finding, as it shows that a majority of surgeons who care for elite-level baseball athletes believe the hamstrings of the drive leg to be more important than the hamstrings of the landing leg. While further studies evaluating muscle activation of the hamstrings of both legs during the baseball pitch are necessary to definitively answer this question, the expert opinion of MLB team physicians can help guide surgeons performing UCLR on baseball players in the interim.

Limitations

As with any survey study, this study has several limitations. These results are level 5 evidence and represent the opinions of team physicians from a single sport. While these physicians take care of athletes from all sports and have expertise in multiple areas, the questions were only aimed at baseball pitchers. Hence, the results may not be translatable to females, softball pitchers, athletes in other overhead sports, and others. There is the small possibility that a survey respondent clicked the wrong answer choice accidentally. This survey did not address all possible technique variables (whether the gracilis or semitendinosus was harvested) or rehabilitation protocols. Furthermore, although an exhaustive search was performed to locate all MLB team physicians, there are some who may have been missed.

CONCLUSION

When performing a UCLR using a hamstring autograft, the majority of MLB team physicians harvest the hamstring from the landing leg as opposed to the drive leg, as they believe the hamstring from the landing leg plays less of a role in generating a forceful pitch than the hamstring from the drive leg.

REFERENCES

1. Bottoni CR, Smith EL, Shaha J, et al. Autograft versus allograft anterior cruciate ligament reconstruction: a prospective, randomized clinical study with a minimum 10-year follow-up. *Am J Sports Med.* 2015;43:2501-2509.
2. Burks RT, Crim J, Fink BP, Boylan DN, Greis PE. The effects of semitendinosus and gracilis harvest in anterior cruciate ligament reconstruction. *Arthroscopy.* 2005;21:1177-1185.
3. Cain EL Jr, Andrews JR, Dugas JR, et al. Outcome of ulnar collateral ligament reconstruction of the elbow in 1281 athletes: results in 743 athletes with minimum 2-year follow-up. *Am J Sports Med.* 2010;38:2426-2434.
4. Campbell BM, Stodden DF, Nixon MK. Lower extremity muscle activation during baseball pitching. *J Strength Cond Res.* 2010;24:964-971.
5. Dines JS, ElAttrache NS, Conway JE, Smith W, Ahmad CS. Clinical outcomes of the DANE TJ technique to treat ulnar collateral ligament insufficiency of the elbow. *Am J Sports Med.* 2007;35:2039-2044.
6. Dugas J, Chronister J, Cain EL Jr, Andrews JR. Ulnar collateral ligament in the overhead athlete: a current review. *Sports Med Arthrosc.* 2014;22:169-182.
7. Dugas JR. Ulnar collateral ligament repair: an old idea with a new wrinkle. *Am J Orthop (Belle Mead NJ).* 2016;45:124-127.
8. Dugas JR, Walters BL, Beason DP, Fleisig GS, Chronister JE. Biomechanical comparison of ulnar collateral ligament repair with internal bracing versus modified Jobe reconstruction. *Am J Sports Med.* 2016;44:735-741.
9. Erickson BJ, Bach BR Jr, Bush-Joseph CA, Verma NN, Romeo AA. Medial ulnar collateral ligament reconstruction of the elbow in Major League Baseball players: where do we stand? *World J Orthop.* 2016;7:355-360.
10. Erickson BJ, Bach BR Jr, Cohen MS, et al. Ulnar collateral ligament reconstruction: the Rush experience. *Orthop J Sports Med.* 2016;4:2325967115626876.
11. Erickson BJ, Chalmers PN, Bush-Joseph CA, Romeo AA. Predicting and preventing injury in Major League Baseball. *Am J Orthop (Belle Mead NJ).* 2016;45:152-156.
12. Erickson BJ, Chalmers PN, Bush-Joseph CA, Verma NN, Romeo AA. Ulnar collateral ligament reconstruction of the elbow: a systematic review of the literature. *Orthop J Sports Med.* 2015;3:2325967115618914.
13. Erickson BJ, Cvetanovich GL, Frank RM, et al. Do clinical results and return-to-sport rates after ulnar collateral ligament reconstruction differ based on graft choice and surgical technique? *Orthop J Sports Med.* 2016;4:2325967116670142.
14. Erickson BJ, Gupta AK, Harris JD, et al. Rate of return to pitching and performance after Tommy John surgery in major League Baseball pitchers. *Am J Sports Med.* 2014;42:536-543.
15. Erickson BJ, Nwachukwu BU, Rosas S, et al. Trends in medial ulnar collateral ligament reconstruction in the United States: a retrospective review of a large private-payer database from 2007 to 2011. *Am J Sports Med.* 2015;43:1770-1774.
16. Erickson BJ, Romeo AA. The ulnar collateral ligament injury: evaluation and treatment. *J Bone Joint Surg Am.* 2017;99:76-86.
17. Erickson BJ, Sgori T, Chalmers PN, et al. The impact of fatigue on baseball pitching mechanics in adolescent male pitchers. *Arthroscopy.* 2016;32:762-771.
18. Escamilla RF, Andrews JR. Shoulder muscle recruitment patterns and related biomechanics during upper extremity sports. *Sports Med.* 2009;39:569-590.
19. Escamilla RF, Barrentine SW, Fleisig GS, et al. Pitching biomechanics as a pitcher approaches muscular fatigue during a simulated baseball game. *Am J Sports Med.* 2007;35:23-33.
20. Fleisig GS, Andrews JR, Dillman CJ, Escamilla RF. Kinetics of baseball pitching with implications about injury mechanisms. *Am J Sports Med.* 1995;23:233-239.
21. Fleisig GS, Barrentine SW, Zheng N, Escamilla RF, Andrews JR. Kinematic and kinetic comparison of baseball pitching among various levels of development. *J Biomech.* 1999;32:1371-1375.
22. Fleisig GS, Kingsley DS, Loftice JW, et al. Kinetic comparison among the fastball, curveball, change-up, and slider in collegiate baseball pitchers. *Am J Sports Med.* 2006;34:423-430.
23. Fleisig GS, Leaddon CE, Laughlin WA, et al. Biomechanical performance of baseball pitchers with a history of ulnar collateral ligament reconstruction. *Am J Sports Med.* 2015;43:1045-1050.
24. Ford GM, Genuario J, Kinkartz J, Githens T, Noonan T. Return-to-play outcomes in professional baseball players after medial ulnar collateral ligament injuries: comparison of operative versus nonoperative treatment based on magnetic resonance imaging findings. *Am J Sports Med.* 2016;44:723-728.
25. Guido JA Jr, Werner SL. Lower-extremity ground reaction forces in collegiate baseball pitchers. *J Strength Cond Res.* 2012;26:1782-1785.
26. Hodgins JL, Vitale M, Arons RR, Ahmad CS. Epidemiology of medial ulnar collateral ligament reconstruction: a 10-year study of New York State. *Am J Sports Med.* 2016;44:729-734.

27. Konrath JM, Vertullo CJ, Kennedy BA, Bush HS, Barrett RS, Lloyd DG. Morphologic characteristics and strength of the hamstring muscles remain altered at 2 years after use of a hamstring tendon graft in anterior cruciate ligament reconstruction. *Am J Sports Med.* 2016;44:2589-2598.
28. Leiter JR, Gourlay R, McRae S, de Korompay N, Macdonald PB. Long-term follow-up of ACL reconstruction with hamstring autograft. *Knee Surg Sports Traumatol Arthrosc.* 2014;22:1061-1069.
29. Letartre R, Isida R, Pommepuy T, Miletic B. Horizontal posterior hamstring harvest. *Orthop Traumatol Surg Res.* 2014;100:959-961.
30. MacWilliams BA, Choi T, Perezous MK, Chao EY, McFarland EG. Characteristic ground-reaction forces in baseball pitching. *Am J Sports Med.* 1998;26:66-71.
31. Milewski MD, Ounpuu S, Solomito M, Westwell M, Nissen CW. Adolescent baseball pitching technique: lower extremity biomechanical analysis. *J Appl Biomech.* 2012;28:491-501.
32. Mohtadi N, Barber R, Chan D, Paolucci EO. Complications and adverse events of a randomized clinical trial comparing 3 graft types for ACL reconstruction. *Clin J Sport Med.* 2016;26:182-189.
33. Montgomery J, Knudson D. A method to determine stride length for baseball pitching. *Appl Res Coach Athl Annu.* 2002;17:75-84.
34. O'Brien DF, O'Hagan T, Stewart R, et al. Outcomes for ulnar collateral ligament reconstruction: a retrospective review using the KJOC assessment score with two-year follow-up in an overhead throwing population. *J Shoulder Elbow Surg.* 2015;24:934-940.
35. Rettig AC, Sherrill C, Snead DS, Mendler JC, Mieling P. Nonoperative treatment of ulnar collateral ligament injuries in throwing athletes. *Am J Sports Med.* 2001;29:15-17.
36. Rohrbough JT, Altchek DW, Hyman J, Williams RJ 3rd, Botts JD. Medial collateral ligament reconstruction of the elbow using the docking technique. *Am J Sports Med.* 2002;30:541-548.
37. Ruffilli A, De Fine M, Traina F, Pilla F, Fenga D, Faldini C. Saphenous nerve injury during hamstring tendons harvest: does the incision matter? A systematic review [published online June 23, 2016]. *Knee Surg Sports Traumatol Arthrosc.* doi:10.1007/S00167-016-4217-8.
38. Savoie FH 3rd, Morgan C, Yaste J, Hurt J, Field L. Medial ulnar collateral ligament reconstruction using hamstring allograft in overhead throwing athletes. *J Bone Joint Surg Am.* 2013;95:1062-1066.
39. Segawa H, Omori G, Koga Y, Kameo T, Iida S, Tanaka M. Rotational muscle strength of the limb after anterior cruciate ligament reconstruction using semitendinosus and gracilis tendon. *Arthroscopy.* 2002;18:177-182.
40. Shelton WR, Fagan BC. Autografts commonly used in anterior cruciate ligament reconstruction. *J Am Acad Orthop Surg.* 2011;19:259-264.
41. Snow BJ, Wilcox JJ, Burks RT, Greis PE. Evaluation of muscle size and fatty infiltration with MRI nine to eleven years following hamstring harvest for ACL reconstruction. *J Bone Joint Surg Am.* 2012;94:1274-1282.
42. Suijkerbuijk MA, Reijman M, Lodewijks SJ, Punt J, Meuffels DE. Hamstring tendon regeneration after harvesting: a systematic review. *Am J Sports Med.* 2015;43:2591-2598.
43. Tashiro T, Kurosawa H, Kawakami A, Hikita A, Fukui N. Influence of medial hamstring tendon harvest on knee flexor strength after anterior cruciate ligament reconstruction. A detailed evaluation with comparison of single- and double-tendon harvest. *Am J Sports Med.* 2003;31:522-529.
44. Vitale MA, Ahmad CS. The outcome of elbow ulnar collateral ligament reconstruction in overhead athletes: a systematic review. *Am J Sports Med.* 2008;36:1193-1205.
45. Yamanouchi T. EMG analysis of the lower extremities during pitching in high-school baseball. *Kurume Med J.* 1998;45:21-25.