



A genomic study on distribution of *human leukocyte antigen (HLA)-A* and *HLA-B* alleles in Lak population of Iran[☆]



Farhad Shahsavari^a, Ali-Mohammad Varzi^{a,*}, Seyyed Amir Yasin Ahmadi^{b,c}

^a Department of immunology, Lorestan University of Medical Sciences, Khorramabad, Iran

^b Research Office for the History of Persian Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

^c Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran

ARTICLE INFO

Article history:

Received 25 September 2016

Received in revised form 3 November 2016

Accepted 9 November 2016

Available online 10 November 2016

Keywords:

HLA class I

Lak population

Anthropology

Immunology

Iran

ABSTRACT

Anthropological studies based on the highly polymorphic gene, *human leukocyte antigen (HLA)*, provide useful information for bone marrow donor registry, forensic medicine, disease association studies, as well as infertility treatment, designing peptide vaccines against tumors, and infectious or autoimmune diseases. The aim of this study was to determine *HLA-A* and *HLA-B* allele frequencies in 100 unrelated Lak/læk/individuals from Lorestan province of Iran. Finally, we compared the results with that previously described in Iranian population. Commercial HLA-Type kits from BAG (Lich, Germany) company were used for determination of the *HLA-A* and *HLA-B* allele frequencies in genomic DNA, based on polymerase chain reaction with sequence specific primer (PCR-SSP) assay. The differences between the populations in distribution of *HLA-A* and *HLA-B* alleles were estimated by chi-squared test with Yate's correction. The most frequent *HLA-A* alleles were *24 (20%), *02 (18%), *03 (12%) and *11 (10%), and the most frequent *HLA-B* alleles were *35 (24%), *51 (16%), *18 (6%) and *38 (6%) in Lak population. *HLA-A**66 (1%), *74(1%) and *HLA-B**48 (1%), *55(1%) were the least observed frequencies in Lak population. Our results based on *HLA-A* and *HLA-B* allele frequencies showed that Lak population possesses the previously reported general features of Iranians but still with unique.

© 2016 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Among the basic medical sciences involving in anthropology, immunology can be considered as a science which deals with the molecular markers playing role in recognition of self and non-self between ethnicities [1]. So biomedicine is not alien to anthropology and offers an ethnographic international classification of ethnicities based on the immunological molecules such as CD markers [2,3]. Hereby the integration between anthropology and immunology as a kind of integration between social and medical sciences could be a strategic way to have a biological information bank of different cultures in order to reach the aims mentioned in the next such as bone marrow transplantation, infertility prognosis and treatment [4,5] and finding the identity of persons not grown with their real parents.

[☆] Authors' contribution: F Shahsavari (data analysis and interpretation and the final revision and approve), AM Varzi (executor of the research proposal and laboratory tests), SAY Ahmadi (interpretation and writing the article). Funding: Present study was performed under the financial support of Lorestan University of Medical Sciences with grant number 1139.

* Corresponding author at: Department of immunology, 3rd floor, Kamalvand branch, Lorestan University of Medical Sciences, Khorramabad, Iran.

E-mail addresses: shahsavari@yaho.com (F. Shahsavari), ali.m.varzi@ums.ac.ir (A.-M. Varzi), yasin_ahmadi73@yahoo.com (S.A.Y. Ahmadi).

In the mankind genome, the *human leukocyte antigen (HLA)* also called as major histocompatibility complex (MHC) [6] with the length of 3600 kb is located on chromosome 6 and includes 239 antigenic loci that about 40% of them are immunogenic [7]. Recently, >2000 alleles for *HLA* class I are known in humans [8]. There are 2 classes of *HLA* and *HLA* class I in turn falls in two categories of classical (*HLA-A* and *B*) and non-classical (*HLA-C*, *G*, *E* and *F*) [4,9–13]. The main role of *HLA* class I is that this biological molecule acts as an identifying card for all nuclear cells of body to be proposed for natural killer cells (CD56CD16 [14]) that different interactions between them results in different outcomes [15]. In addition to the key roles considered by immunologists for *HLA*, such genes have attracted the view point of most developmental biologists because of a high level of allele variety.

Since some alleles of *HLA* are commonplace in specific populations, the alleles are used by anthropologists as markers to determine genetic correlations and interactions in different populations [2]. Clinically, being acquainted with *HLA* distribution is a sine qua non for bone marrow donating centers [16,17], forensic medicine [18], studies of *HLA* related disease such as type 2 diabetes [19–21] or multiple sclerosis [22], designing peptide vaccine and monoclonal antibodies against tumors [23,24], infectious agents [25–27] and autoimmune disease [20,28], as well as infertility treatment and assisted reproductive technologies [29–31].

Race wise, Iran is a variable country that has different ethnicities such as Kurd, Lur, Torkaman, Azeri, Arab and Balouch. Majority of Iranians are Muslims, but Zoroastrians, Christians and Jewish also live in this country [32]. History has it that the majority of Iranians are Aryan but during the history they were attacked by different foreigners such as Alexander and Macedonians, Arabs, Turks and Moguls [32]. As described by a thesis under the supervision of Dr. Maziar Ashrafiyan Bonab [33], Aryans are believed to be one of the early Proto-Indo-European speaking ethnicities migrated toward Iran. They arrived and settled in the north of Afghanistan around 2000–1500 BCE and kept on migrating and headed west settling in Iran and others south into India and Pakistan. It seems that as a result of their vast distribution, the Aryans were the main advocates of the Indo-European languages and promoting their proliferation via cultural and demic diffusion and therefore displacing the indigenous languages. Also Iran has played a key role in connection of different populations via the Silk Road [34, 35]. Thus the living populations in this country might be mixed because of migrations and mentioned relations.

Lak (or Laki) population is an ethnicity living in southwest of Iran and southeast of Iraq that there is still a controversy whether they are Kurd or Lur; because their culture and language are a complex of Kurdish and Luri populations. Kurds, Lurs and Persians are accounted as Indo-Iranian group of Aryans. The geographical condition of Laks (Fig. 1) shows that the majority of Iranian Laks live in the north of Lorestan province and some of them lives in Kermanshah, Ilam and Hamadan [36].

In previous studies, allele frequency of *HLA* class-II were determined in majority of different Iranian populations and compared with each other [18,37]. In the present study, we intend to find allele frequency of *HLA* class-I in Lak/læk/population of Lorestan province (west of Iran) regarding to the lack of this study (on the class-I) in Lak population. Then based on the *HLA-I* profile the genetic relations between this population and total Iranian population [38] is investigated.

2. Material and methods

For the present study, 100 healthy and unrelated Lak individuals living in Lorestan province were randomly chosen by convenient sampling based on the including criteria in 2015. Our including criteria was having the same race (the two recent generations of each sample should be Laks) and having 20–40 years of age, and the excluding criteria was having history of some specific diseases. Complete blood samples were obtained with informed and written consents from the participating individuals. The study was approved by the ethic committee of Lorestan University of Medical Sciences.



Fig. 1. Geographical status of Laks (the red color) (adapted from the reference [36]).

The genomic DNA of individuals was extracted by using kit BAG (Germany). *HLA*-typing kit for polymerase chain reaction with sequence specific primer (PCR-SSP) – a method which has longer and more specific primers for each allele of polymorphisms instead of using restriction enzymes [39,40] – with low resolution (BAG Germany) were used to determine *HLA-A* and *B* alleles via genomic DNA. PCR products got visible in 2% agarose gel including 0.5 mg/ml Ethidium Bromide electrophoresis under an ultra violet light. Since there is a specific primer for each allele in SSP method (and *HLA* is highly polymorphic), we cannot write their sequences in the article and also it's a patent for the company.

Allele frequencies of *HLA-A* and *B* were determined through the direct counting method. The differences between the populations in allele frequencies of *HLA-A* and *B* were estimated by Chi-squared 2 multiplied by 2 test with degree of freedom 1. According to being multiple of the compares, we used Yate's correction to correct the randomized significance. After the correction $p < 0.05$ considered as the level of significance.

3. Results

Allele frequency of *HLA-A* and *B* in Iranian Lak population are given in Table 1. Fifteen allele for *HLA-A* and 23 allele for *HLA-B* were identified. In Lak population, the most frequent *HLA-A* alleles were respectively A*24 (20%), A*02 (18%), A*3 (12%) and A*11 (10%). The most frequent *HLA-B* alleles were respectively B*35 (24%), B*51 (16%), B*18 (6%) and B*38 (6%). The least frequent *HLA-A* alleles were A*66 (1%) and A*74 (1%) and the least frequent *HLA-B* alleles were B*48 (1%) and B*55 (1%).

4. Discussion

In the present study the most frequent *HLA-A* and *HLA-B* alleles in 100 healthy and unrelated Lak individuals were determined with method PCR-SSP. Allele frequency of *HLA-A* and *B* in Lak and total Iranian population [38] is shown in Table 1.

Table 1

Allele frequency of *HLA-A* and *B* in Lak population (our study) in comparison to total Iranian population [38] ($p < 0.05$).

Allele A	Frequency in Lak population	Frequency in total Iranian population	Allele B	Frequency in Lak population	Frequency in total Iranian population
A*01	8	9.25	B*07	4	4.75
A*02	18	18.16	B*08	4	4.25
A*03	12	12.08	B*13	4	3.91
A*11	10	10.41	B*14	2	3.00
A*23	2	2.25	B*15	2	3.16
A*24	20*	16.41	B*18	6	4.33
A*26	8	6.83	B*27	2	2.58
A*29	2	2.5	B*35	24*	21.66
A*30	6	4.66	B*37	2	1.00
A*31	6	6.16	B*38	6	4.33
A*32	2	5.66	B*39	2	0.91
A*33	2	3.66	B*40	4	3.58
A*43	–	0.16	B*41	3	2.75
A*66	1	0.25	B*42	–	0.165
A*68	2	4.16	B*44	2	4.165
A*69	–	0.16	B*45	–	0.165
A*74	1	0.41	B*47	–	0.165
A*80	–	0.16	B*48	1	0.50
			B*49	2	2.50
			B*50	3	3.58
			B*51	16*	13.33
			B*52	3	3.50
			B*53	–	0.25
			B*54	–	0.165
			B*55	1	0.35
			B*56	2	0.50
			B*57	2	1.00
			B*58	3	1.915

Table 2

High frequent alleles in different populations. X = not mentioned in reference [7] or the frequency was <10% in spite of their high frequency.

Ethnicity	High frequent alleles (>10%)	
Laks	A*02,03,11,24	B*35,51
Iranians	A*02,03,11,24	B*35,51
Caucasians (the whites) of the USA	A*01,02,03	B*04,44
The Lebanese	A*01,02,24	B*35
Jordanians	A*02	X
The Irish	A*01,02	B*04,44
The Japanese	A*02,24	B*52
Moroccans	A*01,02	X
Koreans	A*02,11,24	B*15
Africans	A*23	X [B*35,53 [46]]
Bulgarians	A*02,24	B*18,51
The Chinese	A*11,24	B*46
Brazilians	A*02,03,24	X [B*15,35,51 [44]]
Omanis	A*02,11,26,32,68	B*51
The Polish	A*01,02,03,24	X
The Spanish	A*02	X
Armenians [47]	A*01,02,03,24	B*35,51

As it is shown in Table 1, the frequency of the alleles *HLA-A*24*, *HLA-B*35* and *HLA-B*51* in Lak population were significantly more than the frequency of these alleles in Iranian population.

In addition, comparing our findings with the findings of total Iranian population study shows that allele frequency of *HLA-A* and *B* in Lak population has a high similarity to total Iranian population. Recently, *HLA* class I typing has been performed also in the Lur and Kurd populations of western Iran [41]. Based on this study, *A*24* and *B*51* was high frequent in both Lur and Kurd populations of Iran. Because their PCR kit was high resolution, we could not statistically compare this study with the study of ours.

Other than different Iranian populations and ethnicities, these findings are also similar to other studies on different white populations (Table 2). In contrast, these findings are not similar to the findings of Shen et al. in Han population of China [42] and Yao et al. in total Chinese population [43] (except *A*24*, about 12% in the Chinese and 20% in Laks that in the both populations was higher than 10% as the par of high frequency). Of course this fact is reasonable through genetic similarity of Iranians and others Caucasians and genetic difference between the whites and yellow-skinned.

In a Brazilian population the most frequent alleles were *A*02*, *A*03*, *B*15*, *B*35*, *B*51* [44]. Alleles *B*35* and *B*51* high frequency was like Lak population, whereas *B*15* is low frequent in Lak population. The allele frequency of *B*27* – which is associated with Ankylosing spondylitis [45] – is low frequent in Lak population as other populations so. In Africans, alleles *B*35* and *B*53* were high frequent [46] as the Laks respectively are and are not so. These findings suggest that some alleles like *A*24* are widespread in human race. As it's shown in Table 2 the high frequent alleles (>10%) in different ethnicities [7] are compared with Laks. Allele *A*02* seems the most high frequent in different populations (Table 2).

Analyzing frequency of *HLA-A* and *B* alleles enable us to assay genetic relations among various populations in our anthropological studies. As well, determining allele frequency of *HLA-A* and *B* in Lak population could be a good source for other studies in future such as correlations between the alleles and genetic disorders.

5. Conclusions

At the end of the study, our results based on the allele frequencies showed that the population has general features of the Lurs, the Kurds and the total Iranian population reported before. Of course it's obvious that the main conclusion is that establishing immunology-based laboratories is necessary to diagnose and treat diseases through molecular methods. It could be useful for transplantation, assisted reproductive technologies and *HLA*-related disease.

Conflict of interest

Meanwhile we don't have conflict of interest.

Acknowledgements

Finally we take it upon ourselves to appreciate all the participants in the study. Present study was performed under the financial support of Lorestan University of Medical Sciences with grant number 1139.

References

- [1] A.D. Napier, Nonself help: how immunology might reframe the enlightenment. *Cult. Anthropol.* 27 (2012) 122–137.
- [2] A. Cambrosio, P. Keating, A matter of FACS: constituting novel entities in immunology. *Med. Anthropol. Q.* 6 (1992) 362–384.
- [3] E. Martin, Toward an anthropology of immunology: the body as nation state. *Med. Anthropol. Q.* 4 (1990) 410–426.
- [4] M. PrabhuDas, E. Bonney, K. Caron, S. Dey, A. Erlebacher, A. Fazleabas, S. Fisher, T. Golos, M. Matzuk, J.M. McCune, Immune mechanisms at the maternal-fetal interface: perspectives and challenges. *Nat. Immunol.* 16 (2015) 328–334.
- [5] S.A.Y. Ahmadi, F. Shahsavari, S. Akbari, A review on controversies about the role of immune and inflammatory systems in implantation process and durability of pregnancy. *Int. J. Womens Health Reprod. Sc.* 4 (2016) 96–102.
- [6] D.X. Beringer, F.S. Kleijwegt, F. Wiede, A.R. van der Slik, K.L. Loh, J. Petersen, N.L. Dudek, G. Duinkerken, S. Laban, A. Joosten, J.P. Vivian, Z. Chen, A.P. Uldrich, D.I. Godfrey, J. McCluskey, D.A. Price, K.J. Radford, A.W. Purcell, T. Nikolic, H.H. Reid, T. Tiganis, B.O. Roep, J. Rossjohn, T cell receptor reversed polarity recognition of a self-antigen major histocompatibility complex. *Nat. Immunol.* 16 (2015) 1153–1161.
- [7] S. Khansa, R. Hoteit, D. Shammaa, R.A. Khalek, H. El Halas, L. Greige, F. Abbas, R.A.R. Mahfouz, *HLA* class I allele frequencies in the Lebanese population. *Gene* 512 (2013) 560–565.
- [8] S. Mukherjee, J. Warwicker, N. Chandra, Deciphering complex patterns of class-I *HLA*-peptide cross-reactivity via hierarchical grouping. *Immunol. Cell Biol.* 93 (2015) 522–532.
- [9] E.C. Castelli, T.H.A. Lima, R.V. Buttura, M.A. Paz, I.O.P. Porto, J. Ramalho, A.S. Souza, L.C. Veiga-Castelli, C.T. Mendes-Junior, E.A. Donadi, Evaluation of the *HLA-F* variability in Brazil by next generation sequencing. *Hum. Immunol.* 76 (2015) 89 (Supplement).
- [10] E.C. Castelli, C.T. Mendes-Junior, A. Sabbagh, I.O.P. Porto, A. Garcia, J. Ramalho, T.H.A. Lima, J.D. Massaro, F.C. Dias, C.V.A. Collares, V. Jamonneau, B. Bucheton, M. Camara, E.A. Donadi, *HLA-E* coding and 3' untranslated region variability determined by next-generation sequencing in two West-African population samples. *Hum. Immunol.* 76 (2015) 945–953.
- [11] N. Lauterbach, L. Wieten, H.E. Popeijn, C.E.M. Voort, M.G.J. Tilanus, *HLA-E* regulates NKG2C+ natural killer cell function through presentation of a restricted peptide repertoire. *Hum. Immunol.* 76 (2015) 578–586.
- [12] T.D. Veit, J.A.B. Chies, M. Switala, B. Wagner, P.A. Horn, M. Busatto, C.V. Brenol, J.C.T. Brenol, R.M. Xavier, V. Rebmann, The paradox of high availability and low recognition of soluble *HLA-G* by *LILRB1* receptor in rheumatoid arthritis patients. *PLoS One* 10 (2015), e0123838.
- [13] A. Halenius, C. Gerke, H. Hengel, Classical and non-classical MHC I molecule manipulation by human cytomegalovirus: so many targets—but how many arrows in the quiver? *Cell. Mol. Immunol.* 12 (2015) 139–153.
- [14] S.A.Y. Ahmadi, S. Akbari, F. Shahsavari, But the reverse is not true: a critical review on staining protocol of uterine natural killer cells. *J. Chem. Pharm. Sci.* 9 (2016) 2483–2486 [http://jchps.com/issues/jchps%209\(4\)%2014%20Seyyed%20Amir%208%202483-2486.pdf](http://jchps.com/issues/jchps%209(4)%2014%20Seyyed%20Amir%208%202483-2486.pdf).
- [15] T. Mousavi, H. Poormoghimi, M. Moradi, N. Tajik, F. Shahsavari, M. Soofi, Phenotypic study of natural killer cell subsets in ankylosing spondylitis patients. *Iran. J. Allergy Asthma Immunol.* 8 (2009) 193–198.
- [16] Z. Grubic, M.B. Kamenaric, M. Mikulic, K.S. Jankovic, M. Maskalan, R. Zunec, *HLA-A*, *HLA-B* and *HLA-DRB1* allele and haplotype diversity among volunteer bone marrow donors from Croatia. *Int. J. Immunogenet.* 41 (2014) 211–221.
- [17] P.A. Gourraud, M.L. Balere, C. Faucher, P. Loiseau, A. Dormoy, E. Marry, F. Garnier, *HLA* phenotypes of candidates for HSCT: comparing transplanted versus non-transplanted candidates, resulting in the predictive estimation of the probability to find a 10/10 *HLA* matched donor. *Tissue Antigens* 83 (2014) 17–26.
- [18] A.M. Varzi, F. Shahsavari, M.J. Tarrahi, Distribution of *HLA-DRB1* and *HLA-DQB1* alleles in Lak population of Iran. *Hum. Immunol.* 77 (7) (2016) 580–583.
- [19] D.C. Chang, P. Piaggi, R.L. Hanson, W.C. Knowler, J. Bucci, G. Thio, M.G. Hohenadel, C. Bogardus, J. Krakoff, Use of a high-density protein microarray to identify autoantibodies in subjects with type 2 diabetes mellitus and an *HLA* background associated with reduced insulin secretion. *PLoS One* 10 (2015), e0143551.
- [20] N. Tajik, F. Shahsavari, H. Poormoghimi, M. Radjabzadeh, T. Mousavi, A. Jalali, *KIR3DL1* + *HLA-B* Bw4Ile80 and *KIR2DS1* + *HLA-C2* combinations are both associated with ankylosing spondylitis in the Iranian population. *Int. J. Immunogenet.* 38 (2011) 403–409.
- [21] M. Piga, A. Mathieu, Genetic susceptibility to Behcet's disease: role of genes belonging to the MHC region. *Rheumatology* 50 (2011) 299–310.
- [22] F. Shahsavari, S. Mapar, S.A.Y. Ahmadi, Multiple sclerosis is accompanied by lack of *KIR2DS1* gene: a meta-analysis. *Genomics Data* 10 (2016) 75–78.

- [23] S. Matsuoka, Y. Ishii, A. Nakao, M. Abe, N. Ohtsui, S. Momose, H. Jin, H. Arase, K. Sugimoto, Y. Nakauchi, Establishment of a therapeutic anti-pan HLA-class II monoclonal antibody that directly induces lymphoma cell death via large pore formation. *PLoS One* 11 (2016), e0150496.
- [24] W.D. M. Larche, Peptide-based therapeutic vaccines for allergic and autoimmune diseases. *Nat. Med.* 11 (2005) 69–76.
- [25] P.J. McLaren, M. Carrington, The impact of host genetic variation on infection with HIV-1. *Nat. Immunol.* 16 (2015) 577–583.
- [26] T. Mousavi, F. Shahsavari, P. Farnia, N. Tajik, M. Soofi, Study of KIR expression and HLA ligands in CD56+ lymphocytes of drug resistant tuberculosis patients. *Iran. J. Allergy Asthma Immunol.* 10 (2011) 189–194.
- [27] F. Shahsavari, T. Mousavi, A. Azargon, K. Entezami, Association of KIR3DS1 + HLA-B Bw4Ile80 combination with susceptibility to tuberculosis in Lur population of Iran. *Iran. J. Immunol.* 9 (2012) 39–47.
- [28] N. Hussain, G. Jaffery, Distribution of human leukocyte antigen alleles in systemic lupus erythematosus patients with angiotensin converting enzyme insertion/deletion polymorphism. *Bosn. J. Basic Med. Sci.* 13 (2013) 57–62.
- [29] N. Alizadeh, E. Mosaferi, L. Farzadi, J. Majidi, A. Monfaredan, B. Yousefi, B. Baradaran, Frequency of null allele of human leukocyte antigen-G (HLA-G) locus in subjects to recurrent miscarriage. *Int. J. Reprod. BioMed.* 14 (2016) 459.
- [30] M. Fotoohi, N. Ghasemi, S.A. Mirghanizadeh, M. Vakili, M. Samadi, Association between HLA-E gene polymorphism and unexplained recurrent spontaneous abortion (RSA) in Iranian women. *Int. J. Reprod. BioMed.* 14 (2016) 477.
- [31] T. Meuleman, L.E.L.O. Lashley, O.M. Dekkers, J.M.M. van Lith, F.H.J. Claas, K.W.M. Bloemenkamp, HLA associations and HLA sharing in recurrent miscarriage: a systematic review and meta-analysis. *Hum. Immunol.* 76 (2015) 362–373.
- [32] M. Mohebbi, Z. Mohebbi, Demography of race and ethnicity in Iran. *The International Handbook of the Demography of Race and Ethnicity*, Springer 2015, pp. 353–366.
- [33] J.W. Whale, Mitochondrial DNA Analysis of Four Ethnic Groups of Afghanistan. University of Portsmouth, 2012.
- [34] P. Pollegioni, K.E. Woeste, F. Chiocchini, S. Del Lungo, I. Olimpieri, V. Tortolano, J. Clark, G.E. Hemery, S. Mapelli, M.E. Malvolti, Ancient humans influenced the current spatial genetic structure of common walnut populations in Asia. *PLoS One* 10 (2015), e0135980.
- [35] B.I. Boulois, *The Silk Road*. Allen & Unwin, London, 1966.
- [36] E. Anonby, Kurdish or Luri, Laki's Disputed Identity in the Luristan Province of Iran. *Kurdische Studien.* 4, 2004 7–22.
- [37] O. M., S. Farjadian, H. Inoko, A. Ghaderi, The genetic relationship among Iranian ethnic groups: an anthropological view based on HLA class II gene polymorphism. *Mol. Biol. Rep.* 36 (2009) 1943–1950.
- [38] A. Ghashghaie, K. Alimoghaddam, M.R. Ostadali, L. Khansari, M. Sadraee, E. Mirrasekhian, I. Mohyedin, F. Raofi, Z. Noori, H. Yaghmaian, Allele frequencies of HLA class-I loci in the normal Iranian population. *Int. J. Hematol. Oncol. Stem Cell Res.* 3 (2009) 18–20.
- [39] W. De Spiegelaere, J. Philippé, K. Vervisch, C. Verhofstede, E. Malatinkova, M. Kiselinova, W. Trypsteen, P. Bonczkowski, D. Vogelaers, S. Callens, Comparison of methods for in-house screening of HLA-B* 57: 01 to prevent abacavir hypersensitivity in HIV-1 care. *PLoS One* 10 (2015), e0123525.
- [40] N. Tajik, F. Shahsavari, M. Nasiri, M. Radjabzadeh, Compound KIR-HLA genotype analyses in the Iranian population by a novel PCR-SSP assay. *Int. J. Immunogenet.* 37 (2010) 159–168.
- [41] E. Ashouri, P. Norman, L. Guethlein, A. Han, N. Nemat-Gorgani, S. Norberg, A. Ghaderi, P. Parham, HLA class I variation in Iranian Lur and Kurd populations: high haplotype and allotype diversity with an abundance of KIR ligands. *HLA* 88 (2016) 87–99.
- [42] Y.S. Shen, D.F. Cao, Y.L. Li, J.K. Kulski, L. Shi, H.J. Jiang, Q.L. Ma, J.K. Yu, J.X. Zhou, Y.F. Yao, L. Shi, Distribution of HLA-A, -B, and -C alleles and HLA/KIR combinations in Han Population in China. *J. Immunol. Res.* (2014) (<https://www.hindawi.com/journals/jir/2014/565296/>).
- [43] Y. Yao, L. Shi, Y. Tao, J.K. Kulski, K. Lin, X. Huang, H. Xiang, J. Chu, L. Shi, Distinct HLA allele and haplotype distributions in four ethnic groups of China. *Tissue Antigens* 80 (2012) 452–461.
- [44] C. Rodrigues, L.C. Macedo, A.V. Bruder, F.D.C. Quintero, J.B. de Alencar, A.M. Sell, J.E.L. Visentainer, Allele and haplotype frequencies of HLA-A, B, C, DRB1 and DQB1 genes in polytransfused patients in ethnically diverse populations from Brazil. *Int. J. Immunogenet.* 42 (2015) 322–328.
- [45] I. El Mouraghi, A. Ouarour, I. Ghozlan, E. Collantes-Estevez, R. Solana, A. El Maghraoui, Polymorphisms of HLA-A, -B, -Cw and DRB1 antigens in Moroccan patients with ankylosing spondylitis and a comparison of clinical features with frequencies of HLA-B*27. *Tissue Antigens* 85 (2015) 108–116.
- [46] P.J. Norman, J.A. Hollenbach, N. Nemat-Gorgani, L.A. Guethlein, H.G. Hilton, M.J. Pando, K.A. Koram, E.M. Riley, L. Abi-Rached, P. Parham, Co-evolution of human leukocyte antigen (HLA) class I ligands with killer-cell immunoglobulin-like receptors (KIR) in a genetically diverse population of Sub-Saharan Africans. *PLoS Genet.* 9 (2013).
- [47] L. Matevosyan, S. Chattopadhyay, V. Madelian, S. Avagyan, M. Nazaretyan, A. Hyussian, E. Vardapetyan, R. Arutunyan, F. Jordan, HLA-A, HLA-B, and HLA-DRB1 allele distribution in a large Armenian population sample. *Tissue Antigens* 78 (2011) 21–30.