

# Caring for Informal Dementia Caregivers and Their Loved Ones Via the HOMeCARE Exercise and Mindfulness for Health Program (HOMeCARE): A Randomized, Single-Blind, Controlled Trial

Tommy Lang, BAppSc(Hons)<sup>1</sup> , Kenneth Daniel, MND<sup>1</sup>, Michael Inskip, PhD<sup>1,2</sup>, Yorgi Mavros, PhD<sup>1</sup>, and Maria A. Fiatarone Singh, AM, MD<sup>1,3,4</sup>

Gerontology & Geriatric Medicine  
Volume 9: 1–12  
© The Author(s) 2023  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/23337214231203472  
journals.sagepub.com/home/ggm



## Abstract

**Objective:** To investigate the effects of a dyadic intervention of mindfulness-based stress reduction (MBSR) for informal dementia caregivers and home-based balance and progressive resistance training (PRT) for their loved ones. **Methods:** The study was a two arm, randomized, controlled, single-blinded, parallel-group trial. Dyads were randomized to an intervention group: an 8-week MBSR course (daily) and an 8-week PRT and balance training (3 days/week) for their loved ones or a waiting list control group. **Results:** Nine dyads were randomized [caregivers: median age 75 (40–81) years, loved ones: 77 (73–88) years]. The intervention significantly improved caregiver mindfulness [relative effect size (95% confidence interval) 1.35 (−0.10, 2.81);  $p = .009$ ] and functional mobility in their loved ones [mean difference (95% confidence interval) 1.53 (−3.09, 6.14)] with no significant effects on caregiver burden [relative effect size (95% confidence interval) 0.22 (−1.09, 1.54);  $p = .622$ ]. **Conclusion:** The study appeared feasible in the home environment and future large and longer trials should test the efficacy of a more abbreviated MBSR intervention and to optimize adoption and sustain adherence over time.

**Trial registry name:** HOMeCare: Caring for the Dementia Caregiver and their Loved One via the HOMeCare Exercise and Mindfulness for Health Program

**Trial URL:** <https://www.australianclinicaltrials.gov.au/anzctr/trial/ACTRN12617000347369>

**Registration number:** ACTRN12617000347369

## Keywords

dementia, caregiver, mindfulness, exercise, resistance training

**Manuscript received:** June 23, 2023; **final revision received:** September 5, 2023; **accepted:** September 8, 2023.

## Introduction

People with dementia are at risk of transition to residential aged care due to functional decline and caregiver stress (Paradise et al., 2015; Richardson et al., 2013). Frailty contributes to this functional decline (Pedone et al., 2005) and risk of transition to residential aged care, hospitalization (Milte & Crotty, 2014), and premature mortality (Lehmann et al., 2018). Robust anabolic exercise [(progressive resistance training (PRT))] is considered the most potent strategy to treat frailty (Dent et al., 2017), combined with balance training for falls risk (Dent et al., 2017; Singh et al., 2012). The benefits of PRT include cognitive function, as well as many other conditions relevant to dementia: depression, sarcopenia,

osteoporosis, cardiovascular disease, hypertension, diabetes, falls, and other co-morbidities (Escriche-Escuder et al., 2021; Hordern et al., 2011; Sharman et al., 2019;

<sup>1</sup>Sydney School of Health Sciences, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia

<sup>2</sup>College of Healthcare Sciences, Sport and Exercise Science, James Cook University, Townsville, QLD, Australia

<sup>3</sup>Sydney Medical School, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia

<sup>4</sup>The Hinda and Arthur Marcus Institute for Aging Research, Hebrew SeniorLife, Boston, MA, USA

### Corresponding Author:

Maria A. Fiatarone Singh AM, The University of Sydney, Sydney, NSW 2006, Australia.

Email: [maria.fiataronesingh@sydney.edu.au](mailto:maria.fiataronesingh@sydney.edu.au)



Sherrington et al., 2019; Singh et al., 1997; Song et al., 2018; Vanhees et al., 2012).

Recent Australian government policy is targeted toward supporting informal caregivers and their loved ones with dementia to remain living at home for as long as possible, with the overall aims of improving health outcomes and reducing healthcare costs (Australian Health Ministers Advisory Council, 2015). This is not only a government policy but also best practice as supporting “aging in place” for people with dementia is the preferred option for most older adults (Australian Institute of Health Welfare, 2013). Aging in place aims to support older adults to live in their homes in order to delay or prevent transition to residential aged care for as long as possible.

Aging in place often requires informal caregivers who provide unpaid care to their loved ones, in addition to formal services in the home. However, informal dementia caregivers are susceptible to poor health outcomes, isolation, depression, and anxiety due to the challenging behaviors and the functional decline of their loved ones (Mahoney et al., 2005). They are also reported to be significantly more stressed than caregivers for those who are cognitively-intact (Pinquart & Sörensen, 2003) and present with more severe psychological and physical symptoms (Cheng, 2017). Thus, stress-reduction interventions are of particular interest for these individuals. One of the most studied of such interventions for caregivers is mindfulness-based stress reduction (MBSR) developed by Jon Kabat-Zinn in 1979 (Kabat-Zinn, 2003), which has been used in family caregivers of people with dementia (Han, 2022; Kor et al., 2018; Li et al., 2016; Liu et al., 2018), cancer (Birnle et al., 2010), and developmental disabilities (Bazzano et al., 2015). The focus of MBSR is to increase mindfulness which is a skill of being present in the moment, particularly with sustained awareness of perceptible mental states and processes (Grossman et al., 2004). Mindfulness can be practiced through many meditation forms such as body awareness, mindful movements, yoga postures, and being mindful in daily situations. There are promising results reported with MBSR in reducing informal caregiver psychological symptoms such as symptoms of depression, anxiety, stress, and burden (Jaffray et al., 2016; Li et al., 2016; Liu et al., 2018), as well as associations with higher levels of life satisfaction and mindfulness wellbeing with mindfulness practice (Bhattacharyya et al., 2023).

Although exercise has been used in dementia dyads (prescribed either together or separately) with some benefits for both members of the dyad (Doyle et al., 2021), to our knowledge, there are no studies assessing the combined effects of mindfulness training for dementia caregivers in addition to robust exercise for their loved ones. Exercise for older adults with dementia is adaptable to the home environment and is efficacious in delaying functional decline (Doyle et al., 2021; Heyn et al., 2004; Nelson et al., 2004), however this may be limited by the capacity of the caregiver to deliver these interventions effectively without experiencing an increase in burden

and stress. Therefore, a novel and effective program supporting the physical and mental health of older adults with dementia and their informal caregivers could reduce the personal and societal burden of dementia significantly.

We designed “HOMeCARE: Caring for Informal Dementia Caregivers and their Loved Ones via the HOMeCARE Exercise and Mindfulness for Health Program.” The HOMeCARE study implemented MBSR for informal dementia caregivers combined with resistance and balance exercise for people with dementia living at home. The study aimed to not only train the caregiver in the delivery of an effective exercise program to improve the functional capacity of those living with dementia, but to also provide a mindfulness intervention for caregivers to reduce their stress and to increase the sustainability of the intervention. We hypothesized that a dyadic intervention of progressive, home-based resistance and balance training and mindfulness training would improve informal dementia caregivers’ mindfulness state, reduce caregiver burden, and improve their loved one’s functioning mobility.

## Methods

### Study Design

The HOMeCARE study was a two arm, randomized, controlled, single-blinded, parallel-group trial of MBSR for informal dementia caregivers and home-based PRT and balance training for their loved ones. The protocol was prospectively registered with the Australian Clinical Trials Registry (ACTRN12617000347369) and adhered to the Consolidated Standards of Reporting Trials (CONSORT) guidelines. Ethical approval was obtained from the University of Sydney Human Research Ethics Committee on 27/01/2017 and written informed consent was obtained from both members of the dementia dyad. Primary and secondary outcomes were collected at 0, 4, 8, and 12 weeks by a blinded assessor. No follow-up assessments were completed.

### Participants

The participants consisted of a dyad composed of one informal caregiver and one person with dementia. The participants were recruited from dementia carer support groups, volunteer databases from previous trials, advertisements in newsletters, Dementia Australia website, dementia café groups, and dementia care centers.

Informal caregivers were included if they could sufficiently access and navigate the electronic resources developed for the study, were a family member currently living with the participant with dementia, and/or providing some portion of their daily care including activities of daily living in an informal capacity without pay. Their loved ones with dementia were included if they were  $\geq 65$  years in 2016; living in the community with at least one informal caregiver; had a diagnosis of mild-moderate dementia of any type, except for Parkinson’s disease dementia, Mini-mental State Exam

(MMSE; Folstein et al., 1975) score 12 to 24/30; had at least mild deficits in functional mobility, defined as a Short Physical Performance Battery (SPPB; Guralnik et al., 1994) score <10/12. Parkinson's disease dementia was excluded as it may require a different approach to exercise than just addressing frailty and mobility or falls risks as intended with the intervention.

Informal caregivers were excluded if they had prior experience with or current use of MBSR. Their loved ones with dementia were excluded if they were currently participating in exercise at moderate-to-high intensity for three or more days a week. Behavioural and psychological symptoms of dementia were not exclusionary. All participants were excluded if they were unable to commit for the full study duration or had an unstable medical condition.

### Screening

Potential informal caregivers were interviewed on the telephone by the research assistant who assessed eligibility. Once reviewed and approved by the study physician, the participants were invited to the study site at the University of Sydney, Cumberland Campus, Australia. The study physician interviewed all informal caregivers to obtain a complete medical history and performed a physical examination of all participants with dementia at the study site to ensure eligibility. Baseline assessments were then conducted by the research team.

### Interventions

HOMeCARE was a remotely supervised intervention, supported by an initial home visit and weekly video calls. Dyad participants were randomized to either an intervention group or a waiting list group involving usual care. A research assistant demonstrated the intervention in a home visit to each dyad in the intervention group. An iPad was then supplied pre-loaded with mindfulness training material and instructional home-based exercise videos and images, as well as PRT equipment for the exercise component. Comprehensive details of the intervention and control groups are described in Supplemental Materials 1, 2, 3, and 5.

**MBSR.** The 8-week mindfulness training course for the informal caregivers utilized a modified version of the Palouse Mindfulness Based Stress Reduction (MBSR) course (Kabat-Zinn, 2013). The mindfulness practice was performed for the first 4 weeks before their loved ones started their exercise program. The MBSR practice continued until week 8 and self-selected formal and informal practices were continued until week 12. Formal practices were suggested for approximately for 30 minutes each day and informal practices were integrated in the informal caregivers' daily lives.

**PRT and Balance Training.** The person with dementia started their exercise program at week 4, via a home

visit with the research assistant, and subsequently facilitated by their informal caregiver. Informal caregivers were instructed to administer the exercise interventions 3 days/week for at least 45 minutes. Every week, a video call was planned to provide supervision and support to the dyad by the research assistant, while viewing an exercise session, if possible, to give feedback and reinforce appropriate progressions of weights and levels of balance, with additional phone calls as needed.

**Control Group.** The control group dyads were placed on a waiting list to receive the full intervention at the end of the 12-week trial. Both groups continued to receive usual medical care during the intervention.

### Outcomes

At baseline, all outcomes were measured by the same research assistant prior to randomization. All follow-up assessments were performed by a blinded assessor not otherwise involved in the intervention.

Details of the three primary outcomes are presented in Supplemental Material 2. Caregiver burden was assessed via the Zarit Burden Interview (ZBI; Zarit et al., 1980), caregiver mindfulness state via the State Mindfulness Scale (SMS; Tanay & Bernstein, 2013), and the loved one's functional mobility via the SPPB (Guralnik et al., 1994).

### Sample Size

The sample size was calculated based on previous studies using the same primary outcome measures and similar interventions. These calculations were based on (1) reduction in caregiver burden via ZBI (OR 0.18; Lowery et al., 2014) and (2) improvement in function in participants with dementia via the SPPB (ES 0.83; LIFE-P, 2006) with alpha .05 and beta .20. This was estimated at 48 dyads (24 control and 24 experimental).

### Randomization

The participants were randomized after their baseline assessments and were stratified by level of SPPB (<7 and 7–9) and MMSE (12–19 and 20–24). A research assistant not otherwise involved with the study, generated the stratified randomization sequence via a computerized random number generator (accessed at [www.randomization.com](http://www.randomization.com); created by Gerard E. Dallal, Ph.D.) and concealed the allocations in sequentially numbered envelopes for the research assistant to open with the participants after the completion of all baseline testing.

### Blinding

The HOMeCARE study was single blinded (assessor blinded). Both participants and assessors were blinded to the study allocation until completion of the baseline

assessments. It was impossible to blind the participants to the interventions due to awareness of receiving the exercise program and mindfulness training. Participants were asked not to inform the blinded assessor to which group they were allocated.

### Adherence

Adherence was captured on a questionnaire and reviewing the MBSR and exercise logs during the weekly status checks via video calls. Adherence was defined as completing the scheduled formal and informal practices for the MBSR course and the exercise sessions with the planned exercises and intensity.

### Adverse Events

Adverse events were defined *a priori*, including exacerbation of underlying diseases, onset of new musculoskeletal, cardiovascular, or metabolic abnormalities. Adverse events were captured via a questionnaire during the weekly video calls.

### Statistical Methods

Full statistical methods are detailed in Supplemental Material 2. The primary analytic strategy was intention-to-treat, with all randomized participants included, regardless of dropout or adherence level. The primary outcomes data at each time point were inspected for normality through descriptive statistics, histograms, and Shapiro-Wilk tests. Repeated measures linear mixed models with an unstructured repeated covariance type (for the SMS and SPPB) and AR(1; first-order autoregressive; for the ZBI) repeated covariance type were used to determine changes over time in both members of the dyad, in separate mixed models for each primary outcome. Baseline ZBI data was used as a covariate in its mixed model due to the imbalance in baseline scores between groups. Hedges' *g* relative effect sizes (ESs) were determined for each primary outcome using the pooled baseline SD as the denominator and the between group mean difference as the numerator to calculate ES. The analyses were performed on IBM SPSS Statistics version 27 and 28 (IBM Corp. Armonk, NY).

## Results

### Recruitment and Retention

Figure 1 displays the flow of participants through the study. Thirty-four dyads were assessed for eligibility and nine were randomized to the intervention group or the waiting list control group. The less than planned sample size was due to funding limitations and the advent of the COVID-19 restrictions prior to the completion of recruitment. One dyad in the control group did not complete their week 4 and 8 follow up assessments,

there was no week 8 primary outcome data for one informal caregiver in the intervention group and one dyad in the intervention group did not complete their week 12 in-clinic physical performance assessments due to COVID-19. There were no drop-outs in the study.

### Participants

Participant baseline characteristics are reported in Tables 1 to 3; informal caregiver baseline psychological assessments are provided in Supplemental Table 1. The loved ones were primarily female older adults who presented with four chronic conditions and up to eight prescribed medications. Informal caregivers were mainly older adults, who were not experienced in exercise facilitation and were a spouse/partner of the loved one except for one participant who was a daughter aged 40 years. Six informal caregivers presented with controlled chronic conditions.

### Adherence

**Adherence to MBSR.** Adherence to the formal practices was median (range) 36% (0%–55%) and adherence to the informal practices was median 64% (25%–76%).

**Adherence to PRT + Balance Training.** Adherence to the exercise sessions was median (range) 58% (28%–78%).


### Adverse Events

One adverse event was reported for a loved one in the intervention group who had an injurious fall while gardening in her backyard. She was briefly admitted to hospital overnight, had bruising on her face requiring stitches with no musculoskeletal injuries. This was adjudicated as unrelated to the study and her exercise intervention was re-commenced 2 weeks later. There were no adverse events attributed to the MBSR or exercise intervention itself.

### Primary Outcomes

The primary outcomes (ZBI, SMS, and SPPB) are reported in Table 4 and displayed in Supplemental Figures 1 to 3.

**Caregiver Burden: ZBI.** The intervention group reported a higher baseline ZBI score which was therefore used as a covariate in the mixed model analysis. There was no change in the ZBI related to group assignment (small ES = 0.22;  $p = .622$ ) and no significant change over time in the overall cohort ( $F = 0.164$ ,  $p = .92$ ). The mean difference was less than 3% between both groups. We did not find literature reporting the MCID for the ZBI, but in two studies of MBSR in dementia family caregivers that used the ZBI, the ZBI scores were reported to reduce by a mean difference of 5% to 11% (Brown et al., 2016; Epstein-Lubow et al., 2011).



**Figure 1.** CONSORT participant flow diagram.

Note. CONSORT = Consolidated Standards of Reporting Trials; PRT = progressive resistance training; MBSR = mindfulness-based stress reduction.

**Mindfulness State: SMS.** As hypothesized, there was a statistically significant improvement in mindfulness in the intervention group relative to the control group, a mean difference of 16.8 (30.7%), with a large effect size ( $ES=1.35$ ;  $p=.009$ ). We did not find literature reporting the MCID for the SMS. However, an uncontrolled trial (UCT; Bazzano et al., 2015) of MBSR for family caregivers of people with developmental disabilities reported a mindfulness increase of 15%, assessed by a different

mindfulness measure, which is smaller than the increase we saw.

**Physical Performance: SPPB.** As hypothesized, there was a clinically meaningful improvement in the intervention group, a mean difference of 1.53 points, with a moderate effect ( $ES=0.54$ ), and a trend for the group  $\times$  time effect ( $p=.091$ ). A SPPB change greater than 1.0 point indicates a clinically meaningful change (Perera et al., 2006).

**Table 1.** Loved One With Dementia: Baseline Sociodemographic and Health Status.

	Median (range) or <i>n</i> ( <i>n</i> = 9)
<i>Sociodemographic</i>	
Age (years)	77 (73–88)
Sex	
Females, <i>n</i>	6
Ethnicity	
Caucasian, <i>n</i>	8
Asian, <i>n</i>	1
Education	
Primary school, <i>n</i>	1
High school, <i>n</i>	4
Tertiary, <i>n</i>	3
Post-graduate, <i>n</i>	1
Residence type	
House, <i>n</i>	8
Unit/apartment, <i>n</i>	1
Marital status	
Married/de-facto, <i>n</i>	7
Widowed, <i>n</i>	2
Number of persons living in residence	
Alone, <i>n</i>	1
Two persons, <i>n</i>	5
Three or more persons, <i>n</i>	3
<i>Health status</i>	
Duration of dementia diagnosis, years	3 (1–7)
Type of dementia	
Alzheimer's disease, <i>n</i>	1
Vascular dementia, <i>n</i>	1
Frontotemporal dementia, <i>n</i>	1
Lewy Body disease, <i>n</i>	1
Undefined dementia <sup>a</sup> , <i>n</i>	5
Total number of chronic conditions, <i>n</i>	4.22 (1.71)*
Hypertension, <i>n</i>	7
Osteoporosis/Osteopenia, <i>n</i>	3
Osteoarthritis, <i>n</i>	3
Sarcopenia, <i>n</i>	6
Hypercholesterolemia, <i>n</i>	3
Anxiety, <i>n</i>	2
Total number of medications <sup>b</sup> , <i>n</i>	7.89 (2.47)*
Participants prescribed polypharmacy <sup>b</sup> , <i>n</i>	8
Participants with ≥ 1 anticholinergic medication <sup>c</sup> , <i>n</i>	5
Acetylcholinesterase inhibitor and/or memantine, <i>n</i>	6
Participants with nutrition supplements, <i>n</i>	5
Smoking status	
Never, <i>n</i>	6
Current, <i>n</i>	0
Former, <i>n</i>	3
Volume, pack-years <sup>d</sup>	1.25 (0.2–12) <sup>†</sup>

\*Presented as mean (standard deviation).

<sup>†</sup>Data from three participants.<sup>a</sup>Confirmed dementia diagnosis with unconfirmed dementia type.<sup>b</sup>Medications included prescribed, not prescribed, over-the-counter, nutritional, and herbal supplements.<sup>c</sup>Defined as ≥ 5 medications.<sup>d</sup>Estimated volume from packs per day multiplied by years of smoking.**Table 2.** Informal Caregiver Baseline Sociodemographic and Health Status.

	Median (range) or <i>n</i> ( <i>n</i> = 9)
<i>Sociodemographic</i>	
Age (years)	75 (40–81)
Sex	
Females, <i>n</i>	5
Caregiver relationship	
Spouse/partner, <i>n</i>	7
Family member, <i>n</i>	2
Ethnicity	
Caucasian, <i>n</i>	9
Education	
High school, <i>n</i>	4
Tertiary, <i>n</i>	4
Post-graduate, <i>n</i>	1
Marital status	
Married/de-facto, <i>n</i>	8
Single/never married, <i>n</i>	1
Residence type	
House, <i>n</i>	8
Unit/apartment, <i>n</i>	1
Number of persons living in residence	
Two persons, <i>n</i>	5
Three or more persons, <i>n</i>	4
Employment	
Employed, <i>n</i>	1
Unemployed, <i>n</i>	1
Retired, <i>n</i>	7
<i>Health status</i>	
Participants with chronic conditions, <i>n</i>	6
Total number of chronic conditions, <i>n</i>	1 (1–4)
Musculoskeletal, <i>n</i>	3
Hypertension, <i>n</i>	1
Diverticulitis, <i>n</i>	1
Osteoarthritis, <i>n</i>	1
Meniere's disease, <i>n</i>	1
Smoking status	
Current, <i>n</i>	0
Former, <i>n</i>	2
Never, <i>n</i>	7
Volume, pack-years <sup>a</sup>	1.7*
Alcohol consumption status <sup>b</sup>	
None, <i>n</i>	3
Low lifetime risk, <i>n</i>	5
High lifetime risk, <i>n</i>	1
Drinks per week,	7.83 (5.67) <sup>†</sup>

\*Data from two participants.

<sup>†</sup>Presented as mean (standard deviation).<sup>a</sup>Estimated volume from packs per day multiplied by years of smoking.<sup>b</sup>Categories as per The Australian Guidelines to Reduce Health Risks from Drinking Alcohol were used at the time of the study, low lifetime risk is ≤ 2 standard drinks per day or ≤ 14 standard drinks per week and high lifetime risk is ≥ 3 standard drinks per day or ≥ 21 standard drinks per week, lifetime risk regarding lifetime risk of alcohol-related injury (National Health and Medical Research Council, 2009).

**Table 3.** Loved One With Dementia: Baseline Body Composition and Physical Performance.

	Mean (SD) or n (n = 9)	
	Male (n = 3)	Female (n = 6)
<i>Body composition</i>		
Body mass index (kg m <sup>-2</sup> )	31.05 (24.66–31.33) <sup>†</sup>	27.33 (3.28)*
Bioelectrical impedance derived measures <sup>a</sup>		
Skeletal muscle mass (kg)	27.58 (2.43)	17.45 (3.89)
Skeletal muscle mass index (kg m <sup>-2</sup> )	9.85 (0.33)	6.92 (1.09)
Meets sarcopenia low muscle quantity criteria (<10.76 kg m <sup>-2</sup> males, <6.76 kg m <sup>-2</sup> females) <sup>b</sup> , n	3	3
Fat free mass (kg)	60.98 (47.74–62.06) <sup>†</sup>	40.21 (6.67)
Fat mass (kg)	29.79 (14.61–30.17) <sup>†</sup>	29.26 (10.94)
<i>Physical performance</i>		
Short Physical Performance Battery total score <sup>c</sup> , (0–12)		7.56 (2.46)
Balance tests score, (0–4)		4 (0–4) <sup>†</sup>
Gait speed test score, (0–4)		3.33 (0.71)
Chair stand test score, (0–4)		1.1 (0.93)
Habitual gait speed (ms <sup>-1</sup> )		1.02 (0.24)
Meets sarcopenia low performance criteria (≤0.8 ms <sup>-1</sup> ) <sup>d</sup> , n		2
Five chair stands, (seconds)		17.35 (2.44) <sup>‡</sup>
Meets sarcopenia low strength criteria (>15 seconds) <sup>d</sup> , n		8

\*A participant's height was estimated due to an inability to stand upright for a stretch stature measurement.

<sup>†</sup>Presented as median (range).

<sup>‡</sup>Data from three participants were missing for this measure.

<sup>a</sup>Bioelectrical impedance measures derived by the following equations: *skeletal muscle mass* = 0.401 (height in cm<sup>2</sup>/resistance in Ohms) + 3.825 (sex: male = 1; female = 0) + age in years (–0.071) + 5.102<sup>1</sup>, *skeletal muscle index* = skeletal muscle mass/(height in metres<sup>2</sup>), *fat free mass* = –4.03 + 0.734 (height in cm<sup>2</sup>/resistance in Ohms) + 0.116 (body mass in kg) + 0.096 (reactance in Ohms) + 0.984 (sex: male = 1; female = 0)<sup>2</sup>, *fat mass* = body mass in kg – fat free mass.

<sup>b</sup>Values were used from the European Working Group on Sarcopenia in Older People's age-related sarcopenia diagnostic criteria (Cruz-Jentoft et al., 2010) because they were derived from a validated bioelectrical impedance analysis measurement method.

<sup>c</sup>Higher scores indicate better lower extremity physical performance (Guralnik et al., 1994).

<sup>d</sup>Values were used from the updated European Working Group on Sarcopenia in Older People's aged-related sarcopenia diagnostic criteria (Cruz-Jentoft et al., 2019).

## Discussion

We have reported the primary outcomes of HOME CARE, the first remotely-supervised trial of MBSR for informal dementia caregivers combined with provision of PRT and balance training for their loved ones with dementia. As hypothesized, the intervention significantly improved the informal caregiver's mindfulness state and the functional mobility for their loved ones by a clinically meaningful amount. Although it did not improve the informal caregiver's perceived burden, notably it did not worsen it despite the demands of both learning mindfulness and facilitating exercise sessions.

Our first key finding was that the functional mobility of the loved ones with dementia improved meaningfully. Importantly, the informal caregivers had minimal exercise experience and facilitated the exercise intervention to their loved ones as opposed to an exercise professional, with only weekly remote supervision by a research assistant. To our knowledge, only three RCTs of home-based exercise in people with dementia used SPPB as an outcome and one RCT is currently pending results (Cezar et al., 2021). The two completed RCTs (Callahan et al., 2017; Pitkälä et al., 2013) reported no improvements in the SPPB. The first study described approximately 15 minutes of strength and balance

exercises per session for up to 24 sessions over 2 years as part of their occupational therapy intervention, but did not describe the exercise type, intensity, and planned progressions (Callahan et al., 2017). The second study described an hour of endurance, strength, balance, and executive functioning exercises twice a week for 12-months, but did not describe the intensity and planned progressions (Pitkälä et al., 2013). This suggests that the exercise prescriptions were insufficient in terms of progression and/or intensity. Our study progressed the exercise intensities when the participant reached a specific exertion level with safe technique and progressed the balance exercises after mastering their prescribed levels. The provision of exercise by the informal caregivers appeared to be safe as there were no adverse events attributed to the study, although the small sample size precludes any definitive conclusions in this regard. Therefore, the provision of robust, evidence-based exercise to a loved one after physician screening appears adaptable to the home environment. Notably, among the 34 dyads screened, only 1 of the 25 excluded were ineligible because of medical instability, suggesting the potential generalizability of this intervention to patients with dementia and multiple comorbidities living at home with an informal caregiver.

**Table 4.** Repeated Measures Linear Mixed Model Analyses Mean Scores (95% CIs) and Relative Effect Sizes for all Primary Outcomes.

Caregiver psychological assessments outcome measures		Randomization group		MBSR × Time		
	Time point	Control	MBSR	Mean differences* [95% CI]	Relative effect size [95% CI]	p Value
Zarit burden interview	Baseline	29.71 (21.86, 37.55)	26.51 (17.46, 35.57)	2.65 [-14.01, 19.31]	0.22 [-1.09, 1.54]	.622
	Week 4	28.94 (20.72, 37.17)	28.76 (19.71, 37.82)			
	Week 8	28.06 (19.84, 36.29)	32.07 (22.63, 41.51)			
	Week 12	30.31 (22.46, 38.15)	29.76 (20.71, 38.82)			
State mindfulness scale	Baseline	60.95 (51.11, 70.79)	51.25 (40.25, 62.25)	16.80 [-0.70, 34.30]	1.35 [-0.10, 2.81]	.009
	Week 4	69.40 (47.74, 91.06)	59.75 (37.26, 82.25)			
	Week 8	71.11 (42.72, 99.50)	51.20 (19.51, 82.88)			
	Week 12	51.40 (43.52, 65.28)	61.50 (49.34, 73.66)			
Loved one physical performance outcome measure				PRT + Balance training × time		
	Time point	Control	PRT + Balance training	Mean differences* [95% CIs]	Relative effect size [95% CIs]	p Value
Short Physical Performance Battery	Baseline	7.20 (4.89, 9.51)	8.00 (5.37, 10.58)	1.53 [-3.09, 6.14]	0.51 [-0.94, 1.97]	.091
	Week 12	6.00 (3.40, 8.60)	8.325 (5.37, 11.28)			

\*Mean differences of the primary outcomes = (week 12 minus baseline) intervention - (week 12 minus baseline) control. All data were normally distributed, and raw data were used. Mean scores represent the estimated marginal means and 95% confidence intervals (CIs) from repeated measures linear mixed models including all time points, with fixed effects of time, the randomization group, and their interactions. The ZBI mixed model used the AR(1) covariance type while the SMS and SPSS used the unstructured covariance type. Baseline ZBI data was used as a covariate in its mixed model analysis. *p*-Values are from the mixed model type III tests of fixed effects. Relative Hedges' bias corrected ES and 95% CI (Coe, 2022) were calculated for each time point and treatment effect for each primary outcome using estimated marginal means and SD from the mixed models: relative effect size (ES) = (post-test minus baseline) intervention - (post-test minus baseline) control/pooled baseline SD of cohort.

Our second key finding was a significant increase in the informal caregiver's mindfulness state. An increase in caregiver mindfulness state reduces the risk in developing stress, anxiety, and depression (Li et al., 2016). A systematic review of five RCTs implementing MBSR for dementia family caregivers reported reduced symptoms of depression and levels of anxiety (Fjorback et al., 2011), but the RCTs did not report mindfulness as an outcome. However, a systematic review (Li et al., 2016) of MBSR for dementia family caregivers included three UCTs and two RCTs, but mindfulness was not significantly increased in the studies. Our study suggested that MBSR was able to be undertaken in informal dementia caregivers without prior experience, and only one face-to-face training session supported by digital and written materials.

Our third key finding, contrary to our hypothesis, was that caregiver burden did not change differentially in response to the intervention. Over time, there was a

small and non-significant ( $F=0.164$ ,  $p=.92$ ) change of 2.65 points on the ZBI across both groups. We are uncertain whether this change is clinically meaningful as the MCID is not known. Other trials implementing MBSR have reported mixed outcomes for caregiver burden in this cohort. An RCT (Brown et al., 2016) and an uncontrolled trial (Epstein-Lubow et al., 2011) reported a small reduction in caregiver burden, as measured by the ZBI. However a systematic review of three RCTs (Shim et al., 2021) reported an improvement in caregiver burden in only one RCT (Liu et al., 2018), and a review (Liu et al., 2018) of five RCTs was uncertain whether MBSR reduced caregiver burden.

Our novel trial involved the provision of exercise by the informal caregivers in addition to completing the MBSR course, which has never been published to our knowledge. The lack of benefit on perceived burden in the intervention group, despite their improved mindfulness



and improved function of their loved ones, could have been due to the commitment of completing the MBSR intervention in addition to the provision of the exercise program for their loved ones, which included learning how to facilitate the exercise program by older adults with health concerns of their own. Thus, it is an important finding that caregiver burden was *not* differentially increased despite the many demands of learning two entirely new tasks: mindfulness and exercise supervision.

### Limitations

The primary limitation of this study is the smaller than planned sample size of 48 dyads due to limitations of funding and the advent of the COVID-19 restrictions prior to the completion of recruitment. This reduced the statistical power of the primary outcomes and resulted in wide confidence intervals. Post-hoc power analyses (Faul et al., 2009) indicated that a sample size of 55/group would have been needed to show statistical significance for the SPPB, given the ES of 0.54 suggesting the possibility of a type II error.

Second, the exercise adherence was less than expected which reduced the opportunities for exercise progression and full adaptation of strength, balance and mobility. This may have been due to the increase in time and demand for the informal caregivers to facilitate the exercise sessions in addition to attending to their activities of daily living. Exercise adherence may increase if informal caregivers completed the exercises together with their loved ones which will also likely improve informal caregiver psychosocial and physical health (Doyle et al., 2021).

Finally, the adherence to the MBSR program was also sub-optimal, particularly for the formal practices. The higher adherence to informal practices suggests that it was more feasible to integrate into the informal caregiver's activities of daily living than to schedule personal time to complete the formal practices of 30 min/day. Given that the informal caregivers experienced a large and statistical improvement in mindfulness even with their low-to-moderate adherence rate, future trials should test the feasibility and efficacy of a more abbreviated mindfulness intervention. One that could be incorporated *directly* into their caregiving tasks and exercise facilitation, rather than as a separate practice of its own, would be of great relevance.

### Conclusion

The HOMeCARE study was the first trial combining MBSR for informal dementia caregivers and caregiver-provision of PRT and balance training for their loved ones. The study appeared acceptable, feasible and adaptable to the home environment with no safety issues identified, despite the remote supervision. The HOMeCARE program resulted in a statistically significant improvement in informal caregiver mindfulness state which has yet to be reported in this cohort, as well as a clinically meaningful

improvement in functional mobility for their loved ones. There were no significant effects on caregiver burden. However, it is notable that the extensive requirements of the informal caregivers in the intervention group to learn not only how to practice mindfulness, but also how to supervise moderate-high intensity PRT and challenging balance exercises in their frail loved one with dementia for the first time - were substantial. The fact that these requirements did not *increase* burden relative to waiting list control informal caregivers is an extremely novel and important finding. Further large-scale trials are warranted to confirm and extend our findings to optimize adoption, adherence, and adaptation to this dyadic intervention, and ultimately to support ageing in place and quality of life for older adults with dementia and their informal caregivers.

### Author Note

The results were partially presented at the American Geriatrics Society 2023 Annual Scientific Meeting. The analysis and publication of the study was completed in partial fulfillment of Tommy Lang's Honors degree.

### Acknowledgments

The authors would like to thank the caregivers and their loved ones with dementia who contributed their valuable time and efforts to make the HOMeCARE study possible. The authors would also like to thank Prof. Sharon Naismith and Dr. Milena Simic for their contributions to the grant application.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by the Dementia Collaborative Research Centers' (DCRC) national grant round in 2016. The DCRC had no role in the design, methods, subject recruitment, data collections, analysis, and preparation of this paper.

### Institutional Review Board

The University of Sydney Human Research Ethics Committee, approval number 2016/795.

### ORCID iD

Tommy Lang  <https://orcid.org/0000-0001-8957-9028>

### Supplemental Material

Supplemental material for this article is available online.

### References

- Australian Health Ministers Advisory Council. (2015). *National framework for action on dementia 2015–2019*. <https://www.health.gov.au/resources/publications/national-framework-for-action-on-dementia-2015-2019>

- Australian Institute of Health Welfare. (2013). *The desire to age in place among older Australians* (Bulletin no. 114. Cat. no. AUS 169, Issue).
- Bazzano, A., Wolfe, C., Zylowska, L., Wang, S., Schuster, E., Barrett, C., & Lehrer, D. (2015). Mindfulness based stress reduction (MBSR) for parents and caregivers of individuals with developmental disabilities: A community-based approach. *Journal of Child and Family Studies, 24*(2), 298–308. <https://doi.org/10.1007/s10826-013-9836-9>
- Bhattacharyya, K. K., Liu, Y., Gothe, N. P., & Fauth, E. B. (2023). Mind-body practice and family caregivers' subjective well-being: Findings from the midlife in the United States (MIDUS) study. *Gerontology and Geriatric Medicine, 9*, 23337214231185912. <https://doi.org/10.1177/23337214231185912>
- Birnie, K., Garland, S. N., & Carlson, L. E. (2010). Psychological benefits for cancer patients and their partners participating in mindfulness-based stress reduction (MBSR). *Psycho-Oncology (Chichester, England), 19*(9), 1004–1009. <https://doi.org/10.1002/pon.1651>
- Brown, K. W., Coogle, C. L., & Wegelin, J. (2016). A pilot randomized controlled trial of mindfulness-based stress reduction for caregivers of family members with dementia. *Aging & Mental Health, 20*(11), 1157–1166. <https://doi.org/10.1080/13607863.2015.1065790>
- Callahan, C. M., Boustani, M. A., Schmid, A. A., La Mantia, M. A., Austrom, M. G., Miller, D. K., Gao, S., Ferguson, D. Y., Lane, K. A., & Hendrie, H. C. (2017). Targeting functional decline in Alzheimer disease: A randomized trial. *Annals of Internal Medicine, 166*(3), 164–171. <https://doi.org/10.7326/M16-0830>
- Cezar, N. O. C., Ansai, J. H., & de Andrade, L. P. (2021). Home-based multimodal exercise program in older people with Alzheimer disease: Randomized controlled trial protocol [Clinical Trial Protocol]. *Physiotherapy Research International, 26*(2), e1899. <https://doi.org/https://dx.doi.org/10.1002/pri.1899>
- Cheng, S. T. (2017). Dementia caregiver burden: A research update and critical analysis. *Current Psychiatry Reports, 19*(9), 64. <https://doi.org/10.1007/s11920-017-0818-2>
- Coe, R. (2022). *Effect size calculator: A guide to using the spreadsheet*. Accessed September 29, 2022, from <https://www.cem.org/effect-size-calculator>
- Cruz-Jentoft, A. J., Baeyens, J. P., Bauer, J. M., Boirie, Y., Cederholm, T., Landi, F., Martin, F. C., Michel, J.-P., Rolland, Y., Schneider, S. M., Topinková, E., Vandewoude, M., & Zamboni, M. (2010). Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age and Ageing, 39*(4), 412–423. <https://doi.org/10.1093/ageing/afq034>
- Cruz-Jentoft, A. J., Bahat, G., Bauer, J., Boirie, Y., Bruyere, O., Cederholm, T., Cooper, C., Landi, F., Rolland, Y., Sayer, A. A., Schneider, S. M., Sieber, C. C., Topinkova, E., Vandewoude, M., Visser, M., Zamboni, M., & Schols, J. (2019). Sarcopenia: Revised European consensus on definition and diagnosis. *Age and Ageing, 48*(1), 16–31. <https://doi.org/10.1093/ageing/afy169>
- Dent, E., Lien, C., Lim, W. S., Wong, W. C., Wong, C. H., Ng, T. P., Woo, J., Dong, B., de la Vega, S., Hua Poi, P. J., Kamaruzzaman, S. B. B., Won, C., Chen, L. K., Rockwood, K., Arai, H., Rodriguez-Manas, L., Cao, L., Cesari, M., Chan, P., . . . Flicker, L. (2017). The Asia-pacific clinical practice guidelines for the management of frailty. *Journal of the American Medical Directors Association, 18*(7), 564–575. <https://doi.org/10.1016/j.jamda.2017.04.018>
- Doyle, K. L., Toepfer, M., Bradfield, A. F., Noffke, A., Ausderau, K. K., Andreae, S., & Pickett, K. A. (2021). Systematic review of exercise for caregiver–care recipient dyads: What is best for spousal caregivers—exercising together or not at all? *The Gerontologist, 61*(6), e283–e301. <https://doi.org/10.1093/geront/gnaa043>
- Epstein-Lubow, G., McBee, L., Darling, E., Armey, M., & Miller, I. W. (2011). A pilot investigation of mindfulness-based stress reduction for caregivers of frail elderly. *Mindfulness, 2*(2), 95–102. <https://doi.org/10.1007/s12671-011-0047-4>
- Escrache-Escuder, A., Fuentes-Abolafio, I. J., Roldán-Jiménez, C., & Cuesta-Vargas, A. I. (2021). Effects of exercise on muscle mass, strength, and physical performance in older adults with sarcopenia: A systematic review and meta-analysis according to the EWGSOP criteria. *Experimental Gerontology, 151*, 111420. <https://doi.org/10.1016/j.exger.2021.111420>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using GPower 3.1: Tests for correlation and regression analyses. *Behavior Research Methods, 41*(4), 1149–1160. <https://doi.org/10.3758/BRM.41.4.1149>
- Fjorback, L. O., Arendt, M., Ornbol, E., Fink, P., & Walach, H. (2011). Mindfulness-based stress reduction and mindfulness-based cognitive therapy: A systematic review of randomized controlled trials. *Acta Psychiatrica Scandinavica, 124*(2), 102–119. <https://doi.org/10.1111/j.1600-0447.2011.01704.x>
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). “Mini-mental state”: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research, 12*(3), 189–198. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
- Grossman, P., Niemann, L., Schmidt, S., & Walach, H. (2004). Mindfulness-based stress reduction and health benefits: A meta-analysis. *Journal of Psychosomatic Research, 57*(1), 35–43. [https://doi.org/10.1016/S0022-3999\(03\)00573-7](https://doi.org/10.1016/S0022-3999(03)00573-7)
- Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D. G., Scherr, P. A., & Wallace, R. B. (1994). A short physical performance battery assessing lower extremity function: Association with self-reported disability and prediction of mortality and nursing home admission. *Journal of Gerontology (Kirkwood), 49*(2), M85–M94. <https://doi.org/10.1093/geronj/49.2.M85>
- Han, A. (2022). Effects of mindfulness-based interventions on depressive symptoms, anxiety, stress, and quality of life in family caregivers of persons living with dementia: A systematic review and meta-analysis. *Research on Aging, 44*(7–8), 494–509. <https://doi.org/10.1177/01640275211043486>
- Heyn, P., Abreu, B. C., & Ottenbacher, K. J. (2004). The effects of exercise training on elderly persons with cognitive impairment and dementia: A meta-analysis. *Archives of Physical Medicine and Rehabilitation, 85*(10), 1694–1704. <https://doi.org/10.1016/j.apmr.2004.03.019>
- Hordern, M. D., Dunstan, D. W., Prins, J. B., Baker, M. K., Singh, M. A. F., & Coombes, J. S. (2011). Exercise pre-

- scription for patients with type 2 diabetes and pre-diabetes: A position statement from exercise and sport science Australia. *Journal of Science and Medicine in Sport*, 15(1), 25–31. <https://doi.org/10.1016/j.jsams.2011.04.005>
- Jaffray, L., Bridgman, H., Stephens, M., & Skinner, T. (2016). Evaluating the effects of mindfulness-based interventions for informal palliative caregivers: A systematic literature review. *Palliative Medicine*, 30(2), 117–131. <https://doi.org/10.1177/0269216315600331>
- Kabat-Zinn, J. (2003). Mindfulness-based interventions in context: Past, present, and future. *Clinical Psychology: Science and Practice*, 10(2), 144–156. <https://doi.org/10.1093/clipsy.bpg016>
- Kabat-Zinn, J. (2013). *Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness*. Random House.
- Kor, P. P. K., Chien, W. T., Liu, J. Y. W., & Lai, C. K. Y. (2018). Mindfulness-based intervention for stress reduction of family caregivers of people with dementia: A systematic review and meta-analysis. *Mindfulness*, 9(1), 7–22. <https://doi.org/10.1007/s12671-017-0751-9>
- Lehmann, J., Michalowsky, B., Kaczynski, A., Thyrian, J. R., Schenk, N. S., Esser, A., Zwingmann, I., & Hoffmann, W. (2018). The impact of hospitalization on readmission, institutionalization, and mortality of people with dementia: A systematic review and meta-analysis. *Journal of Alzheimer's Disease*, 64, 735–749. <https://doi.org/10.3233/JAD-171128>
- Li, G., Yuan, H., & Zhang, W. (2016). The effects of mindfulness-based stress reduction for family caregivers: Systematic review. *Archives of Psychiatric Nursing*, 30(2), 292–299. <https://doi.org/10.1016/j.apnu.2015.08.014>
- LIFE-P, T. L. S. I. (2006). Effects of a physical activity intervention on measures of physical performance: Results of the lifestyle interventions and independence for elders pilot (LIFE-P) study. *The Journals of Gerontology: Series A*, 61(11), 1157–1165. <https://doi.org/10.1093/gerona/61.11.1157>
- Liu, Z., Sun, Y. Y., Zhong, B. L., & Liu, Z. (2018). Mindfulness-based stress reduction for family carers of people with dementia. *Cochrane Database of Systematic Reviews*, 8(8), CD012791. <https://doi.org/10.1002/14651858.CD012791.pub2>
- Lowery, D., Cerga-Pashoja, A., Iliffe, S., Thune-Boyle, I., Griffin, M., Lee, J., Bailey, A., Bhattacharya, R., & Warner, J. (2014). The effect of exercise on behavioural and psychological symptoms of dementia: The EVIDEM-E randomised controlled clinical trial. *International Journal of Geriatric Psychiatry*, 29(8), 819–827. <https://doi.org/10.1002/gps.4062>
- Mahoney, R., Regan, C., Katona, C., & Livingston, G. (2005). Anxiety and depression in family caregivers of people with alzheimer disease: The LASER-AD study. *The American Journal of Geriatric Psychiatry*, 13(9), 795–801. <https://doi.org/10.1097/00019442-200509000-00008>
- Milte, R., & Crotty, M. (2014). Musculoskeletal health, frailty and functional decline. *Best Practice & Research Clinical Rheumatology*, 28(3), 395–410. <https://doi.org/10.1016/j.berh.2014.07.005>
- National Health and Medical Research Council. (2009). *Australian guidelines to reduce health risks from drinking alcohol*.
- Nelson, M. E., Layne, J. E., Bernstein, M. J., Nuernberger, A., Castaneda, C., Kaliton, D., Hausdorff, J., Judge, J. O., Buchner, D. M., Roubenoff, R., & Fiatarone Singh, M. A. (2004). The effects of multidimensional home-based exercise on functional performance in elderly people. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 59(2), M154–M160. <https://doi.org/10.1093/gerona/59.2.M154>
- Paradise, M., McCade, D., Hickie, I. B., Diamond, K., Lewis, S. J., & Naismith, S. L. (2015). Caregiver burden in mild cognitive impairment. *Aging & Mental Health*, 19(1), 72–78. <https://doi.org/10.1080/13607863.2014.915922>
- Pedone, C., Ercolani, S., Catani, M., Maggio, D., Ruggiero, C., Quartesan, R., Senin, U., Mecocci, P., & Cherubini, A. (2005). Elderly patients with cognitive impairment have a high risk for functional decline during hospitalization: The GIFA study. *The Journals of Gerontology*, 60(12), 1576–1580. <https://doi.org/10.1093/gerona/60.12.1576>
- Perera, S., Mody, S. H., Woodman, R. C., & Studenski, S. A. (2006). Meaningful change and responsiveness in common physical performance measures in older adults. *Journal of the American Geriatrics Society (JAGS)*, 54(5), 743–749. <https://doi.org/10.1111/j.1532-5415.2006.00701.x>
- Pinquart, M., & Sörensen, S. (2003). Differences between caregivers and noncaregivers in psychological health and physical health: A meta-analysis. *Psychology and Aging*, 18(2), 250–267. <https://doi.org/10.1037/0882-7974.18.2.250>
- Pitkälä, K. H., Pöysti, M. M., Laakkonen, M.-L., Tilvis, R. S., Savikko, N., Kautiainen, H., & Strandberg, T. E. (2013). Effects of the Finnish Alzheimer Disease Exercise Trial (FINALEX): A randomized controlled trial. *JAMA Internal Medicine*, 173(10), 1–8. <https://doi.org/10.1001/jamainternmed.2013.359>
- Richardson, T. J., Lee, S. J., Berg-Weger, M., & Grossberg, G. T. (2013). Caregiver health: Health of caregivers of Alzheimer's and other dementia patients. *Current Psychiatry Reports*, 15(7), 367–367. <https://doi.org/10.1007/s11920-013-0367-2>
- Sharman, J. E., Smart, N. A., Coombes, J. S., & Stowasser, M. (2019). Exercise and sport science Australia position stand update on exercise and hypertension. *Journal of Human Hypertension*, 33(12), 837–843. <https://doi.org/10.1038/s41371-019-0266-z>
- Sherrington, C., Fairhall, N. J., Wallbank, G. K., Tiedemann, A., Michaleff, Z. A., Howard, K., Clemson, L., Hopewell, S., Lamb, S. E., & Sherrington, C. (2019). Exercise for preventing falls in older people living in the community. *Cochrane Database of Systematic Reviews*, 1(1), CD012424. <https://doi.org/10.1002/14651858.CD012424.pub2>
- Shim, M., Tilley, J. L., Im, S., Price, K., & Gonzalez, A. (2021). A systematic review of mindfulness-based interventions for patients with mild cognitive impairment or dementia and caregivers. *Journal of Geriatric Psychiatry and Neurology*, 34(6), 528–554. <https://doi.org/10.1177/0891988720957104>
- Singh, N. A., Clements, K. M., & Fiatarone, M. A. (1997). A randomized controlled trial of progressive resistance training in depressed elders. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 52A(1), M27–M35. <https://doi.org/10.1093/gerona/52A.1.M27>
- Singh, N. A., Quine, S., Clemson, L. M., Williams, E. J., Williamson, D. A., Stavrinou, T. M., Grady, J. N., Perry,

- T. J., Lloyd, B. D., Smith, E. U., & Singh, M. A. (2012). Effects of high-intensity progressive resistance training and targeted multidisciplinary treatment of frailty on mortality and nursing home admissions after hip fracture: a randomized controlled trial. *Journal of the American Medical Directors Association, 13*(1), 24–30. <https://doi.org/10.1016/j.jamda.2011.08.005>
- Song, D., Yu, D. S. F., Li, P. W. C., & Lei, Y. (2018). The effectiveness of physical exercise on cognitive and psychological outcomes in individuals with mild cognitive impairment: A systematic review and meta-analysis. *International Journal of Nursing Studies, 79*, 155–164. <https://doi.org/10.1016/j.ijnurstu.2018.01.002>
- Tanay, G., & Bernstein, A. (2013). State mindfulness scale (SMS): Development and initial validation. *Psychological Assessment, 25*(4), 1286–1299. <https://doi.org/10.1037/a0034044>
- Vanhees, L., De Sutter, J., Geladas, N., Doyle, F., Prescott, E., Cornelissen, V., Koudi, E., Dugmore, D., Vanuzzo, D., Börjesson, M., & Doherty, P. (2012). Importance of characteristics and modalities of physical activity and exercise in defining the benefits to cardiovascular health within the general population: Recommendations from the EACPR (Part I). *European Journal of Preventive Cardiology, 19*(4), 670–686. <https://doi.org/10.1177/2047487312437059>
- Zarit, S. H., Reever, K. E., & Bach-Peterson, J. (1980). Relatives of the impaired elderly: Correlates of feelings of burden. *The Gerontologist, 20*(6), 649–655. <https://doi.org/10.1093/geront/20.6.649>