

## Cytotoxic Effect of Some 1, 4-Dihydropyridine Derivatives Containing Nitroimidazole Moiety

Ramin Miri<sup>a,b,\*</sup>, Katayoun Javidnia<sup>a,b</sup>, Zahra Amirghofran<sup>a,c</sup>, Seyyed Hossein Salimi<sup>a,b</sup>, Zahra Sabetghadam<sup>a,b</sup>, Savis Meili<sup>a,b</sup> and Ahmad Reza Mehdipour<sup>a,b</sup>

<sup>a</sup>Medicinal and Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. <sup>b</sup>Department of Medicinal Chemistry, Faculty of Pharmacy, Shiraz University of Medical Sciences, Iran. <sup>c</sup>Department of Immunology, Faculty of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

---

### Abstract

The 1,4-dihydropyridine (DHP) derivatives are a known class of calcium channel blockers. Some derivatives of DHP showed significant cytotoxicity. It was shown that this effect may not be the result of interaction with Ca<sup>2+</sup> channels. In this study, we performed an investigation about the intrinsic cytotoxicity of some derivatives of DHP containing nitroimidazole moiety on their C4 position on four different cancer cell lines (Raji, K562, Fen and HeLa). The result showed that these compounds had moderate-good cytotoxic activity. In addition, QSAR model shows the importance of N atom in cytotoxicity; Ca<sup>2+</sup> channels.

**Keywords:** 1, 4-Dihydropyridines; Cytotoxicity; MTT assay; Nitroimidazole.

---

### Introduction


The 1, 4-dihydropyridine (DHP) derivatives, known as calcium channel antagonists, are used for treatment of cardiovascular diseases such as hypertension and angina pectoris (1-4). It has been discovered that DHPs possess a wide range of other beneficial biological activities such as anticonvulsant analgesic and are also used as a chemical drug delivery systems especially in brain delivery (5-8). Recently, it has been proved that these compounds make a new class of multidrug resistance (MDR) reversals in cancer treatment. Therefore, extensive investigations were made in order to find new DHP derivatives as MDR reversal agents (9-11).

Beside the reversing activity of DHPs, there are several reports about their intrinsic Cytotoxicity. Particularly, some derivatives showed significant cytotoxicity such as dextropropripridine and some dibenzoyl derivatives (12-15). In addition, there are some reports on the effects of DHPs on potentiation of antitumoral and antimetastatic activity of some common cytotoxic drugs (16). Mechanistic investigations were proved that cytotoxicity of DHPs may not be the result of interaction with Ca<sup>2+</sup> channels although it might be related to other pathways in which calcium is involved such as calcium/calmodulin pathway or other cellular pathways like inhibition of topoisomerase I (12, 17, 18).

In this study, we performed an investigation about the intrinsic cytotoxicity of some derivatives of DHP which were bearing a nitroimidazole group on the C<sub>4</sub> position.

---

\* Corresponding author:  
E-mail: mirir@sums.ac.ir

**Table 1.** Structure of DHP derivatives used in this study and their calcium channel antagonist activity.

Compound	R <sub>1</sub>	R <sub>2</sub>	IC <sub>50</sub>
A1	CH <sub>3</sub>	CH <sub>3</sub>	9.03×10 <sup>-12</sup>
A2	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	8.78×10 <sup>-11</sup>
A3	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	5.11×10 <sup>-11</sup>
A4	CH <sub>2</sub> CH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>	4.38×10 <sup>-10</sup>
A5	CH <sub>2</sub> -cyclopropyl	CH <sub>2</sub> -cyclopropyl	2.81×10 <sup>-8</sup>
A6	CH <sub>3</sub>	CH(CH <sub>2</sub> ONO <sub>2</sub> ) <sub>2</sub>	3.86×10 <sup>-12</sup>
A7	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> OCO(1-methyl-pyridinium) Iodide	-----
A8	CH <sub>3</sub>	1-ethyl-4-(4-fluorophenyl)piperazinyl	2.65×10 <sup>-10</sup>
A9	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> OCO(1-methyl-pyridinium) Iodide	-----
A10	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	1.09×10 <sup>-9</sup>
A11	CH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>	CN	2.1×10 <sup>-6</sup>
A12	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> ONO <sub>2</sub>	4.87×10 <sup>-12</sup>
A13	CH <sub>2</sub> CH <sub>2</sub> CN	CN	5.4×10 <sup>-6</sup>
A14	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CN	6.86×10 <sup>-13</sup>
A15	CH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>	1-ethyl-4-(4-fluorophenyl)piperazinyl	2.13×10 <sup>-11</sup>
A16	CH <sub>2</sub> CH <sub>3</sub>	1-ethyl-4-(4-fluorophenyl)piperazinyl	4.88×10 <sup>-11</sup>
A17	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> COCH <sub>3</sub>	9.03×10 <sup>-12</sup>

## Experimental

### *Dihydropyridine compounds*

All compounds used in this study have been synthesized in our chemistry lab and their synthesis process and their Ca<sup>2+</sup> blocking activity were previously reported (8, 19-23). Structures of compounds and their Ca channel blocking activity are presented in Table 1.

### *Cell lines and reagents*

Five cell lines including K-562 (chronic myelogenous leukemia), HeLa (carcinoma of the cervix), Raji (Human, B cell Lymphoma), and Fen (bladder carcinoma cell line) were all obtained from the Iran Pasteur Institute, Tehran, Iran. Cells were cultured in RPMI-1640 (Sigma,

USA) supplemented with 10% fetal bovine serum (Gibco, USA), 100 IU/mL penicillin and 100 µg/mL streptomycin (Biosera, England). Cells were cultured in 50 cm<sup>3</sup> flask (Nunc, Denmark) with 5 mL of culture medium in humidified 5% CO<sub>2</sub> incubator at 37°C.

### *Cytotoxicity evaluation*

Appropriate amount of compounds was mixed with DMSO (dimethyl sulfoxide). Then, four concentration (10, 100, 50, 1000 µM) were made with serial dilution by RPMI-1640. DMSO, as the negative control, was diluted with the same method. For positive control, doxorubicin was mixed with RPMI-1640 to make similar concentrations. MTT [3-(4, 5-dimethylthiazole-2-yl)-2, 5-diphenyl tetrazolium bromide]

**Table 2.** Cell growth inhibitory activity of compounds a-e *in vitro* (IC<sub>15</sub>).

Compound	IC <sub>15</sub> <sup>a</sup> ( $\mu$ M)			
	Raji	K562	Fen	HeLa
A1	< 1	41.8	< 1	38.1
A2	< 1	84.0	< 1	54.7
A3	< 1	6.2	< 1	38.8
A4	< 1	20.2	< 1	45.5
A5	< 1	< 1	< 1	72.4
A6	< 1	< 1	< 1	< 1
A7	< 1	4.1	< 1	< 1
A8	< 1	< 1	< 1	1.2
A9	< 1	< 1	< 1	< 1
A10	< 1	< 1	< 1	< 1
A11	> 100	52.5	51.5	53.5
A12	< 1	< 1	< 1	< 1
A13	< 1	< 1	< 1	< 1
A14	< 1	< 1	< 1	< 1
A15	< 1	< 1	< 1	< 1
A16	< 1	< 1	< 1	< 1
A17	22.4	44.4	44.7	35.5
Doxorubicin	< 1	< 1	< 1	< 1
DMSO	92.4	80.4	71.4	81.1

a: IC<sub>15</sub> is the molar concentration causing 15% growth inhibition of tumor cells.

(Sigma, USA), was dissolved in PBS (phosphate buffered saline) at 5 mg/mL. Each well of the plate was filled with  $1-5 \times 10^4$  cells (depending on the cell line) in 90  $\mu$ L cultured medium. Then, 10  $\mu$ L from the stock solutions of compounds, negative controls and positive controls, was added as triplicate to the wells to reach the final concentration of 1, 10, 50, 100  $\mu$ M. Three wells containing only the same number of cells left in each plate. Plates were kept in a humidified incubator for 48 h. After the incubation period, MTT assay was carried out by the procedure described by Jabbar *et al.* (24).

#### QSAR studies

Chemical structure of the molecules was built using HyperChem software (Version 7, Hypercube Inc., <http://www.hyper.com>, USA) for the structural chemistry. Gaussian98 was

**Table 3.** Cell growth inhibitory activity of compounds a-e *in vitro* (IC<sub>30</sub>).

Compound	IC <sub>30</sub> <sup>a</sup> ( $\mu$ M)			
	Raji	K562	Fen	HeLa
A1	58.3	100 <	33.1	100 <
A2	1 >	100 <	57.8	100 <
A3	32.5	100 <	37.3	100 <
A4	1 >	100 <	58.9	100 <
A5	100 <	100 <	71.1	100 <
A6	1 >	1 >	1 >	1 >
A7	12.3	100 <	10.5	12.6
A8	1.4	1.3	1.4	5.6
A9	1.1	1.1	1.1	5.4
A10	1	1 >	1 >	3.1
A11	100 <	100 <	100 <	100 <
A12	1.4	1.2	1.2	1.7
A13	1 >	1 >	1 >	1 >
A14	1 >	1 >	1 >	1 >
A15	1.1	2.1	1.3	3.1
A16	1	1 >	1 >	3.4
A17	100 <	1 >	100 <	100 <
Doxorubicin	1 >	1 >	1 >	1 >
DMSO	100 <	100 <	100 <	100 <

a: IC<sub>30</sub> is the molar concentration causing 30% growth inhibition of tumor cells.

operated to optimize the molecular structure. The structures were optimized by Ab initio calculations at the level of RHF/ STO-3G. No molecular symmetry constraint was applied; rather full optimization of all bond lengths and angles was carried out. Highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) energies, molecular polarizability (MP) and molecular dipole moment (MDP) were all calculated by Gaussian98. Local charges (LC) at each atom were calculated according to both Mulliken population and natural population analysis methods. Some theoretical QSAR descriptors including physicochemical properties and topological indices were calculated using HyperChem (Hypercube Inc.) and DRAGON ([http://www.taletе.mi.it/dragon\\_exp.htm](http://www.taletе.mi.it/dragon_exp.htm), Italia) software.

**Table 4.** Cell growth inhibitory activity of compounds a-e *in-vitro* (IC<sub>50</sub>).

Compound	IC <sub>50</sub> <sup>a</sup> ( $\mu$ M)			
	Raji	K562	Fen	HeLa
A1	> 100	> 100	> 100	> 100
A2	> 100	> 100	> 100	> 100
A3	> 100	> 100	> 100	> 100
A4	> 100	> 100	> 100	> 100
A5	> 100	> 100	> 100	> 100
A6	4.0	2.8	4.1	5.4
A7	69.3	> 100	56.3	68.3
A8	15.4	14.1	15.1	30.9
A9	7.1	3.0	6.8	29.4
A10	8.1	3.8	4.3	25.3
A11	> 100	> 100	> 100	> 100
A12	14.1	14.3	14.3	14.9
A13	< 1	< 1	< 1	< 1
A14	3.1	2.5	2.4	2.8
A15	8.4	20.2	8.8	23.4
A16	8.2	3.3	7.2	28.5
A17	> 100	> 100	> 100	> 100
Doxorubicin	< 1	< 1	< 1	< 1
DMSO	> 100	> 100	> 100	> 100

a: IC<sub>50</sub> is the molar concentration causing 50% growth inhibition of tumor cells.

The MLR analysis was employed to derive the QSAR models for different cytotoxic activities. In order to make better QSAR models, IC<sub>50</sub> of compounds was transformed into the logarithmic scale (log IC<sub>50</sub>). MLR and correlation analyses were carried out by the statistics software SPSS 11.5 version. Before any MLR analysis, the correlation between the selected descriptors was examined (Table 3) and collinear descriptors ( $r > 0.90$ ) were determined. Among these descriptors, the one which had higher correlation with the dependent variable was retained and the others were removed from the descriptor data matrix. The remaining descriptors were used to construct the MLR model, in accordance with the stepwise selection and elimination method. The better regression models were selected on the basis of the higher R, f-value (a statistic of assessing the overall significance) and the lower

SEE. Cross-validation procedure [leave-one-out ( $Q^2_{LOO}$ )] was applied to measure the predictive capabilities of the models by using MATLAB 6.5 program (25).

## Results and Discussion

The data of cytotoxicity is shown in Tables 2, 3 and 4. IC<sub>15</sub>, IC<sub>30</sub> and IC<sub>50</sub> were calculated for each compound. The most cytotoxic effect was seen on Raji cell line while the lowest activity was observed in K562 and HeLa, since at least 7 compounds had IC<sub>30</sub> and IC<sub>50</sub> of higher than 100  $\mu$ M.

However, synthesized compounds had moderate cytotoxicity. It was obvious that symmetric compounds are significantly less potent since all of them have IC<sub>50</sub> greater than 100  $\mu$ M. On the other hand, the asymmetric derivatives had good cytotoxicity. Among them, compound A13 showed the best activity since its IC<sub>50</sub> in all cell lines were lower than 1  $\mu$ M which was comparable with the reference drug (doxorubicin).

### QSAR studies

MLR analysis with the stepwise selection and elimination of variables was utilized to model the structure-activity relationships with diverse set of descriptors.

In the first step, it was tried to find an appropriate model for cytotoxic activity of all compounds in Raji. The obtained equation is as follows:

$$\log IC_{50} = 5.966 - 0.551 (\pm 0.097) nN - 0.897 (\pm 0.242) C - 040$$

$$N = 17, R_2 = 0.812, Q^2_{LOO} = 0.772, F = 30.1, SE = 0.311$$

In this equation, the values in the parenthesis stand for the standard deviation of the coefficients. N, R<sup>2</sup>, SE, and F are number of components, correlation coefficient, standard error of regression, and Fisher's F-ratio, respectively. Noticeably, this equation has a good statistical feature, which can explain and predict 82% of variances in the cytotoxic activity data.  $Q^2_{LOO}$  represents acceptable correlation coefficient, which is close to that of calibration.

This explains the satisfactory prediction ability and stability of the resulted model. This two-parametric equation, holding the number of nitrogen atoms ( $nN$ ) and an atom center fragment parameters ( $C-040$ ), implies the importance of constitutional parameters in their activity.

The next equation was obtained for cytotoxicity of K562 cell line:

$$\log IC_{50} = 6.002 - 0.571 (\pm 0.142) nN - 0.878 (\pm 0.353) C - 040$$

$N = 17, R_2 = 0.680, Q^2_{LOO} = 0.600, F = 14.88, SE = 0.453$

It is obvious that cytotoxic activity K562 cell line has a great similarity with the activity in Raji cell line, though statistical quality of latter equation is much lower than the first equation. This might imply the similar mechanism of cytotoxicity in both cell lines.

The next equation was obtained for the cytotoxicity of Fen cell line:

$$\log IC_{50} = 4.147 - 0.689 (\pm 0.117) nN - 0.367 (\pm 0.140) MLOGP$$

$N = 17, R_2 = 0.812, Q^2_{LOO} = 0.636, F = 19.1, SE = 0.383$

In this equation as well, the number of nitrogen is the most important factor while, dissimilar to the former equation, the second parameter is  $MLOGP$  which octanol-water partition coefficient. This may suggest that transversing of biological membranes is a key factor in cytotoxic activity in this cell lines, since it was proved that  $\log P$  is a crucial factor in crossing the biological membrane (26).

Finally, an equation for HeLa cell line was obtained as described below:

$$\log IC_{50} = 7.23 - 1.641 (\pm 0.290) nCN - 0.211 (\pm 0.036) SLCDHP + 9.123 (\pm 2.91) nHAcc$$

$N = 17, R_2 = 0.856, Q^2_{LOO} = 0.644, F = 25.8, SE = 0.257$

This three-parametric equation containing the holding number of nitrile groups ( $nCN$ ), sum of local charge on DHP ring ( $SLC_{DHP}$ ) and the number of acceptor atoms for hydrogen bonding

( $nHAcc$ ), shows that cytotoxic activity of DHP derivatives in this cell line probably is slightly different from other cell lines, while the presence of nitrogen (here in nitrile form) is a key factor.

Overall, it can be concluded that presence of nitrogen has a crucial importance in the cytotoxicity DHP derivatives in both total number of nitrogen and also groups containing nitrogen (*i.e.* nitrile group).

On the other hand, modeling the  $Ca^{+2}$  channel blocking activity of these compounds gives the following equation:

$$\log IC_{50} = 9.74 + 3.60 (\pm 0.60) SLCNI - 1.562 (\pm 0.352) H - 046 - 1.459 (\pm 0.569) Mv$$

$N = 15, R_2 = 0.852, Q^2_{LOO} = 0.701, F = 19.1, SE = 0.49$

Where  $SLC_{NI}$  is the sum of local charge on nitroimidazole ring, H-046 is an atom-centered fragment (H attached to C0 ( $sp^3$ )) and  $Mv$  is the mean atomic van der Waals volume. This equation shows the importance of charges in C4 position while volume of molecules has negative effect on  $Ca^{+2}$  channel blocking activity.

## Conclusion

DHPs are one of the major classes of  $Ca^{2+}$  channel blockers. Recently, their MDR reversing activity attracted many researchers. In this way, a large set of new derivatives of DHP were synthesized as MDR reversals, some of which have reached to the upper stage of studies.

In addition, some derivatives showed intrinsic cytotoxicity on cancer cell lines. As a remarkable example, dexniguldipine, a potent MDR reversal, exhibited strong cytotoxicity on various cell lines. On the other hand, it was shown that nitroimidazole and its derivatives possessed appropriate cytotoxic properties (27-29). Therefore, it seems that DHP compounds including nitroimidazole moiety at C4 position may have strong cytotoxicity due to the simultaneous presence of DHP base structure and nitroimidazole substituent. In the present study, cytotoxic activity of some DHP derivatives was evaluated on four different cancer cell lines. Asymmetric derivatives showed good

cytotoxicity while symmetric ones were less potent.

In addition, QSAR studies revealed the significance of N atom and group containing it in cytotoxic activity. This is compatible with previous findings about MDR reversing activity of DHPs which demonstrated the good association of amine groups with MDR reversing effect. For example, Ford's findings showed that tertiary and cyclic amine groups are decisive for MDR reversing activity (30) and in recent years, in most synthesized DHP derivatives as MDR modulators, this rule has been applied (10, 31). This correlation between MDR reversing activity and cytotoxic effect might be the sign of possible similar mechanisms interfering in both phenomena.

On the other hand, QSAR model of Ca<sup>2+</sup> channel blocking activity confirms that key factors in this activity are to some extent different from cytotoxic activity. For example, while local charges of DHP is an important feature in cytotoxic effect, the significant part for Ca<sup>2+</sup> channel blocking activity is nitroimidazole moiety. Besides, the most significant factor in cytotoxic effect (number of nitrogen atoms) does not have any considerable role in Ca<sup>2+</sup> channel blocking activity. Another remarkable point about Ca<sup>2+</sup> channel blocking activity is that the existence of local charge of nitroimidazole moiety in QSAR model is in agreement with previous works which highlighted the importance of the substituent of C4 position (32).

### Acknowledgments

The financial support of the Research Council of Shiraz University of Medical Sciences is acknowledged. This project is a part of Pharm. D theses of SH Salimi and Z. Sabetghadam (Theses Number: 284 and 341).

### References

- (1) Coburn RA, Wierzba M, Suto MJ, Solo AJ, Triggle AM and Triggle DJ. 1, 4-Dihydropyridine antagonist activities at the calcium channel: a quantitative structure-activity relationship approach. *J. Med. Chem.* (1988) 31: 2103-2107.
- (2) Shekarchi M, Binesh Marvasti M, Sharifzadeh M and Shafiee A. Anticonvulsant activities of 7-phenyl-5H-thiazolo[5, 4-e][1, 2, 3, 4]tetrazolo[5, 1-c]pyrrolo[1, 2-a][1, 4]diazepine and 7-phenyl-5H-thiazolo[5, 4-e][1, 3, 4]triazolo[5, 1-c] pyrrolo[1, 2-a][1, 4]diazepines. *Iranian J. Pharm. Res.* (2005) 4: 33-36.
- (3) Miri R, Javidnia K, Sarkarzadeh H and Hemmateenejad B. Synthesis, study of 3D structures, and pharmacological activities of lipophilic nitroimidazolyl-1, 4-dihydropyridines as calcium channel antagonist. *Bioorg. Med. Chem.* (2006) 14: 4842-4849.
- (4) Velazquez C and Knaus EE. Synthesis and biological evaluation of 1, 4-dihydropyridine calcium channel modulators having a diazen-1-ium-1, 2-diolate nitric oxide donor moiety for the potential treatment of congestive heart failure. *Bioorg. Med. Chem.* (2004) 12: 3831-3840.
- (5) Bodor N. Redox drug delivery systems for targeting drugs to the brain. *Ann. NY Acad. Sci.* (1987) 507: 289-306.
- (6) Mortazavi SM. Advances in targeted drug delivery. *Iranian J. Pharm. Res.* (2007) 6 149-150.
- (7) Kaymaz AA, Tan H, Altug T and Buyukdevrim AS. The effects of calcium channel blockers, verapamil, nifedipine and diltiazem, on metabolic control in diabetic rats. *Diabetes Res. Clin. Pract.* (1995) 28: 201-205.
- (8) Miri R, Javidnia K, Kebriaie-Zadeh A, Niknahad H, Shaygani N, Semnianian S and Shafiee A. Synthesis and evaluation of pharmacological activities of 3, 5-dialkyl 1, 4-dihydro-2, 6-dimethyl-4-nitroimidazole-3, 5-pyridine dicarboxylates. *Arch. Pharm. (Weinheim)* (2003) 336: 422-428.
- (9) Kawase M and Motohashi N. New multidrug resistance reversal agents. *Curr. Drug Targets* (2003) 4: 31-43.
- (10) Mehdipour AR, Javidnia K, Hemmateenejad B, Amirghofran Z and Miri R. Dihydropyridine derivatives to overcome atypical multidrug resistance: design, synthesis, QSAR studies, and evaluation of their cytotoxic and pharmacological activities. *Chem. Biol. Drug Des.* (2007) 70: 337-346.
- (11) Zhou XF, Yang X, Wang Q, Coburn RA and Morris ME. Effects of dihydropyridines and pyridines on multidrug resistance mediated by breast cancer resistance protein: *in-vitro* and *in-vivo* studies. *Drug Metab. Dispos.* (2005) 33: 1220-1228.
- (12) Gietzen K, Abdallah F and Bai G. Inhibition of tumor cell growth by a novel dihydropyridine derivative. *Eur. J. Cancer* (1990) 26: 922-923.
- (13) Hahn KA, Legendre AM and Schuller HM. Amputation and dexniguldipine as treatment for canine appendicular osteosarcoma. *J. Cancer Res. Clin. Oncol.* (1997) 123: 34-38.
- (14) Jorjani M, Rastegar H, Roshanzamir F, Varmazyari M, Abdollahi M and Zarghi A. Synthesis and biological evaluation of new 1, 4-dihydropyridines as antihypertensives agents in rats. *Iranian J. Pharm. Res.* (2003) 2: 43-46.
- (15) Morshed SR, Hashimoto K, Murotani Y, Kawase M, Shah A, Satoh K, Kikuchi H, Nishikawa H, Maki

- J and Sakagami H. Tumor-specific cytotoxicity of 3,5-dibenzoyl-1,4-dihydropyridines. *Anticancer Res.* (2005) 25: 2033-2038.
- (16) Fedeli L, Colozza M, Boschetti E, Sabalich I, Aristei C, Guerciolini R, Del Favero A, Rossetti R, Tonato M, Rambotti P and Davis S. Pharmacokinetics of vincristine in cancer patients treated with nifedipine. *Cancer* (1989) 64: 1805-1811.
- (17) Orth SR, Nobiling R, Bonisch S and Ritz E. Inhibitory effect of calcium channel blockers on human mesangial cell growth: evidence for actions independent of L-type  $Ca^{2+}$  channels. *Kidney Int.* (1996) 49: 868-879.
- (18) Straub T, Boesenberg C, Gekeler V and Boege F. The dihydropyridine dextro-gulidipine hydrochloride inhibits cleavage and relegation reactions of eukaryotic DNA topoisomerase I. *Biochemistry* (1997) 36: 10777-10783.
- (19) Miri R, Dehpour AR, Azimi M and Shafiee A. Synthesis and smooth muscle calcium channel antagonist effect of alkyl, aminoalkyl 1, 4-dihydro-2, 6-dimethyl-4-nitroimidazole-3,5 pyridine dicarboxylates. *Daru* (2001) 9: 40-45.
- (20) Miri R, Javidnia K, Mirkhani H, Hemmateenejad H, Kazemi Z, Edraki N and Mehdipour AR. Synthesis and *in-vitro* dual calcium channel antagonist-agonist activity of some 1, 4-dihydro-2, 6-dimethyl-3-nitro and cyano-4-(1-methyl-5-nitro-1H-imidazol-2-yl)-5-pyridinecarboxylates. *Daru* (2008) 16: 263-270.
- (21) Miri R, Niknahad H, Vazin A, Azarpira A and Shafiee A. Synthesis and smooth muscle calcium channel antagonist effects of new derivatives of 1,4-dihydropyridine containing nitroimidazolyl substituent. *Daru* (2002) 10: 130-136.
- (22) Miri R, Niknahad H, Vesal G and Shafiee A. Synthesis and calcium channel antagonist activities of 3-nitrooxyalkyl, 5-alkyl 1,4-dihydro-2,6-dimethyl-4-(1-methyl-5-nitro-2-imidazolyl)-3, 5-pyridinedicarboxylates. *Farmaco.* (2002) 57: 123-128.
- (23) Shafiee A, Miri R, Dehpour AR and Soleymani F. Synthesis and calcium-channel antagonist activity of nifedipine analogues containing nitroimidazolyl substituent in guinea-pig ileal smooth muscle. *Pharm. Sci.* (1996) 2: 541-543.
- (24) Jabbar SA, Twentyman PR and Watson JV. The MTT assay underestimates the growth inhibitory effects of interferons. *Br. J. Cancer* (1989) 60: 523-528.
- (25) Mehdipour AR, Hemmateenejad B and Miri R. QSAR studies on the anesthetic action of some polyhalogenated ethers. *Chem. Biol. Drug Des.* (2007) 69: 362-368.
- (26) Hughes JD, Blagg J, Price DA, Bailey S, Decresceno GA, Devraj RV, Ellsworth E, Fobian YM, Gibbs ME, Gills RW, Greene N, Huang E, Krieger-Burke T, Loesel J, Wager T, Whiteley L and Zhang Y. Physicochemical drug properties associated with *in-vivo* toxicological outcomes. *Bioorg. Med. Chem. Lett.* (2008) 18: 4872-4875.
- (27) Brezden CB, Horn L, McClelland RA and Rauth AM. Oxidative stress and 1-methyl-2-nitroimidazole cytotoxicity. *Biochem. Pharmacol.* (1998) 56: 335-344.
- (28) Lee HH, Palmer BD, Wilson WR and Denny WA. Synthesis and hypoxia-selective cytotoxicity of 2-nitroimidazole mustard. *Bioorg. Med. Chem. Lett.* (1998) 8: 1741-1744.
- (29) Moselen JW, Hay MP, Denny WA and Wilson WR. *N*-[2-(2-methyl-5-nitroimidazolyl) ethyl]-4-(2-nitroimidazolyl)butanamide (NSC 639862), a bisnitroimidazole with enhanced selectivity as a bioreductive drug. *Cancer Res.* (1995) 55: 574-580.
- (30) Ford JM and Hait WN. Pharmacology of drugs that alter multidrug resistance in cancer. *Pharmacol. Rev.* (1990) 42: 155-199.
- (31) Tasaka S, Ohmori H, Gomi N, Lino M, Machida T, Kiue A, Natio S and Kuwano M. Synthesis and structure-activity analysis of novel dihydropyridine derivatives to overcome multidrug resistance. *Bioorg. Med. Chem. Lett.* (2001) 11: 275-257.
- (32) Mohajeri A, Hemmateenejad B, Mehdipour A and Miri R. Modeling calcium channel antagonistic activity of dihydropyridine derivatives using QTMS indices analyzed by GA-PLS and PC-GA-PLS. *J. Mol. Graph. Model.* (2008) 26: 1057-1065.