



# Water oxidation by photosystem II is the primary source of electrons for sustained H<sub>2</sub> photoproduction in nutrient-replete green algae

Sergey Kosourov<sup>a,1</sup>, Valéria Nagy<sup>a</sup>, Dmitry Shevela<sup>b</sup>, Martina Jokel<sup>a</sup>, Johannes Messinger<sup>b,c,1</sup>, and Yagut Allahverdiyeva<sup>a,1</sup>

<sup>a</sup>Molecular Plant Biology, Department of Biochemistry, University of Turku, FI-20520 Turku, Finland; <sup>b</sup>Department of Chemistry, Chemical Biological Centre, Umeå University, 90187 Umeå, Sweden; and <sup>c</sup>Department of Chemistry–Ångström Laboratory, Uppsala University, 75120 Uppsala, Sweden

Edited by Pierre Joliot, Institut de Biologie Physico-Chimique, Paris, France, and approved October 1, 2020 (received for review May 8, 2020)

The unicellular green alga *Chlamydomonas reinhardtii* is capable of photosynthetic H<sub>2</sub> production. H<sub>2</sub> evolution occurs under anaerobic conditions and is difficult to sustain due to 1) competition between [FeFe]-hydrogenase (H<sub>2</sub>ase), the key enzyme responsible for H<sub>2</sub> metabolism in algae, and the Calvin–Benson–Bassham (CBB) cycle for photosynthetic reductants and 2) inactivation of H<sub>2</sub>ase by O<sub>2</sub> coevolved in photosynthesis. Recently, we achieved sustainable H<sub>2</sub> photoproduction by shifting algae from continuous illumination to a train of short (1 s) light pulses, interrupted by longer (9 s) dark periods. This illumination regime prevents activation of the CBB cycle and redirects photosynthetic electrons to H<sub>2</sub>ase. Employing membrane-inlet mass spectrometry and H<sub>2</sub><sup>18</sup>O, we now present clear evidence that efficient H<sub>2</sub> photoproduction in pulse-illuminated algae depends primarily on direct water biophotolysis, where water oxidation at the donor side of photosystem II (PSII) provides electrons for the reduction of protons by H<sub>2</sub>ase downstream of photosystem I. This occurs exclusively in the absence of CO<sub>2</sub> fixation, while with the activation of the CBB cycle by longer (8 s) light pulses the H<sub>2</sub> photoproduction ceases and instead a slow overall H<sub>2</sub> uptake is observed. We also demonstrate that the loss of PSII activity in DCMU-treated algae or in PSII-deficient mutant cells can be partly compensated for by the indirect (PSII-independent) H<sub>2</sub> photoproduction pathway, but only for a short (<1 h) period. Thus, PSII activity is indispensable for a sustained process, where it is responsible for more than 92% of the final H<sub>2</sub> yield.

green algae | hydrogen production | water splitting | carbon dioxide | hydrogenase

Many species of green algae have [FeFe]-hydrogenases (H<sub>2</sub>ases) (1) that catalyze the reversible reduction of protons to molecular hydrogen:



Since [FeFe]-H<sub>2</sub>ases are extremely O<sub>2</sub>-sensitive (2), reaction 1 typically proceeds under anoxic conditions. With respect to H<sub>2</sub> metabolism, *Chlamydomonas reinhardtii* is the most studied alga. This alga possesses two [FeFe]-H<sub>2</sub>ases in the chloroplast, HYDA1 and HYDA2 (3, 4). In the light, they accept electrons from photosynthetically reduced ferredoxin (FDX1) (5), while in the dark electrons come from the activity of pyruvate ferredoxin oxidoreductase (PFR1) (6). PFR1 catalyzes the oxidation of pyruvate to acetyl-CoA, and its activity is linked to H<sub>2</sub>ase via FDX1 (7). Since [FeFe]-H<sub>2</sub>ases interact with the photosynthetic electron transport chain at the level of ferredoxin, they may accept electrons originating both from water oxidation via the photosystem II (PSII)-dependent pathway (“direct water biophotolysis”) and from the degradation of organic substrates via a PSII-independent mechanism (“indirect water biophotolysis” or “indirect pathway”) (8). In the latter case, the reductants are supplied to the plastoquinone (PQ) pool by the type II NADPH dehydrogenase (NDA2), thus bypassing PSII (9, 10).

The release of H<sub>2</sub> leads to a loss of metabolic energy. In healthy, actively growing *C. reinhardtii* cultures, H<sub>2</sub> production is therefore only a temporal phenomenon observed during dark anoxia and upon subsequent onset of illumination (11). In contrast to dark fermentation, H<sub>2</sub> photoproduction is a very efficient process that proceeds for only a short period of time (from a few seconds to a few minutes). Two theories have been developed to explain the short duration. The first is based on the oxygen sensitivity of H<sub>2</sub>ases (12, 13). In the light, algae accumulate O<sub>2</sub> that is produced by water oxidation at PSII (14). As a result, H<sub>2</sub> photoproduction may cease over time (14, 15), and the duration of this process is reported to shorten with increased light intensity (16). Because of the negative correlation between the rates of H<sub>2</sub> photoproduction and O<sub>2</sub> evolution, the inhibition of H<sub>2</sub>ases by O<sub>2</sub> is frequently quoted as the primary reason for the rapid loss in H<sub>2</sub> photoproduction after the onset of illumination (17).

Alternatively, the loss in the H<sub>2</sub> photoproduction efficiency during illumination could be explained by the light-induced induction of competitive pathways, which may drain reducing equivalents away from the [FeFe]-H<sub>2</sub>ase enzyme (18, 19). Candidates for this role are the Mehler-like reaction driven by flavodiiron proteins (FDPs) (15, 20, 21) and the Calvin–Benson–Bassham (CBB) cycle (22). Compelling evidence for the competition between these two

## Significance

Photosynthetic H<sub>2</sub> production in the green alga *Chlamydomonas reinhardtii* is catalyzed by O<sub>2</sub>-sensitive [FeFe]-hydrogenases, which accept electrons from photosynthetically reduced ferredoxin and reduce protons to H<sub>2</sub>. Since the process occurs downstream of photosystem I, the contribution of photosystem II (PSII) in H<sub>2</sub> photoproduction has long been a subject of debate. Indeed, water oxidation by PSII results in O<sub>2</sub> accumulation in chloroplasts, which inhibits H<sub>2</sub> evolution. Therefore, clear evidence for direct water biophotolysis resulting in simultaneous H<sub>2</sub> and O<sub>2</sub> releases in algae has never been presented. This paper demonstrates that sustained H<sub>2</sub> photoproduction in *C. reinhardtii* is directly linked to PSII-dependent water oxidation and brings insights into regulation of PSII activity and H<sub>2</sub> production by CO<sub>2</sub>/HCO<sub>3</sub><sup>−</sup> under microoxic conditions.

Author contributions: S.K., D.S., J.M., and Y.A. designed research; S.K., V.N., D.S., and M.J. performed research; S.K., V.N., D.S., and M.J. analyzed data; S.K., D.S., J.M., and Y.A. wrote the paper; and S.K. and Y.A. conceived the research.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

<sup>1</sup>To whom correspondence may be addressed. Email: serkos@utu.fi, johannes.messinger@kemi.uu.se, or allahve@utu.fi.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2009210117/-DCSupplemental>.

First published November 9, 2020.

pathways and H<sub>2</sub> production has been accumulated in recent studies (23–25). As CO<sub>2</sub> fixation provides the strongest sink for photosynthetic reductants, it should play a major role in the cessation of H<sub>2</sub> photoproduction in algae when the CBB cycle is active (19, 22).

For preventing competition between the [FeFe]-H<sub>2</sub>ases and the CBB cycle, we recently devised a pulse-illumination protocol that allows H<sub>2</sub> production in nutrient-replete algal cultures for up to 3 d (23). To achieve this, we specifically selected the duration of light pulses in the light/dark sequence to avoid activation of the CBB cycle, thus allowing for the redirection of photosynthetic electrons toward the [FeFe]-H<sub>2</sub>ases. Typically, a train of 1- to 6-s light pulses interrupted by 9-s dark periods is sufficient for sustained H<sub>2</sub> photoproduction in *C. reinhardtii* cultures (23, 25). Our protocol thus differs from earlier pulse-illumination approaches that aimed at preventing the accumulation of O<sub>2</sub> in the cultures (26).



While we could demonstrate competition of [FeFe]-H<sub>2</sub>ase with FDPs (25), the origin of reductants for H<sub>2</sub> photoproduction in the pulse-illuminated algae remained unclear. The relatively high efficiency of the process suggests the involvement of water oxidation by PSII, and consequently the simultaneous production of H<sub>2</sub> and O<sub>2</sub>. Although widely proposed in the current literature (8, 24), the presence of the direct water biophotolysis in H<sub>2</sub>-producing green algae has not yet been proven by direct experimental data.

In the present study, we provide clear evidence for the presence of PSII-dependent oxidation of <sup>18</sup>O-labeled water (H<sub>2</sub><sup>18</sup>O) with concomitant evolution of <sup>16</sup>O<sub>2</sub> and <sup>16,18</sup>O<sub>2</sub> during H<sub>2</sub> photoproduction in the pulse-illuminated green alga *C. reinhardtii* under anoxic conditions. O<sub>2</sub> evolution is balanced by light-dependent and light-independent respiration that sustains the anoxic condition. We also demonstrate that the loss of PSII activity in algae can be partly compensated by the PSII-independent H<sub>2</sub> photoproduction pathway. Nevertheless, the activity of PSII is indispensable for the sustained process, where it contributes to more than 92% of the final H<sub>2</sub> yield.


## Results

**Net H<sub>2</sub> Photoproduction, Water Oxidation, and O<sub>2</sub> Exchange.** Employing membrane-inlet mass spectrometry (MIMS), we confirmed the induction of sustained H<sub>2</sub> photoproduction in anoxic *C. reinhardtii* cultures by a train of 1-s light pulses interrupted by 9-s dark periods (Fig. 1A). The signal exhibited a typical sawtooth wave with the H<sub>2</sub> production transients during pulse illumination and a strong H<sub>2</sub> uptake on the shift to darkness, which is in line with previous data (23). The H<sub>2</sub> uptake is the result of H<sub>2</sub> consumption by the MIMS setup and the reversible action of the H<sub>2</sub>ase (as demonstrated in *SI Appendix, Fig. S1*). The response of anoxic algae to the train of light pulses was strain-specific. While in the cell wall-deficient mutant the first H<sub>2</sub> peak appeared already on the first 1-s flash (Fig. 1C, CC-4533), the wild-type CC-124 strain required some time before demonstrating the pronounced H<sub>2</sub> photoproduction yield (Fig. 1A).

The application of <sup>18</sup>O-labeled water (H<sub>2</sub><sup>18</sup>O) to the algae allowed simultaneous monitoring of PSII activity via both nonlabeled and <sup>18</sup>O-labeled O<sub>2</sub> isotopologues (27). Thus, as shown in Fig. 1B, the injection of H<sub>2</sub><sup>18</sup>O into the MIMS cell filled with the algal suspension led to the detection of O<sub>2</sub> evolution at *m/z* 32 (for ambient <sup>16</sup>O<sub>2</sub>) and *m/z* 34 (for singly labeled <sup>16,18</sup>O<sub>2</sub>) signals as a result of the water-oxidizing activity of PSII centers during pulse illumination:




The *m/z* 34 signal showed O<sub>2</sub> evolution at the first 1-s light pulse, and algal cultures reached maximum O<sub>2</sub> production yields after



**Fig. 1.** H<sub>2</sub> photoproduction (A) and O<sub>2</sub> exchange (B) in pulse-illuminated algae. Anoxia was achieved within 3 min after placement of aerobic algae in the O<sub>2</sub>-consuming MIMS chamber under complete darkness. The total (a + b) chlorophyll (Chl) content in the samples was around 9 mg·L<sup>-1</sup>. The photochemical reactions were initiated by a train of 1-s light pulses interrupted by 9-s dark periods. (C) The behavior of the *m/z* 2 and *m/z* 34 signals at better time resolution for the cell wall-deficient strain (CC-4533) in the beginning of pulse illumination and for the wild-type strain (CC-124) in the middle of H<sub>2</sub> photoproduction. One representative result out of three repeats is presented. Additional experimental data can be found in *SI Appendix, Fig. S9*.

10 to 60 pulses (Fig. 1B). Then, the signal slowly declined but kept the sawtooth shape until the end of the pulse-illumination period. The decline in the net O<sub>2</sub> evolution coincided well with the simultaneous increase in the H<sub>2</sub> photoproduction yield (Fig. 1A). Surprisingly, <sup>16</sup>O<sub>2</sub> production (*m/z* 32) of midlog-phase *C. reinhardtii* cultures was significantly delayed by strong light-dependent uptake at the beginning of pulse illumination (Fig. 1B). Since ambient <sup>16</sup>O<sub>2</sub> dominates in the MIMS chamber and the level of <sup>16,18</sup>O<sub>2</sub> is negligible in the beginning of pulse illumination, the pronounced uptake of <sup>16</sup>O<sub>2</sub> in the first few cycles of pulse illumination indicates the domination of light-dependent respiration over water oxidation during this period. On the contrary, the mature (close to the stationary phase) algae did not show any signs of the light-dependent <sup>16</sup>O<sub>2</sub> uptake (*SI Appendix, Fig. S2*). In the latter case, the *m/z* 32 signal simply repeated the shape of the *m/z* 34 curve. Thus, light-dependent O<sub>2</sub> uptake is not caused by the self-shading effect and activation of respiration in dense cultures. This phenomenon is most likely linked to O<sub>2</sub> photoreduction by FDPs that operate during the dark-to-light (or low-light to high-light) transients under oxic and microoxic conditions (25, 28).

The rise of the H<sub>2</sub> (*m/z* 2) signal occurred almost immediately upon the firing of each light pulse. This was followed by a rise in the O<sub>2</sub> (*m/z* 34) signal with ~3-s delay (Fig. 1C). As a result, the H<sub>2</sub> production wave showed maxima at almost the minimums of the *m/z* 34 signal. Since the diffusivity of H<sub>2</sub> in the polydimethylsiloxane membrane used in the MIMS cell (applied to separate the liquid phase of the sample from the high vacuum of the mass spectrometer) is approximately four times faster than for O<sub>2</sub> (29), this



**Fig. 2.** The effect of DCMU and the *psbA* deletion (FuD7) on  $H_2$  (A),  $^{16,18}O_2$  (B), and  $^{16}O_2$  (C) yields in pulse-illuminated algae. The experimental conditions were the same as in Fig. 1, except that the total Chl content was increased to  $\sim 20 \text{ mg}\cdot\text{L}^{-1}$  (to distinguish the difference between  $O_2$  production and  $O_2$  consumption in the control and DCMU-treated samples). DCMU was introduced to aerobic algae at a final concentration of  $30 \mu\text{M}$ . Curves are the mean of two to four independent experiments. The  $H_2$  photoproduction of the FuD7 mutant is a result of one measurement with MIMS but was repeated nine times with a  $H_2$  sensor, producing similar results (*SI Appendix*, Fig. S3). Additional experimental data can also be found in *SI Appendix*, Fig. S10.

behavior of  $H_2$  and  $O_2$  signals is not a physiological response but rather a reflection of the detection method. Therefore, we suggest simultaneous photoproduction of these two gases in *C. reinhardtii* cells.


**The Role of PSII-Dependent and PSII-Independent Pathways.** The addition of  $30 \mu\text{M}$  of DCMU [3-(3,4-dichlorophenyl)-1,1-dimethylurea], a specific inhibitor that binds to the  $Q_B$ -pocket of PSII and blocks electron transport from PSII to the PQ pool, did not affect  $H_2$  photoproduction much during the first 20 min of pulse illumination (Fig. 2A). The  $m/z$  2 signal of the DCMU-treated algae (CC-124 + DCMU) was only slightly reduced as compared to the control (CC-124). This response of pulse-illuminated algae to DCMU was confirmed using an  $H_2$  electrode (*SI Appendix*, Fig. S3). In contrast, the presence of DCMU significantly reduced the  $^{16,18}O_2$ -evolving activity in algae at the start of pulse illumination (Fig. 2B), but inhibition was not complete. Thus, as seen in Fig. 2B, *Inset* a residual water-oxidizing activity ( $<1.5\%$  of the control) was observed in the DCMU-treated samples. At the same time, DCMU-treated algae demonstrated no light-induced evolution of nonlabeled  $^{16}O_2$  ( $m/z$  32) but a strong light-dependent  $^{16}O_2$  uptake (Fig. 2C).

Since the  $m/z$  34 signal represents the net  $^{16,18}O_2$  exchange with a significant expected share of light-induced  $O_2$  uptake (as seen from the  $m/z$  32 signal in Fig. 2C), the contribution of the residual PSII activity in the  $H_2$  photoproduction yield of the DCMU-treated samples remained unclear. Therefore, in the next

experiment, a train of light pulses was applied to the PSII-deficient (FuD7) mutant. Due the absence of fully assembled PSII centers in the thylakoid membrane (30), this strain completely lacks water-oxidizing activity (*SI Appendix*, Fig. S4A). Nevertheless, FuD7 still produced  $H_2$ , albeit at the decreased rate (Fig. 2A and *SI Appendix*, Fig. S3). Similar to DCMU-treated algae, the PSII-deficient mutant showed a strong light-dependent  $^{16}O_2$  uptake (*SI Appendix*, Fig. S4B).

The short-term experiments undertaken with the DCMU-treated wild-type strain and the FuD7 mutant revealed functioning of the PSII-independent  $H_2$  photoproduction pathway when PSII was not available. Therefore, it was worth checking whether this pathway could sustain  $H_2$  photoproduction in the long-term process. It was subsequently observed that the PSII-independent pathway could not substitute the PSII-dependent pathway in the long-term process (Fig. 3A). The  $H_2$  photoproduction yields both in the FuD7 mutant and in the DCMU-treated wild-type strain began decreasing during the first hour (Fig. 3A, *Inset*). This result is in good agreement with those of the short-term experiments (*SI Appendix*, Fig. S3). In the CC-124 and CC-5325 strains (Fig. 3A and B), which possess full PSII activity, DCMU addition stopped  $H_2$  photoproduction after 6 h, while  $H_2$  production of the FuD7 mutant continued, but at a greatly reduced rate. In all cases, final  $H_2$  yields were only slightly higher than in the dark samples.

In contrast to PSII deficiency, the absence of NDA2 (responsible for nonphotochemical reduction of the PQ pool) (9, 10) in *C. reinhardtii* cells was almost completely compensated for by the PSII-dependent pathway. This was particularly evident in the first 4 h of the experiment (Fig. 3B, *Inset*). Here, the input of the PSII-independent pathway was noticeable only by the end of the




**Fig. 3.** Long-term  $H_2$  photoproduction by pulse-illuminated algae. The cultures of the wild-type (CC-124) strain and the PSII-deficient (FuD7) mutant (A), and NDA2-deficient (NDA2) mutant and its wild-type progenitor (CC-5325) (B) were exposed to a train of 1-s light/9-s dark pulses in the absence and in the presence of  $30 \mu\text{M}$  DCMU under an Ar atmosphere. The  $H_2$  production activity of the same strains was also checked under complete darkness. (*Insets*) The same samples in the first 4 h of pulse illumination. Values are mean of 6 to 13 independent replicates ( $\pm$  SD).

experiment, but even in that case the difference was not significant.


**CO<sub>2</sub> Exchange of Algal Cultures Correlates with O<sub>2</sub> Evolution and H<sub>2</sub> Photoproduction.** Previously, we demonstrated that *C. reinhardtii* cells exposed to a train of 1-s light pulses interrupted by 9-s dark periods do not fix CO<sub>2</sub> and as a result do not accumulate biomass (23, 25). These previous experiments were performed at low cell densities (<10 mg total Chl per L) to ensure optimal photosynthetic performance. In the current work, we employed sensitive MIMS methodology at a higher cell density of around 20 mg Chl·L<sup>-1</sup> to improve the resolution of CO<sub>2</sub> analysis. Thus, the culture was observed to release CO<sub>2</sub> during the 1-s light/9-s dark pulse-illumination train (Fig. 4). The ambient (*m/z* 44) CO<sub>2</sub> signal rose within 5 min of engaging the pulse-illumination train and then slowly declined until train cycling was concluded. At this point, the *m/z* 44 signal decayed exponentially to reach the initial state. Along with ambient CO<sub>2</sub> release, *C. reinhardtii* cells also produced <sup>18</sup>O-labeled CO<sub>2</sub> isotopologues: C<sup>16,18</sup>O<sub>2</sub> (*m/z* 46) and C<sup>18,18</sup>O<sub>2</sub> (*m/z* 48) (Fig. 4).

Obviously, the release of all CO<sub>2</sub> isotopologues was light-dependent, but the signals did not fluctuate during pulse illumination (Fig. 4). The typical sawtooth shape appeared only when the duration of the light pulse in the light/dark sequence was extended to 8 s. In this regime, a clear sign of CO<sub>2</sub> fixation appeared 5 min after the start of pulse illumination (Fig. 5A). All three CO<sub>2</sub> signals, *m/z* 44, *m/z* 46, and *m/z* 48, were affected (Fig. 5A and *SI Appendix*, Fig. S5). A similar trend in behavior of *m/z* 44, *m/z* 46, and *m/z* 48 on activation of the CBB cycle indicates that these three signals primarily represent C<sup>16</sup>O<sub>2</sub>, C<sup>16,18</sup>O<sub>2</sub>, and C<sup>18,18</sup>O<sub>2</sub>, but not other possible gases with a similar molecular weight like N<sub>2</sub><sup>16</sup>O and N<sup>16</sup>O<sub>2</sub>. For example, the latter two may appear in *C. reinhardtii* as a result of NO detoxification (31).

As expected, prolonging the light pulse in the light/dark sequence led to enhanced O<sub>2</sub> release, but the burst amounts of O<sub>2</sub> were produced only in the 8-s/9-s regime where the CBB cycle was activated (Fig. 5B; notice the different scale for the 8-s/9-s pulse protocol). By contrast, *C. reinhardtii* cells exposed to the 5-s light/9-s dark protocol yielded almost the same amount of O<sub>2</sub> as the algae exposed to 3-s/9-s pulse illumination, presumably due to a more pronounced O<sub>2</sub> consumption during the dark phases of the same duration. The enhanced O<sub>2</sub> consumption, however, did not correlate with CO<sub>2</sub> release (Fig. 5A). Interestingly, 3-s/9-s and 5-s/9-s pulse-illumination protocols yielded almost the same amounts of H<sub>2</sub> (Fig. 5C). By contrast, the 8-s/9-s protocol released H<sub>2</sub> only in the beginning of pulse illumination. Thus, the activation of the CBB cycle immediately led to H<sub>2</sub> uptake (Fig. 5C). As seen in Fig. 5A and C, the H<sub>2</sub> trace (*m/z* 2) starts to



**Fig. 4.** The release of ambient (*m/z* 44) CO<sub>2</sub> and <sup>18</sup>O-labeled (*m/z* 46, *m/z* 48) CO<sub>2</sub> isotopologues during pulse illumination. The experimental conditions were similar to those presented in Fig. 1. The total Chl content was around 20 mg·L<sup>-1</sup>. A representative experiment is presented. Additional experimental data can be found in *SI Appendix*, Fig. S11.



**Fig. 5.** The effect of light pulse duration in the pulse-illumination sequence on CO<sub>2</sub> exchange (A), O<sub>2</sub> evolution (B), and H<sub>2</sub> photoproduction (C) of algal cultures. In A, only ambient (*m/z* 44) CO<sub>2</sub> signals are presented. For behavior of the *m/z* 46 and *m/z* 48 CO<sub>2</sub> signals see *SI Appendix*, Fig. S5. In B, *m/z* 34 O<sub>2</sub> signal for the 8-s/9-s illumination regime is also shown at low resolution (V) scale. Additional experimental data can be found in *SI Appendix*, Fig. S12.

decline at exactly the time when CO<sub>2</sub> signals establish a pronounced sawtooth wave.

## Discussion

**Clear Evidence for Simultaneous Water Oxidation and H<sub>2</sub> Photoproduction in Green Algae.** The role of PSII in H<sub>2</sub> photoproduction in green algae has long been a subject of debate (32). The PSII-dependent net O<sub>2</sub> release of dark-adapted anoxic algae, when exposed to light, appears only after a period of efficient H<sub>2</sub> production (14). Therefore, direct involvement of PSII in H<sub>2</sub> photoproduction is not obvious, especially because DCMU does not completely inhibit H<sub>2</sub> production (33, 34). In sulfur (S)-deprived *C. reinhardtii* cells, where the recovery of PSII reaction centers is affected by stress (35), the involvement of residual PSII activity in the H<sub>2</sub> photoproduction yield has been proposed based on inhibitory analysis, electron paramagnetic resonance spectroscopy, and fluorescence data (36–38). Nevertheless, Clark-type O<sub>2</sub> electrodes used directly in algal suspensions have not indicated water-splitting activity in S-deprived cells (36), either due to low sensitivity of the technique (39) or due to the absence of net O<sub>2</sub> release in the actively respiring cells (40). The situation improved with application of a high-sensitive MIMS approach (15, 22). The publications showed that the release of O<sub>2</sub> in dark-adapted algae occurs a few seconds after appearance of H<sub>2</sub> in the system, suggesting direct involvement of PSII in H<sub>2</sub> photoproduction.

As shown in Fig. 1, the application of the 1-s light/9-s dark protocol to anoxic algae results in sustained H<sub>2</sub> photoproduction and also leads to the simultaneous evolution of O<sub>2</sub> (detected at *m/z* 32 and *m/z* 34). The appearance of both nonlabeled and <sup>18</sup>O-labeled isotopologues of O<sub>2</sub> upon enrichment with H<sub>2</sub><sup>18</sup>O is well known for monitoring PSII-driven water oxidation reaction (41). The appearance of photosynthetically produced O<sub>2</sub> in algal cultures should indicate a shift in the environment from anoxic to

microoxic. However, the situation is not so simple. This is for two reasons: First, light activates strong O<sub>2</sub> uptake, which is observed in midlog-phase cultures (Fig. 1B, *m/z* 32 signal) in the FuD7 mutant (*SI Appendix*, Fig. S4B) and in the DCMU-treated algae (Fig. 2C, +DCMU trace); second, the release of photosynthetically produced O<sub>2</sub> occurs simultaneously with mitochondrial respiration (which was subtracted from all signals during data processing; *SI Appendix*, Fig. S6A and B). These respiratory processes under normal atmospheric pressure balance photosynthetically produced O<sub>2</sub> to a level undetectable by polarographic techniques (Clark-type O<sub>2</sub> electrode) (23). Nevertheless, the competition of H<sub>2</sub> photoproduction with FDP-driven O<sub>2</sub> photoreduction under such low O<sub>2</sub> levels is still highly possible (25), especially in the beginning of the experiment where extracellular O<sub>2</sub> concentrations are at around 0.1 to 0.5 μM. In this context, the gradually increasing H<sub>2</sub> photoproduction rate (Fig. 1A) may reflect a release of FDPs of such competition. Indeed, the level of extracellular O<sub>2</sub> does decrease in the course of the experiment due to its consumption by the instrument and the culture itself (*SI Appendix*, Fig. S6A).

Surprisingly, the cell wall-deficient CC-4533 strain showed the production of H<sub>2</sub> on the first light pulse (Fig. 1C, CC-4533). The rapid response of cells to the establishment of anoxic conditions (~3 min from fully aerobic environment) suggests the expression of the O<sub>2</sub>-sensitive [FeFe]-H<sub>2</sub>ase in *C. reinhardtii* cells at the time of aerobic cultivation, as has been previously proposed (42). In agreement with this suggestion, transcripts of *hydA1* and *hydA2* hydrogenases and the *hyDEF* and *hydG* maturation factors have been found in aerobic algae (43, 44). Since the HYDA1/HYDA2 proteins are also already available in aerobic *C. reinhardtii* (23), the delay in net H<sub>2</sub> evolution of the CC-124 strain (Fig. 1A) is most likely caused by the limitation of H<sub>2</sub> diffusion through the cell wall. Assuming the reversible nature of the H<sub>2</sub>ase-driven reaction (45, 46), enhanced intracellular levels of H<sub>2</sub> may increase H<sub>2</sub> consumption during dark periods and decrease the final H<sub>2</sub> yield. In accordance with this suggestion, prolongation of the light pulse duration to above 3 s in the pulse-illumination sequence demonstrates the early appearance of the H<sub>2</sub> signal in the same strain (Fig. 5C). Similar to the CC-4533 strain, the periods of H<sub>2</sub> photoproduction in the CC-124 algae coincide well with O<sub>2</sub> evolution (Fig. 1C, CC-124). Simultaneous production of H<sub>2</sub> and O<sub>2</sub> in the light indicates the direct involvement of electrons from water oxidation by PSII in H<sub>2</sub> photoproduction.


**PSII Plays a Major Role in Sustained H<sub>2</sub> Photoproduction.** Reductants for the H<sub>2</sub> photoproduction of *C. reinhardtii* are suggested to be supplied not only by PSII but also by the degradation of stored organic substrates, primarily starch (47). In a similar manner to the PSII-dependent process, the indirect pathway requires active PSI for the donation of electrons to H<sub>2</sub>ase. However, by contrast, the indirect process results in CO<sub>2</sub> release with a maximum molar stoichiometry of 1 CO<sub>2</sub> per 2 H<sub>2</sub>. It has previously been shown that contribution of the indirect pathway to the H<sub>2</sub> photoproduction yield might be significant in S-deprived cells, where the PSII activity is substantially affected by the stress (48).

The results obtained with DCMU-treated algae and the PSII-deficient mutant (Fig. 2A and *SI Appendix*, Fig. S3) clearly show that the PSII-independent H<sub>2</sub> production pathway can partly compensate for the loss of the PSII-dependent H<sub>2</sub> production pathway toward H<sub>2</sub> photoproduction of pulse-illuminated cells. The indirect process, however, could not sustain H<sub>2</sub> production for longer than 6 h (Fig. 3A), and the specific rate of H<sub>2</sub> production had already started to decline during the first hour (Fig. 3A, *Inset*). On the contrary, the elimination of NDA2 in *C. reinhardtii*, which is the main player of the PSII-independent pathway (10), shows almost no effect on H<sub>2</sub> photoproduction under pulse-illumination conditions (Fig. 3B). It has been previously demonstrated that NDA2 deficiency decreases the H<sub>2</sub> photoproduction activity of

S-deprived algae (9, 10). However, S deprivation leads to a significant accumulation of starch reserves in cells during the photosynthetic stage (36). Therefore, it is not surprising that S deprivation increases the contribution of the NDA2-dependent pathway in the H<sub>2</sub> photoproduction yield, especially when the activity of PSII is limited or fully absent (49). Thus, S-deprivation data do not contradict our conclusions, since in actively growing algae, which are used in this study, the starch reserves are limited. In the presence of DCMU, which blocks the electron flow from PSII, the *nda2* deletion mutant produces almost the same amount of H<sub>2</sub> as algae placed in complete darkness. The slightly higher H<sub>2</sub> yield of the DCMU-treated *nda2* deletion mutant might be attributed to 1) the light activation of fermentation, 2) incomplete inhibition of electron flow by DCMU (Fig. 2B), or 3) the presence of other player(s) in the PSII-independent pathway. If any, their input in the total H<sub>2</sub> yield is negligible and limited to the beginning of pulse-illumination (Fig. 3B, *Inset*). A full compensation of algal NDA2 elimination by PSII confirms that H<sub>2</sub> photoproduction during 1-s illumination periods proceeds via the most efficient mechanism of direct water biophotolysis. This conclusion is in agreement with fluorescence data showing that PSII activity in dark-adapted *C. reinhardtii* cells is linearly related to the hydrogenase capacity observed during the first seconds of illumination (50).

In the 1-s/9-s pulse-illumination regime, algae spend most of the time in darkness, where fermentation plays a major role. According to the data presented in Fig. 3A, up to 4% of the H<sub>2</sub> yield in the CC-124 strain is supported by dark fermentation, while 96% is supported by PSII. In the presence of DCMU, the PSII-independent pathway doubles the H<sub>2</sub> yield in pulse-illuminated algae as compared to the dark samples (Fig. 3A). Thus, even if both pathways (PSII-independent and dark) operate in the CC-124 cells during pulse illumination, PSII is still responsible for about 92% of the final H<sub>2</sub> yield. Very similar results (93% of the PSII input) were obtained for the *nda2* deletion mutant, where the contribution of fermentation was close to 7% (Fig. 3B). However, the NDA2 deficiency may also enhance fermentative H<sub>2</sub> production in the mutant. Taking into account all data, we conclude that H<sub>2</sub> production during 1-s light pulses is driven primarily or even exclusively by the PSII-dependent pathway.



**Efficient H<sub>2</sub> Photoproduction in *C. reinhardtii* Occurs before Activation of the CBB Cycle.** The light-dependent water oxidation performed by PSII results in O<sub>2</sub> evolution with a simultaneous release of protons (H<sup>+</sup>) into the thylakoid lumen. In addition, protons are pumped into the lumen by the PQ/plastoquinone cycle driven by PSII and the cytochrome *b<sub>6</sub>f* complex. The accumulation of H<sup>+</sup> in the lumen builds up the Δ*p*H across the thylakoid membrane and ensures adenosine 5'-triphosphate (ATP) biosynthesis. Simultaneously, a relatively small release of CO<sub>2</sub>, which was first detected by MIMS in PSII membranes isolated from spinach (51), could be observed. This CO<sub>2</sub> is known to be released by PSII and is the result of HCO<sub>3</sub><sup>-</sup> reaction with H<sup>+</sup> (produced during water splitting) followed by the subsequent bicarbonate dehydration:





This reaction facilitates the removal of H<sup>+</sup> from PSII. Moreover, the electron-acceptor side of PSII is known to be another source for CO<sub>2</sub> evolution under intensive light illumination (52, 53).

In *C. reinhardtii*, the bicarbonate dehydration reaction (and in general interconversions of inorganic carbon) can be accelerated by a luminal carbonic anhydrase, CrCAH3 (54–56). However, in whole cells under normal conditions, the light-dependent CO<sub>2</sub> release is barely detectable due to the presence of two processes: 1) the consumption of CO<sub>2</sub> by Rubisco (i.e., by CBB cycle) and 2) mitochondrial respiration. Therefore, it is not surprising that we observed CO<sub>2</sub> release in the pulse-illuminated algae, when the CBB cycle is not active (Fig. 4) and mitochondrial respiration

is restricted by an extremely low level of O<sub>2</sub> in the microoxic environment (because of the efficient O<sub>2</sub> consumption by the MIMS setup; *SI Appendix*, Fig. S6A). As shown in Fig. 4, the light-induced rise of *m/z* 44, *m/z* 46, and *m/z* 48 signals occurs simultaneously with O<sub>2</sub> evolution and H<sub>2</sub> photoproduction (Fig. 1). The appearance of *m/z* 46 and *m/z* 48 signals together with ambient CO<sub>2</sub> (*m/z* 44) indicate a CO<sub>2</sub>–water interexchange:



Alternatively, the inclusion of labeled O<sub>2</sub>, which appears as a product of water oxidation (reaction 3), in the organic substrate and its immediate degradation should be expected:



The probability of reaction 7, and especially reaction 8, is much lower than reactions 5 and 6, unless they proceed at the same site as reaction 3. Since the reactions 7 and 8 are expected to occur in the stroma of chloroplasts, they are unlikely to satisfy kinetics of light-dependent release of CO<sub>2</sub> isotopologues during pulse illumination (Fig. 4).

Although the observed CO<sub>2</sub> evolution could be potentially caused by activation of substrate degradation in the light (photofermentation), the positive correlation between O<sub>2</sub> evolution (*SI Appendix*, Fig. S2) and CO<sub>2</sub> release (Fig. 4), and relatively fast recovery of all three CO<sub>2</sub> signals after conclusion of pulse illumination (Fig. 4) strongly suggest direct involvement of PSII in CO<sub>2</sub> evolution. Since the equilibration rate between labeled water and CO<sub>2</sub> occurs with half-times of  $t_{1/2} \approx 30$  s or less (55, 57), any donor-side CO<sub>2</sub> formation or other acidification of the lumen would produce CO<sub>2</sub> isotopologues reflecting the H<sub>2</sub><sup>18</sup>O enrichment of the water (given that measurements were initiated about 5 min after H<sub>2</sub><sup>18</sup>O addition). By contrast, CO<sub>2</sub> release at the acceptor side of PSII, where bicarbonate is bound to the non-heme iron and equilibrates slowly with bulk water, would be predominantly C<sup>16</sup>O<sub>2</sub>. Thus, the different time dependence of C<sup>16</sup>O<sub>2</sub> versus C<sup>16,18</sup>O<sub>2</sub> and C<sup>18</sup>O<sub>2</sub> (Fig. 4, *m/z* 46 and *m/z* 48 signals) suggests that light-induced CO<sub>2</sub> release may originate from both the acceptor and donor sides of PSII, as well as the acidification of the lumen. This is further supported by estimating the total amount of released CO<sub>2</sub>: If just the acceptor side would contribute, we should expect around 2 nmol of CO<sub>2</sub> released per mg Chl [assuming 1.83 mmol of Q<sub>A</sub> per mol of total Chl in photomixotrophic *C. reinhardtii* cells (58)]. Instead, formation of around 4 to 8 nmol CO<sub>2</sub> per mg Chl is typically observed in algal cultures (Figs. 4 and 5 and *SI Appendix*, Fig. S11). These data show PSII-associated CO<sub>2</sub> formation in intact cells, which supports the idea that it has an important regulatory and protective function for PSII *in vivo*.

The prolongation of light pulses to 8 s in the pulse-illumination sequence gradually activates the CBB cycle, leading to a pronounced CO<sub>2</sub> uptake after 5 min of pulse illumination (Fig. 5A and *SI Appendix*, Fig. S5). As shown in Fig. 5, the activation of the CBB cycle immediately results in the burst release of O<sub>2</sub> (Fig. 5B) and inhibition of H<sub>2</sub> evolution followed by H<sub>2</sub> uptake (Fig. 5C). These data support the previously proposed hypothesis that the primary loss of H<sub>2</sub> evolution activity in *C. reinhardtii* cells is caused by competition between ferredoxin-NADP<sup>+</sup> oxidoreductase and H<sub>2</sub>ases for reduced ferredoxin, rather than by the sensitivity of hydrogenase to oxygen (19, 22). Besides the activation of the CBB cycle, increasing O<sub>2</sub> levels inside the chloroplast can also enhance

the flow of photosynthetic electrons toward FDPs. Such competition indeed becomes more pronounced with prolongation of the light pulse in the sequence (25). In long-term experiments, prolongation of the light pulse results in biomass accumulation (25). These experiments show that the pulse-illumination approach allows a fine tuning of algal metabolism between microoxic H<sub>2</sub> photoproduction and aerobic CO<sub>2</sub> fixation, thus enabling PSII-dependent water-oxidation activity to be maintained at the desired level.

## Summary

The illumination of anoxic *C. reinhardtii* cultures with a train of short light pulses interrupted by longer dark periods demonstrates that efficient H<sub>2</sub> photoproduction in algal cells occurs exclusively in the absence of photosynthetic CO<sub>2</sub> fixation, thus when the CBB cycle is not active. The cells produce H<sub>2</sub> via the most efficient mechanism of direct water biophotolysis, where water oxidation at the donor side of PSII provides the electrons for reduction of protons by the [FeFe]-H<sub>2</sub>ase enzyme(s) downstream of PSI. Thus, the two reactions occur simultaneously. However, under normal conditions O<sub>2</sub> is not released by the cells due to its consumption by respiration. During short periods of light illumination, the H<sub>2</sub>ase activity supports a linear photosynthetic electron flow from PSII to PSI, promoting proton translocation across the thylakoid membrane and ensuring efficient water oxidation by PSII. Thus, [FeFe]-H<sub>2</sub>ase, in concert with PSII, creates favorable conditions for O<sub>2</sub> accumulation and ATP biosynthesis to levels sufficient for the activation of mitochondrial respiration and the CBB cycle. If the duration of light pulses is not sufficient to fulfill the above condition, the algae continue to produce H<sub>2</sub>. The long-term production of H<sub>2</sub> in the absence of CO<sub>2</sub> fixation suggests the central role of [FeFe]-H<sub>2</sub>ase in supporting algal photosynthesis and cell fitness under anoxic conditions.

## Materials and Methods

The wild-type *C. reinhardtii* strain CC-124, cell wall-deficient strains CC-4533 and CC-5325, *psbA* deletion (FuD7) mutant CC-4147, and the *nda2* deletion mutant (LMJ.RY0402.257129) were obtained from the Chlamydomonas Resource Center at the University of Minnesota, St. Paul, MN. The *nda2* deletion is characterized as described in *SI Appendix*, Fig. S7. All cultures were maintained, grown, and checked for H<sub>2</sub> photoproduction activity under pulse illumination with H<sub>2</sub> and O<sub>2</sub> microsensors (H2-NP and OX-NP; Unisense A/S) as described in detail by Kosourov et al. (23). Conditions for the long-term experiments were reported in the same publication. All experiments were performed in TAP medium with unstressed, actively growing photomixotrophic algae, which were pipetted into the MIMS chamber or vials just before measurements.

The gas exchange in the suspension of algal cells was studied by time-resolved MIMS setup (*SI Appendix*, Fig. S8) as described previously (59, 60). Briefly, the setup consisted of an isotope ratio mass spectrometer (Delta V Plus; Thermo Fischer Scientific), a cooling trap (−78 °C; dry ice + EtOH), and an in-house-built gas-tight membrane-inlet chamber with 200-μL working volume. Before the measurements, H<sub>2</sub><sup>18</sup>O (97%; Larodan Fine Chemicals AB) was added to the MIMS chamber to a final enrichment of 4%. Analysis of O<sub>2</sub> reactions (evolution/consumption) was based on the *m/z* 32 (<sup>16</sup>O<sub>2</sub>) and *m/z* 34 (<sup>16,18</sup>O<sub>2</sub>) signals, with Faraday cup amplification of 3 × 10<sup>8</sup> and 1 × 10<sup>11</sup>, respectively. The signal *m/z* 36 (<sup>18,18</sup>O<sub>2</sub>) was not considered due to its low evolution at quite low H<sub>2</sub><sup>18</sup>O-enrichment and contamination of this signal by the presence of <sup>36</sup>Ar. Analysis of CO<sub>2</sub> reactions was based on simultaneous monitoring of *m/z* 44 (<sup>12</sup>C<sup>16</sup>O<sub>2</sub>), *m/z* 46 (<sup>12</sup>C<sup>16,18</sup>O<sub>2</sub>), and *m/z* 48 (<sup>12</sup>C<sup>18</sup>O<sub>2</sub>) signals with cup amplification of 1 × 10<sup>9</sup>, 1 × 10<sup>11</sup>, and 1 × 10<sup>12</sup>, accordingly. The argon signal (*m/z* 40) with cup amplification of 1 × 10<sup>11</sup> was used as a control. H<sub>2</sub> photoproduction activity was studied in separate runs by monitoring the *m/z* 2 (<sup>1</sup>H<sub>2</sub>) signal with an amplification of 1 × 10<sup>12</sup>. No H<sub>2</sub><sup>18</sup>O was added for H<sub>2</sub> assays. The white light-emitting diode light pulses (~1,000 μmol photons·m<sup>−2</sup>·s<sup>−1</sup>) were applied using the STM32F103 microcontroller board controlled by the OxyHydrogen software. Before each measurement, microoxic environment inside the MIMS chamber was achieved within ~2 min after sealing by degassing the sample with the vacuum pump of the mass spectrometer. The initial levels of oxygen slightly varied from experiment to experiment from 0.1 to 0.5 μM. The final gas exchange

curves were obtained after correction for the gas consumption by the mass spectrometer during the dark periods in the beginning and in the end of each experiment as demonstrated in *SI Appendix, Fig. S6*. This correction also included the gas diffusion component and consumption/production of the gas by the culture during darkness. The analysis of the collected MIMS signals was done using OriginPro 2019 software. The H<sub>2</sub>, <sup>16</sup>O<sub>2</sub>, and <sup>12</sup>C<sup>16</sup>O<sub>2</sub> yields were determined in moles per milligram of total Chl according to the standard calibration. All other signals were normalized to Chl.

**Data Availability.** All data needed to evaluate conclusions of this paper are present in the paper and in *SI Appendix*.

**ACKNOWLEDGMENTS.** This research was financially supported by the NordForsk NCoE program "NordAqua" (project 82845 to Y.A.), the Academy of Finland (AlgaLEAF, project 322754 to Y.A.), the Kone Foundation (projects 201608799 to S.K. and 201710311 to M.J.), the Nessling Foundation (project 201800011 to V.N.), and Vetenskapsrådet (project 2016-05183 to J.M.).

- V. A. Boichenko, E. Greenbaum, M. Seibert, "Hydrogen production by photosynthetic microorganisms" in *Molecular to Global Photosynthesis: Photoconversion of Solar Energy*, J. Barber, M. D. Archer, Eds. (Imperial College Press, 2004), pp. 397–451.
- S. T. Stripp *et al.*, How oxygen attacks [FeFe] hydrogenases from photosynthetic organisms. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 17331–17336 (2009).
- T. Happe, A. Kaminski, Differential regulation of the Fe-hydrogenase during anaerobic adaptation in the green alga *Chlamydomonas reinhardtii*. *Eur. J. Biochem.* **269**, 1022–1032 (2002).
- M. Forestier *et al.*, Expression of two [Fe]-hydrogenases in *Chlamydomonas reinhardtii* under anaerobic conditions. *Eur. J. Biochem.* **270**, 2750–2758 (2003).
- A. Sawyer, M. Winkler, Evolution of *Chlamydomonas reinhardtii* ferredoxins and their interactions with [FeFe]-hydrogenases. *Photosynth. Res.* **134**, 307–316 (2017).
- A. Dubini, F. Mus, M. Seibert, A. R. Grossman, M. C. Posewitz, Flexibility in anaerobic metabolism as revealed in a mutant of *Chlamydomonas reinhardtii* lacking hydrogenase activity. *J. Biol. Chem.* **284**, 7201–7213 (2009).
- J. Noth, D. Krawietz, A. Hemschemeier, T. Happe, Pyruvate:ferredoxin oxidoreductase is coupled to light-independent hydrogen production in *Chlamydomonas reinhardtii*. *J. Biol. Chem.* **288**, 4368–4377 (2013).
- M. L. Ghirardi *et al.*, "Hydrogen production by water biophotolysis" in *Microbial BioEnergy: Hydrogen Production*, D. Zannoni, R. De Philippis, Eds. (Springer, 2014), pp. 101–135.
- F. Jans *et al.*, A type II NAD(P)H dehydrogenase mediates light-independent plastoquinone reduction in the chloroplast of *Chlamydomonas*. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 20546–20551 (2008).
- E. Mignolet, R. Lecler, B. Ghysels, C. Remacle, F. Franck, Function of the chloroplastic NAD(P)H dehydrogenase Nda2 for H<sub>2</sub> photoproduction in sulphur-deprived *Chlamydomonas reinhardtii*. *J. Biotechnol.* **162**, 81–88 (2012).
- A. A. Tsygankov, "Hydrogen production: Light-driven processes—Green algae" in *Microbial Technologies in Advanced Biofuels Production*, C. P. Hallenbeck, Ed. (Springer, 2012), pp. 29–51.
- T. Happe, J. D. Naber, Isolation, characterization and N-terminal amino acid sequence of hydrogenase from the green alga *Chlamydomonas reinhardtii*. *Eur. J. Biochem.* **214**, 475–481 (1993).
- K. D. Swanson *et al.*, [FeFe]-hydrogenase oxygen inactivation is initiated at the H cluster 2Fe subcluster. *J. Am. Chem. Soc.* **137**, 1809–1816 (2015).
- L. Cournac *et al.*, Limiting steps of hydrogen production in *Chlamydomonas reinhardtii* and *Synechocystis* PCC 6803 as analysed by light-induced gas exchange transients. *Int. J. Hydrogen Energy* **27**, 1229–1237 (2002).
- A. Burlacot *et al.*, Flavodiiron-mediated O<sub>2</sub> photoreduction links H<sub>2</sub> production with CO<sub>2</sub> fixation during the anaerobic induction of photosynthesis. *Plant Physiol.* **177**, 1639–1649 (2018).
- P. J. Aparicio, M. P. Azuara, A. Ballesteros, V. M. Fernández, Effects of light intensity and oxidized nitrogen sources on hydrogen production by *Chlamydomonas reinhardtii*. *Plant Physiol.* **78**, 803–806 (1985).
- M. L. Ghirardi, R. K. Togaasaki, M. Seibert, Oxygen sensitivity of algal H<sub>2</sub> production. *Appl. Biochem. Biotechnol.* **63–65**, 141–151 (1997).
- J. W. Lee, E. Greenbaum, A new oxygen sensitivity and its potential application in photosynthetic H<sub>2</sub> production. *Appl. Biochem. Biotechnol.* **105**, 303–313 (2003).
- D. Godaux, B. Bailleul, N. Berne, P. Cardol, Induction of photosynthetic carbon fixation in anoxia relies on hydrogenase activity and proton-gradient regulation-like1-mediated cyclic electron flow in *Chlamydomonas reinhardtii*. *Plant Physiol.* **168**, 648–658 (2015).
- M. Jokel *et al.*, *Chlamydomonas* flavodiiron proteins facilitate acclimation to anoxia during sulfur deprivation. *Plant Cell Physiol.* **56**, 1598–1607 (2015).
- M. Jokel, X. Johnson, G. Peltier, E.-M. Aro, Y. Allahverdiyeva, Hunting the main player enabling *Chlamydomonas reinhardtii* growth under fluctuating light. *Plant J.* **94**, 822–835 (2018).
- Y. Milrad, S. Schweitzer, Y. Feldman, I. Yacoby, Green algal hydrogenase activity is outcompeted by carbon fixation before inactivation by oxygen takes place. *Plant Physiol.* **177**, 918–926 (2018).
- S. Kosourov, M. Jokel, E.-M. Aro, Y. Allahverdiyeva, A new approach for sustained and efficient H<sub>2</sub> photoproduction by *Chlamydomonas reinhardtii*. *Energy Environ. Sci.* **11**, 1431–1436 (2018).
- V. Nagy *et al.*, Water-splitting-based, sustainable and efficient H<sub>2</sub> production in green algae as achieved by substrate limitation of the Calvin-Benson-Bassham cycle. *Biotechnol. Biofuels* **11**, 69 (2018).
- M. Jokel, V. Nagy, S. Z. Tóth, S. Kosourov, Y. Allahverdiyeva, Elimination of the flavodiiron electron sink facilitates long-term H<sub>2</sub> photoproduction in green algae. *Biotechnol. Biofuels* **12**, 280 (2019).
- J. E. Meuser *et al.*, Phenotypic diversity of hydrogen production in chlorophycean algae reflects distinct anaerobic metabolisms. *J. Biotechnol.* **142**, 21–30 (2009).
- D. Shevela, J. Messinger, Studying the oxidation of water to molecular oxygen in photosynthetic and artificial systems by time-resolved membrane-inlet mass spectrometry. *Front Plant Sci* **4**, 473 (2013).
- Y. Allahverdiyeva, M. Suorsa, M. Tikkanen, E. M. Aro, Photoprotection of photosystems in fluctuating light intensities. *J. Exp. Bot.* **66**, 2427–2436 (2015).
- T. C. Merkel, V. I. Bondar, K. Nagai, B. D. Freeman, I. Pinnau, Gas sorption, diffusion, and permeation in poly(dimethylsiloxane). *J. Polym. Sci. B Polym. Phys.* **38**, 415–434 (2000).
- P. Bennoun *et al.*, Characterization of photosystem II mutants of *Chlamydomonas reinhardtii* lacking the psbA gene. *Plant Mol. Biol.* **6**, 151–160 (1986).
- A. Burlacot, P. Richaud, A. Gosset, Y. Li-Beisson, G. Peltier, Algal photosynthesis converts nitric oxide into nitrous oxide. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 2704–2709 (2020).
- J. Appel, R. Schulz, Hydrogen metabolism in organisms with oxygenic photosynthesis: Hydrogenases as important regulatory devices for a proper redox poising? *J. Photochem. Photobiol. B* **47**, 1–11 (1998).
- J. L. Jurado-Oller, A. Dubini, A. Galván, E. Fernández, D. González-Ballester, Low oxygen levels contribute to improve photohydrogen production in mixotrophic non-stressed *Chlamydomonas* cultures. *Biotechnol. Biofuels* **8**, 149 (2015).
- S. Clowez, D. Godaux, P. Cardol, F. A. Wollman, F. Rappaport, The involvement of hydrogen-producing and ATP-dependent NADPH-consuming pathways in setting the redox poise in the chloroplast of *Chlamydomonas reinhardtii* in anoxia. *J. Biol. Chem.* **290**, 8666–8676 (2015).
- L. Zhang, T. Happe, A. Melis, Biochemical and morphological characterization of sulfur-deprived and H<sub>2</sub>-producing *Chlamydomonas reinhardtii* (green alga). *Planta* **214**, 552–561 (2002).
- S. Kosourov, M. Seibert, M. L. Ghirardi, Effects of extracellular pH on the metabolic pathways in sulfur-deprived, H<sub>2</sub>-producing *Chlamydomonas reinhardtii* cultures. *Plant Cell Physiol.* **44**, 146–155 (2003).
- A. Volgusheva, S. Styring, F. Mamedov, Increased photosystem II stability promotes H<sub>2</sub> production in sulfur-deprived *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 7223–7228 (2013).
- T. K. Antal *et al.*, The dependence of algal H<sub>2</sub> production on Photosystem II and O<sub>2</sub> consumption activities in sulfur-deprived *Chlamydomonas reinhardtii* cells. *Biochim. Biophys. Acta* **1607**, 153–160 (2003).
- D. Shevela, W. P. Schröder, J. Messinger, Liquid-phase measurements of photosynthetic oxygen evolution. *Methods Mol. Biol.* **1770**, 197–211 (2018).
- A. Melis, L. Zhang, M. Forestier, M. L. Ghirardi, M. Seibert, Sustained photobiological hydrogen gas production upon reversible inactivation of oxygen evolution in the green alga *Chlamydomonas reinhardtii*. *Plant Physiol.* **122**, 127–136 (2000).
- D. Shevela, K. Beckmann, J. Clausen, W. Junge, J. Messinger, Membrane-inlet mass spectrometry reveals a high driving force for oxygen production by photosystem II. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 3602–3607 (2011).
- O. Liran *et al.*, Microoxic niches within the thylakoid stroma of air-grown *Chlamydomonas reinhardtii* protect [FeFe]-hydrogenase and support hydrogen production under fully aerobic environment. *Plant Physiol.* **172**, 264–271 (2016).
- A. V. Nguyen *et al.*, Transcriptome for photobiological hydrogen production induced by sulfur deprivation in the green alga *Chlamydomonas reinhardtii*. *Eukaryot. Cell* **7**, 1965–1979 (2008).
- D. González-Ballester *et al.*, RNA-seq analysis of sulfur-deprived *Chlamydomonas* cells reveals aspects of acclimation critical for cell survival. *Plant Cell* **22**, 2058–2084 (2010).
- S. N. Kosourov *et al.*, Maximizing the hydrogen photoproduction yields in *Chlamydomonas reinhardtii* cultures: The effect of the H<sub>2</sub> partial pressure. *Int. J. Hydrogen Energy* **37**, 8850–8858 (2012).
- A. Scoma, A. Hemschemeier, The hydrogen metabolism of sulfur deprived *Chlamydomonas reinhardtii* cells involves hydrogen uptake activities. *Algal Res.* **26**, 341–347 (2017).
- R. P. Gfeller, M. Gibbs, Fermentative metabolism of *Chlamydomonas reinhardtii*: I. Analysis of fermentative products from starch in dark and light. *Plant Physiol.* **75**, 212–218 (1984).
- A. Hemschemeier, S. Fouchard, L. Cournac, G. Peltier, T. Happe, Hydrogen production by *Chlamydomonas reinhardtii*: An elaborate interplay of electron sources and sinks. *Planta* **227**, 397–407 (2008).
- V. Chochois *et al.*, Hydrogen production in *Chlamydomonas*: Photosystem II-dependent and -independent pathways differ in their requirement for starch metabolism. *Plant Physiol.* **151**, 631–640 (2009).
- D. Godaux *et al.*, A novel screening method for hydrogenase-deficient mutants in *Chlamydomonas reinhardtii* based on in vivo chlorophyll fluorescence and photosystem II quantum yield. *Int. J. Hydrogen Energy* **38**, 1826–1836 (2013).

51. S. Koroidov, D. Shevela, T. Shutova, G. Samuelsson, J. Messinger, Mobile hydrogen carbonate acts as proton acceptor in photosynthetic water oxidation. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 6299–6304 (2014).
52. K. Brinkert, S. De Causmaecker, A. Krieger-Liszka, A. Fantuzzi, A. W. Rutherford, Bicarbonate-induced redox tuning in Photosystem II for regulation and protection. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 12144–12149 (2016).
53. D. Shevela, H.-N. Do, A. Fantuzzi, A. W. Rutherford, J. Messinger, Bicarbonate-mediated CO<sub>2</sub> formation on both sides of Photosystem II. *Biochemistry* **59**, 2442–2449 (2020).
54. T. Shutova *et al.*, The photosystem II-associated Cah3 in Chlamydomonas enhances the O<sub>2</sub> evolution rate by proton removal. *EMBO J.* **27**, 782–791 (2008).
55. R. Benlloch *et al.*, Crystal structure and functional characterization of photosystem II-associated carbonic anhydrase CAH3 in Chlamydomonas reinhardtii. *Plant Physiol.* **167**, 950–962 (2015).
56. V. V. Terentyev, A. K. Shukshina, A. V. Shitov, Carbonic anhydrase CAH3 supports the activity of photosystem II under increased pH. *Biochim. Biophys. Acta Bioenerg.* **1860**, 582–590 (2019).
57. J. Clausen, K. Beckmann, W. Junge, J. Messinger, Evidence that bicarbonate is not the substrate in photosynthetic oxygen evolution. *Plant Physiol.* **139**, 1444–1450 (2005).
58. J. E. W. Polle, J. R. Benemann, A. Tanaka, A. Melis, Photosynthetic apparatus organization and function in the wild type and a chlorophyll b-less mutant of Chlamydomonas reinhardtii. Dependence on carbon source. *Planta* **211**, 335–344 (2000).
59. D. Shevela *et al.*, Biogenesis of water splitting by photosystem II during de-etiolation of barley (*Hordeum vulgare* L.). *Plant Cell Environ.* **39**, 1524–1536 (2016).
60. K. Tikhonov, D. Shevela, V. V. Klimov, J. Messinger, Quantification of bound bicarbonate in photosystem II. *Photosynthetica* **56**, 210–216 (2018).