

RESEARCH

Open Access



Tailoring and optimizing fatty acid production by oleaginous yeasts through the systematic exploration of their physiological fitness

Zeynep Efsun Duman-Özdamar^{1,2,4}, Vitor A. P. Martins dos Santos^{1,2,3}, Jeroen Hugenholtz⁴ and Maria Suarez-Diez^{2*}

Abstract

Background: The use of palm oil for our current needs is unsustainable. Replacing palm oil with oils produced by microbes through the conversion of sustainable feedstocks is a promising alternative. However, there are major technical challenges that must be overcome to enable this transition. Foremost among these challenges is the stark increase in lipid accumulation and production of higher content of specific fatty acids. Therefore, there is a need for more in-depth knowledge and systematic exploration of the oil productivity of the oleaginous yeasts. In this study, we cultivated *Cutaneotrichosporon oleaginosus* and *Yarrowia lipolytica* at various C/N ratios and temperatures in a defined medium with glycerol as carbon source and urea as nitrogen source. We ascertained the synergistic effect between various C/N ratios of a defined medium at different temperatures with Response Surface Methodology (RSM) and explored the variation in fatty acid composition through Principal Component Analysis.

Results: By applying RSM, we determined a temperature of 30 °C and a C/N ratio of 175 g/g to enable maximal oil production by *C. oleaginosus* and a temperature of 21 °C and a C/N ratio of 140 g/g for *Y. lipolytica*. We increased production by 71% and 66% respectively for each yeast compared to the average lipid accumulation in all tested conditions. Modulating temperature enabled us to steer the fatty acid compositions. Accordingly, switching from higher temperature to lower cultivation temperature shifted the production of oils from more saturated to unsaturated by 14% in *C. oleaginosus* and 31% in *Y. lipolytica*. Higher cultivation temperatures resulted in production of even longer saturated fatty acids, 3% in *C. oleaginosus* and 1.5% in *Y. lipolytica*.

Conclusions: In this study, we provided the optimum C/N ratio and temperature for *C. oleaginosus* and *Y. lipolytica* by RSM. Additionally, we demonstrated that lipid accumulation of both oleaginous yeasts was significantly affected by the C/N ratio and temperature. Furthermore, we systematically analyzed the variation in fatty acids composition and proved that changing the C/N ratio and temperature steer the composition. We have further established these oleaginous yeasts as platforms for production of tailored fatty acids.

Keywords: Oleaginous yeasts, Microbial oil, Response surface methodology, Carbon to nitrogen ratio, Lipid accumulation

*Correspondence: maria.suarezdiez@wur.nl

² Laboratory of Systems and Synthetic Biology, Wageningen University & Research, 6708 WE Wageningen, The Netherlands
Full list of author information is available at the end of the article



Introduction

The use of plant-derived oils, especially palm oil, is increasing at an alarming rate. This is happening in part as a replacement for fossil foils, but mostly as they are cheap sources of many useful components. The oils and fatty acids derived from palm trees are used in food, feed, chemical, personal care, and cosmetic products for health benefits, sensorial reasons (texture, flavor), to extend shelf-life, and as surfactants or emulsifiers [1–3]. As a result, palm tree groves are rapidly replacing the original tropical forests, and other original and traditional vegetation in many Asian, South American, and African countries. This replacement is not only threatening the local ecosystem but is also having a major effect on the local livelihoods, as it causes deforestation and contributes to climate change [4, 5]. Despite some responsible actions that have been taken among them fighting against deforestation driven by RSPO (Roundtable on Sustainable Palm Oil), the use of palm oil remains controversial [6]. To that end, developing a sustainable alternative to fatty acids and oils is urgent and of utmost interest.

Oil-producing yeasts, referred to as oleaginous yeasts, have strong potential as sustainable alternatives for lipid production in various industrial applications [7]. *Yarrowia lipolytica* and *Cutaneotrichosporon oleaginosus* also known as *Apiotrichum curvatum*, *Cryptococcus curvatus*, *Trichosporon cutaneum*, *Trichosporon oleaginosus*, and *Cutaneotrichosporon curvatum* are reported among the top five most well-known oil-producing yeasts. *C. oleaginosus* and *Y. lipolytica* can accumulate oils up to 70% and 40% of their biomass respectively [8–10]. Lipid accumulation in oleaginous yeasts is induced by limiting specific nutrients such as nitrogen, phosphate, and sulphur. Nitrogen limitation or, in other words, a high C/N ratio in the growth medium, has been observed to be the most effective lipid induction strategy [9]. As reported by Ykema et al., after passing the critical C/N of 11 g/g, the oleaginous yeast starts to accumulate oils by re-routing the excess carbon to be stored as lipids [11, 12]. Under nitrogen limiting conditions, the produced fatty acid composition has been reported to be 25% palmitic acid (C16:0), 10% stearic acid (C18:0), 57% oleic acid (C18:1), and 7% linoleic acid (C18:2) by *C. oleaginosus* [13] and 15% C16:0, 13% C18:0, 51% C18:1, and 21% C18:2 by *Y. lipolytica* [14]. This composition is comparable to that of palm oil. Furthermore, these yeasts can use a broad range of carbon sources, such as glucose, xylose, glycerol, sucrose, and lactose. They can also use more complex and inexpensive side streams such as crude glycerol from bioethanol production or whey permeate as a feedstock, which is significant to reduce raw materials cost [15–17]. Moreover, *Y. lipolytica* is non-pathogenic and regarded as food-grade yeast, thus its oil can be used for food-related

applications [18, 19]. Due to these advantages, oleaginous yeasts are flagged as attractive microbial-cell factories to sustain a bio-based circular economy for industrial implementation. However, from an economic point of view, the lipid production process of oleaginous yeasts still requires substantial optimization for industrial purposes. Koutinas et al. [20] reported that implementation of microbial oil in industrial applications is strongly dependent on the final microbial oil concentrations and lipid productivity [19]. For *C. oleaginosus*, it has been calculated that the process is only economically feasible if lipid accumulation reaches approximately 85% (w/w).

Identifying and designing optimal production conditions is a challenging step in developing bioconversion systems since these cultivation conditions play a crucial role in productivity [21]. For at least two decades, efforts have been made to design the optimum growth medium and fermentation conditions to boost the lipid accumulation as well as to sustain the growth of oleaginous yeasts [22–24]. When Ochoa-Estopier et al. revealed the effect of the C/N ratio in a cultivation medium is critical for the regulation of lipid accumulation [25], Kuttiraja et al. reported that glycerol concentration has affected lipid accumulation more than the concentration of trace elements and inoculum size for *Y. lipolytica* [26]. One of the most efficient strategies to systematically identify optimum production conditions is through the Response Surface Methodology (RSM). This method decreases experimental time and laborious work compared to the one-factor-at-a-time (OFAT). Whereas OFAT only allows changing one of the considered factors in each of the experiments, RSM provides a design on which multiple parameters are changed at each experimental run. Moreover, RSM aims to predict the observed response, by reliably estimating the experimental variability [27]. For instance, Awad et al. assessed the effect of various carbon and nitrogen sources on the physiology of *C. oleaginosus* via RSM [28]. In another example, Cui et al. focused on the effect of temperature and pH on lipid content and the growth on crude glycerol by maintaining C/N ratio [29]. Additionally, Canonico et al. reported optimum C/N ratio and time to maximize lipid production of *Y. lipolytica* [17]. However, no studies have focused on the synergistic effect between the C/N ratio of a defined medium and cultivation parameters.

In addition to the optimized lipid productivity of oleaginous yeasts, tailoring the composition of produced lipids is important for increasing economic competitiveness. For instance, the longer fatty acids, such as oleic acid, lauric acid, and palmitic acid are heavily used in home and personal care products due to their cleaning/surfacting activities. Polyunsaturated fatty acids (PUFAs) such as linolenic acid supply various health properties. A part

of the previous studies on *Y. lipolytica* enabled switching the fatty acid composition by metabolic engineering strategies. These approaches were mainly focused on the manipulation of the elongation and desaturation pathway [30, 31]. Additionally, Ochsenreither et al. [19] reported that temperature and the composition of the medium lead to variation in fatty acid composition. However, there are no studies focused on a systematic evaluation of the effect of these parameters on variation in fatty acids produced by oleaginous yeasts. Therefore, analyzing the fatty acid compositions under different cultivation conditions will provide valuable information that will enable us to devise strategies to tailor fatty acid compositions.

Thus, in this study, we aimed to further develop *C. oleaginosus* and *Y. lipolytica* as microbial cell factories for the improved production of lipids and lipids with higher content of specific fatty acids. The first set of experiments was designed to assess the variance between the fatty acid composition produced by *C. oleaginosus* and *Y. lipolytica* at various C/N ratios and temperatures. We extensively studied the variance in fatty acids composition via PCA. Additionally, we performed another set of experiments to broaden the current experimental region for RSM and determined the optimized C/N ratio and temperature for *C. oleaginosus* and *Y. lipolytica*.

Materials and methods

Yeast strains and pre-culture preparation

Cutaneotrichosporon oleaginosus ATCC 20509 and *Yarrowia lipolytica* DSM 1345 were maintained on Yeast extract Peptone Dextrose (YPD) agar plates containing 10 g/L yeast extract, 20 g/L peptone, 20 g/L glucose, 20 g/L agar. The maintained cultures were stored at 4 °C for up to a week. The inoculum was prepared by transferring a single colony of the oleaginous yeasts into 10 mL YPD broth (10 g/L yeast extract, 20 g/L peptone, 20 g/L glucose, 20 g/L) in 50 mL tubes and incubated at 30 °C, 250 rpm for 18 h in a shaking incubator.

Cultivation conditions

C. oleaginosus and *Y. lipolytica* were grown in defined media consisting of glycerol as a carbon source and urea as a nitrogen source. The medium was adapted from Meester et al. [32] with modifications. In addition to the variation in carbon to nitrogen ration, the medium contained 2.7 g/L KH_2PO_4 , 1.79 g/L $\text{NaH}_2\text{PO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g/L EDTA with the pH 5.5 as well as the trace elements: 40 mg/L $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 5.5 mg/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 5.2 mg/L citric acid- H_2O , 1 mg/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.76 mg/L $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 10 $\mu\text{L/L}$ H_2SO_4 (36 N).

Experimental design

The C/N ratio varied from 30:1 (g/g) to 300:1 (g/g) by mixing 4–40 g carbon/L with 0.13 g nitrogen/L for both oleaginous yeasts. After that, prepared cultures were incubated at different temperatures (15 °C, 25 °C, 30 °C, and 35 °C), 250 rpm for 96 or 144 h. For the experimental design, the C/N ratio (X_1) and temperature (X_2) were selected as variables. Low levels and high levels for C/N ratio and temperature are C/N 30, C/N 120, and 15 °C, 35 °C. The coded values of independent variables were calculated by considering the high and low levels together with the real values:

$$\text{Coded value} = \frac{\text{real value} - \left(\frac{\text{high level value} + \text{low level value}}{2} \right)}{\frac{1}{2}x(\text{high value} - \text{low value})}$$

and are presented in Table 1.

Determination of biomass

The growth of oleaginous yeasts was monitored every 24 h by measuring the absorbance at 600 nm (OD_{600}). A calibration curve was plotted with the absorbance versus the dry cell weight for *C. oleaginosus* and *Y. lipolytica* (Additional file 1: Figure S1). The dry cell was obtained as follows: (1) 5 mL of culture was centrifuged at 3200g for 15 min, (2) the cells were washed twice with 10 mL of deionized water (3) and freeze-dried.

Identification of lipids and fatty acid composition

The total fatty acids and the fatty acid composition were determined quantitatively. The samples were prepared by mixing 20–25 mg freeze-dried yeast cells with 2 mL of 15% H_2SO_4 in methanol and 2 mL of chloroform containing methyl pentanoate as an internal standard. The samples were incubated for 4 h at 85–95 °C and cooled on ice for 5 min, 1 mL of distilled water was added. Following

Table 1 Levels of two independent variables employed in RSM in terms of real and coded values for *C. oleaginosus* and *Y. lipolytica*

Levels and coded values	Real values	
	C/N ratio (g/g) X_1	Temperature (°C) X_2
−1.0	30	15
−0.3	60	−
0.0	75	25
0.5	−	30
1.0	120	35
3.7	240	−
5.0	300	−

the phase separation by centrifugation at 2200g for 5 min, the organic phase was collected from the bottom of the tube and dried with NaSO₄. Subsequently, The fatty acid methyl esters (FAME) were analyzed with a gas chromatograph (Brand, City, Country) equipped with a Zebron ZB-FAME column (30 m × 0.25 mm × 0.20 μm; Phenomenex, Torrance, CA, The US). The yeast's oil content was calculated from the internal standard.

Computational analysis

All computational analysis was performed with R version 4.0.2 [33].

Principal Component Analysis (PCA) was carried out on the fatty acid profiles by the statistics function `prcomp` within R [34]. The correlation biplots of the principal component scores and the loading vectors were plotted through R `ggplot2` package [35].

Response Surface Methodology (RSM) was performed by the `rsm` package, and the contour plots were generated with the R `pers` or `cont` functions [36]. The relationship between the responses and factors was expressed by the second-order polynomial equation:

$$Y = \beta_0 + \sum \beta_i x_i + \sum \beta_{ii} x_i^2 + \sum \beta_{ij} x_i x_j.$$

Y represents the predicted response, β_0 is the interception coefficient, β_i is the linear coefficient, β_{ii} is the quadratic coefficient and β_{ij} is the interaction coefficient, X_i is the independent variable, X_i^2 is the squared effect, and $X_i X_j$ is the interaction effect. The quality of the regression equations was assessed according to the coefficient of determination (R^2) and lack of fit F-test. Statistical analysis of the model was performed using Analysis of Variance (ANOVA) and $p < 0.05$ was considered significant. Optimal levels of the C/N ratio and temperature were given as stationary points via RSM.

Results

In total, 10 experiments were conducted for *C. oleaginosus* and 9 experiments for *Y. lipolytica*. The biomass concentration of *C. oleaginosus* and *Y. lipolytica* varied from 1.24 g/L to 5.54 g/L and 2.34 to 4.52 g/L, lipid content ranged from 3.62 to 47.41% and 3.35% and 18.51% for all tested conditions (Tables 2 and 3). While the highest lipid content was obtained at C/N 120 at 30 °C for *C. oleaginosus*, it reached the maximum point at C/N 140 at 25 °C in *Y. lipolytica*. Incubating *C. oleaginosus* at 15 °C slightly decreased the biomass and lipid content compared to the other tested temperatures. On the other hand, decreasing the C/N ratio reduced the biomass concentration of *Y. lipolytica*.

Table 2 Biomass density and lipid content of *C. oleaginosus* at different C/N ratios and temperatures

C (g/L)	C/N (g/g)	Temperature (°C)	Biomass (g/L)	Lipid content (% g/g)
4	30	15	1.24 ± 0.07	10.71 ± 0.93
16	120	15	2.61 ± 0.24	18.09 ± 1.38
32	240	15	2.36 ± 0.21	3.62 ± 0.32
4	30	30	5.43 ± 0.08	31.16 ± 2.54
8	60	30	5.18 ± 0.08	38.48 ± 1.61
16	120	30	5.54 ± 0.16	47.41 ± 0.70
32	240	30	4.43 ± 0.16	44.12 ± 1.06
40	300	30	5.33 ± 0.21	37.83 ± 1.99
4	30	35	2.56 ± 0.03	28.95 ± 1.08
16	120	35	2.54 ± 0.16	38.92 ± 0.73

Table 3 Biomass density and lipid content of *Y. lipolytica* at different C/N ratios and temperatures

C (g/L)	C/N (g/g)	Temperature (°C)	Biomass (g/L)	Lipid content (% g/g)
4	30	15	2.62 ± 0.08	7.91 ± 0.49
16	120	15	4.40 ± 0.08	15.91 ± 0.07
4	30	30	2.45 ± 0.16	4.00 ± 0.24
16	120	30	4.52 ± 0.14	14.26 ± 1.09
4	30	35	2.34 ± 0.11	3.35 ± 0.33
16	120	35	3.12 ± 0.13	9.25 ± 0.06
9.75	75	25	3.01 ± 0.13	15.29 ± 0.41
9.75	75	25	3.06 ± 0.09	15.01 ± 0.17
18.2	140	25	4.18 ± 0.14	18.51 ± 0.96

Development of regression models and ANOVA

C/N ratio and temperature as coded variables, and lipid content and biomass as responses were analyzed with RSM. The relation was fitted by second-order polynomial equations to obtain the regression equation models (Table 4). These models represent the empirical relationships between the biomass density, lipid content of cells, and the variables (C/N ratio (X_1) and temperature (X_2)) in coded units. These regression models suggest that both linear and quadratic effects of C/N ratio and temperature significantly affected the lipid content of *C. oleaginosus* and *Y. lipolytica* (Table 4). However, interaction of C/N ratio and temperature has no significant effect on any considered responses of *Y. lipolytica*. When the growth of *C. oleaginosus* is not significantly affected by the C/N ratio, the combined effect of the C/N ratio and temperature has a significant effect. On the other hand, only the linear effect of

Table 4 Regression equations, statistics of regression equations for lipid content and biomass of *C. oleaginosus* and *Y. lipolytica*

<i>C. oleaginosus</i>			<i>Y. lipolytica</i>		
Dependent variable: Lipid content % (w/w)					
$Y(\text{Lipidcontent}) = 39.26 + 4.85X_1 + 10.08X_2 + 2.59X_1X_2 - 1.38X_1^2 - 14.01X_2^2$			$Y(\text{Lipidcontent}) = 14.47 + 4.30X_1 - 3.12X_2 - 0.28X_1X_2 - 1.50X_1^2 - 4.03X_2^2$		
Source	Estimate	p-values	Source	Estimate	p-values
Model	39.26	< e-15***	Model	14.47	< e-15***
X ₁	4.85	< e-08***	X ₁	4.30	< e-10***
X ₂	10.08	< e-12***	X ₂	- 3.12	< e-07***
X ₁ :X ₂	2.59	< e-07***	X ₁ :X ₂	- 0.28	0.480
X ₁ ²	- 1.38	< e-09***	X ₁ ²	- 1.50	0.011*
X ₂ ²	- 14.01	< e-08***	X ₂ ²	- 4.03	< e-05***
Dependent variable: Biomass (g/L)					
$Y(\text{Biomass}) = 6.12 + 0.11X_1 + 0.36X_2 - 0.17X_1X_2 - 0.02X_1^2 - 3.91X_2^2$			$Y(\text{Biomass}) = 3.17 + 0.71X_1 - 0.33X_2 - 0.13X_1X_2 - 0.14X_1^2 - 0.12X_2^2$		
Source	Estimate	p-values	Source	Estimate	p-values
Model	6.12	< e-15***	Model	3.17	< e-15***
X ₁	0.11	0.210	X ₁	0.71	< e-09***
X ₂	0.36	0.005**	X ₂	- 0.33	0.005**
X ₁ :X ₂	- 0.17	0.005**	X ₁ :X ₂	- 0.13	0.238
X ₁ ²	- 0.02	0.323	X ₁ ²	- 0.14	0.344
X ₂ ²	- 3.91	< e-13***	X ₂ ²	- 0.12	0.500

Table 5 Evaluating the significance of regression models by ANOVA

<i>C. oleaginosus</i>						<i>Y. lipolytica</i>				
Source	Degrees of Freedom	Sum Square	Mean Square	F-Value	p-Value	Degrees of Freedom	Sum Square	Mean Square	F-Value	p-Value
<i>Lipid content % (w/w)</i>										
FO(X ₁ , X ₂)	2	4181.0	2090.50	356.93	< 2.2e-16	2	562.80	281.40	141.65	6.459e-13
TWI (X ₁ , X ₂)	1	497.2	497.20	84.89	2.374e-09	1	1.15	1.15	0.58	0.4561
PQ (X ₁ , X ₂)	2	1022.5	511.27	87.29	9.709e-12	2	128.23	64.11	32.27	3.936e-07
Residual	24	140.6	5.86			21	41.72	1.99		
Lack of fit	4	82.5	20.63	7.11	0.000992	2	29.90	14.95	24.04	6.237e-06
Pure error	20	58.1	2.90			19	11.81	0.62		
Dependent variable = Lipid content % (w/w); R ² = 0.9759; Adj R ² = 0.9709, p-value = < e-15						Dependent variable = Lipid content % (w/w); R ² = 0.9432; Adj R ² = 0.9296, p-value = < e-11				
<i>Biomass (g/L)</i>										
FO(X ₁ , X ₂)	2	23.91	11.97	69.94	1.128e-10		14.03	7.01	49.52	1.122e-08
TWI (X ₁ , X ₂)	1	0.31	0.31	1.80	0.1919	1	0.24	0.24	1.70	0.2063
PQ (X ₁ , X ₂)	2	42.61	21.30	122.84	2.467e-13	2	0.16	0.08	0.55	0.5834
Residual	24	4.16	0.17			21	2.97	0.14		
Lack of fit	4	3.43	0.86	23.58	2.483e-07	2	2.37	0.19	37.48	2.539e-07
Pure error	20	0.73	0.04			19	0.60	0.03		
Dependent variable = Biomass (g dcw/L); R ² = 0.9414; Adj R ² = 0.9292, p-value = < e-13						Dependent variable = Biomass (g dcw/L); R ² = 0.8291; Adj R ² = 0.7884, p-value = < e-06				

FO, TWI, and PQ refer to the linear function, two-way interactions, and quadratic terms in the model formula of RSM respectively.

investigated factors significantly affected the growth of *Y. lipolytica* (Table 5).



ANOVA was performed to assess the significance and adequacy of response surface quadratic models. The quality of the model fit can be evaluated by the coefficient

of determination (R^2), which provides a measure of how much variability in the observed response values can be explained by the experimental factors and their interactions. The R^2 value is always between 0 and 1 and the closer the R^2 value is to 1, the stronger the model is and the better it predicts the response. For the developed models of *C. oleaginosus* and *Y. lipolytica*, the determination confidence coefficients (R^2) are 97.58% and 94.32% for lipid content, 94.14% and 82.91% for biomass. These R^2 values represented that sample variation of the regression models described the experimental data accurately. The model is regarded as significant if the p-value is lower than 0.05. In other words, ‘Model F-value’ could occur because of noise with only a 5% chance [37]. Therefore p-value of the models, for *C. oleaginosus* $P_{lipid\ accumulation} = < e-15$, $P_{biomass} = < e-13$, and for *Y. lipolytica*

$P_{lipid\ accumulation} = < e-11$, and $P_{biomass} = < e-6$ suggested the coefficients are significant. The Lack of Fit P-values showed that the Lack of Fit F-value could occur due to noise with the possibility of almost 0% for the second model for *C. oleaginosus* and 0% for all models for *Y. lipolytica*, and 0.0992% for the first model of *C. oleaginosus*.

Response surface analysis

The factors, C/N ratio of the growth medium, and temperature were selected to be optimized for *C. oleaginosus* and *Y. lipolytica*. Three-dimensional surface responses were plotted to illustrate the relationships between the responses and variables for *C. oleaginosus* (Fig. 1) and *Y. lipolytica* (Fig. 2). When the temperature was around 30 °C and the C/N ratio was between C/N 75 to C/N 220,



the lipid accumulation of *C. oleaginosus* was improved (Fig. 1). On the other hand, the biomass density increased within the range of C/N 60 to C/N 240 and from 25 °C to 30 °C (Fig. 1). Additionally, lipid accumulation of *Y. lipolytica* was enhanced starting from C/N 100 and between 18 °C to 25 °C (Fig. 2).

The optimal values for the investigated dependence factors were predicted from these 3D response surface plots (Figs. 1 and 2). Accordingly, the maximum predicted responses for *C. oleaginosus* were 47.67% lipid accumulation and 6.26 g yeast dry cell weight/L. While the optimum C/N ratio and temperature were approximately C/N 175 and 30 °C for predicted lipid accumulation, C/N 185 and 25 °C were suggested to achieve the predicted maximum biomass density. The suggested optimum values for lipid accumulation resulted in $51.17\% \pm 0.66$ lipid accumulation, and 5.28 ± 0.08 g dry yeast cells/L which confirmed the predictions of developed regression models (Table 6). Moreover, suggestions of RSM improved lipid accumulation by 9% in *C. oleaginosus*. Regression models of *Y. lipolytica* predicted the optimum conditions as C/N 140 and 21 °C for maximum lipid production, which predicted 18.33% lipid accumulation, and 4.72 g yeast dry cell weight/L (Table 6). These predicted optimum conditions provided $19.03\% \pm 0.37$ lipid accumulation, and 5.07 ± 0.16 g dry yeast cells/L for *Y. lipolytica*. Additionally, model predictions were validated via experiments at C/N 45 and 30 °C for *C. oleaginosus* and C/N 180 at 21 °C for *Y. lipolytica* (Table 6).

Analysis of the fatty acid profile

Principal Component Analysis (PCA) was conducted to clarify the variation in the fatty acid profile at different C/N ratios and temperatures. Produced the fatty acid compositions of *C. oleaginosus* and *Y. lipolytica* are represented in Tables 7 and 8. The PCA showing the variation in the fatty acid profile is shown in Fig. 3. As seen in Fig. 3 for *C. oleaginosus* the variance explained in PC1 and PC2 were 43% and 22% respectively and for *Y. lipolytica* was

56% and 20%. As observed from Fig. 3, the highest temperature (35 °C) and the combination of C/N 30 with the other tested temperatures (15 °C, 30 °C) caused a higher content of saturated and longer chain fatty acids (C20:0, C22:0, C24:0). *C. oleaginosus* produced higher content of unsaturated fatty acids (C18:1, C18:2, C18:3) at the lowest temperature, 15 °C. On the other hand, we observed the same effect on the fatty acid profile of *Y. lipolytica* in terms of saturation level and chain length by changing the temperature. *Y. lipolytica* produced higher content of C14:0, C16:0, C18:0, C20:0, and C24:0 when it was incubated at 35 °C, and C14:1, C16:1, C18:1, and C20:1 at 15 °C.

Discussion

The lipid accumulation and lipid productivity in oleaginous yeasts are strongly dependent on the composition of the cultivation medium and the operational conditions. Therefore, this study sought to assess the importance of such factors in lipid production. To this end, we performed an RSM analysis and represented the optimum C/N ratio and temperature for the maximization of lipid content, and biomass density. The optimum C/N ratio and temperature are C/N 175 at 30 °C for *C. oleaginosus* and C/N 140 at 21 °C for *Y. lipolytica*. Moreover, we demonstrated that a C/N ratio and temperature cause variations in the fatty acid composition of oleaginous yeasts.

In the experiments performed for RSM, we used glycerol as a carbon source as this is efficiently utilized by *C. oleaginosus* and *Y. lipolytica* [39, 40] and urea as a nitrogen source as it provides higher biomass yields compared to the ammonium salts [28]. RSM is one of the most preferred methods to optimize operational conditions and medium composition in biotechnology [41, 42]. This method facilitates obtaining more information with a fewer number of experiments by changing multiple factors at a time because RSM reflects on the complex nonlinear relationships between independent variables and measured responses of the system. In contrast, the

Table 6 Validation of RSM models by suggested optimum conditions and additional experiments

C/N (g/g)	Temperature (°C)	Predicted Values		Measured/Calculated Values	
		Lipid Accumulation (%) g/g)	Biomass (g/L)	Lipid Accumulation (%), g/g)	Biomass (g/L)
<i>C. oleaginosus</i>					
45	30	36.08	5.30	38.48 ± 1.61	5.28 ± 0.08
175	30	47.67	4.97	51.17 ± 0.66	5.35 ± 0.28
<i>Y. lipolytica</i>					
140	21	18.33	4.72	19.03 ± 0.37	5.07 ± 0.16
180	21	17.20	5.81	17.72 ± 0.16	4.96 ± 0.16

Table 7 Fatty acid profile of *C. oleginosus* at various C/N ratios and temperatures

C/N (g/g)	Temperature (°C)	Fatty Acid Profile (%)															
		C14:0	C16:0	C16:1	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C24:0	Saturated FAs	MUFAs	PUFAs	Long chain FAs	Very long chain FAs
30	15	-	24.62 ± 1.47	-	43.92 ± 0.53	26.58 ± 1.09	4.88 ± 1.96	-	-	-	-	-	68.54 ± 1.47	26.58 ± 1.09	4.88 ± 1.96	100 ± 0.0	-
30	30	-	15.44 ± 3.06	-	21.09 ± 4.22	46.70 ± 1.41	14.27 ± 1.93	-	-	-	2.50 ± 0.54	39.03 ± 1.05	46.7 ± 1.41	14.27 ± 1.93	97.5 ± 0.54	2.5 ± 0.54	
35	35	-	9.33 ± 0.58	-	28.14 ± 0.80	42.52 ± 0.86	12.50 ± 0.81	1.90 ± 0.17	-	1.27 ± 0.06	4.34 ± 0.06	44.98 ± 0.05	42.52 ± 0.86	12.50 ± 0.81	94.39 ± 0.0	5.61 ± 0.0	
60	30	-	21.00 ± 1.67	1.09 ± 0.07	18.96 ± 2.74	51.06 ± 0.87	8.25 ± 0.91	-	-	-	-	39.96 ± 2.26	52.15 ± 1.36	8.25 ± 0.91	100 ± 0.0	-	
120	15	-	18.32 ± 0.54	1.79 ± 0.43	10.74 ± 0.63	56.16 ± 1.64	10.74 ± 0.62	2.26 ± 0.24	-	-	-	29.05 ± 0.24	57.95 ± 1.21	13.00 ± 0.97	100 ± 0.0	-	
30	30	0.39 ± 0.03	30.33 ± 2.14	2.71 ± 0.23	7.39 ± 0.49	46.50 ± 1.98	11.93 ± 0.50	0.44 ± 0.01	-	-	0.30 ± 0.02	38.85 ± 2.57	49.22 ± 2.14	11.93 ± 0.50	99.7 ± 0.02	0.30 ± 0.02	
35	35	0.40 ± 0.01	14.61 ± 0.11	0.71 ± 0.01	22.09 ± 0.22	47.85 ± 0.04	8.54 ± 0.18	1.75 ± 0.06	-	1.13 ± 0.03	3.09 ± 0.00	43.34 ± 0.43	48.55 ± 0.03	8.54 ± 0.18	95.95 ± 0.08	4.22 ± 0.04	
240	15	-	24.89 ± 2.20	-	-	59.08 ± 2.55	16.01 ± 0.82	-	-	-	-	24.89 ± 2.20	59.08 ± 2.55	16.01 ± 0.82	100 ± 0.0	-	
30	30	0.38 ± 0.03	29.30 ± 1.46	2.88 ± 0.10	6.82 ± 0.19	47.23 ± 1.49	13.10 ± 0.33	-	-	-	0.29 ± 0.03	36.79 ± 1.62	50.11 ± 1.51	13.10 ± 0.33	100 ± 0.0	0.29 ± 0.03	
Palm oil (range) [38]		0.3 - 1.1	32.6-46.0	0.0 -0.5	1.7-7.6	35.1 - 53.4	7.0 - 15.0	0.2-0.6	0.1-0.5	0.1 - 0.2	-	36.7 - 51.5	35.3 - 53.8	7.3 - 15.5	100	-	

Fatty acid profile of palm oil added for direct comparison


MUFAs: Monounsaturated fatty acids, PUFAs: Polyunsaturated fatty acids

Table 8 Fatty acid profile of *Y. lipolytica* at various C/N ratios and temperatures

C/N (g/g)	Temperature (°C)	Fatty Acid Profile (%)																
		C10:0	C14:0	C14:1	C16:0	C16:1	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C24:0	Saturated FAs	MUFAs	PUFAs	Long chain FAs	Very long chain FAs
30	15	1.27 ± 0.15	-	0.68 ± 0.01	8.56 ± 0.34	13.41 ± 0.37	6.27 ± 0.70	46.54 ± 1.17	21.41 ± 0.01	-	-	1.84 ± 0.32	-	16.11 ± 1.20	62.48 ± 1.22	21.41 ± 0.01	98.72 ± 0.16	-
30	30	2.53 ± 0.31	-	-	10.59 ± 0.35	11.47 ± 0.68	5.57 ± 0.38	43.97 ± 0.88	24.67 ± 0.44	1.18 ± 0.10	-	-	-	19.88 ± 0.27	55.44 ± 0.21	24.67 ± 0.44	97.47 ± 0.31	-
35	35	-	1.02 ± 0.31	-	18.77 ± 0.27	6.47 ± 0.18	14.94 ± 1.37	33.83 ± 0.13	19.14 ± 0.71	3.60 ± 0.27	-	-	2.22 ± 0.04	40.56 ± 1.02	40.30 ± 0.31	19.14 ± 0.71	97.77 ± 0.04222 ± 0.04	-
120	15	-	-	0.30 ± 0.00	10.26 ± 0.10	19.98 ± 0.66	5.38 ± 0.25	50.21 ± 0.43	11.38 ± 0.20	-	-	1.81 ± 0.03	0.68 ± 0.01	16.32 ± 0.40	72.29 ± 0.52	11.38 ± 0.20	99.32 ± 0.10068 ± 0.01	-
30	30	-	-	-	23.31 ± 0.58	11.40 ± 1.37	7.81 ± 0.66	35.73 ± 1.06	21.74 ± 1.39	-	-	-	-	31.12 ± 0.08	47.14 ± 1.30	21.75 ± 1.38	100 ± 0.00	-
35	35	-	0.29 ± 0.01	-	20.97 ± 0.05	6.79 ± 0.24	22.14 ± 1.06	32.75 ± 1.23	12.72 ± 0.58	2.23 ± 0.14	-	2.11 ± 0.13	47.73 ± 0.87	39.54 ± 1.42	12.72 ± 0.58	97.88 ± 0.132.11 ± 0.13	-	
Palm oil (range) [38]	-	-	0.3 - 1.1	-	32.6 - 46.0	0.0 - 0.5	1.7 - 7.6	35.1 - 53.4	7.0 - 15.0	0.2 - 0.6	0.1 - 0.5	0.1 - 0.2	-	36.7 - 51.5	35.3 - 53.8	7.3 - 15.5	100	-

Fatty acid profile of palm oil added for direct comparison

MUFAs: Monounsaturated fatty acids, PUFAs: Polyunsaturated fatty acids



OFAT approach allows changing only one of the factors for each of the experiments. Whereas around 15 experiments are required to follow the OFAT approach with the same factors and same levels, the number of experiments decreased by approximately 40% via the DoE approach. The statistics tables of developed regression models in this study represented that both the C/N ratio and the temperature have a significant effect on the lipid content of cells, and biomass density of *Y. lipolytica*. Although

they utilized crude glycerol as a cultivation medium, the effect of the C/N ratio on the behavior of biomass and lipid content is comparable to the findings of Canonico et al. [17]. In another study, Ochoa-Estopier reported that the C/N ratio has a critical role in the regulation of lipid metabolism of *Y. lipolytica* as lipid accumulation was controlled by concentration of carbon source without citrate excretion [25]. Therefore, this approach is consistent for different cultivation mediums. In this study, we

investigated a broader range of C/N ratios, the effect of temperature, and their combinatorial effect which has not been tested on total lipid production and growth of *Y. lipolytica* to our knowledge. Optimum temperature predicted by our model (21 °C) for lipid accumulation and growth for *Y. lipolytica* was surprisingly lower than previously reported value (28 °C) [43, 44]. On contrary to *Y. lipolytica*, in this study, the growth of *C. oleagnosus* was significantly affected by only temperature and the combined effect of temperature and C/N ratio. These findings are consistent with the report of Cui et al. even though they tested a narrower temperature range between 27 °C to 33 °C, whereas in our design, it was extended from 15 °C to 35 °C [29]. This extension was applied not only in temperature range but also in C/N ratio compared to Awad et al. which provided different optimum C/N ratios that resulted in ~3% improvement in lipid accumulation compared to their findings [28]. Although the maximum lipid accumulation of *Y. lipolytica* is much less than *C. oleagnosus*, it is corresponding to the amounts reported for the wild-type strain [45, 46]. Gao et al. reported lipid accumulation of *Y. lipolytica* CICC 31,596 up to 30% (g/g) when it grew on volatile fatty acids [43]. On the other hand, *Y. lipolytica* ACA-DC 50,109 produced 20% (g/g) lipids on a glycerol-based cultivation medium [39]. These findings show lipid accumulation of *Y. lipolytica* is strain-dependent. Our findings provided an optimum C/N ratio and temperature as these predicted values provided higher lipid contents and biomass for both oleaginous yeasts. These optimum values identified via RSM in this study will potentially contribute to solving other optimization problems such as revealing the effect of other factors, and finding optimum conditions for other strains and engineered strains. In addition to optimization of cultivation conditions, lipid accumulation ability of *Y. lipolytica* and *C. oleagnosus* can be further improved by strain engineering as suggested in other reports [47].

Fatty acids with different chain lengths and saturation levels are utilized in various products. Engineering of elongation and desaturation of fatty acids has been studied for *Y. lipolytica* however limiting genetic engineering strategies and genetic elements restricts the applicability of that strategy for other oleaginous yeast such as *C. oleagnosus* [48, 49]. In our work, we demonstrated that the C/N ratio and temperature affected the fatty acids composition of *C. oleagnosus* and *Y. lipolytica* in addition to the lipid content and biomass. In previous studies, when Awad et al. mentioned the effect of selected N-source and C-source on produced fatty acid composition [28], Moon et al. reported that 15 °C as a growth temperature shifted the fatty acid profile of *C. oleagnosus* to more

unsaturated fatty acids [50]. Moreover, Hackenschmidt et al. claimed that there was only a slight variation in the fatty acid profile of *Y. lipolytica* from 25 to 35 °C [51]. However, a systematic evaluation of the effect of low and high levels of temperature, C/N ratio, and their combination on fatty acid composition had not been performed before, to our knowledge. In this study, PCA allowed us to evaluate the variation in produced fatty acids composition by reducing the noise and creating uncorrelated components from the analyzed data. When the growth temperature of oleaginous yeasts was increased from optimum to 35 °C and the C/N ratio decreased to C/N 30, the saturation level and the chain length of fatty acids were increased. Therefore we showed that the C/N ratio can be used to manipulate fatty acid composition in addition to the temperature and N-source. On the other hand, fatty acids produced at 15 °C were slightly more unsaturated (C14:1, C16:1, C18:1, and C20:1 for *Y. lipolytica*, C18:1, C18:2, and C18:3 for *C. oleagnosus*) than they were at the optimum temperature. This variety can be explained by the adaptation of an organism to maintain lipid fluidity at different temperatures [50]. Because the saturation level and the chain length directly influence the melting point of fatty acids. The melting points are much lower for unsaturated and shorter chain length of fatty acids whereas it is higher for saturated and longer chain fatty acids [52]. While low incubation temperatures increased unsaturation levels on both oleaginous yeasts, it decreased the growth and lipid productivity of *C. oleagnosus*. Unexpectedly, low incubation temperatures positively affected lipid accumulation and biomass production in *Y. lipolytica*. This situation demonstrated that *Y. lipolytica* is promising to produce fatty acids, especially with higher content of unsaturated fatty acids as growth and lipid accumulation were enhanced by lower temperature. Tezaki et al. related functions of some genes in *Y. lipolytica* with the adaptation ability of this organism at low temperatures [53], still the changes in the cellular metabolic network leading to these changes are not fully understood nor the regulatory and signalling pathways orchestrating this response. Therefore, elucidation of the adaptation mechanism of oleaginous yeasts to low temperatures or too high temperatures could contribute to achieving higher productivity and expanding the application potential of microbial oils.

Conclusion

In this study, we sought to determine the major operational factors affecting physiological fitness toward fatty acid production by oleaginous yeasts. We aimed to enhance lipid accumulation as well as to enable *C.*

oleaginosus and *Y. lipolytica* to attain higher content of particular fatty acids by changing operating conditions such as the available C/N ratio and temperature. We applied a thorough DoE method (RSM) and developed second-order polynomial equations to identify the optimum C/N ratios and temperatures for both oleaginous yeasts. The predictions of RSM improved the lipid accumulation by approximately 71% for *C. oleaginosus* and about 66% for *Y. lipolytica* compared to the average lipid accumulation in the tested conditions. While the lipid accumulation was significantly affected by the C/N ratio and temperature, the growth of *C. oleaginosus* was mainly affected by temperature. Additionally, changing the C/N ratio in the cultivation medium and temperature resulted in variations in fatty acid profile, which we observed switches from saturated to unsaturated fatty acids, unsaturated to saturated fatty acids, and shorter chain to longer chain fatty acids. These switches in fatty acid compositions might be the result of maintaining the fluidity of lipid bodies within oleaginous yeasts as a response to low and high temperatures, still the molecular and regulatory changes in the underlying cellular metabolic network remain to be elucidated. Altogether, these findings helped strengthen the basis to deploy these oleaginous yeasts as platforms for tailored fatty acid production and thereby contribute to the development of processes substituting palm oil that are more sustainable.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12934-022-01956-5>.

Additional file 1. Calibration curves to calculate dry cell weight of *C. oleaginosus* and *Y. lipolytica* from OD₆₀₀.

Acknowledgements

We thank Sara Moreno Paz for her valuable contribution to design of experiment and for providing the initial code for RSM. We thank Janine Verbokkem for her valuable contribution to the analysis of fatty acid composition.

Author contributions

All authors conceived and designed the study. ZEDÖ and MSD performed the data analysis. ZEDÖ drafted the manuscript and performed the experiments. VAPMds, JH, and MSD acquired project funding, conceived and supervised the research. All authors reviewed and edited the study. All authors read and approved the final manuscript.

Funding

This research was financed by the Dutch Ministry of Agriculture through the “TKI-toeslag” project LWV19221 “Tailor-made microbial oils and fatty acids”.

Declarations

Competing interests

JH has interests in NoPalm Ingredients BV and VAPMds has interests in LifeGlimmer GmbH.

Author details

¹Bioprocess Engineering, Wageningen University & Research, 6708 PB Wageningen, The Netherlands. ²Laboratory of Systems and Synthetic Biology, Wageningen University & Research, 6708 WE Wageningen, The Netherlands. ³LifeGlimmer GmbH, 12163 Berlin, Germany. ⁴Wageningen Food & Biobased Research, Wageningen University & Research, 6708 WE Wageningen, The Netherlands.

Received: 22 June 2022 Accepted: 13 October 2022

Published online: 03 November 2022

References

- Rustan AC, Drevon CA. Fatty Acids: Structures and Properties. *Encycl Life Sci*. 2005. <https://doi.org/10.1038/npg.els.0003894>.
- Holley RA, Patel D. Improvement in shelf-life and safety of perishable foods by plant essential oils and smoke antimicrobials. *Food Microbiol*. 2005. <https://doi.org/10.1016/j.fm.2004.08.006>.
- Desbois P, Potential A. Applications of antimicrobial fatty acids in medicine, agriculture and other industries. *Recent Pat Antiinfect Drug Discov*. 2012. <https://doi.org/10.2174/157489112801619728>.
- Abubakar A, Ishak MY, Makmom AA. Impacts of and adaptation to climate change on the oil palm in Malaysia: a systematic review. *Environ Sci Pollut Res*. 2021. <https://doi.org/10.1007/s11356-021-15890-3>.
- Snashall GB, Poulos HM. Oreos versus orangutans: The need for sustainability transformations and nonhierarchical polycentric governance in the global palm oil industry. *Forests*. 2021. <https://doi.org/10.3390/f12020252>.
- Cazzolla Gatti R, Liang J, Velichevskaya A, Zhou M. Sustainable palm oil may not be so sustainable. *Sci Total Environ*. 2019. <https://doi.org/10.1016/j.scitotenv.2018.10.222>.
- Patel A, Karageorgou D, Rova E, Katapodis P, Rova U, Christakopoulos P, et al. An overview of potential oleaginous microorganisms and their role in biodiesel and omega-3 fatty acid-based industries. *Microorganisms*. 2020. <https://doi.org/10.3390/microorganisms8030434>.
- Görner C, Redai V, Bracharz F, Schrepfer P, Garbe D, Brück T. Genetic engineering and production of modified fatty acids by the non-conventional oleaginous yeast *Trichosporon oleaginosus* ATCC 20509. *Green Chem*. 2016;18:2037–46.
- Beopoulos A, Nicaud JM, Gaillardin C. An overview of lipid metabolism in yeasts and its impact on biotechnological processes. *Appl Microbiol Biotechnol*. 2011. <https://doi.org/10.1007/s00253-011-3212-8>.
- Xue SJ, Chi Z, Zhang Y, Li YF, Liu GL, Jiang H, et al. Fatty acids from oleaginous yeasts and yeast-like fungi and their potential applications. *Crit Rev Biotechnol*. 2018. <https://doi.org/10.1080/07388551.2018.1428167>.
- Ykema A, Verbree EC, van Verseveld HW, Smit H. Mathematical modelling of lipid production by oleaginous yeasts in continuous cultures. *Antonie Van Leeuwenhoek*. 1986;52:491–506.
- Carsanba E, Papanikolaou S, Erten H. Production of oils and fats by oleaginous microorganisms with an emphasis given to the potential of the nonconventional yeast *Yarrowia lipolytica*. *Crit Rev Biotechnol*. 2018;38:1230–43. <https://doi.org/10.1080/07388551.2018.1472065>.
- Pham N, Reijnders M, Suarez-Diez M, Nijssse B, Springer J, Eggink G, et al. Genome-scale metabolic modeling underscores the potential of *Cutaneotrichosporon oleaginosus* ATCC 20509 as a cell factory for biofuel production. *Biotechnol Biofuels*. 2021. <https://doi.org/10.1186/s13068-020-01838-1>.
- Papanikolaou S, Aggelis G. Lipid production by *Yarrowia lipolytica* growing on industrial glycerol in a single-stage continuous culture. *Bioresour Technol*. 2002. [https://doi.org/10.1016/S0960-8524\(01\)00149-3](https://doi.org/10.1016/S0960-8524(01)00149-3).
- Dobrowolski A, Drzymała K, Miłuta P, Mirończuk AM. Production of tailor-made fatty acids from crude glycerol at low pH by *Yarrowia lipolytica*. *Bioresour Technol*. 2020;314:10–3.
- Ykema A, Verbree EC, Verwoert IIGS, van der Linden KH, Nijkamp HJJ, Smit H. Lipid production of revertants of Ufa mutants from the oleaginous yeast *Apiotrichum curvatum*. *Appl Microbiol Biotechnol*. 1990. <https://doi.org/10.1007/BF00176521>.
- Canonico L, Ashoor S, Taccari M, Comitini F, Antonucci M, Truzzi C, et al. Conversion of raw glycerol to microbial lipids by new *Metschnikowia* and

- Yarrowia lipolytica* strains. *Ann Microbiol.* 2016. <https://doi.org/10.1007/s13213-016-1228-0>.
18. Zinjarde SS. Food-related applications of *Yarrowia lipolytica*. *Food Chem.* 2014. <https://doi.org/10.1016/j.foodchem.2013.11.117>.
 19. Amalia L, Zhang YH, Ju YH, Tsai SL. Enhanced Lipid Production in *Yarrowia lipolytica* Po1g by Over-expressing Iro1 Gene under Two Different Promoters. *Appl Biochem Biotechnol.* 2020. <https://doi.org/10.1007/s12010-020-03226-9>.
 20. Koutinas AA, Chatzifragkou A, Kopsahelis N, Papanikolaou S, Kookos IK. Design and techno-economic evaluation of microbial oil production as a renewable resource for biodiesel and oleochemical production. *Fuel.* 2014. <https://doi.org/10.1016/j.fuel.2013.08.045>.
 21. Singh V, Haque S, Niwas R, Srivastava A, Pasupuleti M, Tripathi CKM. Strategies for fermentation medium optimization: An in-depth review. *Front Microbiol.* 2017. <https://doi.org/10.3389/fmicb.2016.02087>.
 22. Ykema A, Verbree EC, Kater MM, Smit H. Optimization of lipid production in the oleaginous yeast *Apiotrichum curvatum* in whey permeate. *Appl Microbiol Biotechnol.* 1988. <https://doi.org/10.1007/BF00251704>.
 23. Seo Y, Han S, Han JI. Economic biodiesel production using algal residue as substrate of lipid producing yeast *Cryptococcus curvatus*. *Renew Energy.* 2014. <https://doi.org/10.1016/j.renene.2014.03.062>.
 24. Back A, Rossignol T, Krier F, Nicaud JM, Dhulster P. High-throughput fermentation screening for the yeast *Yarrowia lipolytica* with real-time monitoring of biomass and lipid production. *Microb Cell Fact.* 2016. <https://doi.org/10.1186/s12934-016-0546-z>.
 25. Ochoa-Estopier A, Guillouet SE. D-stat culture for studying the metabolic shifts from oxidative metabolism to lipid accumulation and citric acid production in *Yarrowia lipolytica*. *J Biotechnol Elsevier.* 2014;170:35–41.
 26. Kuttiraja M, Douha A, Valéro JR, Tyagi RD. Elucidating the Effect of Glycerol Concentration and C/N Ratio on Lipid Production Using *Yarrowia lipolytica* SKY7. *Appl Biochem Biotechnol.* 2016. <https://doi.org/10.1007/s12010-016-2189-2>.
 27. Sarabia LA, Ortiz MC, Sánchez MS. Response Surface Methodology. *Compr Chemom.* 2020. <https://doi.org/10.1016/B978-0-12-409547-2.14756-0>.
 28. Awad D, Bohnen F, Mehlmer N, Brueck T. Multi-factorial-guided media optimization for enhanced biomass and lipid formation by the oleaginous yeast *Cutaneotrichosporon oleaginosus*. *Front Bioeng Biotechnol.* 2019. <https://doi.org/10.3389/fbioe.2019.00054>.
 29. Cui Y, Blackburn JW, Liang Y. Fermentation optimization for the production of lipid by *Cryptococcus curvatus*: use of response surface methodology. *Biomass Bioenerg.* 2012. <https://doi.org/10.1016/j.biombioe.2012.09.017>.
 30. Gao Q, Yang JL, Zhao XR, Liu SC, Liu ZJ, Wei LJ, et al. *Yarrowia lipolytica* as a metabolic engineering platform for the production of very-long-chain wax esters. *J Agric Food Chem.* 2020. <https://doi.org/10.1021/acs.jafc.0c04393>.
 31. Konzock O, Matsushita Y, Zaghen S, Sako A, Norbeck J. Altering the fatty acid profile of *Yarrowia lipolytica* to mimic cocoa butter by genetic engineering of desaturases. *Microb Cell Fact.* 2022. <https://doi.org/10.1186/s12934-022-01748-x>.
 32. Meesters PAEP, Huijberts GNM, Eggink G. High-cell-density cultivation of the lipid accumulating yeast *Cryptococcus curvatus* using glycerol as a carbon source. *Appl Microbiol Biotechnol.* 1996;45:575–9.
 33. R Core Team. R software: Version 4.0.2. R Found Stat Comput. 2020.
 34. Martins TG. Computing and visualizing PCA in R. R-bloggers. 2013.
 35. Wickham H. ggplot2. Wiley Interdiscip Rev Comput Stat. 2011. <https://doi.org/10.1002/wics.147>.
 36. Lenth RV. Response-surface methods in R, using RSM. *J Stat Softw.* 2009. <https://doi.org/10.18637/jss.v032.i07>.
 37. Tan HT, Lee KT, Mohamed AR. Pretreatment of lignocellulosic palm biomass using a solvent-ionic liquid [BMIM]Cl for glucose recovery: An optimisation study using response surface methodology. *Carbohydr Polym.* 2011. <https://doi.org/10.1016/j.carbpol.2010.10.052>.
 38. Montoya C, Lopes R, Flori A, Cros D, Cuellar T, Summo M, et al. Quantitative trait loci (QTLs) analysis of palm oil fatty acid composition in an interspecific pseudo-backcross from *Elaeis oleifera* (HBK) Cortés and oil palm (*Elaeis guineensis* Jacq). *Tree Genet Genomes.* 2013. <https://doi.org/10.1007/s11295-013-0629-5>.
 39. Makri A, Fakas S, Aggelis G. Metabolic activities of biotechnological interest in *Yarrowia lipolytica* grown on glycerol in repeated batch cultures. *Bioresour Technol.* 2010;101:2351–8. <https://doi.org/10.1016/j.biortech.2009.11.024>.
 40. Pham N, Reijnders M, Suarez-Diez M, Nijssse B, Springer J, Eggink G, et al. Genome-scale metabolic modeling underscores the potential of *Cutaneotrichosporon oleaginosus* ATCC 20509 as a cell factory for biofuel production. *Biotechnol Biofuels.* 2021;14:1–17. <https://doi.org/10.1186/s13068-020-01838-1>.
 41. Xu YX, Li YL, Xu SC, Liu Y, Wang X, Tang JW. Improvement of xylanase production by *Aspergillus niger* XY-1 using response surface methodology for optimizing the medium composition. *J Zhejiang Univ Sci B.* 2008. <https://doi.org/10.1631/jzus.B0820038>.
 42. Xu Y, Yang L, Zhao S, Wang Z. Large-scale production of tauroursodeoxycholic acid products through fermentation optimization of engineered *Escherichia coli* cell factory. *Microb Cell Fact.* 2019. <https://doi.org/10.1186/s12934-019-1076-2>.
 43. Gao R, Li Z, Zhou X, Bao W, Cheng S, Zheng L. Enhanced lipid production by *Yarrowia lipolytica* cultured with synthetic and waste-derived high-content volatile fatty acids under alkaline conditions. *Biotechnol Biofuels.* 2020. <https://doi.org/10.1186/s13068-019-1645-y>.
 44. Qiao K, Wasylenko TM, Zhou K, Xu P, Stephanopoulos G. Lipid production in *Yarrowia lipolytica* is maximized by engineering cytosolic redox metabolism. *Nat Biotechnol.* 2017. <https://doi.org/10.1038/nbt.3763>.
 45. Fontanille P, Kumar V, Christophe G, Nouaille R, Larroche C. Bioconversion of volatile fatty acids into lipids by the oleaginous yeast *Yarrowia lipolytica*. *Bioresour Technol.* 2012;114:443–9. <https://doi.org/10.1016/j.biortech.2012.02.091>.
 46. Jia YL, Wang LR, Zhang ZX, Gu Y, Sun XM. Recent advances in biotechnological production of polyunsaturated fatty acids by *Yarrowia lipolytica*. *Crit Rev Food Sci Nutr.* 2021. <https://doi.org/10.1080/10408398.2021.1937041>.
 47. Carsamba E, Papanikolaou S, Fickers P, Erten H. Lipids by *Yarrowia lipolytica* strains cultivated on glucose in batch cultures. *Microorganisms.* 2020. <https://doi.org/10.3390/microorganisms8071054>.
 48. Gao S, Tong Y, Wen Z, Zhu L, Ge M, Chen D, et al. Multiplex gene editing of the *Yarrowia lipolytica* genome using the CRISPR-Cas9 system. *J Ind Microbiol Biotechnol.* 2016. <https://doi.org/10.1007/s10295-016-1789-8>.
 49. Wen Z, Al Makishah NH. Recent advances in genetic technology development of oleaginous yeasts. *Appl Microbiol Biotechnol.* 2022. <https://doi.org/10.1007/s00253-022-12101-y>.
 50. Moon NJ, Hammond EG. Oil production by fermentation of lactose and the effect of temperature on the fatty acid composition1. *J Am Oil Chem Soc.* 1978;55:683–8.
 51. Hackenschmidt S, Bracharz F, Daniel R, Thürmer A, Bruder S, Kabisch J. Effects of a high-cultivation temperature on the physiology of three different *Yarrowia lipolytica* strains. *FEMS Yeast Res.* 2019;19:1–11.
 52. Fasina OO, Craig-Schmidt M, Colley Z, Hallman H. Predicting melting characteristics of vegetable oils from fatty acid composition. *LWT - Food Sci Technol.* 2008. <https://doi.org/10.1016/j.lwt.2007.09.012>.
 53. Tezaki S, Iwama R, Kobayashi S, Shiwa Y, Yoshikawa H, Ohta A, et al. $\Delta 12$ -fatty acid desaturase is involved in growth at low temperature in yeast *Yarrowia lipolytica*. *Biochem Biophys Res Commun Elsevier Ltd.* 2017;488:165–70. <https://doi.org/10.1016/j.bbrc.2017.05.028>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.