

Glycemic outcomes among rural patients in the type 1 diabetes T1D Exchange registry, January 2016–March 2018: a cross-sectional cohort study

Arashpreet Gill,¹ M David Gothard,² Kathaleen Briggs Early ³

To cite: Gill A, Gothard MD, Briggs Early K. Glycemic outcomes among rural patients in the type 1 diabetes T1D Exchange registry, January 2016–March 2018: a cross-sectional cohort study. *BMJ Open Diab Res Care* 2022;**10**:e002564. doi:10.1136/bmjdr-2021-002564

AG and KBE contributed equally.

Received 26 August 2021
Accepted 6 December 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹College of Osteopathic Medicine, Pacific Northwest University, Yakima, Washington, USA

²BIOSTATS, East Canton, Ohio, USA

³College of Osteopathic Medicine, Department of Biomedical Sciences, Pacific Northwest University, Yakima, Washington, USA

Correspondence to

Dr Kathaleen Briggs Early; kearly@pnwu.edu

ABSTRACT

Introduction Does rural status influence glycemic outcomes among participants in the type 1 diabetes T1D Exchange clinic registry?

Research design and methods Data from the T1D Exchange clinic registry between January 2016 and March 2018 were identified by rural–urban status and stratified by age and hemoglobin A1c (HbA1c). Multivariable regression modeling was performed to isolate HbA1c differences. A full model including all significant ($p < 0.05$ via two-sided testing) differential factors was determined with an additional indicator for rural status, and adjusted for duration of diabetes, use of continuous glucose monitoring device, age, race/ethnicity, and private insurance status. The model was reduced using backwards elimination stepwise procedures until only significant factors remained.

Results Mean HbA1c levels for all rural participants were significantly higher (8.71%; 72 mmol/mol) compared with the urban group (8.48%; 69 mmol/mol), $p < 0.001$. For youth under 13 years of age, rural participants had a higher mean HbA1c (8.65%; 71 mmol/mol) compared with urban (8.45% 69 mmol/mol), $p = 0.022$. Rural youth (13–<18 years) had a higher mean HbA1c (9.39%; 79 mmol/mol) than urban youth (9.14%; 76 mmol/mol), $p < 0.001$. Rural young adults (18–<26 years) had a higher mean HbA1c (9.07%; 76 mmol/mol) than urban young adults (8.88%; 74 mmol/mol), $p = 0.042$. Rural adults (≥ 26 years; $n = 589$) were the only group that did not have a higher mean HbA1c (7.76%, 61.3 mmol/mol) than urban adults ($n = 4770$; 7.72%, 60.9 mmol/mol), $p = 0.503$. Rural locale was highly significant ($\beta = 0.175$, $p < 0.001$) despite controlling for potentially confounding differences between rural and urban groups.

Conclusions Among this T1D Exchange cohort, there is a pattern of higher mean HbA1c being associated with rural status, even after adjustment for characteristic differences, most strikingly among those under 26 years of age. This disparity and contributing factors need to be more thoroughly studied to provide effective solutions.

INTRODUCTION

One in five Americans (about 60 million people) live in a rural area,¹ and they also carry a higher burden of chronic disease compared with their urban counterparts.²

Significance of this study

What is already known about this subject?

- ▶ Diabetes prevalence and mortality are higher among rural Americans.
- ▶ Limited information is known about type 1 diabetes among rural Americans.

What are the new findings?

- ▶ Despite data for this study being collected via endocrinology services, this analysis of the type 1 diabetes T1D Exchange registry found that rural participants had worse glycemic control than participants from urban areas.
- ▶ Rural youth under age 26 years had worse mean hemoglobin A1c (HbA1c) values than urban youth of the same age.
- ▶ For adults aged 26 years and over, there were no differences in HbA1c between rural and urban groups.

How might these results change the focus of research or clinical practice?

- ▶ Rural residence and associated barriers encountered among rural-residing persons with type 1 diabetes should be anticipated and addressed in care planning.
- ▶ Researchers and providers working in rural areas should strive to provide creative care delivery solutions that can reach patients and their families where they reside.

Incidence of type 1 diabetes among rural Americans has been reported to be 2.28 times (95% CI 2.08 to 2.50) that of persons living in high-density areas,³ and it is well recognized that prevalence of type 1 diabetes in the USA and worldwide is increasing.⁴ While there have been positive changes in diabetes-related mortality over time, urban mortality rates have decreased almost three to five times more than the rates observed in rural areas.⁵

Multiple barriers, including those related to socioeconomic and transportation issues, contribute to rural diabetes disparities.

Additionally, there is often a culture of self-reliance and hesitancy to use urban-based medical resources.^{2 6} Limited access to endocrinology specialty care in rural communities is another barrier to rural diabetes outcome improvement.^{7 8} These significant and ongoing diabetes health disparities throughout rural America^{5 9 10} have instigated discussions about how rural health equity can be improved.¹¹ Moreover, type 1 diabetes outcomes have not been thoroughly studied in the context of rural residence.

The type 1 diabetes T1D Exchange clinic registry was established in 2010 with more than 35 000 adult and youth participants of all ages. While not a nationally representative sample of persons with type 1 diabetes, the registry includes patients from 81 pediatric and adult endocrinology clinics from more than 35 states.¹² Beck *et al*¹³ have published a complete description of the T1D Exchange clinic registry including participant recruitment, inclusion and exclusion criteria, and overall baseline characteristics. Analyses of the registry have covered topics varying from device use to disparities in treatments and outcomes.^{12 14} To our knowledge, however, there has been no prior investigation of the T1D Exchange clinic registry comparing rural–urban participant glycemetic observations. Additionally, there is limited understanding of the effects of rural status on type 1 diabetes outcomes.^{15–18}

The purpose of this study was to investigate the cross-sectional mean hemoglobin A1c (HbA1c) and other diabetes-related characteristics among the rural and urban participants in the T1D Exchange. We hypothesized that disparities in glycemia, as indicated by mean HbA1c, would diminish significantly because all T1D Exchange registry participants receive diabetes care via endocrinology services. We also believed it would be of interest to look at the T1D Exchange clinic registry data through the lens of rural residence stratified by age-specific groupings (all ages; youth below 13 years; youth 13–<18 years; young adults 18–<26 years; and adults 26 years of age and older), and predetermined HbA1c ranges (<7% (<53 mmol/mol), 7%–<9% (53–<75 mmol/mol), and ≥9% (≥75 mmol/mol)).

RESEARCH DESIGN AND METHODS

Study data for this retrospective cohort analysis were obtained from the T1D Exchange registry of participants between January 1, 2016 and March 31, 2018. Year 5 data were provided with registrant zip codes by the Jaeb Center for Health Research T1D Exchange clinic registry team.^{12 19} Year 5 data include registrants who were followed for 5 years and completed the baseline and year 5 questionnaires.

Variables for this study were obtained through medical records and patient questionnaires completed by patients and/or their guardians, which is explained more thoroughly by Foster *et al*.¹² Age, duration of

diabetes, body mass index, insulin pump and continuous glucose monitoring (CGM) device use, non-insulin glucose-lowering medication use, and HbA1c levels were collected via medical records. Self-monitoring of blood glucose habits were reported as meter downloads when available, or self-report. Occurrences of severe hypoglycemia (SH; defined as loss of consciousness or seizure) and diabetic ketoacidosis (DKA; defined as requiring overnight hospitalization) were reported if they occurred in the 3 months before the participant/caregiver completed the year 5 questionnaire.¹² The analysis for participant diabetes control is based on the calculated average prior 12 months' HbA1c from the year 5 questionnaire.

Data were imported into SPSS V.25.0 software and the cohort was stratified by rural location status (Yes/No) based on zip code as determined by the Centers for Medicaid and Medicare Services (CMS)²⁰ definitions of rurality for payment, which defines 'rural' by zip code based on urbanized areas and clusters.²¹ We recognize there are many ways to define rural,²² but we chose to use the CMS rurality for payment definition as it was feasible and efficient for our small research team's limited resources, and it is also based on a real-world and clinically meaningful application (eg, payment for medical care/equipment). For simplicity, we will use the terms 'rural' and 'urban' to describe the two groups reported here.

The age group cut-offs and HbA1c categories used to frame our analysis were based on commonly used ranges from the literature^{23 24} and adjusted to reflect the Affordable Care Act coverage available to those up to age 26 years. For age ranges, we used all ages; youth below 13 years; youth 13–<18 years; young adults 18–<26 years; and adults 26 years of age and older. For the predetermined HbA1c ranges, we used <7% (<53 mmol/mol), 7%–8.99% (53–<75 mmol/mol), and ≥9% (≥75 mmol/mol). Looking specifically at mean HbA1c groupings across several age strata would allow us to more effectively investigate the youth-specific concerns, which is widely recognized as a difficult time period for optimal diabetes control,^{12 25} and understudied in rural youth.

Statistical analysis

Demographic data were summarized both overall for the entire cohort and by rural–urban status. Numeric data were summarized using means, SDs, and minimum to maximum range values. Comparisons between rural and urban groups for mean equality were performed via independent sample t-tests. Categorical data were summarized overall and by rural–urban status using frequencies and percentages. Depending on the cell sample sizes and ordinality of the data, either Pearson χ^2 tests, Fisher's exact tests, linear association χ^2 tests, or exact linear association tests were performed to compare rural with urban location status for distributional equality.

Table 1 Demographic characteristics

	Urban, n (%)	Rural, n (%)	P value
Total participants, n (%)	12 476 (100)	1837 (100)	0.040
Age in years, mean (SD)	28.9 (18.06)	27.9 (19.33)	0.040
Child (<13 years)	1188 (9.5)	224 (12.2)	<0.001
Adolescent (13–<18 years)	3567 (28.6)	609 (33.2)	
Young adult (18–<26 years)	2951 (23.7)	415 (22.6)	
Adult (26+ years)	4770 (38.2)	589 (32.1)	
Gender, n (%)			0.316
Female	6375 (51.1)	964 (52.5)	
Male	6097 (48.9)	872 (47.5)	
Transgender	4 (0.0)	1 (0.1)	
Race/ethnicity, n (%)			<0.001
White non-Hispanic	10 399 (83.4)	1697 (92.4)	
Black non-Hispanic	565 (4.5)	25 (1.4)	
Hispanic or Latino	982 (7.9)	60 (3.3)	
Other	530 (4.2)	55 (3.0)	
Annual income, n (%)			<0.001
Missing/unreported	5881 (47.1)	883 (48.1)	
<\$50 000	1726 (26.2)	338 (35.4)	
\$50 000–\$75 000	934 (14.2)	203 (21.3)	
>\$75 000	3935 (59.7)	413 (43.3)	
Education, n (%)			<0.001
Missing/unreported	3493 (28.0)	557 (30.3)	
Less than HS diploma	2362 (26.3)	416 (32.5)	
HS diploma/GED	2500 (27.8)	402 (31.4)	
Associate's degree	565 (6.3)	118 (9.2)	
Bachelor's degree	2047 (22.8)	226 (17.7)	
Master's degree	1092 (12.2)	94 (7.3)	
Professional/doctoral degree	417 (4.6)	24 (1.9)	
Insurance status, n (%)			<0.001
Private	9477 (77.1)	1240 (69.2)	
Other	2728 (22.2)	533 (29.7)	

GED, General Educational Diploma; HS, high school.

Treatment-related and crude outcome data were similarly summarized and analyzed. Due to characteristic differences between the rural and urban strata for demographic and/or treatment-related data, a multivariable linear regression model was employed to adjust for these differences. A full multivariable regression model including all significant ($p < 0.05$ via two-sided testing) differential factors (duration of diabetes, age, use of CGM, race/ethnicity, and private insurance status) was determined, with an additional indicator factor for rural status. Two factors, education and income, were significant but not included in the model due to missing data ranging from 28% to over 48% of the rural and urban sample, which could have biased the model. Therefore, we used insurance status as a proxy measure

for socioeconomic status, rather than education and income. Finally, the multivariable model was reduced using backwards elimination stepwise procedures until only significant ($p < 0.05$ via two-sided testing) factors were included.

For the primary outcome variables, mean HbA1c and number of HbA1c measurements, the percentage of non-missing, subject-level data were determined to be above 98% for each strata. Similarly for the multivariable regression model, the percentage of complete subject data including non-missing data for all predictive factors was determined to be 93.9%. Due to the completeness of the data, missing data were not included in the analyses and no imputation was performed. All data are based on the 12-month time frame within the data collection

Table 2 Diabetes-related health characteristics

	Urban, n (%)	Rural, n (%)	P value
Age at diagnosis (years), n (%)	12 476 (100)	1837 (100)	0.663
Mean (SD)	11.6 (11.19)	11.5 (11.70)	
Duration of diabetes (years), n (%)	12 475 (>99.9)	1837 (100)	0.004
Mean (SD)	17.3 (12.40)	16.4 (12.60)	
Use of CGM, n (%)	3684 (30.5)	420 (23.6)	<0.001
Use of insulin pump, n (%)	8056 (65.7)	1224 (67.4)	0.142
BMI (m ² /kg), n (%)	11 316 (90.7)	1705 (92.8)	0.731
Mean (SD)	25.4 (5.70)	25.4 (5.71)	
Total daily insulin (units per kg), n (%)	7270 (58.3)	1151 (62.7)	0.089
Mean (SD)	0.78 (0.30)	0.79 (0.30)	
Use of non-insulin medications for blood sugar control, n (%)			0.874
Missing/unreported	2619 (21.0)	361 (19.7)	
Yes	836 (8.5)	127 (8.6)	
No	9021 (91.5)	1349 (91.4)	
General health (patient reported), n (%)			0.644
Missing/unreported	3262 (26.1)	525 (28.6)	
Excellent	1602 (17.4)	227 (17.3)	
Very good	3523 (38.2)	487 (37.1)	
Good	2945 (32.0)	443 (33.8)	
Fair	976 (10.6)	136 (10.4)	
Poor	168 (1.8)	19 (1.4)	
Smoke at least 1 cigarette/week, n (%)			0.127
Missing/did not wish to answer	5055 (40.5)	822 (44.7)	
Yes	304 (4.1)	52 (5.1)	
No	7117 (95.9)	963 (94.9)	
Days per week exercising, n (%)*	8493 (68.0)	1184 (64.5)	0.383
Mean (SD)	4.0 (2.11)	4.0 (2.19)	
≥1 DKA events in past 3 months, n (%)			0.517
Missing	3262 (26.1)	525 (28.6)	
Yes	252 (2.7)	40 (3.0)	
No	8962 (97.3)	1272 (97.0)	
≥1 severe hypoglycemic events† in past 3 months, n (%)			0.937
Missing	3262 (26.1)	525 (28.6)	
Yes	567 (6.2)	80 (6.1)	
No	8647 (93.8)	1232 (93.9)	
Average patient-reported SMBG/day, n (%)	9055 (72.6)	1287 (70.1)	0.629
Mean (SD)	4.88 (2.42)	4.91 (2.33)	
Average glucose tests/day from meter download, n (%)	8635 (69.2)	1382 (75.2)	0.619
Mean (SD)	4.19 (2.47)	4.23 (2.49)	

*Exercise was only reported for youth and young adults below age 26 years.

†Severe hypoglycemic events were defined as an episode of documented or presumed low blood glucose that resulted in seizure or loss of consciousness per Cengiz *et al.*³⁹

BMI, body mass index; CGM, continuous glucose monitoring; DKA, diabetic ketoacidosis; SMBG, self-monitoring of blood glucose.




Figure 1 Mean HbA1c by age group and rural status. *Significant difference between urban and rural participants. All ages: $p<0.001$; below 13 years of age: $p=0.022$; 13–below 18 years of age: $p=0.001$; 18–26 years of age: $p=0.042$; 26 years of age and older: $p=0.503$. HbA1c, hemoglobin A1c.

period (January 2016–March 2018) when participants or caregivers completed their year 5 questionnaire.

RESULTS

Demographics

A total of 14 313 registrants were included in this analysis (1837 rural, 13%; and 12 476 urban, 87%). The rural and urban participants in this analysis differed significantly across multiple demographic variables, summarized in [table 1](#). Briefly, rural participants differed significantly from the urban group by age distribution ($p=0.04$), but both groups were primarily non-Hispanic white (rural, 92.4%; urban, 83.4%). Overall, rural participants reported lower income and educational attainment than urban participants ($p<0.001$). Both groups reported having primarily private insurance, but more urban participants had private insurance compared with those in the rural group ($n=9477$, 77.1%; and $n=1240$, 69.2%, respectively, $p<0.001$).

Diabetes-related health characteristics

The two groups varied in some, but not all, diabetes-related health characteristics ([table 2](#)). Age at diagnosis for type 1 diabetes onset was not different between the urban and rural groups (11.6 years, 11.19 SD; and 11.5 years, 11.70 SD, respectively), however, the urban group had a significantly longer duration of diabetes (17.3 years, 12.4 SD) than the rural group (16.4 years, 12.60 SD; $p=0.004$).

Diabetes device use was also different between the groups. Use of CGM was significantly higher among participants in the urban compared with rural groups ($n=3684$, 30.5%; $n=420$, 23.6%, respectively; $p<0.001$), but insulin pump use was not significantly different ($n=8056$, 65.7% for the urban group; $n=1224$, 67.4% for the rural group; $p=0.142$).

No other significant differences were found in other diabetes-related health characteristics including body

mass index, total daily dose of insulin per kilogram body weight (rural group mean 0.79 units/kg, SD 0.30; urban group mean 0.78 units/kg, 0.30 SD, $p=0.089$), use of non-insulin medications, general self-rated health, smoking status, or reported days per week of exercise. There were also no differences in reported instances of DKA (urban mean affirming DKA episode $n=252$, 2.7%; rural mean $n=40$, 3.0%; $p=0.517$), or severe hypoglycemic (SH) episodes (urban mean affirming SH episodes $n=567$, 6.2%; rural mean $n=80$, 6.1%; $p=0.937$). There were also no differences in the frequency of self-reported monitoring of blood glucose (mean tests per day for rural 4.91, SD 2.33; urban 4.88, SD 2.42; $p=0.629$), or the average number of tests downloaded from glucometers (rural 4.23, SD 2.49; urban 4.19, SD 2.47; $p=0.619$).

HbA1c: number of measurements and mean values

Across all age strata, there were significantly more HbA1c measurements taken within the past year of the data collection period among the rural versus urban group (3.18 mean tests, 1.25 SD; 3.08 mean tests, 1.37 SD, respectively, $p=0.003$). Rural adults over the age of 26 years had significantly more HbA1c tests (2.90 tests in the past 12 months, 1.19 SD) compared with those in the urban group (2.67 tests in the past 12 months, 1.31 SD; $p<0.001$).

[Figure 1](#) and [table 3](#) present the mean HbA1c findings across all age strata by rural–urban status. Briefly, rural participants had significantly higher mean HbA1c (8.71% or 72 mmol/mol, 1.66 SD; $p<0.001$) compared with urban participants (8.48%, 69 mmol/mol; 1.63 SD). Fewer rural youth 13–<18 years of age had an HbA1c between 7% (53 mmol/mol) and 8.99% (75 mmol/mol) compared with urban youth ($n=273$; 45.1% and $n=1772$; 50.1%, respectively, $p=0.002$). More rural youth between 13 and <18 years of age had an average HbA1c 9% (75 mmol/mol) and over compared with the urban group ($n=315$; 52.1% and $n=1596$; 45.1%, respectively, $p<0.002$).

Multivariable linear regression

After adjustment for characteristic differences (duration of diabetes, use of CGM, age, race/ethnicity, and private insurance), rural location was a statistically significant factor in the reduced model for predicting average HbA1c value (beta=0.175, $p<0.001$) ([table 4](#)). Rural locale was associated with a significantly increased HbA1c even after identifying and adjusting for significant characteristic differences in potentially confounding factors between the rural and urban strata.

DISCUSSION

After examination of all age groups, this analysis of the 2016–2018 T1D Exchange registry cohort found that rural participants under 26 years of age had significantly higher average HbA1c values than urban participants. Moreover, when looking at the T1D Exchange sample stratified by both age and HbA1c, we continued to find that the HbA1c generally worsens among the younger

Table 3 HbA1c outcomes by age and rural status

	HbA1c	Urban, n (%)	Rural, n (%)	P value	
All ages	<7%	1725 (14.1)	179 (9.9)	<0.001	
	7%–8.99%	6833 (55.8)	1002 (55.3)		
	≥9%	3689 (30.1)	631 (34.8)		
	Average				<0.001
	n (%)	12247 (98.2)	1812 (98.6)		
	Mean (SD)	8.48 (1.63)	8.71 (1.66)		
<13 years	<7%	81 (6.9)	6 (2.7)	0.059	
	7%–8.99%	771 (65.7)	151 (68.0)		
	≥9%	322 (27.4)	65 (29.3)		
	Average				0.022
	n (%)	1174 (98.8)	222 (99.1)		
	Mean (SD)	8.45 (1.19)	8.65 (1.17)		
13–<18 years	<7%	170 (4.8)	17 (2.8)	0.002	
	7%–8.99%	1772 (50.1)	273 (45.1)		
	≥9%	1596 (45.1)	315 (52.1)		
	Average				0.001
	n (%)	3538 (99.2)	60 (99.3)		
	Mean (SD)	9.14 (1.69)	9.39 (1.73)		
18–<26 years	<7%	269 (9.2)	30 (7.3)	0.115	
	7%–8.99%	1506 (51.7)	200 (48.7)		
	≥9%	1140 (39.1)	181 (44.0)		
	Average				0.042
	n (%)	2915 (98.8)	411 (99.0)		
	Mean (SD)	8.88 (1.78)	9.07 (1.77)		
≥26 years	<7%	1205 (26.1)	126 (22.0)	0.033	
	7%–8.99%	2784 (60.3)	378 (65.9)		
	≥9%	631 (13.7)	70 (12.2)		
	Average				0.503
	n (%)	4620 (96.9)	574 (97.5)		
	Mean (SD)	7.72 (1.21)	7.76 (1.15)		

HbA1c, hemoglobin A1c.

population and improves with age, illustrated by rural youth having worse mean HbA1c levels than their urban counterparts. Similarly, but not significantly different due to the small convenience sample size, Stumetz *et al*²⁶ also reported higher HbA1c levels among their sample of 61 youth with type 1 diabetes (mean age 13.3 years) for rural (9%; 75 mmol/mol) versus urban (8.5%; 69 mmol/mol) participants.

Frequency of HbA1c monitoring, which can contribute to overall glycemic control, was higher in the rural cohort reported herein. The American Diabetes Association Standards of Medical Care in Diabetes recommends testing HbA1c levels more often among those adults with HbA1c above 7%.²⁷ The importance of frequent HbA1c monitoring among those with type 1 diabetes was established with the Diabetes Control and Complications

Trial²⁸ and others.²⁹ More recently, a study with over 15 000 Austrian and German patients with type 1 diabetes found both mean HbA1c (different between rural and urban in our study) and self-monitoring of blood glucose (not different between rural and urban in our study) were associated with achieving target HbA1c.³⁰ Specifically, among the rural adults aged 26 years and over in our study, we found an increased average number of HbA1c tests reported, which could contribute to the improved HbA1c in adults—both rural and urban. Previous research has found that while rural physicians do indeed order HbA1c tests, many rural patients do not have optimal control of their diabetes,^{31 32} which aligns with our findings.

The significantly higher usage of CGM among urban participants is likely related to the differences in

Table 4 Multivariable regression model for factors influencing HbA1c outcomes

Outcome/factors	Factor effect—full model			Factor effect—reduced model		
	Beta (SE)	95% CI	P value	Beta (SE)	95% CI	P value
Mean HbA1c in past 12 months n=13 436 (93.9%)						
Intercept	10.112 (0.068)	9.979 to 10.245	<0.001	10.149 (0.044)	10.062 to 10.237	<0.001
Duration of diabetes	-0.006 (0.002)	-0.010 to -0.003	<0.001	-0.006 (0.002)	-0.010 to -0.003	<0.001
Use of CGM	-0.574 (0.028)	-0.630 to -0.519	<0.001	-0.574 (0.028)	-0.630 to -0.519	<0.001
Age	-0.024 (0.001)	-0.026 to -0.022	<0.001	-0.024 (0.001)	-0.027 to -0.022	<0.001
Race—white	-0.323 (0.064)	-0.449 to -0.198	<0.001	-0.359 (0.041)	-0.439 to -0.278	<0.001
Race—black	0.723 (0.089)	0.549 to 0.896	<0.001	0.686 (0.073)	0.543 to 0.830	<0.001
Race—Hispanic	0.057 (0.078)	-0.096 to 0.209	0.464			
Private insurance	-0.535 (0.031)	-0.596 to -0.475	<0.001	-0.537 (0.031)	-0.597 to -0.476	<0.001
Rural location	0.176 (0.038)	0.101 to 0.251	<0.001	0.175 (0.038)	0.100 to 0.250	<0.001

Model was reduced using stepwise backwards elimination procedure with $p < 0.05$ required for reduced model inclusion. CGM, continuous glucose monitoring; HbA1c, hemoglobin A1c.

income and insurance status observed in this cohort. Previous research supports the association between greater CGM use, and socioeconomic status, as we found in the urban group in this study.^{33–35} For rural families who have youth with type 1 diabetes, there can also be barriers to quality of care such as appointment adherence and communication between patients and providers.²⁶

Our multivariable model controlled for many of the confounding factors already well recognized to influence glycemia in persons with type 1 diabetes: duration of diabetes, CGM use, race/ethnicity, and private insurance. It is important to note that black race, CGM status, and private insurance had larger effects on HbA1c than rurality in the multivariable regression model; however, those factors are already widely acknowledged as important predictors of disparities in type 1 diabetes outcomes.^{7,12,36} Therefore, we believe it is noteworthy that rural status persisted as a significant predictor for HbA1c among this group with type 1 diabetes obtaining care at endocrinology clinics. While rurality has previously been reported as a negative influence on HbA1c, most of this work has been done among the adult population with type 2 diabetes.^{18,37}

As these data were collected and analyzed prior to the COVID-19 pandemic, there is no way to know the impact of the pandemic's widespread adoption of virtual medical visits and telemedicine on the T1D Exchange participant sample we reported on here, especially those living in rural communities. Technology-based approaches to increase and democratize³⁸ diabetes care access among socioeconomically and geographically at-risk rural communities can be effective. Researchers and clinicians should work towards building flexible solutions that meet a wide diversity of needs for rural patient populations with type 1 diabetes.

Limitations

Despite our large, multistate sample, there are limitations. This sample does not include all persons with type 1 diabetes throughout the USA. The data are from those participants who obtained care at endocrinology clinics, consented to participate in the T1D Exchange registry, and therefore does not capture patients with type 1 diabetes who obtained healthcare from other providers (eg, internal medicine, pediatricians, or family practice providers), or patients who chose not to participate in the registry. The data on insurance coverage are very generalized (only reported as private, public, or other), so we were unable to determine more specific information concerning insurance coverage, underinsured, and uninsured status in this sample. Due to the lack of detail about participant education and income, the linear regression model could be biased. We chose to use the CMS methodology for defining rural, but there are many ways to define rural, so these findings should be interpreted with some caution.²² We were not able to obtain psychosocial data (eg, depression, quality of life) for this cohort beyond general health status. Despite these limitations, we believe this study sheds light on an understudied group of rural Americans living with type 1 diabetes.

CONCLUSIONS

In this analysis of a 2-year span of the T1D Exchange clinic registry participants, all receiving care at endocrinology clinics, we found significantly different levels of glycemic control among rural versus urban participants, especially in youth. The demographic differences reported here showed the rural group had lower levels of education, income, and less private insurance coverage. More importantly, our model showed that the disparities in glycemic control in this sample remained significantly associated with rural participant location even after controlling

for other more significantly associated factors including duration of diabetes, CGM use, age, race/ethnicity, and private insurance status. To our knowledge, this is the first study to look specifically at rurality and HbA_{1c} levels among a large sample of the American population with type 1 diabetes. With approximately 20% of Americans living in rural communities, there is still much more to be done to fully address the commonly recognized health disparities observed in rural America, and those living with type 1 diabetes.

Acknowledgements We want to express our gratitude to the Leona M and Harry B Helmsley Charitable Trust and the Jaeb Center for Health Research (JCHR) in Tampa, Florida, USA for providing us with the T1D Exchange registry data. We also appreciate the peer reviews provided prior to submitting for publication by Dr Josh Neumiller and Dr Malcom Cutchin.

Contributors KBE (author guarantor) conceived the research idea and obtained the data from Jaeb Center for Health Research. MDG planned and executed the statistical analysis and generated the tables. All authors contributed to interpreting the results. AG created and presented the research poster for this paper (presented at 2020 ADA Scientific Sessions), drafted the manuscript, and created the tables and figure for the paper. All authors contributed to revisions, and reviewed and approved the final submission. KBE submitted the paper on behalf of the research team.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval All research procedures were reviewed and approved by the Pacific Northwest University (PNWU) Institutional Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. All data relevant to the study are included in the article.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Kathaleen Briggs Early <http://orcid.org/0000-0003-2224-2548>

REFERENCES

- 1 U.S. Census Bureau. New Census Data Show Differences Between Urban and Rural Populations [press release]. American Community Survey: 2011-2015, 2016. Available: <https://www.census.gov/newsroom/press-releases/2016/cb16-210.html> [Accessed Dec 08].
- 2 Coughlin SS, Clary C, Johnson JA, et al. Continuing challenges in rural health in the United States. *J Environ Health Sci* 2019;5:90-2.
- 3 Rogers MAM. Onset of type 1 diabetes mellitus in rural areas of the USA. *J Epidemiol Community Health* 2019;73:1136-8.
- 4 Patterson CC, Karuranga S, Salpea P, et al. Worldwide estimates of incidence, prevalence and mortality of type 1 diabetes in children and adolescents: results from the International diabetes Federation diabetes atlas, 9th edition. *Diabetes Res Clin Pract* 2019;157:107842.
- 5 Callaghan T, Ferdinand AO, Akinlotan MA, et al. The changing landscape of diabetes mortality in the United States across region and rurality, 1999-2016. *J Rural Health* 2020;36:410-5.
- 6 Brundisini F, Giacomini M, DeJean D. Chronic disease patients' experiences with accessing health care in rural and remote areas: a systematic review and qualitative meta-synthesis. *Ont Health Technol Assess Ser* 2013;13:1-33.
- 7 Johnston KJ, Wen H, Joynt Maddox KE. Lack of access to specialists associated with mortality and preventable hospitalizations of rural Medicare beneficiaries. *Health Aff* 2019;38:1993-2002.
- 8 Lu H, Holt JB, Cheng YJ, et al. Population-Based geographic access to endocrinologists in the United States, 2012. *BMC Health Serv Res* 2015;15:541.
- 9 Rutledge SA, Masalovich S, Blacher RJ, et al. Diabetes Self-Management Education Programs in Nonmetropolitan Counties - United States, 2016. *MMWR Surveill Summ* 2017;66:1-6.
- 10 O'Connor MR, Carlin K, Coker T, et al. Disparities in insulin pump therapy persist in youth with type 1 diabetes despite rising overall pump use rates. *J Pediatr Nurs* 2019;44:16-21.
- 11 National Academies of Sciences, Engineering, and Medicine. Achieving rural health equity and well-being: proceedings of a workshop 2018.
- 12 Foster NC, Beck RW, Miller KM, et al. State of type 1 diabetes management and outcomes from the T1D exchange in 2016-2018. *Diabetes Technol Ther* 2019;21:66-72.
- 13 Beck RW, Tamborlane WV, Bergenstal RM, et al. The T1D exchange clinic registry. *J Clin Endocrinol Metab* 2012;97:4383-9.
- 14 Chalew S, Gomez R, Vargas A, et al. Hemoglobin A1c, frequency of glucose testing and social disadvantage: metrics of racial health disparity in youth with type 1 diabetes. *J Diabetes Complications* 2018;32:1085-90.
- 15 Kirkbride K, Wallace N. Rural health clinics and diabetes-related primary care for Medicaid beneficiaries in Oregon. *J Rural Health* 2009;25:247-52.
- 16 Krishna S, Gillespie KN, McBride TM. Diabetes burden and access to preventive care in the rural United States. *J Rural Health* 2010;26:3-11.
- 17 Myers CA, Slack T, Broyles ST, et al. Diabetes prevalence is associated with different community factors in the diabetes belt versus the rest of the United States. *Obesity* 2017;25:452-9.
- 18 Hunt KJ, Davis M, Pearce J, et al. Geographic and racial/ethnic variation in glycemic control and treatment in a national sample of veterans with diabetes. *Diabetes Care* 2020;43:2460-8.
- 19 Beck RW, Miller KM, Foster NC. The T1D exchange clinic network and registry: 10 years of enlightenment on the state of type 1 diabetes in the United States. *Diabetes Technol Ther* 2019;21:310-2.
- 20 Centers for Medicare & Medicaid Services. Details for DME_Rural Zip and Formats. In: *Durable medical equipment PO, and supplies fee schedule*. CMS.gov, 2016. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/DMEPOSFeeSched/DMEPOS-Fee-Schedule-Items/DME-Rural-Zip-and-Formats.html>
- 21 Martino SC, Elliott MN, Dembosky JW. Rural-Urban disparities in health care in Medicare, 2020. Available: <https://www.cms.gov/About-CMS/Agency-Information/OMH/equity-initiatives/rural-health>
- 22 Bennett KJ, Borders TF, Holmes GM, et al. What is rural? challenges and implications of definitions that inadequately encompass rural people and places. *Health Aff* 2019;38:1985-92.
- 23 Geifman N, Cohen R, Rubin E. Redefining meaningful age groups in the context of disease. *Age* 2013;35:2357-66.
- 24 Mercado CI, McKeever Bullard K, Gregg EW, et al. Differences in U.S. rural-urban trends in diabetes ABCs, 1999-2018. *Diabetes Care* 2021;44:1766-73.
- 25 Petitti DB, Klingensmith GJ, Bell RA, et al. Glycemic control in youth with diabetes: the search for diabetes in youth study. *J Pediatr* 2009;155:668-72.
- 26 Stumetz KS, Yi-Frazier JP, Mitrovich C, et al. Quality of care in rural youth with type 1 diabetes: a cross-sectional pilot assessment. *BMJ Open Diabetes Res Care* 2016;4:e000300.
- 27 American Diabetes Association. 6. Glycemic Targets: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S73-84.
- 28 Diabetes Control and Complications Trial Research Group, Nathan DM, Genuth S, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977-86.
- 29 Larsen ML, Horder M, Mogensen EF. Effect of long-term monitoring of glycosylated hemoglobin levels in insulin-dependent diabetes mellitus. *N Engl J Med* 1990;323:1021-5.
- 30 Schwandt A, Best F, Biester T, et al. Both the frequency of HbA_{1c} testing and the frequency of self-monitoring of blood glucose predict metabolic control: A multicentre analysis of 15 199 adult type 1 diabetes patients from Germany and Austria. *Diabetes Metab Res Rev* 2017;33. doi:10.1002/dmrr.2908. [Epub ahead of print: 14 06 2017].
- 31 Coon P, Zulkowski K. Adherence to American diabetes association standards of care by rural health care providers. *Diabetes Care* 2002;25:2224-9.
- 32 Grant JS, Steadman LA. Barriers to diabetes self-management among rural individuals in the workplace. *Workplace Health Saf* 2016;64:243-8.
- 33 Sheikh K, Bartz SK, Lyons SK, et al. Diabetes device use and glycemic control among youth with type 1 diabetes: a single-center, cross-sectional study. *J Diabetes Res* 2018;2018:1-6.

- 34 DeSalvo DJ, Miller KM, Hermann JM, *et al*. Continuous glucose monitoring and glycemic control among youth with type 1 diabetes: international comparison from the T1D exchange and DPV initiative. *Pediatr Diabetes* 2018;19:1271–5.
- 35 Messer LH, Tanenbaum ML, Cook PF, *et al*. Cost, hassle, and on-body experience: barriers to diabetes device use in adolescents and potential intervention targets. *Diabetes Technol Ther* 2020;22:760–7.
- 36 Agarwal S, Schechter C, Gonzalez J, *et al*. Racial-Ethnic disparities in diabetes technology use among young adults with type 1 diabetes. *Diabetes Technol Ther* 2021;23:306–13.
- 37 N C Fu H, Skolnick VG, S Carlin C, *et al*. The effect of depression and rurality on diabetes control. *J Am Board Fam Med* 2020;33:913–22.
- 38 Walker AF, Cuttriss N, Haller MJ, *et al*. Democratizing type 1 diabetes specialty care in the primary care setting to reduce health disparities: project extension for community healthcare outcomes (echo) T1D. *BMJ Open Diabetes Res Care* 2021;9:e002262.
- 39 Cengiz E, Xing D, Wong JC, *et al*. Severe hypoglycemia and diabetic ketoacidosis among youth with type 1 diabetes in the T1D exchange clinic registry. *Pediatr Diabetes* 2013;14:447–54.