



The Tenebrionidae of California: A Time Sensitive Snapshot Assessment

Rolf L. Aalbu¹, Aaron D. Smith²

I Department of Entomology, California Academy of Sciences, 55 Music Concourse Dr., Golden Gate Park, San Francisco, California, 94118 **2** Department of Biological Sciences, Northern Arizona University, PO Box 5640, Flagstaff, AZ, 86011-5640, USA

Corresponding author: Rolf L. Aalbu (raalbu@comcast.net)

Academic editor: P. Bouchard | Received 29 October 2013 | Accepted 1 February 2014 | Published 12 June 2014

http://zoobank.org/D530808E-29A0-444C-B87C-CD01AF10DDF7

Citation: Aalbu RL, Smith AD (2014) The Tenebrionidae of California: A Time Sensitive Snapshot Assessment. In: Bouchard P, Smith AD (Eds) Proceedings of the Third International Tenebrionoidea Symposium, Arizona, USA, 2013. ZooKeys 415: 9–22. doi: 10.3897/zookeys.415.6523

Abstract

Due to a diversity of habitats and its geologic history, the US state of California hosts a spectacular assemblage of darkling beetle species (Coleoptera: Tenebrionidae). In addition to being part of the California Floristic Province, one of 34 global biodiversity hotspots identified by Conservation International, California also has additional areas which are parts of the Great Basin, Mojave, and Sonoran deserts. California is divided into nine floristic regions. Each region is assessed in terms of faunal composition and endemism. A "snapshot" of our present knowledge of the Tenebrionidae indicates that 447 currently recognized species, representing 108 genera, occur in California of which one hundred and ninety are endemic. California is compared to other nearby regions in diversity and endemism. An analysis of currently valid species vs a more realistic species account based on unpublished records of likely synonyms and known species yet to be described in the scientific literature is presented. The California Floristic Region, rather than other more arid parts of California, has the highest number of total and endemic species. Because of their high diversity and endemism, tenebrionids could potentially provide a valuable tool for monitoring the environment for conservation purposes.

Keywords

California, Floristic Regions, Tenebrionidae, Biodiversity, Hotspots, Conservation

Introduction

The state of California is part of the California Floristic Province, one of 34 global biodiversity hotspots identified by Conservation International¹. Over 50 percent of the world's plant species and 42 percent of all terrestrial vertebrate species are endemic to these 34 biodiversity hotspots, a total area which covers only 2.3 percent of the Earth's land surface. The California Floristic Province includes most of western California and a small section of Baja California and Southwestern Oregon. On Conservation International's California Floristic Province website, although numbers of endemic plants, birds, mammals and amphibians are listed, nothing is mentioned concerning insects². California also includes areas not considered to be part of the California floristic province. These areas contain aspects of the Great Basin, Mojave, and Sonoran deserts.

In 2010 the Essig Museum of Entomology at University of California, Berkeley began CalBug (NSF-DBI: 0956389), a collaborative project among nine California museums with a goal to digitize and geographically reference over one million specimens from target groups and localities³. Tenebrionidae was one of the focus groups in Coleoptera. However, to date, few tenebrionids (2%) have been digitized and georeferenced, all at Santa Barbara Museum of Natural History (SBMNH).

In 2005 Mike Caterino, formerly at SBMNH, solicited the author's help in contributing to a web accessible list of "Beetles of California". This was followed by a visit to the SBMNH in 2007 to provide additional identifications of beetles in the collection. The list, last updated in 2009, is posted on http://www.sbnature.org/collections/invert/entom/cbphomepage.php [accessed on December 9, 2013]. An updated list is present here (Fig. 1) that reflects a current "snapshot" of our knowledge of this fauna. It is also available online (http://insectbiodiversitylab.org/CaliforniaDarklingBeetles. html). To account for active research and our growing understanding of the California fauna, the list includes a separate column assessing the potential that each species will be synonymized in future works (see below). Both the current valid species list and a list excluding likely synonymous species, but including known undescribed species, are analyzed based on each species' known occurrence in each of California's nine floristic provinces to assess number of tenebrionid species in each province and their endemicity.

Materials and methods

Sources of Information other than the SBMNH list above include publications from early workers (LeConte, Horn, Motschulsky, Casey, Blaisdell, and others), modern workers

See http://www.conservation.org/where/priority_areas/hotspots/Pages/hotspots_main.aspx [accessed on December 9, 2013].

See http://www.conservation.org/WHERE/PRIORITY_AREAS/HOTSPOTS/NORTH_CENTRAL_AMER-ICA/CALIFORNIA-FLORISTIC-PROVINCE/Pages/default.aspx [accessed on December 9, 2013].

³ See http://calbug.berkeley.edu/data.html [accessed on December 9, 2013].

(Doyen, Triplehorn, Somerby, Brown, Smith, and others) ⁴, and modern revisions: Parts of the Coniontini (Doyen 1984), Cnodalonini (Doyen 1973), Amphidorini (Aalbu et al. 2012, Triplehorn and Thomas 2011), Edrotini (Pape et al. 2007), Stenosini (Papp 1981) and Asidini (Brown and Doyen 1991, Smith 2013) as well as complete revisions of the Cryptoglossini (Aalbu 2005) and Anepsiini (Doyen 1987). Other major sources of information include the Species Database of the California Academy of Sciences and information from the author's personal collection (the Rolf L. Aalbu Collection – RLAC), as well as visits to all major beetle collections in California and many others outside of the state. Information for potential future species synonymies and undescribed species come from the authors' research, discussions with other tenebrionid workers, and currently unpublished studies by the authors, Ron Somerby, and Charles Triplehorn.

To account for the many groups in which data has been accumulated but no recent revision has been published, the Tenebrionidae records from California were categorized in the following status groups based on their current and future status: 0), Known new but undescribed species; 1), Currently projected valid species and subspecies⁵; 2) Most likely synonyms, but synonymy not determined without further study; and 3), Known but unpublished synonyms. Published synonyms were omitted. The assessment was then divided into two categories: A.) Described Species Count: All species currently valid in the literature including known synonyms (groups 1, 2, and 3 above). B), Realistic Species Count: (groups 0, 1, and 2 above). Endemism was calculated on a strict basis (species endemic to specific regions which include parts of adjacent areas not in California were not considered).

For the purpose of this study, California is divided into nine floristic regions modified from a map by the Jepson Herbarium⁶ (Fig. 1). Four of these are not considered parts of the California Floristic Region. These are: Region 1, The Northern Great Basin Province, including the Warner Mountains and Modoc Plateau; Region 2, The Southern Great Basin Province, including the White and Inyo Mountains and intermountain valleys east of the Sierras Nevada's and White Mountains; Region 3, The Mojave Desert and associated desert mountains; and Region 4, The Sonoran (Colorado) Desert and associated desert mountains.

Regions belonging to the California Floristic Region include: Region 5. The South Coast, including the Transverse and Peninsular Ranges and Channel Islands; Region 6, The Sierra Nevada Mountains; Region 7, The Central Valley; Region 8, The Central Coast, including the San Francisco Bay area and Coast Ranges; and Region 9, The Northern Coast, including the Cascade and Klamath Ranges as well as the Northern Coast Ranges. In these regions we examined species occurrence and regional endemism. Regional endemism was also calculated on a strict basis as described above.

⁴ All publications prior to 2002 are listed in Aalbu et al. 2002. Newer pertinent publications are listed in the reference section below.

Some of the early described species as Edrotines etc., described by early workers as Casey may potentially be synonyms.

Geographic subdivisions of California, Jepson Flora Project (eds.) [2013] Jepson eFlora, http://ucjeps.berkeley. edu/IJM.html [accessed on December 9, 2013].

Results and discussion

It is important to keep in mind that this study represents a snapshot in time and thus is subject to change as new information becomes available. However, this assessment is also a balance between future synonymies from previous descriptions (Casey and other early workers: *Coniontis*, various genera of edrotines) on one side and new species discoveries, as well as new foreign introductions, on the other. At present, we know of at least eight distinct new species.

A list of all described species is presented in phylogenetic order (Fig. 1). Differences in group numbers and endemics are presented in Table 1. Differences in species count categories (numbers, endemics and percent endemism) are shown in Table 2. It is notable that despite the differences in numbers, both analyses (described vs realistic) indicate a very similar percent endemism. Since this study is intended as a "snapshot" of our current knowledge, species counts and analysis, unless otherwise specified, include only groups 0, 1, and 2 (Realistic Species Count). This tenebrionid inventory of California thus includes 34 tribes, 118 genera and subgenera, 447 species and subspecies (including known new species). Of these, 190 are endemic to California. The present SBMNH web list includes 471 species from California. Of these, 10 are collection data errors. These included *Argoporis alutacea* Casey; *Asidopsis consentanea* Casey; *Asidopsis planata* (Horn); *Cryptoglossa variolosa* Horn; *Eleodes alticola* Blaisdell; *Eleodes subnitens* LeConte; *Neatus tenebrioides* Beauvois; *Platydema micans* Zimmerman; and *Stenomorpha obovatus* (LeConte) none of which are known to occur in California. Others are known but unpublished synonymies (status group 3).

The fauna is composed of the following subfamilies in descending species number: Pimeliinae (204), Tenebrioninae (168), Alleculinae (33), Diaperinae (23), Stenochiini (11), Lagriinae (7), and Phrenapatinae (1). California is clearly a center of diversity for the family Tenebrionidae, representing 38% of all U.S. species. The most abundant tribes and genera in terms of species numbers are: Amphidorini (73 species), Edrotini (71 species), Coniontini (53 species), Alleculini (33 species), Opatrini (26 species), Asidini (25 species), and Helopini (21 species); and genera such as *Eleodes* (64 species), *Stenomorpha* (19 species), *Coniontis* (38 species), and *Metoponium* and *Helops* each with 21 species. A number of tribes such as Amphidorini, Coniontini, and Nyctoporini, and genera such as *Eleodes*, *Coelocnemis*, *Nyctoporis*, *Asbolus*, *Coniontis*, and *Alaudes* also exhibit their greatest diversity in genera/species in California.

Compared to other known nearby geographical regions, California also has a high species per area diversity (1.05 per 1000 square miles) which is higher than the U.S. as a whole⁷ (.12) or even Mexico⁷ (.68), but not Baja California⁷ which has a species diversity of 5.47 (see Table 3). California shares species with the following adjoining areas in descending order: 1. Southwest U.S.: (including Arizona, 101, Nevada, 76; New Mexico, 23; and Utah, 42). 2. Mexico (mainland 32, Baja California, 68) and 3. Northwest U.S. (including Oregon, 56; Washington, 33; and Idaho, 32. A number of

Numbers probably 5–8 years old.

Table 1. Status Groups and Endemicity. Group 0: Known new but undescribed species; Group 1: currently projected valid species and subspecies; Group 2: most likely synonyms, but synonymy not determined without further study; and Group 3: known but unpublished synonyms.

Status group	Non endemic species	Endemic species	Total
0	2	8	10
1	249	155	404
2	6	27	33
3	22	17	39
	279	207	486

Table 2. California Species, Described vs Realistic. Species counts for the state and % endemicity based on current valid species (A) and a realistic estimation of actual species counts (B).

Category	Status groups	Species	Endemics	Total	% Endemic
A: Described	1, 2 & 3	277	199	476	41.81%
B: Realistic	0, 1, & 2	257	190	447	42.51%

Table 3. Comparison of currently valid species/endemics per area for various regions.

Region	Number of species	Number of endemics	% Endemism	Area (km²)	Species diversity per 1000 km ²
California*	447**	190	43%	423970	1.05
USA***	1184	?	>60%	9827000	0.12
Mexico***	1340	723	54%	1973000	0.68
Baja California***	404	225	56%	73909	5.47

^{*} Bordered by 3 states and Baja California.

species are known only from the type and have undetermined California localities (16). Twenty species are cosmopolitan pests. See Fig. 1 for additional locality information.

The distribution of California tenebrionids can be divided into six patterns: 1), Widespread species, 2), Restricted but not especially hard to collect species (Caves, single canyons (Eschatomoxys andrewsi Aalbu & Thomas, Eleodes (Caverneleodes) microps Aalbu et al.), 3), Restricted but very difficult to collect species (Eleodimorpha, Oxygonodera), 4), Historically abundant but now difficult to collect species (Eleodes (Melaneleodes) quadricollis Eschscholtz), 5) Introduced species composed of standard stored product pests as well as other introductions not associated with stored products (Opatroides punctulatus Brullé and Gonocephalum sp.) and 6) species only known form the type material with specific locality unknown. California also has some unusual darkling beetle occurrences and absences compared to the rest of North America. One is the presence of two species from the Asian tribe Laenini, which is otherwise absent on the continent. Another is the absence of the genus Strongylium, a species-rich genus found worldwide including in Arizona (2 species) and most of the rest of the United States.

^{** 34%} of all U.S. species.

^{***} numbers probably 5–8 years old.

Regional analysis

For the purpose of this study, California was into 9 floristic regions (Fig. 2) to examine species occurrence and regional endemism. Regional endemism was also calculated on a strict basis as mentioned above. A list of all regional endemics is presented as well as total species numbers for the region and percent endemism (Fig. 3). These areas are ranked in Table 4. Adding the above data suggests that over 62% (62.11) of the endemic species in California are regional endemics while 43% (42.60) of all tenebrionids are endemic in terms of being regional endemics or multiple region endemics.

One may note that, somewhat surprisingly, subregions within the California Floristic Region have more regional endemic species (87) as well as California endemic species (124) despite the common association of tenebrionids with desert habitats, where they are always abundant (see Table 5). On this table, "all endemics" in the "unknown...." region refer to species where the type locality is simply listed as "California". This "snapshot" assessment emphasizes how much remains to be done in this area, especially in revising tribes or genera which have not been looked at since their description, as well as rediscovering species of "unknown" California localities. Additional new species, as well as new introductions, will undoubtedly be discovered as well. It is hoped that this type of assessment can be useful in environment monitoring and conservation studies.

Tabl	e 4.	Com	parison	of	regional	enc	lemics	and	all	end	lemics	for	Califor	rnia.

Region	Endemic species	All species	% Endemic	% of all California Endemics
5. South Coast & Islands	42	171	24.56%	35.59%
8. Central Coast & Bay	20	110	18.18%	16.95%
6. Sierra Nevada	16	100	16.00%	13.56%
4. Sonoran Desert	13	113	11.50%	11.02%
3. Mojave Desert	12	112	10.71%	10.17%
8. Central Valley	5	76	6.58%	4.24%
2. South Great Basin	5	55	9.09%	4.24%
9. North Coast	4	73	5.48%	3.39%
1: North Great Basin	1	29	3.45%	0.85%

Table 5. Comparison of species endemicity for California Floristic affinities.

Floristic Region	All Endemics	Non Endemic	All Species
Desert Areas	37	94	131
California Floristic Province	124	81	205
Both Areas	16	60	76
Unknown California locality, cosmopolitan or introduction	13	22	35

Figures

Endemic Status

group Pimeliinae Cnemeplatiini 1 Alaudes alternata Fa 3 Alaudes fallax Fall,

Pimeliinae Cryptoglossini

Pimeliinae Nyctoporini

Pimeliinae Asidini

1 Typhleusechus singularis Linell, 1897

1 Asbolus mexicanus angularis (Horn), 1894

1 Asbolus papillosus (Triplehorn), 1964

1 Cryptoglossa muricata (LeConte) 1851

1 Schizillus nunenmacheri Blaisdell, 1921

1 Nyctoporis aequicollis Eschscholtz,1831

1 Nyctoporis carinatus LeConte, 1851

1 Nyctoporis cristata Eschscholtz,1831

3 Nyctoporis maura Casey, 1907

3 Nyctoporis pullata Casey, 1907

1 Nyctoporis sponsa Casey, 1907

3 Nyctoporis tetrica Casey, 1907

1 Nyctoporis vandykei Blaisdell, 1931

1 Microschatia championi Horn, 1893

1 Philolithus actuosus (Horn) 1870

2 Philolithus jaegeri (Papp), 1961

1 Philolithus opimus Casey, 1912

3 Philolithus porcatus (Papp), 1961

3 Philolithus rugosus (Papp), 1961

1 Stenomorpha (Asidina) confluens (LeConte), 1851

1 Philolithus carinatus LeConte, 1851

1 Microschatia inaequalis LeConte. 1851

1 Cryptoglossa spiculifera (LeConte), 1861

1 Asbolus verrucosus LeConte, 1851

1 Asbolus laevis LeConte, 1851

1 Schizillus laticeps Horn, 1874

Pimeliinae Stenosini

CA

3

6

10

12

13

14

15

16

17

18 19

20

21

22

23

25

26

27

28

29

30

32

33

35

36

37

39

40

42

43

44

46

47

•

group	Species	Distribution	#	CA	group	Species	Distribution
nae Cnemeplatiin	i		50		1 Stenomorpha	(Asidina) parallela (LeConte), 1851	3 4 AZ
1 Alaudes alternata	Fall 1929	5	51		1 Stenomorpha	(Asidina) semilaevis (Horn), 1870	3 4 NV AZ ME
3 Alaudes fallax Fa		5	52		1 Stenomorpha	(Asidina) wickami (Horn), 1894	4 AZ
O Alaudes n.sp. Col	•	3 4 5	53	✓	1 Stenomorpha	(Pycnomorpha) angulata (LeConte), 1851	58
Alaudes n.sp. Fre		57	54		1 Stenomorpha 1870	(Pycnomorpha) californica (Motschulsky),	5 BC
O Alaudes n.sp. Moj	ave	23	55		1 Stenomorpha	(Pycnomorpha) cressoni (Blaisdell), 1933	4 AZ
 Alaudes setigera 	Blaisdell, 1919	5678BC	56	H		(Pycnomorpha) gibicollis (LeConte), 1851	5 BC
 Alaudes singularis 	Horn, 1870	234568 NV AZ ID	57	~	3 Stenomorpha	(Stenomorpha) amplicollis (Casey), 1912	678
3 Alaudes testacea	Blaisdell, 1919	8	58	V	1 Stenomorpha	(Stenomorpha) captiosa (Horn), 1870	678
 Lepidocnemeplati 	a n.sp.	4	59	V	3 Stenomorpha	(Stenomorpha) compressa (Horn), 1870	7
 Lepidocnemeplati 	a sericea (Horn), 1870	67958AZ OR ID ME	60	V	3 Stenomorpha	(Stenomorpha) crassa (Casey), 1912	678
		NV UT +	61	~	3 Stenomorpha	(Stenomorpha) deceptor (Casey), 1912	678
nae Stenosini			62	~	3 Stenomorpha	(Stenomorpha) evanescens (Casey), 1912	23497
1 Araeoschizus and	rewsi Papp, 1981	4	63	✓	3 Stenomorpha	(Stenomorpha) gravidipes (Casey), 1912	678
1 Araeoschizus arm	atus Horn, 1870	3 2 NV	64	✓	3 Stenomorpha	(Stenomorpha) integer (Casey), 1912	678
1 Araeoschizus cos	tipennis LeConte, 1851	5 4	65	✓	3 Stenomorpha	(Stenomorpha) lecontei (Horn), 1866	23497
1 Araeoschizus doy	eni Papp, 1981	5	66	~	3 Stenomorpha	(Stenomorpha) lecontella Blaisdell, 1936	23497
1 Araeoschizus exig	uus Casey, 1907	5	67		1 Stenomorpha	(Stenomorpha) marginata (LeConte), 1851	3 AZ NM
1 Araeoschizus hard	dyi Papp, 1981	4	68	✓	3 Stenomorpha	(Stenomorpha) maritima (Casey), 1912	678
1 Araeoschizus hys	trix Papp, 1981	4	69	✓	1 Stenomorpha	(Stenomorpha) mckittricki (Pierce), 1944	7
1 Araeoschizus kas:	zabi Papp, 1981	53	70	✓	3 Stenomorpha	(Stenomorpha) neutralis (Casey), 1912	23497
1 Araeoschizus lariv	versi Papp, 1981	3	71	✓	3 Stenomorpha	(Stenomorpha) speculata Blaisdell, 1936	23497
1 Araeoschizus sim	ulans Casey, 1907	U type only	72	✓	3 Stenomorpha	(Stenomorpha) spurcans (Casey), 1912	23497
1 Araeoschizus sulc	icollis disjunctus Papp, 1981	8	73	✓	3 Stenomorpha	(Stenomorpha) subelegans (Casey), 1912	
1 Araeoschizus sulc	icollis sulcicollis Horn, 1870	3 NV	74	✓	3 Stenomorpha	(Stenomorpha) tularensis (Casey), 1912	678
1 Typhleusechus ch	emehuevii Aalbu & Andrews, 1985	3	75		3 Stenomorpha	(Stethasida) adumbrata Casey, 1912	5 BC AZ

77

78

80

81

83

84

85

87

AA

89

91

92

94

95

97

QR

101

102

104

103 🗹

Pimeliinae Coniontini

4 BC AZ

4 BC AZ

783452AZNV ME

7534A7NV MF

23457BC

23 NV AZ UT

56798NV

69875BC

6789

6

9

976

6758

4 BC

5 BC

3 AZ

3

3

3

3 AZ

34AZNV

3 4

4 ME

3 Stenomorpha (Stethasida) angustula (Casey), 1890

3 Stenomorpha (Stethasida) convergens Casey, 1912

3 Stenomorpha (Stethasida) brevines Casev 1912

3 Stenomorpha (Stethasida) discreta Casey, 1912

3 Stenomorpha (Stethasida) longula Casey, 1912

3 Stenomorpha (Stethasida) pertinax Casey, 1912

3 Stenomorpha (Stethasida) socors Casey, 1912

3 Stenomorpha (Stethasida) stricta Casey, 1912

3 Stenomorpha (Stethasida) tarsalis Casev. 1912

3 Stenomorpha (Stethasida) tenax Casey, 1912

3 Stenomorpha (Stethasida) torpida Casey, 1912

1 Stenomorpha (Stethasida) unica Casey, 1912

3 Stenomorpha (Stethasida) vergrandis Casey, 1912

1 Stenomorpha (Trichiasida) hirsuta (LeConte), 1851

1 Stenomorpha (Trichiasida) impetrata (Horn), 1894

1 Stenomorpha (Trichiasida) luctata (Horn), 1870

1 Coelus ciliatus Eschscholtz, 1829

1 Coelus globosus LeConte, 1851

1 Coniontis abdominalis LeConte, 1859

1 Coelus gracilis Blaisdell 1939

1 Coniontis affinis LeConte, 1851

1 Conjontis blaisdelli Casev. 1908

1 Coelus pacificus Fall, 1897

1 Stenomorpha (Trichiasida) hispidula (LeConte), 1851

3 Stenomorpha (Trichiasida) lineatopilosa Casey, 1912

1 Stenomorpha (Trichiasida) acerba (Horn), 1878

3 Stenomorpha (Stethasida) laevigatus (Papp), 1961

1 Stenomorpha (Stethasida) muricatula (LeConte), 1851

1 Stenomorpha (Stethasida) obsoleta (LeConte), 1851

5 BC AZ

5 BC

5 BC

5 BC

5 BC

5 BC

5 BC

5 BC AZ

5 BC AZ

5 BC AZ

3 4 AZ UT

34 AZ UT

5 8 9 OR WA CA

4 AZ

5 8 BC

5

58

896

6789AZ

Figure 1. Checklist of the California Tenebrionidae species with distributions and likelihood for future
synonymy. Distribution numbers refer to California regions (Fig. 2) and the following: ME (Mexico)
BC (Baja California) NV (Nevada) AZ (Arizona) ID (Idaho) UT (Utah) NM (New Mexico) OR (Oregon)
WA (Washington) CA (Canada) U (unknown California distribution) C (refers to cosmopolitan pest),
ASIA SA (South America), and OW (Old World).

Endemic	Endemic Status Endemic Status						
# CA	group	Species	Distribution		CA	group Species	Distribution
105 🔲	1 Coniontis c	allida Casey, 1908	6 9 CA WA OR NV	159		1 Chilometopon brachystomum Doyen, 1982	3 4 NV BC
106 🗸		atalinae Casey, 1908	5	160		1 Chilometopon helopoides Horn, 1874	2 3 4 AZ ID NV NM UT
107 🗸		ostulata Casey, 1908	8	404		4 Obiles stores wisses Mark sables 8 Oless 4000	BC
108 🗸		lliptica Casey, 1884	4567	161	✓	1 Chilometopon microps MacLachlan & Olson, 1990	34
109 🗸		longata Casey, 1890	4567891	162	Ш	1 Chilometopon pallidum Casey, 1890	3 4 AZ NV NM TX ME BC
110 🗸	1 Coniontis e	schscholtzi Mannerheim, 1840	U	163	П	1 Cryptadius inflatus LeConte, 1851	5 BC
111 🔲		xtricata Casey, 1908	589 OR	164	H	1 Edrotes arens La Rivers, 1947	5 AZ
112 🗸		arallonica Casey, 1895	8	165	H	1 Edrotes ventricosus LeConte, 1851	3 4 5 AC BC ME ID NV
113 🗸		enitiva Casey, 1890	9	166	~	1 Eurymetopon ochraceum Eschscholtz,1831	U type only
114 🗸	-	lobulina Casey, 1895	5	167	Ĭ	1 Eurymetopon rufipes Eschscholtz, 1831	U type only
115 🗸		oppingi Blaisdell, 1918	6	168	~	1 Hylocrinus blaisdelli Casey, 1890	76
116 🗸		nteger Casey, 1908	57	169	•	1 Hylocrinus depressulus Casey, 1907	U
117 🗸		amentabilis Blaisdell, 1924	587	170		1 Hylocrinus filitarsis Casey, 1907	5 4 8
118 🗌		assenica Casey, 1908	9 1 5 NV ID	171		1 Hylocrinus longulus LeConte, 1851	56358AZ
119 🗸		ata LeConte, 1866	8	172		1 Hylocrinus oblongulus Casey, 1907	4 BC
120 🗌		nicrosticta Casey, 1908	8 ID		~	1 Hylocrinus piceus Casey, 1890	5
121 🔽		nusculus Blaisdell, 1918	U	174		2 Melanastus aequicollis Casey,1907	9 NV
122 🗌		emoralis Eschscholtz, 1840	8567OR	175	V	1 Melanastus ater (LeConte), 1851	98
123		besa LeConte, 1851	12569	176		2 Melanastus crassicornis (Casey), 1907	9 OR
124		paca Horn, 1870	2 NV WA	177	V	2 Melanastus exoletus Casey,1907	5
125 🗌	1 Coniontis o	valis LeConte, 1851	12679AZCACO ID NV OR WA UT	178		2 Melanastus lucidulus Casey,1907	8
126 🗸	1 Conjuntis n	allidicornis Casey, 1890	5	179	П	2 Melanastus moestus Casey,1907	8 UT
127		arallela Casey, 1890	U	180	$\overline{\Box}$	1 Melanastus obesus (LeConte), 1851	587BC
128		arviceps Casey, 1890	5 8 7 BC	181	~	1 Melanastus obtusus (LeConte), 1866	587
129 🔽		ectoralis Casey, 1908	U	182	V	2 Melanastus otiosus Casey,1907	8
130		uncticollis LeConte, 1851	56789	183	~	2 Melanastus sterilis Casey,1907	3
131		emnans Pierce, 1945	5	184	~	2 Melanastus thoracicus (Casey), 1907	5
132		obusta Horn, 1870	5678	185	✓	1 Melanastus vergrandis Casey,1907	6
133		anfordi Blaisdell, 1895	689	186		1 Metoponium abnorme LeConte, 1851	4 ME
134 🗸		antarosae Blaisdell, 1921	5	187		1 Metoponium bicolor Horn, 1870	7 3 AZ
135		etosa Casey, 1890	6 OR WA CA ID UT	188	V	2 Metoponium concors Casey,1907	3
136		ubpubescens LeConte, 1851	83569 OR BC	189		2 Metoponium convexicolle LeConte, 1851	7 5 3 AZ
137		mida Casey, 1908	8	190	~	2 Metoponium cylindricum Casey, 1890	3
138 🔽		andykei Blaisdell, 1921	5	191	✓	2 Metoponium dubium Casey, 1884	4
139 🔽		iatica Eschscholtz, 1929	8	192	~	2 Metoponium edax Casey,1907	5
140	1 Eusattus co	ostatus Hom, 1870	5 BC	193	~	2 Metoponium egregium Casey,1907	4
141	1 Eusattus di	fficilis LeConte, 1851	3 4 5 7 ME AZ NV	194	✓	2 Metoponium extensum Casey,1907	U
142	1 Eusattus di	latatus LeConte, 1851	4 3 AZ ME	195	✓	2 Metoponium faustum Casey,1907	3 4
143	1 Eusattus du	ubius arizonensis Doyen, 1984	3 4 AZ NV	196		2 Metoponium fusculum Casey, 1890	4 AZ
144	1 Eusattus du	ubius dubius LeConte, 1851	3 4 2 NV AZ UT	197	✓	2 Metoponium gravidum Casey,1907	3
145	1 Eusattus m	uricatus muricatus LeConte, 1851	123495 AZ NM TX	198	✓	2 Metoponium gulosum Casey,1907	7
			CO UT NV ID OR WA	199		2 Metoponium hebes Casey,1907	4 AZ
146		hreatophilus Doyen, 1984	3 2 NV	200	✓	2 Metoponium insulare Casey,1907	5
147 🗸		olitus Horn, 1883	5	201	✓	2 Metoponium integer Casey,1907	4
148 🗌		roductus LeConte, 1858	4 ME	202	•	2 Metoponium molestum Casey,1907	4
149 🗌		ticulatus (Say), 1824	4 AZ	203	•	2 Metoponium opacipenne Casey,1907	3 4
150 🗸		bustus LeConte,1866	5	204	•	2 Metoponium probatum Casey, 1907	5
Pimeliina	ae Edrotini			205	✓	2 Metoponium tersum Casey,1907	5
151 🔽	1 Auchmobiu	s angelicus Blaisdell. 1934	5	206	✓	2 Metoponium testaceum Casey,1907	3
152 🗸		s parvicollis Blaisdell. 1934	8	207		1 Micromes maritimus (Casey), 1891	5 BC
153		s picipes Blaisdell. 1934	8	208		1 Micromes ovipennis (Horn), 1870	5 BC
154		s sanfordi Blaisdell. 1934	5	209		O Oxygonodera n.sp.	2
155 🔽		s slevini Blaisdell. 1934	8	210	✓	2 Stibia blairi Blaisdell, 1936	U type only (BMNH)
156		s subboreus Blaisdell. 1934	8	211		1 Stibia imperialis Blaisdell, 1936	4 BC
157 🗸		s sublaevis LeConte, 1851	2	212		1 Stibia puncticollis Horn, 1870	5 BC
158		on abnorme (Horn), 1870	3 4 AZ NV ME	213		3 Stibia tanneri Blaisdell, 1936	6
				214	✓	1 Telabis fidelis Casey, 1907	4

Figure I. Continue.

Endemic Status	1 chebi ioniu	Endemic Status	
# CA group Species	Distribution	# CA group Species	Distribution
215 1 Telabis incisa Casey, 1907	4 NV	Diaperinae Diaperini	
216 🕡 1 Telabis opacella Casey, 1907	5	257 1 Alphitophagus bifasciatus (Say), 1824	С
217 1 Telabis punctulata LeConte, 1866	3 AZ	258 1 Cynaeus angustus (LeConte), 1851	5 AZ UT ME
218 1 Telabis serrata (LeConte), 1866	3 4 BC ME AZ UT NV	259 1 Cynaeus depressus Horn, 1870	5 AZ BC
219 1 Telabis sodalis Horn, 1870	3 4 AZ	260 1 Diaperis californica Blaisdell, 1929	6
220 1 Triorophus laevis LeConte, 1851	3 4 AZ NV ME UT BC	261 1 Pentaphyllus californicus Horn, 1870	6
221 1 Triorophus rugiceps LeConte, 1851	237865NV	262 1 Platydema americanum Laporte & Brulle, 1831	6 AZ ID CA NM TX UT
222 1 Triphalopsis californicus Doyen, 1982	4 BC		WA +
Pimeliinae Epitragini		263 🗹 1 Platydema neglectum Triplehorn, 1965	1652
223 1 Bothrotes plumbeaus tenebriosus Casey, 1907	5 AZ	264 1 Platydema oregonense LeConte, 1857	5679 CA OR WA
224 1 Metopoloba pruinosa pruinosa (Horn), 1870	3 4 AZ NV ME BC	265 1 Sitophagus hololeptoides Laporte, 1840	С
225 1 Polemiotus submetallicus (LeConte), 1854	4 AZ ME	Diaperinae Crypticini	
Pimeliinae Anepsiini		266 1 Gondwanocrypticus platensis (Fairmaire), 1883	C5SA
226 1 Anchomma costatum LeConte, 1858	3567NV	Diaperinae Myrmechixenini	
227 1 Anepsius delicatulus LeConte, 1851	8 5 7 2 3 4 BC AZ UT	267 1 Myrmechixenis lathridioides Crotch, 1873	68 TX +
	NV	Diaperinae Hypophloeini	
228 1 Batuliodes obesus Doyen, 1987	3 4	. ,,,	
229 1 Batuliodes rotundicollis (LeConte), 1851	3 4 BC AZ	268 1 Corticeus hatchi Boddy, 1957	9 OR
230 1 Batuliodes spatulatus Doyen, 1987	3 4 BC AZ	269 1 Corticeus opaculus (LeConte), 1878	8 5 AZ
231 1 Batuliodes wasbaueri Doyen, 1987	4 BC	270 1 Corticeus praetermissus (Fall), 1926	12456789AZBC NM UT NV ID OR WA
232 1 Batuliomorpha comatus Doyen, 1987	3 BC		CA+
233 1 Batuliomorpha imperialis Doyen, 1987 234 1 Batulius setosus LeConte, 1851	4 3 4 BC AZ	271 1 Corticeus strublei Blaisdell, 1934	369 WA ID UT AZ NIV
Pimeliinae Vacronini	0 4 DO AZ	270 - 4 0-4	+ 40056700MEA7
		272 1 Corticeus substriatus (LeConte), 1852	12356789 ME AZ NM UT NV ID WA OR
235 1 Alaephus gracilis Fall, 1905	853 AZ		CA+
236 🕡 1 Alaephus longicornis Casey, 1924	5	273 1 Corticeus tenuis (LeConte), 1878	6 OR WA CA AZ ID +
237 1 Alaephus macilentus Casey, 1891	3 ME NM AZ	Tenebrioninae Bolitophagini	
238 1 Alaephus maderensis Casey, 1924	5	274 1 Eleates explanatus Casey, 1890	679 OR WA CA ID
239 1 Alaephus pallidus Horn, 1870	762UT	275 1 Eleates occidentalis Casey, 1886	68 WA
240 1 Alaephus puberulus Fall, 1907	5 UT	276 1 Megeleates sequoiarum Casey,1895	96753BC OR
241 1 Eupsophulus castaneus Horn, 1870 Pimeliinae Cnemodini	3 4 ME BC AZ	Tenebrioninae Opatrini	
	2447	277 1 Blapstinus brevicollis LeConte, 1851	2345678AZ
242 1 Cnemodinus testaceus Horn, 1870	3 4 AZ	278 1 Blapstinus castaneus Casey, 1890	5 AZ
Lagriinae Lagriini		279	5
243 🕡 1 Statira dumalis Parsons,1973	4	280 1 Blapstinus dilatatus LeConte, 1851	2345678 AZ NV UT
244 1 Statira latitator Parsons,1973	4 BC	281 1 Blapstinus discolor Horn, 1870	987654321NV OF
245 1 Statira pluripunctata Horn, 1888	4 AZ NM TX ME		ID UT WA CA
Lagriinae Goniaderini		282	4578 AZ NV
246 O Eschatoporis n.sp. Aalbu m.s.	9	 283 1 Blapstinus lecontei Mulsant & Rey, 1859 	5
247 1 Eschatoporis nunenmacheri Blaisdell, 1906	8	284 1 Blapstinus pimalis Casey, 1885	58 AZ NM TX CO UT
Lagriinae Laenini	0	285 1 Blapstinus pulverulentus Mannerheim, 1843	NV 9 OR WA
		286 1 Blapstinus substriatus Champion, 1885	196 AR NV NM TX CC
248 1 Eschatomoxys andrewsi Aalbu & Thomas, 2007	4	200 T biapsinus substitutus Onumpon, 1000	UT WY ID OR WA MT
249 1 Eschatomoxys wagneri Blaisdell, 1935	3 4 AZ UT		CA
Phrenapatinae Penetini		287 1 Blapstinus sulcatus LeConte, 1851	2345678 AZ NV
250 1 Clamoris americana (Horn), 1874	9 4 OR	288 1 Blapstinus vandykei Blaisdell, 1942	2345AZ
Diaperinae Phaleriini		289 1 Cheirodes californica (Horn), 1870	2347 AZ NV
		. 290 1 Conibiosoma elongatum Horn, 1870	3574
251 1 Phaleria debilis LeConte, 1866	4 ME	291 O Conibius n.sp.	7 OR
252	5 8 BC	292 1 Conibius seriatus LeConte, 1851	34578BC
253 1 Phaleromela globosa (LeConte), 1857	89 CA OR WA	293 1 Gonocephalum sp.	5 OW
254 1 Phaleromela humeralis (Laporte), 1840	3 ASIA	294 1 Mecysmus angustus (LeConte), 1851	3 4 5 8 BC
255 1 Phaleromela prohumeralis Triplehorn, 1961	UOR	295 1 Mecysmus tenuis Casey,1890	5 BC
256 1 Phaleromela variegata Triplehorn, 1961	9 4 CA OR WA	296 1 Nocibiotes crassipes (Casey), 1890	4 BC
		297 1 Nocibiotes granulatus (LeConte), 1851	3 4 AZ

Figure I. Continue.

Ender		Status		Enden	ic Status		
# CA	١.	group Species	Distribution	# CA	group	Species	Distribution
298	7 1	Notibius laticeps Casey, 1890	4 AZ BC	349	1 Eleodes (Lithel	leodes) corvinus Blaisdell 1921	1
299	ī 1	Notibius puberulus LeConte, 1851	3 4 AZ	350	1 Eleodes (Lithel	leodes) granulatus LeConte 1857	9 OR
300 🗀	ī 1	Notibius puncticollis LeConte, 1851	753 NV	351	1 Eleodes (Lithel	leodes) letcheri Blaisdell 1909	1 2 NV ID
301 🗀	ī 1	Opatroides punctulatus Brullé, 1832	7 OW	352	1 Eleodes (Lithel	leodes) papillosus Blaisdell 1917	9
302 🗀	ī 1	Tonibius sulcatus LeConte, 1851	35BC	353	1 Eleodes (Lithel	leodes) vandykei Blaisdell 1909	19 OR WA NV AR
303	ī 1	Trichoton sordidum (LeConte), 1851	4 AZ BC ME NM	354	1 Eleodes (Mela	neleodes) carbonarius (Say) 1823	3 4 5 ME AR UT NV B
304		I Ulus crassus LeConte, 1851	857 BC ME ZA NM TX		,	(,,	TX NM ID
	_	ninae Amphidorini		355	,	neleodes) humeralis LeConte 1857	9 CA WA OR NV CO L
			2	356 🗸	,	neleodes) quadricollis Eschscholtz 1833	8
305	_	Eleodes (Blapylis) aristatus Somerby 1977	_	357		neleodes) rileyi Casey 1891	2 NV AZ UT NM ID OF
306			5	358 🗸		blapylis) aalbui Triplehorn 2007	2
307	_	Eleodes (Blapylis) blanchardii Blaisdell 1909	56AZ	359	1 Eleodes (Metal	blapylis) californicus Blaisdell 1929	35AZ
308] 1	Eleodes (Blapylis) brunnipes Casey 1890	6 9 AZ NM CO UT NV OR WA ID WY	360	1 Eleodes (Metal	blapylis) dissimilis Blaisdell 1909	3 ME NV AZ
309 🗆	7 4	I Eleodes (Blapylis) caseyi Blaisdell 1909	25 NV	361 🗸	O Eleodes (Metal	blapylis) n.sp.	5
	_		58	362	1 Eleodes (Metal	blapylis) nigrinus LeConte 1858	12 CA AZ CO ID MT
310	_	I Eleodes (Blapylis) clavicornis Eschscholtz 1833	568				NV NM OR UT WA WY
311		I Eleodes (Blapylis) consobrinus LeConte 1851		363		deleodes) granosus LeConte 1866	2 3 4 NV ID
312 🗸		Eleodes (Blapylis) cooperi Somerby & Doyen 1976	6	364		deleodes) inyoensis Tanner 1961	2 NV
313 _	_	Eleodes (Blapylis) cordatus Eschscholtz 1863	9685OR	365 🗸	1 Eleodes (Stene	eleodes) gigantea Mannerheim 1843	6789
314 🗸	_	l Eleodes (Blapylis) fuchsii Blaisdell 1909	58	366	1 Eleodes (Trich	eleodes) hirsutus LeConte 1861	23 NV
315	_	l Eleodes (Blapylis) hoppingi Blaisdell 1909	6 2 NV AR	367	1 Eleodes (Trich	eleodes) obesus Doyen 1985	9 1 OR
316 🗸	_	l Eleodes (Blapylis) hornii Blaisdell 1909	621	368	1 Eleodes (Trich	eleodes) pilosus Horn, 1870	1239 NV UT
317 🗸	1	Eleodes (Blapylis) incultus LeConte 1861	5	369 🗸	1 Eleodimorpha	bolcan Blaisdell, 1909	5
318 🗸	1	Eleodes (Blapylis) kaweanus Blaisdell 1933	6	370	1 Embaphion de	pressum LeConte, 1858	43 AZ
319 🗸	1	Eleodes (Blapylis) lariversi Somerby & Doyen 1976	6	371	1 Embaphion eld	ingatum Horn, 1870	2 NV
320 🗸	1	Eleodes (Blapylis) littoralis Eschscholtz 1831	5678	372	O Lariversius n.s	p.	3 NV
321] 1	Eleodes (Blapylis) neotomae Blaisdell 1909	5 5 BC	373		alis Blaisdell, 1947	2 NV
322	1	Eleodes (Blapylis) nigropilosa (LeConte) 1851	58BC	374	1 Neobaphion el	ongatum Blaisdell, 1933	2 NV
323 🔽	1	Eleodes (Blapylis) panamintensis Somerby 1977	3	375		status LeConte. 1879	3 2 NV
324	1	Eleodes (Blapylis) parvicollis Eschscholtz 1829	6	376		status mayhewi Papp, 1961	3 NV
325		Eleodes (Blapylis) pimelioides Mannerheim 1843	1358 CA AZ CO ID	377		status pappi Kulzer, 1960	3
	_		OR UT WA WY	•	oninae Cereno		v
326 _] 1	Eleodes (Blapylis) propinquus Blaisdell 1918	1 OR	Tellebi		<u> </u>	
327 🗸	9 1	Eleodes (Blapylis) scabripennis LeConte 1859	56	378	 Cerenopus cor 	ncolor LeConte, 1851	3 4 ME AZ
328 🗸	1	Eleodes (Blapylis) scabriventris Blaisdell 1933	678	Tenebri	oninae Eulabii	ni	
329] 1	Eleodes (Blapylis) scabrosus Eschscholtz 1833	8 9 OR WA	379	1 Ansona harbar	ae Blaisdell, 1932	5
330 🗸	1	Eleodes (Blapylis) schlingeri Somerby & Doyen 1976	6	•		ae Bialsdell, 1932 cornis (Casey), 1890	5
331 🔽	1	Eleodes (Blapylis) spilmani Somerby & Doyen 1976	9	•		· //·	
332	1	Eleodes (Blapylis) subtestitus Blaisdell 1939	5	381		(LeConte), 1866	5
333		Eleodes (Blapylis) tenebrosus (Horn) 1870	126 NV UT OR ID	382 🗸		nis (Casey), 1891	5
334	ī 1	Eleodes (Blapylis) versatilis Blaisdell 1921	9 OR	383	1 Apsena leachi		6
335		Eleodes (Blapylis) volcanensis Somerby 1977	9 OR	384		cens (LeConte),1851	5 BC
_	_		2	385 🗸	 Apsena rufipes 	rufipes (Eschscholtz), 1829	5678
336							
-		I Eleodes (Caverneleodes) microps Aalbu et al., 2012		386		simplex Blaisdell, 1932	8
337	<u> </u>	Eleodes (Cratidus) osculans (LeConte) 1851	58BC			simplex Blaisdell, 1932 alis californica Berry, 1851	8 4 AZ ME
337 <u> </u>] 1	Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833	58BC 658	386	1 Argoporis apic		•
337 = 338 2 339 =] 1] 1	Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858	5 8 BC 6 5 8 6 7 OR NV	386 🔽 387 🗌	1 Argoporis apica 1 Argoporis bicol	alis californica Berry, 1851	4 AZ ME
337	1 9 1 1 1	Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851	5 8 BC 6 5 8 6 7 OR NV 8 5 BC	386 ✓ 387 □ 388 □	Argoporis apica Argoporis bicol Epantius obscu	alis californica Berry, 1851 lor (LeConte), 1851	4 AZ ME 3 4 AZ
337	1 9 1 1 1	Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858	58BC 658 67 OR NV 85BC 782345AZ NV BC	386	Argoporis apica Argoporis bicol Epantius obscu	alis californica Berry, 1851 lor (LeConte), 1851 urus LeConte, 1851 ata Eshscholtz, 1829	4 AZ ME 3 4 AZ 5 8 9 BC
337	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851	5 8 BC 65 8 67 OR NV 8 5 BC 7 8 2 3 4 5 AZ NV BC ME BC	386 387	Argoporis apice Argoporis bicol Epantius obscu Eulabis bicarin Oninae Ulomir	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshscholtz, 1829 ii	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851	58BC 658 67 OR NV 85BC 782345AZ NV BC	386 387 388 389 390 390 391 391	Argoporis apica Argoporis bicol Epantius obscu Eulabis bicarin Oninae Ulomir Uloma longula	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshschottz, 1829 ii LeConte, 1861	4 AZ ME 3 4 AZ 5 8 9 BC
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851	5 8 BC 6 5 8 6 7 OR NV 8 5 BC 7 8 2 3 4 5 AZ NV BC ME BC 9 8 7 6 1 5 OR	386 387 388 389 390 390 391 391	Argoporis apice Argoporis bicol Epantius obscu Eulabis bicarin Oninae Ulomir	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshschottz, 1829 ii LeConte, 1861	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851	58BC 658 67 OR NV 85BC 782345AZ NV BC ME BC 987615OR 53824 NV AZ NM T>	386 387 388 389 390 390 391 391	Argoporis apica Argoporis bicol Epantius obscu Eulabis bicarin Coninae Ulomir Uloma longula Oninae Helopiu	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshschottz, 1829 ii LeConte, 1861	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851	5 8 BC 65 8 67 OR NV 85 BC 7 8 2 3 4 5 AZ NV BC ME BC 98 7 6 15 OR 5 3 8 2 4 NV AZ NM T> UT CO 98 7 AR CA NM TX UT	386 387 388 389 390 390 391 Tenebri	Argoporis apic: Argoporis bicol Epantius obsci Eulabis bicarin Oninae Ulomir Uloma longula Oninae Helopi Helops angust	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshschottz, 1829 ii LeConte, 1861	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851 Eleodes (Eleodes) dentipes Eschscholtz 1833 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) gracilis LeConte 1858	5 8 BC 6 5 8 6 7 OR NV 8 5 BC 7 8 2 3 4 5 AZ NV BC ME BC 9 8 7 6 1 5 OR 5 3 8 2 4 NV AZ NM T> UT CO 8 5 4 BC 9 8 7 AR CA NM TX U' WA OR NV CO ID +	386 387 388 389 390 390 391 Tenebri 392 393 393 393	Argoporis apic: Argoporis bicol Epantius obsci Eulabis bicarin Oninae Ulomir Uloma longula Oninae Helopi Helops angust	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshscholtz, 1829 ii LeConte, 1861 ni us LeConte, 1859 ttus (LeConte), 1859 ttus (LeConte), 1859	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7 5 9 6 7 OR
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) aculicaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851	5 8 BC 65 8 67 OR NV 85 BC 7 8 2 3 4 5 AZ NV BC ME BC 98 7 6 15 OR 5 3 8 2 4 NV AZ NM T> UT CO 98 7 AR CA NM TX UT	386 387 388 389 390 390 391 391 392 393 394 394 394	1 Argoporis apic. 1 Argoporis bicol 1 Epantius obset 1 Eulabis bicarin oninae Ulomir 1 Uloma longula oninae Helopi 1 Helops angust 1 Helops attenue 1 Helops bachei	alis californica Berry, 1851 or (LeConte), 1851 rurus LeConte, 1851 ata Eshscholtz, 1829 ii LeConte, 1861 ni us LeConte, 1859 ttus (LeConte), 1859 ttus (LeConte), 1851 LeConte, 1861	4 AZ ME 3 4 AZ 58 9 BC 89 6 7 59 6 7 OR 5 3 4 NV AZ 5
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) acuticaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851 Eleodes (Eleodes) dentipes Eschscholtz 1833 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) gracilis LeConte 1858	5 8 BC 65 8 67 OR NV 85 BC 7 8 2 3 4 5 AZ NV BC ME BC 98 7 6 1 5 OR 53 8 2 4 NV AZ NMT > UT CO 98 7 AR CA NM TX U' WA OR NV CO ID + 5 BC 1 2 AZ ME CO ID NV	386 387 388 389 390 390 391 391 392 393 394 395 395	Argoporis apic. Argoporis bicol Epantius obsci Eulabis bicarin Coninae Ulomir Uloma longula Oninae Helopi Helops angust Helops attenus Helops blaisde Helops blaisde	alis californica Berry, 1851 or (LeConte), 1851 urus LeConte, 1851 ata Eshschottz, 1829 ii LeConte, 1861 ni us LeConte, 1859 attus (LeConte), 1851 LeConte, 1861 leConte, 1861	4 AZ ME 3 4 AZ 5 8 9 BC 8 9 6 7 5 9 6 7 OR 5 3 4 NV AZ 5 5 5 BC
337		Eleodes (Cratidus) osculans (LeConte) 1851 Eleodes (Discogenia) marginatus Eschscholtz 1833 Eleodes (Discogenia) scabriculus LeConte 1858 Eleodes (Eleodes) aculicaudus LeConte 1851 Eleodes (Eleodes) aculicaudus LeConte 1851 Eleodes (Eleodes) armatus LeConte 1851 Eleodes (Eleodes) dentipes Eschscholtz 1833 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) gracilis LeConte 1858 Eleodes (Eleodes) prandicollis Mannerheim 1843 Eleodes (Eleodes) hispitabris Say 1824	58 BC 658 67 OR NV 85 BC 78 23 45 AZ NV BC ME BC 98 76 15 OR 53 82 4 NV AZ NM T> UT CO 98 7 AR CA NM TX U* WA OR NV CO ID + 5 BC	386 387 388 389 390 390 391 391 392 393 394 394 394	Argoporis apic. Argoporis bicol Epantius obsci Epantius obsci Eulabis bicarin Ulomir Uloma longula Oninae Helopi Helops angust Helops attenus Helops bachei Helops bachei Helops californ Helops californ	alis californica Berry, 1851 or (LeConte), 1851 rurus LeConte, 1851 ata Eshscholtz, 1829 ii LeConte, 1861 ni us LeConte, 1859 ttus (LeConte), 1859 ttus (LeConte), 1851 LeConte, 1861	4 AZ ME 3 4 AZ 58 9 BC 89 6 7 59 6 7 OR 5 3 4 NV AZ 5

Figure I. Continue.

		Californi	a Tenebrionic					
Endemic # CA	Status	Species	Distribution	E #		ic Status group	Species	Distribution
	8r				CA	8 r	-	
399 🔲		ardsi Horn, 1870	35 OR 7	449	•		orus inquilinus Casey, 1891	6
400 🗹 401 🗆		oensis Blaisdell, 1931 s LeConte, 1857	7 8 WA	450 451	•	,	orus irritus Fall, 1931 orus jacobinus Fall, 1931	58 U
		s Lecone, 1657 sangula Blaisdell, 1921	5 8	451	•		orus jacobinus iraii, 1931 orus montivagus Fall, 1931	U
402 🗸 403 🗸		us LeConte, 1859	19657	453	•		orus parvus Fall, 1931	3
404		tipennis LeConte, 1866	6	454	•		orus protibialis Fall, 1931	3 4 AZ
405		es (LeConte), 1851	5 BC	455		-	orus punctatissimus LeConte, 1866	3 4 NM TX UT AZ
406		ollis LeConte, 1866	5	456		,	orus punctulatus (LeConte), 1859	6
407		losus LeConte, 1866	5796	457	_		orus rufohumeralis Campbell, 1982	8
408	1 Helops simul	lator Blaisdell, 1921	6	458	\Box	1 Hymeno	orus sinuatus Fall, 1931	9 OR
409	1 Helops spret	us Horn, 1880	2 NV	459		1 Hymeno	orus thoracicus Fall, 1931	4 AZ
410 🗸	1 Helops stend	otrichoides Blaisdell, 1895	68	460	✓	1 Hymeno	orus ulomoides Fall, 1931	U
411 🗸		collis Horn, 1885	3 4	461	~	1 Hymeno	orus uniseriatus Casey, 1891	U
412	1 Helops tume	scens LeConte, 1866	3 4 AZ	462		1 Isomira	comstocki Papp, 1956	769851CA ME ID
Tenebrio	oninae Tribo	liini		463		1 leomira	damnata Marshall, 1970	OR UT WA 67
413 🖂	1 Gnathocerus	cornutus (Fabricius) 1801	С	464			luscitiosa Casey, 1891	586
414	1 Gnathocerus	maxillosus (Fabricius) 1801	С	465			monticola Casey, 1891	U
415	1 Latheticus or	ryzae Chittenden, 1880	С	466	•		variabilis (Horn),1875	8965
416	1 Latheticus pr	rosopis Chittenden, 1904	С	467			chara marshalli Campbell, 1978	7
417	1 Lyphia tetrap	ohylla (Fairmaire,1856)	C	468	•		chara procera Casey, 1891	9 6 CA AZ ID OR WA
418	1 Mycotrogus a	angustus Horn, 1870	4 AZ	469			chara pubipennis (LeConte), 1878	5
419	1 Mycotrogus p	piceus Horn, 1870	4 BC	470			cistela opaca LeConte, 1859	98653
420 🗌		eburgii (Wissnmann), 1848	С	471		1 Pseudoo	cistela pacifica Hopping, 1933	9 CA OR WA
421		depressus Wollaston, 1864	С	472		1 Pseudoo	cistela pinguis LeConte, 1866	5 WA CA OR
422		itiosus LeConte, 1866	С	473		1 Stenoch	idius cyanescens LeConte, 1859	9 8 6 OR ID
423		dax Uyttenboogsart, 1933	С	474	~		nidius gracilis LeConte, 1851	658
424		evicorne (LeConte), 1859	С	475	\checkmark	1 Stenoch	nidius robustus Schaeffer, 1911	8976
425		staneum Herbst, 1797	C C	Ste	enoch	niinae Cn	odalonini	
426 427		nfusum Jacquelin du Val, 1868 structor Uyttenboogsart, 1933	C	476		1 Alobates	s pennsylvanicus (DeGeer), 1775	5 BC
427 428		adens (Charpentier), 1825	C	477	~	1 Cibdelis	bachei LeConte, 1861	9876
	oninae Apoc		O	478		1 Cibdelis	blaschkei Mannerheim, 1843	8
		•		479		1 Coelocn	nemis californica Mannerheim, 1843	5 8 9 6 CA ME ID NV
429 🗸		inthicoides Eschscholtz, 1831	98576	400		4.0		OR UT
430 🗌		livinoides Horn, 1870	2 NV	480 481			nemis lucia Doyen, 1973	8 58962AZNM
431 🔽		etosa Doyen & Kityama, 1980	8	482			nemis magna LeConte, 1851 nemis punctata LeConte, 1855	123 NV UT AZ
Tenebrio	oninae Alphi	toblini		• 483			nemis rugulosa Doyen, 1973	10R
432	1 Alphitobius d	fiaperinus (Panzer), 1797	С	484			nemis sulcata Casey, 1895	23 NV AZ UT
433	1 Alphitobius la	aevigatus Fabricius, 1781	С	485			inus laevissimus (Casey), 1890	56931OR
434		arginalis Horn, 1870	568179BC	486	_		inus serratus Mannerheim, 1843	1965 AZ WY WA UT
435		s LeConte, 1851	7 8 2 NV BC				,	OR NM MT ID CO
436 🗸	O Neatus n.sp.		68					
437		litor Linnaeus, 1758	С					
438 🔲		scurus Fabricius,1792	С					
renebrio	oninae Centr	ronopini						
439	1 Scotobaenus	s parallelus LeConte,1859	96 OR					
440 🗸		s punctatus (Blaisdell), 1933	6					
441		s simplex (Blaisdell), 1937	6					
442		s wagneri (Blaisdell), 1933	6					
	nae Alleculin							
443 🔲		apacheanus Casey, 1891	3 4 A Z					
444 🗹		discrepans Casey, 1891	5					
445		exiguus Casey, 1891	4 AZ					
446	,	fusculus Casey, 1891	56					
447 🗹 448 🗸		fusicornis Casey, 1891	5 5					
448 🔽	- mymenorus I	infuscatus Casey, 1891	J					

Figure 1. Continue.




Figure 2. Geographic subdivisions of California from http://ucjeps.berkeley.edu/cguide.html#Map with Unit Boundaries with regions 1–9 outlined.

Regional Endemic California Tenebrionidae (January 2014)

REGION Species present, Percent endemic	REGION	Species present, Percent	endemic F	REGION	Species present, Percent e	ndemi
1: The Northern Great Basin 29: 3.45%		catalinae Casey, 1908			ius parvicollis Blaisdell. 1934	
Province	15 Coniontis	globulina Casey, 1895		5 Auchmob	ius picipes Blaisdell. 1934	
including the Warner Mountains and Modoc Plateau	16 Coniontis	pallidicornis Casey, 1890		6 Auchmob	ius slevini Blaisdell. 1934	
1 Eleodes (Litheleodes) corvinus Blaisdell 1921	_	remnans Pierce, 1945			ius subboreus Blaisdell. 1934	
2: The Southern Great Basin 55: 9.09%		santarosae Blaisdell, 1921			olaschkei Mannerheim, 1843	
Province		vandykei Blaisdell, 1921			emis lucia Doyen, 1973	
including the White and Inyo Mountains and		Blapylis) bishopensis Somerby & I	-		costulata Casey, 1908	
intermountain valleys east of the Sierras Nevada's and		Blapylis) incultus LeConte 1861			farallonica Casey, 1895	
White Mountains 1 Auchmobius sublaevis LeConte, 1851		Blapylis) subtestitus Blaisdell 1939			hoppingi Blaisdell, 1918	
2 Eleodes (Blapylis) aristatus Somerby 1977		Metablapylis) n.sp.			lata LeConte, 1866	
3 Eleodes (Caverneleodes) microps Aalbu et al., 2012		rpha bolcan Blaisdell, 1909			timida Casey, 1908	
4 Eleodes (Metablapylis) aalbui Triplehorn 2007		politus Horn, 1883			viatica Eschscholtz, 1929	
5 Oxygonodera n.sp.		robustus LeConte,1866			(Melaneleodes) quadricollis Eschsch	oltz 1833
		ngustus LeConte, 1859			oris nunenmacheri Blaisdell, 1906	
3: The Mojave Desert 112: 10.71%		achei LeConte, 1861			us rufohumeralis Campbell, 1982	
and associated desert mountains		igicollis LeConte, 1866			us lucidulus Casey,1907	
1 Araeoschizus Iariversi Papp, 1981		s piceus Casey, 1890		20 Melanast	us otiosus Casey,1907	
2 Eleodes (Blapylis) panamintensis Somerby 1977	,	us discrepans Casey, 1891				73: 5.489
3 Hymenorus parvus Fall, 1931		us fusicornis Casey, 1891			ne Cascade and Klamath Ranges as	well as
4 Melanastus sterilis Casey,1907		us infuscatus Casey, 1891			n Coast Ranges	
5 Metoponium concors Casey,1907		us exoletus Casey,1907			genitiva Casey, 1890	4070
6 Metoponium cylindricum Casey, 1890		us thoracicus (Casey), 1907			(Blapylis) spilmani Somerby & Doyen	
7 Metoponium gravidum Casey,1907		um edax Casey,1907		,	Litheleodes) papillosus Blaisdell 191	/
Metoponium testaceum Casey,1907 Philolithus jaegeri (Papp), 1961		um insulare Casey,1907		4 Eschatop	oris n.sp. Aalbu m.s.	
		um probatum Casey,1907				
10 Philolithus opimus Casey, 1912 11 Trogloderus costatus pappi Kulzer, 1960		um tersum Casey,1907 ara pubipennis (LeConte), 1878				
12 Typhleusechus chemehuevii Aalbu & Andrews, 1985	,	rpha (Trichiasida) luctata (Horn), 18	770			
	_	pacella Casey, 1907	570			
4: The Sonoran (Colorado) Desert 113: 11.50%						
and associated desert mountains	6: The Sie	erra Nevada Mountains	100: 16.00%			
1 Araeoschizus andrewsi Papp, 1981						
2 Araeoschizus hardyi Papp, 1981		eachi Blaisdell, 1932				
3 Araeoschizus hystrix Papp, 1981 4 Batuliomorpha imperialis Doven, 1987		californica Blaisdell, 1929	- 4070			
4 Batuliomorpha imperialis Doyen, 1987 5 Eschatomoxys andrewsi Aalbu & Thomas, 2007		Blapylis) cooperi Somerby & Doyer	1 1976			
· · · · · · · · · · · · · · · · · · ·		Blapylis) kaweanus Blaisdell 1933	- 4070			
6 Lepidocnemeplatia n.sp. 7 Metoponium dubium Casey, 1884		Blapylis) lariversi Somerby & Doye				
7 Metoponium dubium Casey, 1884 8 Metoponium egregium Casey, 1907		Blapylis) parvicollis Eschscholtz 18				
9 Metoponium integer Casey,1907		Blapylis) schlingeri Somerby & Doy unctipennis LeConte, 1866	reii 1970			
10 Metoponium molestum Casey,1907		mulator Blaisdell, 1921				
11 Statira dumalis Parsons,1973						
12 Stenomorpha (Trichiasida) hirsuta (LeConte), 1851		us inquilinus Casey, 1891 us punctulatus (LeConte), 1859				
13 Telabis fidelis Casey, 1907		us purictulatus (LeConte), 1659 us vergrandis Casey,1907				
*		llus californicus Horn, 1870				
5: The South Coast 170: 24.12%		nus punctatus (Blaisdell), 1933				
including the Transverse and Peninsular Ranges and Channel Islands		nus simplex (Blaisdell), 1937				
1 Alaephus longicornis Casey, 1924		nus wagneri (Blaisdell), 1933				
2 Alaephus maderensis Casey, 1924			70. 0.500/			
3 Alaudes alternata Fall, 1928	7 : The Ce	ntral Valley	76: 6.58%			
4 Apsena barbarae Blaisdell, 1932	4 01	iii- Di-id-II 4000				
5 Apsena crassicomis (Casey), 1890	-	racilis Blaisdell, 1939				
6 Apsena grossa (LeConte), 1866		esnoensis Blaisdell, 1931				
7 Apsena laticornis (Casey), 1891		um gulosum Casey,1907				
8 Araeoschizus doyeni Papp, 1981		ara marshalli Campbell, 1978) 4044			
9 Araeoschizus exiguus Casey, 1907	5 Stenomo	rpha (Stenomorpha) mckittricki (Pie	rce), 1944			
10 Auchmobius angelicus Blaisdell. 1934	8: The Ce	ntral Coast	110: 18.18%			
11 Auchmobius sanfordi Blaisdell. 1934		e San Francisco Bay area and Co	oast Ranges			
12 Blapstinus lecontei Mulsant & Rey, 1859	1 Apocryph	a setosa Doyen & Kityama, 1980				
13 Coelus pacificus Fall, 1897	2 Apsena r	ufipes simplex Blaisdell, 1932				
	3 Araeosch	izus sulcicollis disjunctus Papp, 198				

Figure 3. Regional Endemic California Tenebrionidae.

Acknowledgements

We would like to thank the Jepson Herbarium at UC Berkeley for use of their map, Mike Caterino for encouraging interest in California beetles and regional endemism within the state, Drs. Ronald Somerby and Charles Triplehorn for sharing their ongoing research, and the NSF ARTS program (DEB-1258154) for support of this study.

References

- Aalbu RL (2005) The pimeliine tripe Cryptoglossini: classification, biology and inferred phylogeny (Coleoptera: Tenebrionidae). Annales Zoologici 55: 677–756.
- Aalbu RL, Smith AD, Triplehorn CA (2012) A Revision of the *Eleodes* Subgenus *Caverneleodes* with new species and notes on cave breeding *Eleodes* (Tenebrionidae: Amphidorini). Annales Zoologici (Warszawa) 62(2): 199–216. doi: 10.3161/000345412X652729
- Aalbu RL, Kanda K, Steiner WE Jr. (2009) Opatroides punctulatus Brullé now esablished in California (Coleoptera: Tenebrionidae). Pan-Pacific Entomol. 85(2):38–42. doi: 10.3956/2008-24.1
- Aalbu RL, Triplehorn CA, Campbell JM, Brown KW, Somerby R, Thomas DB (2002) 106. Tenebrionidae. In: Volume 2 of Arnett R et al. The Beetles of the United States. CRC press, New York, 463–509.
- Brown KW, Doyen JT (1991) Review of the genus *Microschatia* (Solier) (Tenebrionidae: Coleoptera). J. New York. Entomol. Soc. 99: 539–582.
- Doyen JT (1973) Systematics of the Genus *Coelocnemis*: A Quantitative Study of Variation. Univ. of Calif. publications in entomology 73: 1–110.
- Doyen JT (1984) Systematics of *Eusattus* and *Conisattus* (Coleoptera; Tenebrionidae; Coniontini; Eusatti). Occ. Pap. Calif. Acad. Sci. No. 141, 104 pp.
- Doyen JT (1987) Review of the tenebrionid tribe Anepsiini (Coleoptera). Proceedings of the California Academy of Sciences 44: 343–371
- Geographic subdivisions of California, Jepson Flora Project (Eds) [2013] Jepson eFlora, http://ucjeps.berkeley.edu/IJM.html [accessed on December 9, 2013]
- Pape RB, Aalbu RL, Thomas DB (2007) A Revision of the Genus *Eschatomoxys* Blaisdell (Tenebrionidae: Pimeliinae: Edrotini)) with notes on the biology of a new species. Coleop. Bull. 61(4): 519–540. doi: 10.1649/0010-065X(2007)61[519:AROTGE]2.0.CO;2
- Papp CS (1981) Revision of the genus Araeoschizus LeConte (Coleoptera: Tenebrionidae). Ent. Arb. Mus. Frey 29: 273–420.
- Smith AD (2013) Phylogenetic revision of the North American Asidini (Coleoptera: Tenebrioni-dae). Sys. Entomol.
- Thomas DB (2005) Blaisdell's formae and homonyms in the genus *Eleodes* Eschscholtz (Coleoptera; Tenebrionidae: Embaphionini). Annales Zoologici 55(4): 549–560.
- Triplehorn CA (2007) New species of *Eleodes* (Coleoptera: Tenebrionidae). Proc. Entomol. Soc. Wash. 109(3): 628–642.
- Triplehorn CA, Thomas DB (2011) Studies in the Genus *Eleodes* Eschscholtz with a Revision of the Subgenus *Melaneleodes* Blaisdell and *Omegeleodes*, new subgenus (Coleoptera: Tenebrionidae). Trans. American Entomol. Soc. 137(3+4): 251–281.