




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original article

Emotional Impact on Health Personnel, Medical Students, and General Population Samples During the COVID-19 Pandemic in Lima, Peru



Bruno Pedraz-Petrozzi^{a,*}, Hever Krüger-Malpartida^{b,c}, Martin Arevalo-Flores^{b,c}, Frine Salmavides-Cuba^{c,d}, Victor Anculle-Arauco^b, Mauricio Dancuart-Mendoza^b

^a Cognitive Neurosciences, Center for Psychiatry and Psychotherapy, Justus-Liebig University, Giessen, Hessen, Germany

^b Department of Neuropsychiatry, Cayetano Heredia Hospital, San Martin de Porres, Lima, Peru

^c Faculty of Medicine, Cayetano Heredia Peruvian University, San Martin de Porres, Lima, Peru

^d Department of Tropical Medicine and Infectious Diseases, Cayetano Heredia Hospital, San Martin de Porres, Lima, Peru

ARTICLE INFO

Article history:

Received 17 December 2020

Accepted 21 April 2021

Available online 5 June 2021

Keywords:

COVID-19

Mental health

Pandemics

Latin America

Medical students

Health personnel

Peru

ABSTRACT

Background: The aim of the study is to compare the emotional effects of COVID-19 among three different groups, namely: health personnel, medical students, and a sample of the general population.

Methods: 375 participants were recruited for this study, of which 125 were medical students (preclinical studies, 59; clinical studies, 66), 125 were health personnel (COVID-19 frontline personnel, 59; personnel not related with COVID-19, 66), and 125 belonged to the general population. The PHQ-9, GAD-7, and CPDI scales were used to assess the emotional impact. A multinomial logistic regression was performed to measure differences between groups, considering potential confounding factors.

Results: Regarding CPDI values, all other groups showed reduced values compared to COVID-19 frontline personnel. However, the general population, preclinical and clinical medical students showed increased PHQ-9 values compared to COVID-19 frontline personnel. Finally, confounding factors, gender and age correlated negatively with higher CPDI and PHQ-9 scores.

Conclusions: Being frontline personnel is associated with increased COVID-19-related stress. Depression is associated, however, with other groups not directly involved with the treatment of COVID-19 patients. Female gender and younger age correlated with COVID-19-related depression and stress.

© 2021 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

* Autor para correspondencia.

E-mail address: bruno.pedraz@med.uni-giessen.de (B. Pedraz-Petrozzi).

<https://doi.org/10.1016/j.rcp.2021.04.006>

0034-7450/© 2021 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Impacto emocional en personal de salud, estudiantes de Medicina y población general durante la pandemia de COVID-19 en Lima, Perú

R E S U M E N

Palabras clave:

COVID-19
Salud mental
Pandemias
América Latina
Estudiantes de Medicina
Personal de salud
Perú

Introducción: El objetivo del estudio es comparar los efectos emocionales de la COVID-19 entre 3 grupos diferentes, a saber: personal de salud, estudiantes de Medicina y una muestra de la población general.

Métodos: Se incluyó a 375 participantes en este estudio, de los que 125 eran estudiantes de Medicina (estudios preclínicos, 59; estudios clínicos, 66), 125 eran personal de salud (personal de primera línea contra la COVID-19, 59; personal no relacionado con la COVID-19, 66) y 125 pertenecían a la población general. Las escalas PHQ-9, GAD-7 y CPDI se utilizaron para evaluar el impacto emocional. Se realizó una regresión logística multinomial para medir las diferencias entre grupos, considerando posibles factores de confusión.

Resultados: Con respecto a los valores del CPDI, todos los demás grupos mostraron valores reducidos en comparación con el personal de primera línea contra la COVID-19. Sin embargo, la población general y los estudiantes de Medicina preclínica y clínica mostraron un aumento de los valores del PHQ-9 en comparación con el personal en primera línea. Por último, los factores de confusión, sexo y edad se correlacionaron negativamente con puntuaciones más altas del CPDI y el PHQ-9.

Conclusiones: Ser personal de primera línea se asocia con más estrés relacionado con la COVID-19. Sin embargo, la depresión está asociada con otros grupos que no están directamente involucrados en el tratamiento de los pacientes con COVID-19. Las mujeres y los participantes más jóvenes se correlacionaron con la depresión y el estrés relacionados con la COVID-19.

© 2021 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U.

Todos los derechos reservados.

Introduction

Since the first case reports, the coronavirus disease of 2019 (COVID-19) has spread rapidly and has caused diverse negative changes in the world population. These involved not only social restrictions, but also negative consequences for the mental health of society. Different groups, who work actively with COVID-19¹ as well as those who have to comply with the COVID-19 lockdown rules,² are exposed to higher stress, anxiety, and depression rates. For instance, two cross-sectional studies revealed that during the COVID-19 pandemic, college students reported high levels of mental health distress and academic difficulties,^{3,4} causing a significant negative impact on their mental health.^{3,4} Another study in Pakistan, which involved active COVID-19 frontline health personnel, found higher prevalence rates with moderate-to-severe anxiety and depression scores.⁵ Similar results have been found in the study of AlAteeq et al. This study revealed that depression and anxiety symptoms are prevalent among COVID-19 health personnel.⁶ Depression and anxiety are not only prevalent in health personnel, but also in medical students, who are health personnel in training. In this case, a study with medical students reported that almost two-thirds of the medical students showed moderate-to-severe symptoms of depression.⁷ Furthermore, the same study found that half of these students showed higher anxiety symptoms. In this case, and similar to the health personnel, medical students are affected severely by mental health issues related to the COVID-19 pandemic.⁷⁻⁹ These negative consequences of the COVID-19 pandemic

have also affected the general population not related directly to COVID-19. A Peruvian study showed that almost half of the participants showed mild to severe scores related to COVID-19 related stress.¹⁰ Other non-healthcare workers have reported that processing COVID-19 information is correlated positively with distress levels.¹¹ Another example with the general population not related to health professions is the study of Zhang et al. Higher levels of depression were found among the general population during the COVID-19 pandemic, mostly related to stress and hopeless feelings.¹²

Pandemics (e.g., COVID-19) are mostly related to higher rates of depression,¹³⁻¹⁵ anxiety^{16,17} and stress^{18,19} reported in general population, health personnel and medical students. Despite the numerous reports, few studies compare the aforementioned groups. The information obtained regarding this matter will establish better intervention policies to prioritize the most affected population groups, offering medical intervention, social and therapeutic support. To this end, the main objective of this study is to find out differences between medical students, COVID-19 frontline health personnel, health personnel not exposed to COVID-19, and the general population, regarding depression, anxiety, and distress scores.

Material and methods

Study design and selection criteria

For this study, 375 participants between 18 and 80 years old were recruited voluntarily from Lima (Peru) for an online




Figure 1

survey. The volunteers included medical students from the faculty of Medicine of Cayetano Heredia University (UPCH) and health personnel from Lima's health centers. The participants were recruited between 20th August and 20th November 2020 and allocated into 5 groups: general population (GP; n = 125), medical students in preclinical studies (MS-pre; n = 59), medical students in clinical studies (MS-cli; n = 66), COVID-19 frontline health personnel (HP-COVID; n = 59) and health personnel not involved with COVID-19 patients (HP; n = 66). A complete description is shown in "Results". A flowchart of the study is presented in Figure 1.

Participants younger than 18 years old, with insufficient knowledge of Spanish and medical difficulties that could restrict the participation (i.e., learning difficulties, especially illiteracy as well incomplete school studies, or blindness in the online survey were not included in this study.

All participants were fully informed of the study and gave their consent to participate. This study was approved by the ethics committee from the Faculty of Medicine of the Peruvian University Cayetano Heredia and carried out in accordance with the Helsinki Declaration and the ethical standards of the APA.

Data collection

Online survey

The information of this study comes from a database of a study project that englobes emotional impact and COVID-19 pandemics. Part of this information was published recently by Krüger-Malpartida et al.,¹⁰ and some methodological aspects of this article were followed also in this study.

For the data collection, an online survey was carried out. Due to the restrictive policies for avoiding COVID-19 contagions, all instruments and questions were digitalized and programmed in using a free internet survey program (Google

Forms). Afterwards, the online survey was distributed through different social media (i.e., Facebook, Twitter, Instagram, LinkedIn, etc.) and using the principles of the snowballing sampling for the data recollection. In case of the medical students, we distributed also the online survey by using the e-mail program of the university and to different social groups of the UPCH medical faculty. The questions included: informed consent, general information (i.e., age, gender, district, confession/faith, and occupation), previous medical diagnosis and medication intake, and the COVID-19 peritraumatic distress index (CPDI) for the COVID-19 pandemic, GAD-7, and PHQ-9 instruments. Before answering the questions, each participant must accept its participation through the informed consent and declare that the participant is full age (i.e., 18 years old).

Finally, additional questions were asked, as follows: "In the last 14 days, did you have a cough, difficulty breathing, sore throat, and fever?" (COVID.1); "Do you have positive results for any COVID-19 test?" (COVID.2); "Have you been hospitalized (or are you hospitalized at the moment) due to COVID-19?" (COVID.3); "Do you have relatives with positive results for any COVID-19 test?" (COVID.4); "Do you have relatives who were hospitalized due to COVID-19?" (COVID.5), and "Do you have relatives who have passed away due to COVID-19?" (COVID.6).

COVID-19 peritraumatic distress index

The authors of this study followed the definition of COVID-19 related stress used in a recent study of Krüger-Malpartida et al., published elsewhere¹⁰. Some methodological aspects of this article were followed also in this study.

The CPDI was first applied in China²⁰ and recently validated in other countries.^{20,21} In addition, the Spanish version of this instrument was also validated for its use in Peru.²² This instrument was designed for a populational evaluation of changes related to mood, behavior, cognitive skills, circadian rhythm, and other somatic symptoms due to the COVID-19 pandemic.

This instrument consists of 24 items, with a four-factor design: negative mood, cognition, behavioral change, somatization, and hyperarousal/exhaustion. Each item was evaluated by using Likert elements (from 0 to 4: never, occasionally, sometimes, often, and most of the time).¹⁰ The sum of each value per question results in the raw score. The displayed score is obtained by adding 4 to the raw score and used to calculate the CPDI severity degrees. For this reason, this instrument defines different categories for peritraumatic stress due to the COVID-19 pandemic: normal (0 to 28 display points), mild (29 to 52 display points, and severe (53 to 100 display points).¹⁰

Depressive and anxiety symptoms

The authors of this study followed the definition of depression and anxiety used in a recent study of Krüger-Malpartida et al., published elsewhere.¹⁰ Some methodological aspects of this article were followed also in this study.

The Peruvian version of the PHQ-9²³ was used to assess the severity of depressive symptoms. The PHQ-9 delivers values in the range between 0 and 27. The highest value indicates a higher depression score. This instrument was validated in Peru with a representative sample (n = 30 446) and showed significant internal consistency (Cronbach's $\alpha = .87$). This inventory defines different categories for depression scores: minimal (1 to 4 points), mild (5 to 9 points), moderate (10 to 14 points), and severe (15 to 27 points).²³

For anxiety symptoms, the Peruvian version of the GAD-7²⁴ was used to assess the severity of anxiety symptoms. The GAD-7 delivers values in the range between 0 and 21 points. The highest value indicates a higher anxiety score. This instrument was also validated in Peru with a representative sample (n = 2978), showing significant internal consistency (Cronbach's $\alpha = .89$). This inventory also defines different categories for anxiety scores: minimal (0 to 4 points), mild (5 to 10 points), moderate (11 to 15 points, and severe (16 to 21 points).²⁴

Statistical analysis

Statistical analyses were performed using SPSS version 26.0 (Statistical Package for the Social Sciences, IBM Corp., United States of America) and jamovi 1.2.5.0²⁵.

Descriptive data were managed with count data and percentages. To improve readability, the information is presented in tables. Quantitative variables approximately fitting a normal distribution are specified in the text as the mean \pm standard deviation, and those with a non-normal distribution are expressed as the median (Me) [interquartile range]. Categorical variables were specified with count data and percentages. Data were rounded to 2 decimals. Values smaller than 0.001 were shown as $<.001$ and values greater than one million were expressed in scientific notation.

Multinomial logistic regression was computed for comparing the 5 groups, considering the group status (GP, MS-pre, MS-cli, HP or HP-COVID) as a dependent variable. Regarding predictor variables, the following were used as predictor variables and added in 4 different blocks, as follows:

Block 1: CPDI scores, GAD-7 scores, PHQ-9 scores.

Block 2: age, gender, CPDI scores, GAD-7 scores, PHQ-9 scores.

Block 3: COVID_1, COVID_2, COVID_3, COVID_4, COVID_5, COVID_6, age, gender, CPDI scores, GAD-7 scores, PHQ-9 scores.

Block 4: residence, religion/faith, medication intake, previous medical disease, COVID_1, COVID_2, COVID_3, COVID_4, COVID_5, COVID_6, age, gender, CPDI scores, GAD-7 scores, PHQ-9 scores.

The "block" that best explained the data was chosen by using the Akaike information criterion (AIC).

The results of this statistical modeling are presented in "Results". The odds ratio (OR) was flagged as "significant" if the 2-tailed P-value was $<.05$. The 95% confidence intervals (95%CI) were calculated for this model.

Results

General sample descriptions, including medication intake and previous clinical diagnoses of the 5 groups included in the study, are listed in Table 1. Previous information regarding COVID-19 infection, including COVID-19 infection of relatives, is represented in Table 1. Information regarding the most frequent previous clinical diagnosis and the most frequent prescribed medication is represented in Table 2. Descriptive data regarding anxiety scores, depression scores, and CPDI scores are presented in Table 3.

Statistical modeling for emotional impact after COVID-19 lockdown

This study's objective was to find out differences between the 5 established groups (i.e., GP, MS-pre, MS-cli, HP and HP-COVID) regarding depression, anxiety, and distress scores. Variables were introduced stepwise in 4 different blocks. The block which best explained the data was chosen using a smaller AIC. In this case, Block 2 was chosen as the model that could best explain the data of the 5 groups (Block 1, $\chi^2_{(df=12)} = 63.10$, AIC = 1139; Block 2, $\chi^2_{(df=20)} = 408.40$, AIC = 809; Block 3, $\chi^2_{(df=44)} = 442.40$, AIC = 823; Block 4, $\chi^2_{(df=56)} = 462.10$, AIC = 828). Block 2 included the variables age, gender, CPDI scores, GAD-7 scores, PHQ-9 scores as predictors.

Different correlations were found between the 5 groups. These are presented in Table 4. In summary, being part of the GP correlated positively with higher PHQ-9 values, when compared to the HP-COVID group (GP vs. HP-COVID, OR = 1.15; 95%CI, 1.03-1.29; $P = .02$). Similar positive correlations were observed in the MS-pre (MS-pre vs. HP-COVID, OR = 1.23; 95%CI, 1.02-1.47; $P = .03$). Moreover, being part of the MS-cli group correlated with higher PHQ-9 values, when compared to HP-COVID (MS-cli vs. HP-COVID, OR = 1.17; 95%CI, 1.01-1.35; $P = .03$). Within those comparisons, significant correlations were found between PHQ-9 scores, gender and age (Table 4).

On the other hand, higher CPDI values correlated negatively in the comparison GP vs. HP-COVID (OR = 0.92; 95%CI, 0.88-0.96; $P < .001$). Similar negative correlations were found also between MS-pre vs. HP-COVID (OR = 0.90; 95%CI, 0.84-0.97; $P = 0.01$), MS-cli vs. HP-COVID (OR = 0.91; 95%CI,

Table 1 – General socio-demographic data and COVID-19 contact information.

	GP (n = 125)	MS-pre (n = 59)	MS-cli (n = 66)	HP (n = 66)	HP-COVID (n = 59)
Age (years)	41.10 ± 14.30 41 [54-29]	19.70 ± 2.15 19 [20 -19]	23.30 ± 1.74 23 [24-22]	39.60 ± 11.90 38 [49.00-31.30]	35.00 ± 9.41 32 [37.50-29.00]
Gender					
Female	78 (62.40)	28 (47.46)	40 (60.61)	43 (65.15)	25 (42.37)
Male	47 (37.60)	31 (52.54)	26 (39.39)	23 (34.85)	34 (57.63)
Domicile					
Lima – peripheral	35 (28.00)	25 (42.37)	23 (34.85)	14 (21.21)	19 (32.20)
Lima – center	90 (72.00)	34 (57.63)	43 (65.15)	52 (78.79)	40 (67.80)
Faith/religious beliefs					
Absent	26 (20.80)	30 (50.85)	29 (43.94)	19 (28.79)	14 (23.73)
Present	99 (79.20)	29 (49.15)	37 (56.06)	47 (71.21)	45 (76.27)
Medication intake					
No	54 (43.20)	35 (59.32)	35 (53.03)	42 (63.64)	37 (62.71)
Yes	71 (56.80)	24 (40.68)	31 (46.97)	24 (36.36)	22 (37.29)
Presence of past medical diseases					
No	83 (66.40)	48 (81.36)	35 (53.03)	47 (71.21)	38 (64.41)
Yes	42 (33.60)	11 (18.64)	31 (46.97)	19 (28.79)	21 (35.59)
COVID_1					
No	115 (92.00)	51 (86.44)	57 (86.36)	58 (87.88)	50 (84.75)
Yes	10 (8.00)	8 (13.56)	9 (13.64)	8 (12.12)	9 (15.25)
COVID_2					
No	119 (95.20)	59 (100)	62 (93.94)	60 (90.91)	49 (83.05)
Yes	6 (4.80)	0	4 (6.06)	6 (9.09)	10 (16.95)
COVID_3					
No	124 (99.20)	59 (100)	66 (100)	64 (96.97)	58 (98.31)
Yes	1 (0.80)	0	0	2 (3.03)	1 (1.69)
COVID_4					
No	89 (71.20)	25 (42.37)	30 (45.45)	40 (60.61)	28 (47.46)
Yes	36 (28.80)	34 (57.63)	36 (54.55)	26 (39.39)	31 (52.54)
COVID_5					
No	110 (88.00)	44 (74.58)	49 (74.24)	57 (86.36)	46 (77.97)
Yes	15 (12.00)	15 (25.42)	17 (25.76)	9 (13.64)	13 (22.03)
COVID_6					
No	112 (89.60)	43 (72.88)	54 (81.82)	58 (87.88)	50 (84.75)
Yes	13 (10.40)	16 (27.12)	12 (18.18)	8 (12.12)	9 (15.25)

GP: general population; HP: health personnel (not COVID-19 frontline health personnel); HP-COVID: COVID-19 frontline health personnel; MS-cli: medical students, clinical studies; MS-pre: medical students, pre-clinical studies.

COVID_1: in the last 14 days, did you have cough, difficulty breathing, sore throat and fever?; COVID_2: do you have positive results for any sort of COVID-19 test?; COVID_3: have you been hospitalized (or are you hospitalized at the moment) due to COVID-19?; COVID_4: do you have relatives with positive results for any sort of COVID-19 test?; COVID_5: do you have relatives who were hospitalized due to COVID-19?; COVID_6: do you have relatives who have passed away?

Data shown as n (%), mean ± standard deviation, or median [interquartile range].

0.86-0.96; $P < .001$), and HP vs. HP-COVID (OR = 0.94; 95%CI, 0.90-0.99; $P = .02$). Within those comparisons, significant correlations were found between CPDI scores, gender and age (Table 5).

Discussion

This study results in 2 different outcomes concerning differences on the emotional impact due to the COVID-19 pandemic between the five groups (i.e., GP, MS-pre, MS-cli, HP, and HP-COVID). On the one hand, this study showed that higher CPDI values correlate with the COVID-19 frontline health personnel compared with the other remaining four groups. On the other hand, the groups GP, MS-pre and MS-cli showed greater

PHQ-9 values than COVID-19 frontline health personnel. No correlations were found with the GAD-7 between groups.

Our findings reveal 2 important points. First of all, higher psychological distress reflected in higher CPDI values. Secondly, higher depression values and reduced functionality reflected in higher PHQ-9 values. Moreover, our results revealed that higher psychological distress is mostly related to the COVID-19 frontline health personnel group. The current pandemic resulted for many health personnel overwhelming, mostly because of the different factors that daily they have to confront (i.e., long shifts, increased working hours, altered circadian rhythm, fears, death of patients and colleagues, constant increase of hospitalized patients, etc.). Most people tend to restrict or repress during a constant stress situation

Table 2 – Most frequent diagnoses and prescript medication for each group.

	GP	MS-pre	MS-cli	HP	HP-COVID
Most frequent diagnoses for each group ^a	Arterial hypertension (14.29%), diabetes mellitus (11.11%), metabolic syndrome (9.52%)	Bronchial asthma (26.67%), hypothyroidism (13.33%), atopy (13.33%), depression (13.33%)	Bronchial asthma (17.74%), allergic rhinitis (11.29%), polycystic ovary syndrome (8.06%)	Bronchial asthma (22.22%), arterial hypertension (13.89%), obesity (8.33%)	Bronchial asthma (36.36%), hypothyroidism (18.18%), migraine (6.06%)
Most frequent prescript medication for each group ^b	Nutritional supplements (8.61%), vitamin supplements (7.95%), antihypertensive agents (7.28%), oral antidiabetics (5.96%)	Nutritional supplements (21.62%), vitamin supplements (16.22%), antihistaminic agents (8.11%), asthma therapy (β2-agonists + corticoids, 8.11%)	Asthma therapy (β2-Agonists + Corticoids, 16.39%), oral contraceptives (14.75%), antidepressants (9.83%), paracetamol (8.20%)	Asthma therapy (β2-Agonists + Corticoids, 11.76%), antihypertensive agents (9.80%), antidepressants (mostly SSRI, 9.80%), antihistaminic agents (7.84%)	L-thyroxine (20.00%), vitamin supplements (16.67%), antidepressants (13.33%), bronchial asthma (β2-agonists + corticoids, 8.11%)

GP: general population; HP: health personnel (not COVID-19 frontline health personnel); HP-COVID: COVID-19 frontline health personnel; MS-cli: medical students, clinical studies; MS-pre: medical students, pre-clinical studies.

^a The percentages described above represented the frequency of the medical diagnoses in participants who had a medical condition.

^b The percentages described above represented the frequency of the prescribed medication in participants who were taking any medication at the time of the interview.

Table 3 – CPDI, PHQ-9 and GAD-7 descriptive data.

	GP (n = 125)	MS-pre (n = 59)	MS-cli (n = 66)	HP (n = 66)	HP-COVID (n = 59)
CPDI					
Mean ± SD	27.50 ± 13.00	33.60 ± 12.90	28.70 ± 14.20	26.10 ± 15.10	30.30 ± 13.20
Median [Q3-Q1]	26 [36-17]	33 [43.50-23.50]	27 [36.80-16.50]	22.50 [35-14.30]	29 [39-21]
Skewness	0.63	0.20	0.75	1.30	0.30
Categories					
Normal	71	22	37	42	29
Mild	47	34	25	22	26
Severe	7	3	4	2	4
PHQ-9					
Mean ± SD	5.22 ± 5.28	9.59 ± 6.20	7.32 ± 5.48	4.14 ± 4.66	4.73 ± 4.40
Median [Q3-Q1]	4 [8-1]	8 [13-5]	6 [11-3]	3 [6-0.25]	4 [7-1]
Skewness	1.24	0.81	0.78	1.66	1.18
Categories					
Minimal	67	12	21	43	30
Mild	37	23	26	14	21
Moderate	13	12	10	6	7
Severe	8	12	9	3	1
GAD-7					
Mean ± SD	4.46 ± 3.68	6.66 ± 4.44	5.45 ± 4.54	3.76 ± 3.60	4.32 ± 3.74
Median [Q3-Q1]	4 [7-1]	7 [8.50-3.50]	5 [7-2]	3 [6-1]	4 [6.00-1.50]
Skewness	1.04	0.78	1.26	1.36	1.48
Categories					
Minimal	66	19	32	39	35
Mild	52	32	26	24	22
Moderate	4	4	5	2	0
Severe	3	4	3	1	2

CPDI: COVID-19 peritraumatic distress index; GAD-7: general anxiety disorder questionnaire 7; GP: general population; HP: health personnel (not COVID-19 frontline health personnel); HP-COVID: COVID-19 frontline health personnel; MS-cli: medical students, clinical studies; MS-pre: medical students, pre-clinical studies; PHQ-9: patient health questionnaire 9; Q1: 25th percentile; Q3: 75th percentile; SD: standard deviation.

emotions and feelings in order to achieve tasks,²⁶ that in this case involve the life of high-risk patients. This could be why the group of the COVID-19 frontline health personnel correlated higher psychological distress values. Finally, our

results revealed that higher depression scores correlated with other groups not directly involved with COVID-19. Following the stress-vulnerability model,²⁷ it is possible that some individuals could develop mental health issues due to an stressful

Table 4 – Multinomial logistic regression.

	Predictor	P	OR	95%CI Lower	95%CI Upper
GP (1) vs. MS-pre (0) (reference level: MS-pre)	CPDI scores	.58	1.02	0.95	1.10
	GAD-7 scores	.70	0.96	0.77	1.20
	PHQ-9 scores	.47	0.94	0.79	1.11
	Age*	<.001	3.57	2.64	4.82
	Gender	.27	0.52	0.16	1.67
GP (1) vs. MS-cli (0) (reference level: MS-cli)	CPDI scores	.56	1.02	0.96	1.07
	GAD-7 scores	.89	0.99	0.84	1.17
	PHQ-9 scores	.83	0.99	0.87	1.12
	Age*	<.001	1.59	1.39	1.81
	gender	.84	1.09	0.47	2.57
GP (1) vs. HP (0) (reference level: HP)	CPDI scores	.28	0.98	0.94	1.02
	GAD-7 scores	.42	1.07	0.91	1.25
	PHQ-9 scores	.16	1.08	0.97	1.21
	age	.13	1.02	0.99	1.05
	gender	.50	1.25	0.66	2.39
GP (1) vs. HP-COVID (0) (reference level: HP-COVID)	CPDI scores*	<.001	0.92	0.88	0.96
	GAD-7 scores	.09	1.15	0.98	1.34
	PHQ-9 scores*	.02	1.15	1.03	1.29
	Age*	<.001	1.06	1.02	1.09
	Gender*	.01	0.41	0.21	0.82
MS-pre (1) vs. MS-cli (0) (reference level: MS-cli)	CPDI scores	.88	1.00	0.94	1.06
	GAD-7 scores	.74	1.03	0.86	1.24
	PHQ-9 scores	.48	1.05	0.91	1.21
	Age*	<.001	0.45	0.34	0.58
	gender	.13	2.11	0.80	5.56
MS-pre (1) vs. HP (0) (reference level: HP)	CPDI scores	.27	0.96	0.89	1.03
	GAD-7 scores	.39	1.11	0.87	1.42
	PHQ-9 scores	.13	1.16	0.96	1.39
	Age*	<.001	0.29	0.21	0.39
	gender	.16	2.41	0.71	8.18
MS-pre (1) vs. HP-COVID (0) (reference level: HP-COVID)	CPDI scores*	.01	0.90	0.84	0.97
	GAD-7 scores	.14	1.20	0.94	1.52
	PHQ-9 scores*	.03	1.23	1.02	1.47
	Age*	<.001	0.30	0.22	0.40
	gender	.72	0.80	0.24	2.68
MS-cli (1) vs. HP (0) (reference level: HP)	CPDI scores	.19	0.96	0.91	1.02
	GAD-7 scores	.44	1.08	0.89	1.31
	PHQ-9 scores	.21	1.10	0.95	1.27
	Age*	<.001	0.64	0.56	0.73
	gender	.78	1.14	0.46	2.86
MS-cli (1) vs. HP-COVID (0) (reference level: HP-COVID)	CPDI scores*	<.001	0.91	0.86	0.96
	GAD-7 scores	.12	1.16	0.96	1.40
	PHQ-9 scores*	.03	1.17	1.01	1.35
	Age*	<.001	0.67	0.58	0.76
	Gender*	.04	0.38	0.15	0.94
HP (1) vs. HP-COVID (0) (reference level: HP-COVID)	CPDI scores*	.02	0.94	0.90	0.99
	GAD-7 scores	.43	1.08	0.90	1.29
	PHQ-9 scores	.36	1.06	0.93	1.21
	Age*	.04	1.04	1.00	1.07
	Gender*	.01	0.33	0.15	0.71

CPDI: COVID-19 peritraumatic distress index; GAD-7: general anxiety disorder questionnaire 7; GP: general population; HP: health personnel (not COVID-19 frontline health personnel); HP-COVID: COVID-19 frontline health personnel; MS-cli: medical students, clinical studies; MS-pre: medical students, pre-clinical studies; PHQ-9: patient health questionnaire 9.

event, such as the COVID-19 pandemics and its consequences on the society. The long-lasting isolation, the restriction of free movement and the limitation of different activities (e.g., studying, working, interacting with other people, etc.) could influence, in those vulnerable individuals, the occurrence of depressive symptoms through different biological factors.

To the best of our knowledge, there is information on this line of evidence that could be comparable to our results. For

instance, Naser et al. published a cross-sectional study comparing the adverse effects of the COVID-19 pandemic in three different groups: university students, health care professionals, and the general population. They found that mental health problems (i.e., anxiety and depression) were more prevalent in university students.²⁸

In this study, the groups of medical students showed correlations with PHQ-9 values compared to the HP-COVID group.

Table 5 – PHQ-9 and CPDI values, and their correlations with gender and age.

PHQ-9 values, correlations with gender and age		
	Gender	Age
GP vs. HP-COVID	$r_s = -.23, P = .001^*$	$r_s = -.28, P < .001^*$
MS-pre vs. HP-COVID	$r_s = -.15, P = .11$	$r_s = -.42, P < .001^*$
MS-cli vs. HP-COVID	$r_s = -.26, P = .004^*$	$r_s = -.29, P = .001^*$
CPDI values, correlations with gender and age		
	Gender	Age
GP vs. HP-COVID	$r_s = -.25, P = .001^*$	$r_s = -.13, P = .07$
MS-pre vs. HP-COVID	$r_s = -.30, P = .001^*$	$r_s = -.19, P = .04^*$
MS-cli vs. HP-COVID	$r_s = -.32, P < .001^*$	$r_s = 0.04, P = .65$
HP vs. HP-COVID	$r_s = -.23, P = .01^*$	$r_s = -.13, P = .16$

CPDI: COVID-19 peritraumatic distress index; GAD-7: general anxiety disorder questionnaire 7; GP: general population; HP: health personnel (not COVID-19 frontline health personnel); HP-COVID: COVID-19 frontline health personnel; MS-cli: medical students, clinical studies; MS-pre: medical students, pre-clinical studies; PHQ-9: patient health questionnaire 9.

However, no correlations were seen for anxiety scores (Table 4). These outcomes revealing higher depression values among medical students could be due to the social restrictions that obligate many medical students to abandon their studies.^{7,9,29} The switch to distance education and the suspension of clinical activities or night shifts could have negatively influenced this group's mental health, leading mostly to adverse mood effects.^{7,9,29}

Krishnamoorthy et al. published a meta-analysis indicating that during pandemic, health professionals have a higher prevalence of psychological morbidities than the general population, characterized mostly by psychological distress and poor sleep quality.³⁰ The results of this study indeed support that higher values of psychological distress are correlated to the COVID-19 frontline health personnel group. However, being part of the general population is mostly correlated with higher depression scores. Frontline professionals are faced most of the time with fears of getting infected or infecting others.³⁰ Other concerns, like personal protection, death of their own colleagues, or excess working hours, could have influenced the obtained results.

On the other hand, the general population is exposed to isolation due to the sanitary restrictions.^{10,29,31} People belonging to the general population are confined in their homes and have their transit restricted.^{10,29,31} This could lead to higher values of depression, poor sleep quality, hopeless feelings, and suicidal thoughts.^{32,33}

Also remarkable are the results between HP and HP-COVID (Table 4). In this case, being a COVID-19 frontline health professional is correlated with higher values of psychological distress due to COVID-19 (CPDI). Our results could be comparable with the findings of Cai et al.³⁴ In this mentioned study, COVID-19 frontline medical workers have higher rates of mental health problems of any kind compared to non-frontline medical workers.³⁴ Moreover, Cai et al. found correlations between anxiety symptoms, insomnia and depressive mood and being a frontline health professional.³⁴

Finally, the variables age and gender are essential to mention. Significant correlations with the variables age and gender were found in the comparisons (Table 4 and Table 5). This could mean that age and gender could also have influenced in the correlations found by this multinomial logistic regression. In

this case, younger age or female gender correlated with higher scores of the aforementioned psychological instruments (i.e., PHQ-9 and CPDI). These obtained results coincided with other studies that have found associations with female gender and age of the participant.^{21,28,35,36}

Although these results contribute to the actual COVID-19 pandemic's panorama, concerning mostly adverse effects on mental health as an emotional impact, the reader must consider some limitations. Firstly, the sampling procedure (in this case the snowballing method) could contravene the principle of randomization. However, the social restrictions due to COVID-19 pandemic made it difficult to have direct contact with other persons, making the online distribution of the survey one of the most suitable methods for recruiting participants and recollecting data. Moreover, the sample size could be larger to generalize the results beyond the study's scope. However, the power obtained from this study with 375 participants and 5 groups was $1-\beta=.93$, a value that overcame the $1-\beta=.80$ threshold. Therefore, the sample size for the study design should be sufficient to examine the expected effects. Also, the higher total of women compared to men could have influenced the results. However, studies related to COVID-19 lockdown distress have also reported a higher proportion of female participants, which is also reflected in this study. As expected, there were many very low CPDI, PHQ-9 and GAD-7 scores in the evaluated participants, which led to skewed distributions. Multinomial logistic regression was computed to overcome this limitation because there is no consideration regarding skewed statistical distributions. Finally, medication intake, previous medical conditions, and district of residence could affect COVID-19 distress scores. All these variables were included in the model. However, these variables did not appear to affect the results of the current study.

In conclusion, being a COVID-19 frontline health personnel is associated with more psychological distress, involving harmful behavioral and emotional components. This could be related to the fact that frontline health professionals confront excess working hours, their colleagues' death, fear of being infected, among others. On the other hand, higher depression scores were associated mostly with other groups not directly involved with COVID-19 patients' treatment. This could be related to the sanitary emergency restrictions, the long-lasting

isolation, the restriction of free movement, and the limitation of different activities, like studying and accomplishing essential duties (in medical students). Finally, female gender and younger age correlated with depression and stress related to COVID-19, also playing an important role in these findings.

Future studies and public health authorities may consider these obtained results for generating intervention and support. First of all, intervention studies and health policies should consider higher psychological distress in COVID-19 frontline health personnel for creating intervention strategies, such as group or individual psychotherapy. Furthermore, Balint groups and related interventions could also help to comprehend different situations with COVID patients as well to reduce the stress originated from the patient-health personnel relationships. Finally, future studies and public health interventions should consider severe stress cases and their management, especially a combined management with psychotherapy and pharmacotherapy. For the other groups, intervention studies and health policies should consider higher depression values. The intervention and assessment of the other groups (i.e., students, general population and health personnel not related to COVID-19) should be based on telepsychotherapeutic interventions or telepsychiatric consultations. In most severe depression cases, such as extreme agitation or suicide attempts, public health authorities should consider a 24-hour telephone support line for the population.

Author contributions

Bruno Pedraz-Petrozzi: First author, wrote the introduction, methods, results, discussion and conclusions. Corrected the manuscript, did the data analysis and the literature search. Hever Krüger-Malpartida: First and corresponding author; PI of this working group, HKM contributed with the introduction, discussion and conclusions, corrected the manuscript and did the literature search. Martin Arévalo-Flores: contributed with the literature search, introduction and discussion, helped with the paper proofreading, corrections and the interpretation of the results in the discussion. Frine Samalvides-Cuba: Corrected the manuscript and helped with the proofreading and paper mentoring. Victor Anculle-Arauco: Helped with the data recollection and contributed with the proofreading. Mauricio Dancuart-Mendoza: Helped with the data recollection and contributed with the proofreading.

Conflict of interests

The authors declare no conflict of interests.

Acknowledgements

The authors of this work would like to thank Dr. Natalie Tinney, M.D. (physician and English native speaker) and Dr. Cecilia Clement, M.D. (The University of Texas-Medical Branch, Galveston, Texas) for supporting with the proofreading, spelling and grammar correction of the English language.

REFERENCES

1. Teng Z, Huang J, Qiu Y, Tan Y, Zhong Q, Tang H, et al. Mental health of front-line staff in prevention of coronavirus disease 2019. *Zhong nan da xue xue bao. Yi xue ban = J. Cent. South Univ. Med. Sci.* 2020;45:613-9.
2. Zhou Y, Macgeorge EL, Myrick JG. Mental health and its predictors during the early months of the COVID-19 pandemic experience in the United States. *Int J Environ Res Public Health.* 2020;17:6315, <http://dx.doi.org/10.3390/ijerph17176315>.
3. Kecojevic A, Basch CH, Sullivan M, Davi NK. The impact of the COVID-19 epidemic on mental health of undergraduate students in New Jersey, cross-sectional study. *PLoS One.* 2020;15:e0239696, <http://dx.doi.org/10.1371/journal.pone.0239696>.
4. Hubner C, von K, Bruscatto ML, Lima RD. Distress among Brazilian university students due to the COVID-19 pandemic: survey results and reflections. *medRxiv.* 2020.
5. Amin F, Sharif S, Saeed R, Durrani N, Jilani D. COVID-19 pandemic- knowledge, perception, anxiety and depression among frontline doctors of Pakistan. *BMC Psychiatry.* 2020;20:459, <http://dx.doi.org/10.1186/s12888-020-02864-x>.
6. AlAteeq DA, Aljhani S, Althiyabi I, Majzoub S. Mental health among healthcare providers during coronavirus disease (COVID-19) outbreak in Saudi Arabia. *J. Infect. Public Health.* 2020;13:1432-7, <http://dx.doi.org/10.1016/j.jiph.2020.08.013>.
7. Sartorao Filho CI, et al. Impact of covid-19 pandemic on mental health of medical students: a cross-sectional study using GAD-7 and PHQ-9 Questionnaires. *medRxiv.* 2020.
8. Komer L. COVID-19 amongst the Pandemic of Medical Student Mental Health. *Int J Med Students.* 2020;8:56-7, <http://dx.doi.org/10.5195/ijms.2020.501>.
9. Chandratte S. Medical Students and COVID-19: Challenges and Supportive Strategies. *J Med Educ Curric Dev.* 2020;7, <http://dx.doi.org/10.1177/2382120520935059>, 2382120520935059.
10. Krüger-Malpartida H, Pedraz-Petrozzi B, Arevalo-Flores M, Samalvides-Cuba F, Anculle-Arauco V, Dancuart-Mendoza M, et al. Effects on Mental Health After the COVID-19 Lockdown Period: Results From a Population Survey Study in Lima. *Peru. Clin. Med. Insights Psychiatry.* 2020;11, <http://dx.doi.org/10.1177/1179557320980423>, 1179557320980423.
11. Ruiz-Frutos C, Ortega-Moreno M, Dias A, Bernardes JM, Garcia-Iglesias JJ, Gómez-Salgado J, et al. Information on COVID-19 and psychological distress in a sample of non-health workers during the pandemic period. *Int J Environ Res Public Health.* 2020;17:6982, <http://dx.doi.org/10.3390/ijerph17196982>.
12. Zhang W, Yang X, Zhao J, Yang F, Jia Y, Cui C, et al. Depression and Psychological-Behavioral Responses Among the General Public in China During the Early Stages of the COVID-19 Pandemic: Survey Study. *J. Med. Internet Res.* 2020, <http://dx.doi.org/10.2196/22227>.
13. Ho CS, Chee CY, Ho RC. Mental Health Strategies to Combat the Psychological Impact of COVID-19 Beyond Paranoia and Panic. *Ann. Acad. Med. Singapore.* 2020;49:155-60.
14. Liu X, Kakade M, Fuller C, Fan B, Fang Y, Kong J, et al. Depression after exposure to stressful events: Lessons learned from the severe acute respiratory syndrome epidemic. *Compr. Psychiatry.* 2012;53:15-23, <http://dx.doi.org/10.1016/j.comppsy.2011.02.003>.
15. Hawryluck L, Gold WL, Robinson S, Porgoski S, Galea S, Styra R. SARS control and psychological effects of quarantine, Toronto. *Canada. Emerg. Infect. Dis.* 2004;10:1206-12, <http://dx.doi.org/10.3201/eid1007.030703>.

16. Jeong H, Yim HW, Song Y-J, Ki M, Min J-A, Cho J, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol. Health.* 2016;38:e2016048, <http://dx.doi.org/10.4178/epih.e2016048>.
17. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open.* 2020;3:e203976.
18. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int. J. Environ. Res. Public Health.* 2020;17:1729, <http://dx.doi.org/10.3390/ijerph17051729>.
19. Mihashi M, Otsubo Y, Yinjuan X, Nagatomi K, Hoshiko M, Ishitake T, et al. Predictive Factors of Psychological Disorder Development During Recovery Following SARS Outbreak. *Heal. Psychol.* 2009;28:91-100, <http://dx.doi.org/10.1037/a0013674>.
20. Afshar Jahanshahi A, Mokhtari Dinani M, Nazarian Madavani A, Li J, Zhang SX. The distress of Iranian adults during the Covid-19 pandemic - More distressed than the Chinese and with different predictors. *medRxiv.* 2020;87:124-5.
21. Zhang S, Yifei W, Afshar Jahanshahi A, Jia J, Haensel-Schmitt V. First study on mental distress in Brazil during the COVID-19 crisis. *medRxiv.* 2020, <http://dx.doi.org/10.1101/2020.04.18.20070896>.
22. Pedraz-Petrozzi B, Arévalo-Flores M, Krüger-Malpartida H, Anculle-Arauco V. Validación por expertos del Índice de Distrés Peri-traumático relacionado con la Enfermedad por Coronavirus 2019 para su uso en el Perú. *Rev Neuropsiquiatr.* 2021;83:228-35, <http://dx.doi.org/10.20453/rmp.v83i4.3888>.
23. Villarreal-Zegarra D, Copez-Lonzoy A, Bernabé-Ortiz A, Melendez-Torres GJ, Bazo-Alvarez JC. Valid group comparisons can be made with the Patient Health Questionnaire (PHQ-9): A measurement invariance study across groups by demographic characteristics. *PLoS One.* 2019;14:e0221717, <http://dx.doi.org/10.1371/journal.pone.0221717>.
24. Zhong QY, Gelaye B, Zaslavsky AM, Fann JR, Rondon MB, Sánchez SE, et al. Diagnostic validity of the generalized anxiety disorder - 7 (GAD-7) among pregnant women. *PLoS One.* 2015;10:e0125096, <http://dx.doi.org/10.1371/journal.pone.0125096>.
25. Love J, Dropmann D, Selker R, Gallucci M, Jentschke S, Balci S et al. The jamovi project. [accessed Jun 22 2020]. Available from: <https://www.jamovi.org>
26. Piqueras Rodríguez JA, Ramos Linares V, Martínez González AE, Oblitas Guadalupe LA. Emociones negativas y su impacto en la salud mental y física. *Suma Psicol.* 2009;16(2):85-112.
27. Goh C, Agius M. The stress-vulnerability model how does stress impact on mental illness at the level of the brain and what are the consequences? *Psychiatr Danub.* 2010;22:198-202, [http://dx.doi.org/10.1016/s0924-9338\(10\)71572-8](http://dx.doi.org/10.1016/s0924-9338(10)71572-8).
28. Naser AY, Dahmash EZ, Al-Rousan R, Alwafi H, Alrawashdeh HM, Ghoul I, et al. Mental health status of the general population, healthcare professionals, and university students during 2019 coronavirus disease outbreak in Jordan: A cross-sectional study. *Brain Behav.* 2020;10:e01730, <http://dx.doi.org/10.1002/brb3.1730>.
29. Alkhamees AA, Alrashed SA, Alzunaydi AA, Almohimeed AS, Aljohani MS. The psychological impact of COVID-19 pandemic on the general population of Saudi Arabia. *Compr Psychiatry.* 2020;102:152192, <http://dx.doi.org/10.1016/j.comppsy.2020.152192>.
30. Krishnamoorthy Y, Nagarajan R, Saya GK, Menon V. Prevalence of psychological morbidities among general population, healthcare workers and COVID-19 patients amidst the COVID-19 pandemic: A systematic review and meta-analysis. *Psychiatry Res.* 2020;293:113382.
31. Qiu J, Shen B, Zhao M, Wang Z, Xie B, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *Gen. Psychiatry.* 2020;33:e100213.
32. Gunnell D, Appleby L, Arensman E, Hawton K, John A, Kapur N, et al. Suicide risk and prevention during the COVID-19 pandemic. *Lancet Psychiatr.* 2020;7:468-71, [http://dx.doi.org/10.1016/S2215-0366\(20\)30171-1](http://dx.doi.org/10.1016/S2215-0366(20)30171-1).
33. Kawohl W, Nordt C. COVID-19, unemployment, and suicide. *Lancet Psychiatr.* 2020;7:389-90, [http://dx.doi.org/10.1016/S2215-0366\(20\)30141-3](http://dx.doi.org/10.1016/S2215-0366(20)30141-3).
34. Cai Q, Feng H, Huang J, Wang M, Wang Q, Lu X, et al. The mental health of frontline and non-frontline medical workers during the coronavirus disease 2019 (COVID-19) outbreak in China: A case-control study. *J. Affect. Disord.* 2020;275:210-5.
35. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: Immediate psychological responses and associated factors. *Int. J. Environ. Res. Public Health.* 2020;17:3165, <http://dx.doi.org/10.3390/ijerph17093165>.
36. Losada-Baltar A, Jiménez-Gonzalo L, Gallego-Alberto L, Pedroso-Chaparro M, Fernandes-Pires J, Márquez-González M. 'We're staying at home'. Association of self-perceptions of aging, personal and family resources and loneliness with psychological distress during the lock-down period of COVID-19. *J. Gerontol B Psychol Sci Soc Sci.* 2020;76:e10-6, <http://dx.doi.org/10.1093/geronb/gbaa048>.