

ORIGINAL RESEARCH

Angiotensin II stimulates superoxide production by nitric oxide synthase in thick ascending limbs

Agustin Gonzalez-Vicente^{1,2,*}, Jagannath H. Saikumar^{3,*}, Katherine J. Massey³, Nancy J. Hong¹, Fernando P. Dominici^{2,4}, Oscar A. Carretero³ & Jeffrey L. Garvin^{1,3}

1 Department of Physiology and Biophysics, School of Medicine, Case Western Reserve University, Cleveland, Ohio

2 Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Ciudad Autónoma de Buenos Aires, Argentina

3 Department of Internal Medicine, Hypertension and Vascular Research Division, Henry Ford Hospital, Detroit, MI

4 Instituto de Química y Físicoquímica Biológicas, CONICET, Ciudad Autónoma de Buenos Aires, Argentina

Keywords

Kidney, NADPH oxidase, protein kinase C, reactive oxygen species, uncoupling.

Correspondence

Agustin Gonzalez-Vicente, Case Western Reserve University School of Medicine, Department of Physiology and Biophysics, 10900 Euclid Avenue, MS 4970, Cleveland, OH 44106-4970.

Tel: +216 368 1687

Fax: +216 368 5586

E-mail: agustin.gonzalezvicente@case.edu

Present addresses

Katherine J. Massey, Wayne State University, Detroit, Michigan

Jagannath H. Saikumar, Division of Nephrology, The University of Tennessee Health Science Center, Memphis, Tennessee

Funding Information

This work was supported by grants from the National Institutes of Health (HL 28982 and HL 70985) to J.L. Garvin.

Received: 14 September 2015; Revised: 17 December 2015; Accepted: 8 January 2016

doi: 10.14814/phy2.12697

Physiol Rep, 4 (4), 2016, e12697,
doi: 10.14814/phy2.12697

*These authors contributed equally to this manuscript.

Introduction

The reactive oxygen species nitric oxide (NO) and superoxide (O_2^-) have opposing physiological effects in the

Abstract

Angiotensin II (Ang II) causes nitric oxide synthase (NOS) to become a source of superoxide (O_2^-) via a protein kinase C (PKC)-dependent process in endothelial cells. Ang II stimulates both NO and O_2^- production in thick ascending limbs. We hypothesized that Ang II causes O_2^- production by NOS in thick ascending limbs via a PKC-dependent mechanism. NO production was measured in isolated rat thick ascending limbs using DAF-FM, whereas O_2^- was measured in thick ascending limb suspensions using the lucigenin assay. Consistent stimulation of NO was observed with 1 nmol/L Ang II ($P < 0.001$; $n = 9$). This concentration of Ang II-stimulated O_2^- production by 50% (1.77 ± 0.26 vs. 2.62 ± 0.36 relative lights units (RLU)/s/ μ g protein; $P < 0.04$; $n = 5$). In the presence of the NOS inhibitor L-NAME, Ang II-stimulated O_2^- decreased from 2.02 ± 0.29 to 1.10 ± 0.11 RLU/s/ μ g protein ($P < 0.01$; $n = 8$). L-arginine alone did not change Ang II-stimulated O_2^- (2.34 ± 0.22 vs. 2.29 ± 0.29 RLU/s/ μ g protein; $n = 5$). In the presence of Ang II plus the PKC α/β_1 inhibitor Gö 6976, L-NAME had no effect on O_2^- production (0.78 ± 0.23 vs. 0.62 ± 0.11 RLU/s/ μ g protein; $n = 7$). In the presence of Ang II plus apocynin, a NADPH oxidase inhibitor, L-NAME did not change O_2^- (0.59 ± 0.04 vs. 0.61 ± 0.08 RLU/s/ μ g protein; $n = 5$). We conclude that: (1) Ang II causes NOS to produce O_2^- in thick ascending limbs via a PKC- and NADPH oxidase-dependent process; and (2) the effect of Ang II is not due to limited substrate.

kidney. In the thick ascending limb of the loop of Henle, a nephron segment that is important in salt, water and acid/base homeostasis (Burg 1982; Greger 1985; Mount 2014), NO and O_2^- regulate ion transport (Garvin and

Ortiz 2003). NO promotes natriuresis and diuresis (Majid et al. 1993; Eitle et al. 1998; Plato et al. 1999; Ortiz et al. 2004b), whereas O₂⁻ enhances Na reabsorption (Zou et al. 2001; Majid and Nishiyama 2002; Juncos and Garvin 2005; Juncos et al. 2006; Silva et al. 2006) and water retention. Imbalances between these two factors in the kidney can lead to renal pathophysiological conditions such as hypertension (Sedeeq et al. 2003; Kopkan and Majid 2005; Kopkan et al. 2007; Majid and Kopkan 2007; Ramseyer et al. 2015) and progressive renal injury.

The thick ascending limb is one of the predominant sources of O₂⁻ in the kidney (Zou et al. 2001; Li et al. 2002). In this segment, most of the O₂⁻ is generated by NADPH oxidase 4 (Hong and Garvin 2012; Massey et al. 2012), a process involving several steps. Ultimately, an electron from NADPH is transferred to the oxidase sub-domain and then to molecular O₂, reducing it to O₂⁻. Thick ascending limb NADPH oxidase activity is enhanced by a number of stimuli including luminal flow (Hong and Garvin 2012), ion delivery (Hong and Garvin 2007), and angiotensin II (Ang II) (Li et al. 2002). We have previously shown that Ang II stimulates NADPH oxidase-dependent O₂⁻ production by binding to Ang II type 1 (AT1) receptors and activating protein kinase C (PKC) (Herrera et al. 2010). We have also shown that once O₂⁻ is generated, it can stimulate PKC activity and exert physiological effects including increasing Na reabsorption by thick ascending limbs (Silva et al. 2006). Although NADPH oxidase is the primary source of O₂⁻ after Ang II stimulation, short hairpin RNAs against and inhibitors of NADPH oxidase only reduce Ang II-enhanced O₂⁻ by about 70% suggesting that other sources are involved. Potential sources include NO synthase (NOS), cyclooxygenase, and the mitochondria (Zou et al. 2001; Li et al. 2002).

Thick ascending limbs express all three isoforms of NOS (Mount and Power 2006). NO synthesis involves several steps. Ultimately, electrons are transferred from NADPH via FAD and FMN to the oxidase domain of NOS, which contains a heme group. These electrons are then passed from the enzyme to the substrate, L-arginine, causing its reduction to L-citrulline and release of NO. If substrate or cofactors are lacking, under some circumstances NOS can produce O₂⁻ rather than NO because the electrons are transferred to molecular O₂ rather than L-arginine (Andrew and Mayer 1999). The balance between NO and O₂⁻ production by NOS can be regulated by phosphorylation by a number of kinases including PKC (Forstermann et al. 1994; Fleming 2010; Herrera and Garvin 2010; Forstermann and Sessa 2012; Ramseyer et al. 2015). The NOS inhibitor L-NAME blocks both NO and O₂⁻ generation by NOS. Several stimuli can enhance NOS activity in thick ascending limbs including Ang II

(Cabral et al. 2010, 2012; Herrera and Garvin 2010; Peng et al. 2014).

Similar to thick ascending limbs, Ang II stimulates both O₂⁻ production by NADPH oxidase and NO production by NOS in endothelial cells. Ang II also causes NOS to produce O₂⁻ (Mollnau et al. 2002; Oak and Cai 2007; Lobysheva et al. 2011; Lee et al. 2013; Galougahi et al. 2014; Saura et al. 2014) in endothelial cells. Thus, we hypothesized that Ang II induces O₂⁻ production by NOS in thick ascending limbs via a PKC- and NADPH oxidase-dependent process.

Materials and Methods

Animals

All protocols in this study were approved by the Case Western Reserve University and the Henry Ford Hospital Institutional Animal Care and Use Committees. All experiments were conducted in accordance with the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals. Male Sprague–Dawley rats (Charles River Breeding Laboratories, Kalamazoo, MI) were fed a diet containing 0.22% Na⁺ and 1.1% K⁺ for at least 6 days prior to the experiments. For terminal surgery, animals were anesthetized with ketamine (100 mg/kg bw IP) and xylazine (20 mg/kg bw IP), and given 2 IU heparin (IP). Animals were killed while still under anesthesia.

Drugs and buffers

Unless specified, all drugs and reagents were obtained from Sigma-Aldrich (St Louis, MO). The cell-permeant NO-selective fluorescent dye DAF-FM-diacetate was purchased from Invitrogen (Grand Island, NY). Coomassie Plus Protein Assay Reagent was obtained from Thermo-Scientific (Rockford, IL).

HEPES-buffered physiological saline contained (in mmol/L): 10 HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) (pH 7.5 at 22°C), 130 NaCl, 4 KCl, 2.5 NaH₂PO₄, 1.2 MgSO₄, 5.5 glucose, 6.0 DL-alanine, 2.0 Ca(lactate)₂, and 1.0 Na₃citrate. Osmolality was adjusted to 300 ± 5 mOsmol/L with mannitol.

Thick ascending limb isolation

Single medullary thick ascending limbs were isolated as previously described (Cabral et al. 2010, 2012). Briefly, animals weighing 120–150 g were anesthetized and the abdominal cavity opened. The left kidney was bathed in ice-cold 150 mmol/L NaCl, and then, immediately removed and placed in HEPES-buffered physiological saline at 4°C. Coronal slices were cut and individual thick

ascending limbs isolated from the outer medulla under a stereomicroscope at 4–8°C. Tubules ranging from 0.5–1.0 mm were transferred to a temperature-regulated chamber and maintained at 37 ± 1°C. The bath was exchanged at 0.6 ml/min.

Thick ascending limb suspensions

Suspensions were prepared as described before (Gonzalez-Vicente and Garvin 2013; Gonzalez-Vicente et al. 2014). Briefly, kidneys were perfused retrograde *via* the abdominal aorta with cold HEPES-buffered physiological saline containing 2 USP/ml heparin and 0.1% of Type 4 collagenase. Perfused kidneys were removed, coronal slices cut, and outer medullary tissue dissected and minced. Minced tissue was digested in 0.1% collagenase for 30 min at 37°C. During digestion, tissue was agitated and gassed with 100% O₂ every 5 min. The sample was then centrifuged (100 × g, 2 min, 4°C), and the resulting pellet of tubules resuspended in fresh HEPES-buffered physiological saline and stirred on ice for 30 min. After stirring, the sample was filtered through a 250 μm nylon mesh, and the filtered tubules were collected and rinsed at 4°C. This preparation resulted in a 95% pure suspension of thick ascending limbs.

NO production measurement

NO production was measured by fluorescence microscopy as previously described (Ramseyer et al. 2015). Briefly, manually dissected thick ascending limbs were transferred to a temperature-controlled chamber (37 ± 1°C) on the stage of an inverted microscope and fixed in place using glass pipettes for live cell microscopy (Diaphot TMJ, Nikon, Japan). The bath contained 100 μmol/l L-arginine, but tubules were not submitted to luminal flow. Under these conditions, NO production is negligible (Ortiz et al. 2001, 2004a). Tubules were loaded with the NO-sensitive fluorescent dye DAF-FM (2 μmol/L) for 15 min and then washed with physiological saline for 5 min. For imaging, a 40X oil immersion objective was used. The dye was excited by a xenon arc lamp with a 488 nm band pass filter. The fluorescence emitted by the NO-bound dye (>500 nm) was measured using Metafluor software (Molecular Devices, Sunnyvale CA). Results were expressed as arbitrary fluorescent units (AFU)/min.

Superoxide production measurement

O₂⁻ production was measured with bis-N-methylacridinium nitrate (lucigenin). Lucigenin is a validated luminescent probe to measure O₂⁻ (Skatchkov et al. 1999; Ohashi et al. 2015), and it has been used in our

laboratory before (Ortiz and Garvin 2002; Herrera et al. 2010; Silva and Garvin 2010; Massey et al. 2012). Briefly, for each experiment, lucigenin (5 μmol/L final concentration) was added to each of two plastic tubes containing 800 μl of warm-oxygenated physiological saline and kept in the dark at 37°C. Then, 200-μl aliquots of freshly prepared thick ascending limb suspension containing 100 to 200 μg of protein were added to each tube. The control tube was placed in a luminometer (FB12/Sirius; Zylux Oak Ridge, TN) and basal luminescence recorded at 4.8 sec intervals for 10 min using the provided software. Then, the O₂⁻ scavenger 4,5-dihydroxy-1,3-benzenedisulfonic acid (Tiron) was added to a final concentration of 10 mmol/L, and measurements were continued for 5 min. The average luminescence of the last 2 min following the addition of Tiron was then subtracted from the steady-state luminescence before Tiron. Tubules were recovered by centrifugation, and luminescence normalized for protein content.

In the experiments using Ang II, it was added before placing the tubes in the luminometer. In some experiments, we used the lipid-dependent PKC activator phorbol 12-myristate 13-acetate (PMA) instead of Ang II, which was also added before placing the tubes in the luminometer. For each set of experiments, the order of the control and treatment tubes was randomized.

For the experiments involving either L-arginine or L-NAME, the chemical was added during the generation of the suspensions. When Gö 6976 or apocynin was used, it was added 3 min before the addition of Ang II.

Statistical analysis

Results are expressed as the arithmetic mean ± the standard error of the mean. The Student t-tests was used to compare means. The *P*-values were calculated using two-tailed tests in all cases, and paired or unpaired test were used where appropriate. *P* < 0.05 was considered significant.

Results

To show that Ang II activates thick ascending limb NOS and stimulates NO production, and to study which concentration of Ang II should be used in subsequent experiments, we measured the effect of different concentrations (10⁻¹³, 10⁻¹¹, 10⁻⁹ mol/L) of Ang II on NO production by isolated thick ascending limbs. We found that within 5 min, 10⁻⁹ mol/L Ang II consistently increased NO production (*P* < 0.001; *n* = 9), while lower concentrations gave a highly variable response (Fig. 1).

Next, we tested whether a 5-min treatment with 10⁻⁹ mol/L Ang II increased O₂⁻ production in thick




Figure 1. Effect of acute angiotensin II (Ang II) treatment on NO production by isolated thick ascending limbs. Tubules were manually dissected, held with pipettes and NO production measured with DAF-FM. $P < 0.001$ versus basal; $n = 9$ for 1 nmol/L Ang II. AFU, arbitrary fluorescence units.

ascending limb suspensions in the absence of general NOS substrate L-arginine, to avoid any potential interference from endogenously produced NO. We found that within 5 minutes, 10⁻⁹ mol/L Ang II increased O₂⁻ production by about 50% from 1.77 ± 0.26 to 2.62 ± 0.36 relative light units (RLU)/s/μg protein ($P < 0.04$; $n = 5$; Fig. 2). Given that 5-min treatments with 10⁻⁹ mol/L Ang II are capable of stimulating both NO and O₂⁻, we used this timeframe and concentration in all other experiments.

Ang II stimulates O₂⁻ production by NOS in endothelial cells (Lobysheva et al. 2011). Thus, we next evaluated the contribution of NOS-derived O₂⁻ by measuring Ang II-induced O₂⁻ in the presence and absence of the general NOS inhibitor L-NAME (1 mmol/L). These experiments were conducted in the absence of L-arginine to avoid interference by NO. We found that Ang II-induced O₂⁻ production in tubules treated with vehicle was




Figure 2. Effect of 1 nmol/L Ang II on O₂⁻ production by thick ascending limb suspensions ($P < 0.04$; $n = 5$). Lucigenin plus Tiron were used to detect O₂⁻ production in the presence or absence of Ang II.

2.02 ± 0.29 RLU/s/μg protein, while in those treated with L-NAME, it was 1.10 ± 0.11 RLU/s/μg ($P < 0.01$; $n = 8$; Fig. 3). Thus, blocking NOS reduces Ang II-stimulated O₂⁻ more than 40%.

The results presented in Figure 3 indicate that Ang II stimulates O₂⁻ production by NOS. This can be due to a direct effect or due to lack of substrate, that is, L-arginine. To test whether Ang II-stimulated NOS-dependent O₂⁻ production was a result of lack of L-arginine, we repeated the experiments in the presence and absence of 100 μmol/L L-arginine. We found that Ang II-induced O₂⁻ production was not different in the presence (2.34 ± 0.22 RLU/s/μg protein) or absence (2.29 ± 0.29 RLU/s/μg protein) of L-arginine ($n = 5$; Fig. 4). These data indicate that Ang II-stimulated O₂⁻ production by NOS is not only a consequence of the lack of L-arginine to accept electrons, and thus other effects may be involved.

Previous reports indicate that PKC activation causes NOS to produce O₂⁻. Ang II can stimulate PKC activity in thick ascending limbs both directly (Herrera et al. 2010) and indirectly via O₂⁻ (Silva et al. 2006; Herrera et al. 2010). Therefore, we next tested whether PKC activation was required for Ang II to increase NOS-derived O₂⁻, by measuring Ang II-induced O₂⁻ in the presence and absence of L-NAME, while inhibiting PKCα/β₁ with 100 nmol/L Gö 6976 (Fig. 5). We found that in the presence of Ang II and Gö 6976, O₂⁻ production was unaltered by L-NAME (0.78 ± 0.23 vs. 0.62 ± 0.11 RLU/s/μg protein; $n = 7$). These data indicate that PKC activation is necessary for Ang II to stimulate O₂⁻ production by NOS.

Ang II can also stimulate PKC indirectly by increasing O₂⁻ derived from NADPH oxidase (Silva et al. 2006; Herrera et al. 2010). Apocynin reduces Ang II-stimulated O₂⁻ from NADPH oxidase (Herrera et al. 2010). Therefore, to test whether NADPH oxidase was involved in the




Figure 3. Effect of the NOS inhibitor L-NAME (1 mmol/L) on Ang II-induced O₂⁻ in thick ascending limb suspensions ($P < 0.01$; $n = 8$). Lucigenin plus Tiron were used to detect Ang II-stimulated O₂⁻.




Figure 4. Effect of the NOS substrate L-arginine (100 μ mol/L) on Ang II-induced O_2^- in thick ascending limb suspensions ($n = 5$). Lucigenin plus Tiron were used to detect Ang II-stimulated O_2^- .




Figure 5. Effect of the NOS inhibitor L-NAME (1 mmol/L) on Ang II-stimulated O_2^- in thick ascending limb suspensions in the presence of the PKC α/β_1 inhibitor Gö 6976 (100 nmol/L) ($n = 7$). Lucigenin plus Tiron were used to detect Ang II-stimulated O_2^- .

ability of Ang II to stimulate O_2^- by NOS, we measured Ang II-induced O_2^- in the presence and absence of L-NAME, when NADPH oxidase was blocked with 10 μ mol/L apocynin (Fig. 6). We found that in the presence of Ang II and apocynin, O_2^- production was not affected by L-NAME (0.59 ± 0.04 vs. 0.61 ± 0.08 RLU/s/ μ g protein; $n = 5$). These data indicate that NADPH oxidase activation is a necessary event for Ang II to increase O_2^- production by NOS in thick ascending limbs.

Finally, to test whether direct activation of lipid-dependent PKC isoforms was able to stimulate O_2^- production by NOS in our system, we measured the effect of PMA alone on O_2^- production, and the effect of L-NAME on PMA-stimulated O_2^- production in the presence of apocynin. We found that PMA-stimulated O_2^- production from 1.5 ± 0.3 to 12.8 ± 2.8 RLU/s/ μ g protein ($P < 0.05$, $n = 5$). L-NAME reduced PMA-stimulated O_2^- production in the presence of apocynin by




Figure 6. Effect of the NOS inhibitor L-NAME (1 mmol/L) on Ang II-stimulated O_2^- in thick ascending limb suspensions in the presence of the NADPH oxidase inhibitor apocynin (10 μ mol/L) ($n = 5$). Lucigenin plus Tiron were used to detect Ang II-stimulated O_2^- .

$49 \pm 15\%$ ($P < 0.05$, $n = 4$). These data indicate that lipid-dependent PKC isoform activation increases O_2^- production by NOS in the absence of NADPH oxidase-derived O_2^- in thick ascending limbs.

Discussion

We hypothesized that Ang II causes NOS to become a source of O_2^- and that this effect is dependent on PKC and NADPH oxidase. This implies that Ang II stimulates both O_2^- and NO production. Therefore, we began our study by directly measuring the effect of different concentrations of Ang II on NO production by rat thick ascending limbs. We found that 1 nmol/L Ang II stimulates thick ascending limb NO production within 5 minutes. Using this concentration of Ang II, we next tested whether it also stimulates O_2^- production. We found that 1 nmol/L Ang II stimulated O_2^- by approximately 50% in the same time frame as it stimulates NO. Taken together, these data suggest that there is a concomitant production of NO and O_2^- upon Ang II stimulation in the thick ascending limbs. The O_2^- experiments were performed in the absence of L-arginine, the substrate for NOS, because NO and O_2^- can react to form peroxynitrite if they are generated in the same compartment at the same time (Pryor and Squadrito 1995; Squadrito and Pryor 1995) and we have shown that NO can inhibit O_2^- production via a cGMP-dependent mechanism. The results showing that Ang II stimulates both NO and O_2^- in the same time frame are consistent with our previous studies in which we found that Ang II stimulates NO (Herrera and Garvin 2010) and O_2^- (Herrera et al. 2010; Massey et al. 2012). They are also consistent with findings

of other investigators (Mori and Cowley 2003; Lobysheva et al. 2011).

Although it is generally accepted that Ang II stimulates O₂⁻ in the thick ascending limb primarily by NADPH oxidase, this enzyme is not the only source of Ang II-stimulated O₂⁻ in this nephron segment. We and others have shown that other sources account for 10 (Mori and Cowley 2003; Herrera et al. 2010) to 40% (Massey et al. 2012) of the O₂⁻ produced by thick ascending limbs upon Ang II stimulation. However, in the previous studies, the identity of such sources was not determined. Therefore, we next investigated the contribution of NOS to the increase in O₂⁻ production caused by Ang II by studying whether L-NAME, a NOS inhibitor, alters the ability of Ang II to enhance O₂⁻ production. We found that blocking NOS reduced Ang II-stimulated O₂⁻ by 40%. These data suggest that Ang II stimulates NOS to produce O₂⁻.

Our results showing that Ang II causes NOS to produce O₂⁻ are similar to reports in other cell types (Lobysheva et al. 2011; Lee et al. 2013; Galougahi et al. 2014; Kossmann et al. 2014; Saura et al. 2014). NOS has been shown to be a source of O₂⁻ after Ang II treatment in endothelial cells (Lobysheva et al. 2011). Further support of this concept comes from studies in Ang II-induced hypertensive animals. For instance, NOS3 was upregulated at both the RNA and protein levels in the myocardium (Tambascia et al. 2001) and vessels (Mollnau et al. 2002; Sullivan et al. 2002) of Ang II-treated rats as compared to controls. However, these animals presented a marked decrease in vascular NO bioavailability as well as endothelial dysfunction, which is associated with elevated O₂⁻. Thus, elevated NOS does not necessarily lead to an increase in NO, and NOS-derived O₂⁻ may represent a mechanism whereby Ang II causes prolonged oxidative stress (Oak and Cai 2007).

An alternative explanation for our data could be that Ang II-stimulated NO production is required to activate O₂⁻ production from another source such as xanthine oxidase, cyclooxygenase, or the mitochondria. This seems unlikely because neither xanthine oxidase nor cyclooxygenase activities have been shown to be affected by NO or cGMP. While mitochondrial respiration has been shown to be regulated by NO, it reduces oxygen consumption by this organelle rather than increasing it (Boveris et al. 2003).

In our experiments, we showed that Ang II causes NOS to produce O₂⁻ in the absence of exogenously added L-arginine. Lack of substrate causes the active enzyme to transfer electrons not to L-arginine but to molecular oxygen thereby generating O₂⁻. Thus, we tested whether Ang II-stimulated O₂⁻ was altered in the presence or absence of L-arginine. We found that Ang II-induced O₂⁻ production was not significantly altered by L-arginine. These

data indicate that a simple lack of NOS substrate is not the main mechanism mediating the increase in O₂⁻ in our experimental conditions.

Since NO inhibits O₂⁻ production and that NO and O₂⁻ can react to form peroxynitrite in isolated perfused tubules, one might ask why adding L-arginine to the suspension media did not alter Ang II-stimulated O₂⁻ production via one of these pathways. The explanation is that unlike the isolated, perfused tubule preparation, where the bath is >10⁶ times the volume of the tubule and L-arginine is being washed out continuously, the thick ascending limb suspensions are always in a restricted space. This allows the tubules to retain enough endogenous L-arginine to support NO production during the experiment, given that the K_{1/2} for L-arginine is only about 2 μmol/L, and at the time of isolation, the thick ascending limb cells may contain as much as 300 μmol/L L-arginine.

A similar consideration should be made for the cofactor tetrahydrobiopterin (BH₄). Even though reductions in BH₄ can cause O₂⁻ production by NOS (Bec et al. 1998; Andrew and Mayer 1999; Adak et al. 2000), this seems unlikely in our experimental conditions. Addition of the BH₄ precursor sepiapterin to thick ascending limb suspensions has no effect on NO production (Herrera et al. 2006). This indicates that BH₄ is not a limiting factor in our preparation. Thus, although BH₄ levels are affected in models of chronic hypertension (Landmesser et al. 2003), including those dependent on Ang II (Kang et al. 2011), there is no evidence that acute exposure to Ang II *per se* reduces BH₄ levels. Thus, reduced cofactors availability is not likely the cause of NOS-derived O₂⁻ in this case.

Because Ang II activates PKC both directly and indirectly, and PKC activation can increase O₂⁻ production from NOS in other cells (Chen et al. 2014), we next tested whether PKC was necessary for Ang II to increase O₂⁻ production from NOS in thick ascending limbs. We found that when PKC was blocked, L-NAME had no effect on Ang II-stimulated O₂⁻. The importance of PKC as a mediator of O₂⁻ production is in agreement with our previous studies in thick ascending limbs (Silva et al. 2006; Herrera et al. 2010; Hong et al. 2010). However, this study is the first to identify a role in NOS-derived O₂⁻ production.

Ang II can indirectly activate PKC by stimulating NADPH oxidase activity. To test whether NADPH oxidase is required for Ang II to stimulate O₂⁻ production by NOS, we used apocynin. We found that apocynin prevented Ang II from enhancing O₂⁻ production by NOS. These data indicate that NADPH oxidase activity is required for Ang II's effect on NOS.

When taken together with published studies, the current PKC, apocynin, and PMA data suggest two possible

pathways by which Ang II treatment can lead to O_2^- production by NOS. Ang II binds AT1 receptors which activate PKC α (Herrera et al. 2010). PKC α then increases NADPH oxidase activity (Herrera et al. 2010; Hong et al. 2010; Massey et al. 2012). The O_2^- thus produced either: (1) further activates the same pool of PKC α which increases NOS phosphorylation; or (2) activates a different pool of PKC α/δ (Silva et al. 2006). PKC α/δ then phosphorylates NOS causing it to produce O_2^- . Therefore, according to this model, the PKC α directly activated by Ang II does not cause NOS to produce O_2^- because either: (1) it is in a different cellular compartment than the one that phosphorylates NOS; or (2) Ang II may simply not increase PKC α sufficiently to affect NOS. The proposed model is represented in Figure 7.

Another open question that cannot yet be answered is which NOS isoform is responsible from O_2^- production in response to Ang II; however, some conclusions can




Figure 7. Ang II-stimulated O_2^- production by NOS requires at least one of these pathways involving NADPH oxidase: (1) NADPH oxidase-derived O_2^- exerts a positive feedback over the PKC α pool stimulated by Ang II (dashed lines); or (2) NADPH oxidase-derived O_2^- stimulating a different pool of PKC α/δ (solid lines). Both pathways converge in the final step which is NOS phosphorylation by PKC.

be drawn based on published studies. First, NOS2 is mainly regulated at the transcriptional level and its abundance is at the limit of detection under nonstimulating conditions in the rat kidney (Zhang et al. 2000; Stumm et al. 2002), thereby making it unlikely to mediate any effect in acute experiments. Second, NOS1 is not phosphorylated by PKC (Okada 1996); PKC instead affects its sensitivity for calcium indirectly (Okada 1995), and exerts inhibitory rather than stimulatory effects (Riccio et al. 1996). Thus, NOS1 is not a likely candidate either. Finally, NOS3 is directly phosphorylated by PKC (Fleming et al. 2001; Chen et al. 2014) causing it to produce O_2^- (Lin et al. 2003; Chen et al. 2014). Taking all this into account, our data suggest that NOS3 mediates NOS-derived O_2^- in response to acute Ang II stimulation. Studies with double knockouts that would provide a definitive answer to this question are far out of the scope of this paper.

In summary, we found that Ang II acutely causes O_2^- production by NOS in thick ascending limbs. This process is dependent on PKC and NADPH oxidase and remains present in excess of L-arginine. Stimulation of NO and O_2^- by such factors as Ang II is important in regulating renal water and salt reabsorption. Understanding the processes involved in maintaining the balance between NO and O_2^- may give insight into the pathogenesis and treatment of diseases associated with oxidative stress and Na retention.

Conflict of Interest

None declared.

References

- Adak, S., Q. Wang, and D. J. Stuehr. 2000. Arginine conversion to nitroxide by tetrahydrobiopterin-free neuronal nitric-oxide synthase. Implications for mechanism. *J. Biol. Chem.* 275:33554–33561.
- Andrew, P. J., and B. Mayer. 1999. Enzymatic function of nitric oxide synthases. *Cardiovasc. Res.* 43:521–531.
- Bec, N., A. C. Gorren, C. Voelker, B. Mayer, and R. Lange. 1998. Reaction of neuronal nitric-oxide synthase with oxygen at low temperature. Evidence for reductive activation of the oxy-ferrous complex by tetrahydrobiopterin. *J. Biol. Chem.* 273:13502–13508.
- Boveris, A., L. B. Valdez, S. Alvarez, T. Zaobornyj, A. D. Boveris, and A. Navarro. 2003. Kidney mitochondrial nitric oxide synthase. *Antioxid. Redox Signal.* 5:265–271.
- Burg, M. B. 1982. Thick ascending limb of Henle's loop. *Kidney Int.* 22:454–464.
- Cabral, P. D., N. J. Hong, and J. L. Garvin. 2010. Shear stress increases nitric oxide production in thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 299:F1185–F1192.

- Cabral, P. D., N. J. Hong, and J. L. Garvin. 2012. ATP mediates flow-induced NO production in thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 303:F194–F200.
- Chen, F., S. Kumar, Y. Yu, S. Aggarwal, C. Gross, Y. Wang, et al. 2014. PKC-dependent phosphorylation of eNOS at T495 regulates eNOS coupling and endothelial barrier function in response to G + -toxins. *PLoS ONE* 9:e99823.
- Eitle, E., S. Hiranyachattada, H. Wang, and P. J. Harris. 1998. Inhibition of proximal tubular fluid absorption by nitric oxide and atrial natriuretic peptide in rat kidney. *Am. J. Physiol.* 274:C1075–C1080.
- Fleming, I. 2010. Molecular mechanisms underlying the activation of eNOS. *Pflugers Arch.* 459:793–806.
- Fleming, I., B. Fisslthaler, S. Dimmeler, B. E. Kemp, and R. Busse. 2001. Phosphorylation of Thr(495) regulates Ca²⁺/calmodulin-dependent endothelial nitric oxide synthase activity. *Circ. Res.* 88:E68–E75.
- Forstermann, U., and W. C. Sessa. 2012. Nitric oxide synthases: regulation and function. *Eur. Heart J.* 33:837a–837d.
- Forstermann, U., E. I. Closs, J. S. Pollock, M. Nakane, P. Schwarz, I. Gath, et al. 1994. Nitric oxide synthase isozymes. Characterization, purification, molecular cloning, and functions. *Hypertension* 23:1121–1131.
- Galougahi, K. K., C. C. Liu, C. Gentile, C. Kok, A. Nunez, A. Garcia, et al. 2014. Glutathionylation mediates angiotensin II-induced eNOS uncoupling, amplifying NADPH oxidase-dependent endothelial dysfunction. *J. Am. Heart Assoc.* 3:e000731.
- Garvin, J. L., and P. A. Ortiz. 2003. The role of reactive oxygen species in the regulation of tubular function. *Acta Physiol. Scand.* 179:225–232.
- Gonzalez-Vicente, A., and J. L. Garvin. 2013. Angiotensin II-induced hypertension increases plasma membrane Na pump activity by enhancing Na entry in rat thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 305:F1306–F1314.
- Gonzalez-Vicente, A., P. D. Cabral, and J. L. Garvin. 2014. Resveratrol increases nitric oxide production in the rat thick ascending limb via Ca²⁺/calmodulin. *PLoS ONE* 9:e110487.
- Greger, R. 1985. Ion transport mechanisms in thick ascending limb of Henle's loop of mammalian nephron. *Physiol. Rev.* 65:760–797.
- Herrera, M., and J. L. Garvin. 2010. Angiotensin II stimulates thick ascending limb NO production via AT(2) receptors and Akt1-dependent nitric-oxide synthase 3 (NOS3) activation. *J. Biol. Chem.* 285:14932–14940.
- Herrera, M., G. Silva, and J. L. Garvin. 2006. A high-salt diet dissociates NO synthase-3 expression and NO production by the thick ascending limb. *Hypertension* 47:95–101.
- Herrera, M., G. B. Silva, and J. L. Garvin. 2010. Angiotensin II stimulates thick ascending limb superoxide production via protein kinase C(alpha)-dependent NADPH oxidase activation. *J. Biol. Chem.* 285:21323–21328.
- Hong, N. J., and J. L. Garvin. 2007. Flow increases superoxide production by NADPH oxidase via activation of Na-K-2Cl cotransport and mechanical stress in thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 292:F993–F998.
- Hong, N. J., and J. L. Garvin. 2012. NADPH oxidase 4 mediates flow-induced superoxide production in thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 303:F1151–F1156.
- Hong, N. J., G. B. Silva, and J. L. Garvin. 2010. PKC-alpha mediates flow-stimulated superoxide production in thick ascending limbs. *Am. J. Physiol. Renal. Physiol.* 298:F885–F891.
- Juncos, R., and J. L. Garvin. 2005. Superoxide enhances Na-K-2Cl cotransporter activity in the thick ascending limb. *Am. J. Physiol. Renal. Physiol.* 288:F982–F987.
- Juncos, R., N. J. Hong, and J. L. Garvin. 2006. Differential effects of superoxide on luminal and basolateral Na⁺/H⁺ exchange in the thick ascending limb. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 290:R79–R83.
- Kang, K. T., J. C. Sullivan, F. T. Spradley, L. V. d'Uscio, Z. S. Katusic, and J. S. Pollock. 2011. Antihypertensive therapy increases tetrahydrobiopterin levels and NO/cGMP signaling in small arteries of angiotensin II-infused hypertensive rats. *Am. J. Physiol. Heart Circ. Physiol.* 300:H718–H724.
- Kopkan, L., and D. S. Majid. 2005. Superoxide contributes to development of salt sensitivity and hypertension induced by nitric oxide deficiency. *Hypertension* 46:1026–1031.
- Kopkan, L., Z. Huskova, Z. Vanourkova, M. Thumova, P. Skaroupkova, L. Cervenka, et al. 2007. Superoxide and its interaction with nitric oxide modulates renal function in prehypertensive Ren-2 transgenic rats. *J. Hypertens.* 25:2257–2265.
- Kossmann, S., H. Hu, S. Steven, T. Schonfelder, D. Fraccarollo, Y. Mikhed, et al. 2014. Inflammatory monocytes determine endothelial nitric-oxide synthase uncoupling and nitro-oxidative stress induced by angiotensin II. *J. Biol. Chem.* 289:27540–27550.
- Landmesser, U., S. Dikalov, S. R. Price, L. McCann, T. Fukai, S. M. Holland, et al. 2003. Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J. Clin. Invest.* 111:1201–1209.
- Lee, D. Y., F. Wauquier, A. A. Eid, L. J. Roman, G. Ghosh-Choudhury, K. Khazim, et al. 2013. Nox4 NADPH oxidase mediates peroxynitrite-dependent uncoupling of endothelial nitric-oxide synthase and fibronectin expression in response to angiotensin II: role of mitochondrial reactive oxygen species. *J. Biol. Chem.* 288:28668–28686.
- Li, N., F. X. Yi, J. L. Spurrier, C. A. Bobrowitz, and A. P. Zou. 2002. Production of superoxide through NADH oxidase in thick ascending limb of Henle's loop in rat kidney. *Am. J. Physiol. Renal. Physiol.* 282:F1111–F1119.
- Lin, M. I., D. Fulton, R. Babbitt, I. Fleming, R. Busse, K. A. Jr Pritchard, et al. 2003. Phosphorylation of threonine 497 in endothelial nitric-oxide synthase coordinates the coupling of

- L-arginine metabolism to efficient nitric oxide production. *J. Biol. Chem.* 278:44719–44726.
- Lobysheva, I., G. Rath, B. Sekkali, C. Bouzin, O. Feron, B. Gallez, et al. 2011. Moderate caveolin-1 downregulation prevents NADPH oxidase-dependent endothelial nitric oxide synthase uncoupling by angiotensin II in endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* 31:2098–2105.
- Majid, D. S., and L. Kopkan. 2007. Nitric oxide and superoxide interactions in the kidney and their implication in the development of salt-sensitive hypertension. *Clin. Exp. Pharmacol. Physiol.* 34:946–952.
- Majid, D. S., and A. Nishiyama. 2002. Nitric oxide blockade enhances renal responses to superoxide dismutase inhibition in dogs. *Hypertension* 39:293–297.
- Majid, D. S., A. Williams, and L. G. Navar. 1993. Inhibition of nitric oxide synthesis attenuates pressure-induced natriuretic responses in anesthetized dogs. *Am. J. Physiol.* 264:F79–F87.
- Massey, K. J., N. J. Hong, and J. L. Garvin. 2012. Angiotensin II stimulates superoxide production in the thick ascending limb by activating NOX4. *Am. J. Physiol. Cell Physiol.* 303:C781–C789.
- Mollnau, H., M. Wendt, K. Szocs, B. Lassegue, E. Schulz, M. Oelze, et al. 2002. Effects of angiotensin II infusion on the expression and function of NAD(P)H oxidase and components of nitric oxide/cGMP signaling. *Circ. Res.* 90:E58–E65.
- Mori, T., and A. W. Jr Cowley. 2003. Angiotensin II-NAD(P)H oxidase-stimulated superoxide modifies tubulovascular nitric oxide cross-talk in renal outer medulla. *Hypertension* 42:588–593.
- Mount, D. B. 2014. Thick ascending limb of the loop of henle. *Clin. J. Am. Soc. Nephrol.* 9:1974–1986.
- Mount, P. F., and D. A. Power. 2006. Nitric oxide in the kidney: functions and regulation of synthesis. *Acta Physiol.* 187:433–446.
- Oak, J. H., and H. Cai. 2007. Attenuation of angiotensin II signaling recouples eNOS and inhibits nonendothelial NOX activity in diabetic mice. *Diabetes* 56:118–126.
- Ohashi, K., E. Munetsuna, H. Yamada, Y. Ando, M. Yamazaki, N. Taromaru, et al. 2015. High fructose consumption induces DNA methylation at PPARalpha and CPT1A promoter regions in the rat liver. *Biochem. Biophys. Res. Commun.* 468:185–189.
- Okada, D. 1995. Protein kinase C modulates calcium sensitivity of nitric oxide synthase in cerebellar slices. *J. Neurochem.* 64:1298–1304.
- Okada, D. 1996. Differential effects of protein kinase C on neuronal nitric oxide synthase activity in rat cerebellar slices and in vitro. *J. Chem. Neuroanat.* 10:213–220.
- Ortiz, P. A., and J. L. Garvin. 2002. Superoxide stimulates NaCl absorption by the thick ascending limb. *Am. J. Physiol. Renal. Physiol.* 283:F957–F962.
- Ortiz, P. A., N. J. Hong, and J. L. Garvin. 2001. NO decreases thick ascending limb chloride absorption by reducing Na (+)-K(+)-2Cl(-) cotransporter activity. *Am. J. Physiol. Renal. Physiol.* 281:F819–F825.
- Ortiz, P. A., N. J. Hong, and J. L. Garvin. 2004a. Luminal flow induces eNOS activation and translocation in the rat thick ascending limb. *Am. J. Physiol. Renal. Physiol.* 287:F274–F280.
- Ortiz, P. A., N. J. Hong, and J. L. Garvin. 2004b. Luminal flow induces eNOS activation and translocation in the rat thick ascending limb. II. Role of PI3-kinase and Hsp90. *Am. J. Physiol. Renal. Physiol.* 287:F281–F288.
- Peng, X. L., W. Qu, L. Z. Wang, B. Q. Huang, C. J. Ying, X. F. Sun, et al. 2014. Resveratrol ameliorates high glucose and high-fat/sucrose diet-induced vascular hyperpermeability involving Cav-1/eNOS regulation. *PLoS ONE* 9:e113716.
- Plato, C. F., B. A. Stoos, D. Wang, and J. L. Garvin. 1999. Endogenous nitric oxide inhibits chloride transport in the thick ascending limb. *Am. J. Physiol.* 276:F159–F163.
- Pryor, W. A., and G. L. Squadrito. 1995. The chemistry of peroxynitrite: a product from the reaction of nitric oxide with superoxide. *Am. J. Physiol.* 268:L699–L722.
- Ramseyer, V. D., A. Gonzalez-Vicente, O. A. Carretero, and J. L. Garvin. 2015. Angiotensin II-induced hypertension blunts thick ascending limb NO production by reducing NO synthase 3 expression and enhancing threonine 495 phosphorylation. *Am. J. Physiol. Renal. Physiol.* 308:F149–F156.
- Riccio, A., E. Esposito, and M. L. Eboli. 1996. Modulation by protein kinase C of nitric oxide and cyclic GMP production in cultured cerebellar granule cells. *Brain Res.* 718:159–164.
- Saura, M., S. Marquez, P. Reventun, N. Olea-Herrero, M. I. Arenas, R. Moreno-Gomez-Toledano, et al. 2014. Oral administration of bisphenol A induces high blood pressure through angiotensin II/CaMKII-dependent uncoupling of eNOS. *FASEB J.* 28:4719–4728.
- Sedeek, M. H., M. T. Llinas, H. Drummond, L. Fortepiani, S. R. Abram, B. T. Alexander, et al. 2003. Role of reactive oxygen species in endothelin-induced hypertension. *Hypertension* 42:806–810.
- Silva, G. B., and J. L. Garvin. 2010. Rac1 mediates NaCl-induced superoxide generation in the thick ascending limb. *Am. J. Physiol. Renal. Physiol.* 298:F421–F425.
- Silva, G. B., P. A. Ortiz, N. J. Hong, and J. L. Garvin. 2006. Superoxide stimulates NaCl absorption in the thick ascending limb via activation of protein kinase C. *Hypertension* 48:467–472.
- Skatchkov, M. P., D. Sperling, U. Hink, A. Mulsch, D. G. Harrison, I. Sindermann, et al. 1999. Validation of lucigenin as a chemiluminescent probe to monitor vascular superoxide as well as basal vascular nitric oxide production. *Biochem. Biophys. Res. Commun.* 254:319–324.
- Squadrito, G. L., and W. A. Pryor. 1995. The formation of peroxynitrite in vivo from nitric oxide and superoxide. *Chem. Biol. Interact.* 96:203–206.

- Stumm, M. M., D. D'Orazio, L. T. Sumanovski, P. Y. Martin, J. Reichen, and C. C. Sieber. 2002. Endothelial, but not the inducible, nitric oxide synthase is detectable in normal and portal hypertensive rats. *Liver* 22:441–450.
- Sullivan, J. C., D. M. Pollock, and J. S. Pollock. 2002. Altered nitric oxide synthase 3 distribution in mesenteric arteries of hypertensive rats. *Hypertension* 39:597–602.
- Tambascia, R. C., P. M. Fonseca, P. D. Corat, H. Jr Moreno, M. J. Saad, and K. G. Franchini. 2001. Expression and distribution of NOS1 and NOS3 in the myocardium of angiotensin II-infused rats. *Hypertension* 37:1423–1428.
- Zhang, C., L. M. Walker, and P. R. Mayeux. 2000. Role of nitric oxide in lipopolysaccharide-induced oxidant stress in the rat kidney. *Biochem. Pharmacol.* 59:203–209.
- Zou, A. P., N. Li, and A. W. Jr Cowley. 2001. Production and actions of superoxide in the renal medulla. *Hypertension* 37:547–553.