

Short report

Open Access

Quantitative expression analysis of HHV-6 cell receptor CD46 on cells of human cord blood, peripheral blood and G-CSF mobilised leukapheresis cells

Stefanie Thulke*¹, Aleksandar Radonić², Andreas Nitsche³ and Wolfgang Siegert⁴

Address: ¹Charité-Universitätsmedizin Berlin, CCM – Medizinische Klinik m.S. Onkologie/Hämatologie, Charitéplatz 1, 10117 Berlin, Germany, ²Charité-Universitätsmedizin Berlin, CCM – Medizinische Klinik m.S. Onkologie/Hämatologie, Charitéplatz 1, 10117 Berlin, Germany, ³Robert Koch Institut, ZBS 1, Nordufer 20, 13353 Berlin, Germany and ⁴Charité-Universitätsmedizin Berlin, CCM – Medizinische Klinik m.S. Onkologie/Hämatologie, Charitéplatz 1, 10117 Berlin, Germany

Email: Stefanie Thulke* - stefanie.thulke@charite.de; Aleksandar Radonić - aleksandar.radonic@charite.de; Andreas Nitsche - NitscheA@rki.de; Wolfgang Siegert - wolfgang.siegert@t-online.de

* Corresponding author

Published: 19 September 2006

Received: 20 January 2006

Virology Journal 2006, 3:77 doi:10.1186/1743-422X-3-77

Accepted: 19 September 2006

This article is available from: <http://www.virologyj.com/content/3/1/77>

© 2006 Thulke et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Human herpesvirus-6 (HHV-6) can infect blood cells and thereby may inhibit hematopoietic stem and progenitor cell expansion and differentiation. In this context, it has been discussed if early progenitor cells can be infected by HHV-6. CD46 was identified as one possible cellular surface receptor for HHV-6. The study presented here had been done to get insight into the susceptibility of various leukocyte subpopulations to HHV-6 (including early hematopoietic progenitors) by determining the amount of CD46 molecules expressed on their surfaces. Human cord blood cells, peripheral blood cells and G-CSF mobilised progenitor cells were analysed by flow cytometry. CD46 molecule number per cell was determined and compared to calibration beads conjugated with known ratio of PE per bead. Highest CD46 expression was detected on B- lymphocytes, whereas T-lymphocytes only showed about half of the amount found on B cells. Hematopoietic progenitors also carried CD46 at intermediate levels. Unexpectedly, CD46 expression on progenitors from G-CSF mobilised leukapheresis products was approximately 20% of that found on comparable cells from untreated cord blood. In conclusion, hematopoietic progenitor cells express CD46 on their surface, thereby fulfilling a basic requirement for the susceptibility of HHV-6 infection.

Findings

Human herpesvirus-6 (HHV-6) was first isolated in 1986 [1]. To date all HHV-6 isolates can be differentiated into the variants HHV-6A and HHV-6B. In early childhood HHV-6B infection causes *exanthema subitum* and febrile illness. After primary infection, HHV-6 persists life-long in host cells and may be reactivated under conditions of

immunosuppression, thereby causing various, in some extend life-threatening diseases, including mononucleosis, lymphoid/hematopoietic diseases, myelosuppression, encephalitis, pulmonitis and hepatitis [2]. HHV-6 induced myelosuppression, as occurring after stem cell transplantation, is recognised by leuko- and thrombocytopenia. Moreover, we showed that early HHV-6B infec-

tions may contribute to delayed platelet engraftment after stem cell transplantation [3]. There have been evidences for latently HHV-6 infected hematopoietic progenitors reactivating HHV-6 replication within the graft [4,5]. Our own studies showed that HHV-6A as well as HHV-6B are able to infect cord blood (CB) derived mononuclear cells and thereby inhibit *in vitro* expansion of the total cell number and of BFU-e, CFU-GM, as well as CD34⁺ or CD33⁺ cells. Contrariwise we could show only less HHV-6 mediated inhibition of CD34⁺ cell expansion of MACS separated CB CD34⁺ cells [6]. So far we have had no success to show an infected CD34⁺ cell. In order to clarify the role of HHV-6 in early hematopoietic stem cell development, we were interested in determining the level of differentiation when blood cells, especially early hematopoietic progenitor cells, became susceptible for HHV-6 infection. We believed that this question might be answered by quantifying HHV-6 membrane receptor CD46. CD46 was identified as cellular surface receptor for HHV-6 in 1999 [7] by interaction with viral glycoprotein complex gH-gL-gQ [8]. CD46 is the cellular receptor for further pathogens: Measles virus, group B adenoviruses [9] and other pathogenic microorganisms [10,11]. It is also known to act as a membrane cofactor for factor-I proteolytic cleavage of C3b and C4b in complement activation. CD46 also affects various cellular activities in response to pathogen or complement binding, and thus influences the host response to infection [12]. Recently, analysis of the short consensus repeat (SCR) regions that comprise most of the extracellular domain of CD46, was shown to have an essential role of SCR 2 and 3 in HHV-6 receptor activity [13].

We analysed samples of peripheral blood (PB) of healthy adults, CB, G-CSF mobilised peripheral blood progenitor cells collected as leukapheresis product (LP), and the HHV-6 infectable cell lines KG-1 and CRF-HSB-2 with regard to CD46 expression. In addition, we analysed CD34⁺ hematopoietic precursor cells purified by MACS separation (Miltenyi Biotec GmbH, Bergisch Gladbach, Germany). Heparinised blood was diluted 1:10 in FACS lysing Solution (Becton Dickinson, Heidelberg, Germany) to lyse erythrocytes and cells were washed twice in PBS. The cells were stained with the following monoclonal antibodies (mAB) for 15 min at room temperature: R-PE conjugated anti-CD46 (Cymbus Biotechnology LTD, Chandlers Ford, UK) and PerCP conjugated anti-CD45 (Becton Dickinson). To characterise different blood cell types, cells were stained with the following FITC conjugated mAB: anti-CD3, anti-CD8, anti-CD13, anti-CD15, anti-CD19, anti-CD22, anti-CD28, anti-CD33, anti-CD38, anti-CD45, anti-CD65 (Beckman Coulter GmbH, Unterschleissheim-Lohhof, Germany), anti-CD4, anti-CD14, (Becton Dickinson), anti-CD34 (Miltenyi Biotec GmbH). All FACS analyses were performed using the FAC-

Scalibur (Becton Dickinson). Levels of CD46 expression were determined in reference to calibration beads conjugated with a known ratio of PE per bead (QuantiBRITE PE conjugated beads, Becton Dickinson).

Levels of CD46 obtained on T and B-lymphocytes of CB, PB and LP are shown in figure 1. CD46 expression on B-lymphocytes (CD22⁺, CD19⁺ cells) was significantly higher than on T-lymphocytes from CB and PB. Highest CD46 expression levels were detected on CD22⁺ B-cells, the median number of molecules per CD22⁺ cell in CB, PB and LP was 11,650 (range, 10,702–14,158), 17,490 (range, 13,772–19,997) and 9,618 (range, 4,339–10,774), respectively. On CD19⁺ B-cells the median number of CD46 molecules per cell in CB, PB and LP was 9190 (range, 5,807–11,437), 11,060 (range, 10,378–




Figure 1

Detection of CD46 molecules on cell membranes of mature leukocytes from cord blood (CB), [A, squares], peripheral blood of adult donors (PB), [B, circles] and leukapheresis products (LP) from G-CSF mobilised precursor cells [C, triangles]. Statistical analysis was performed by paired t-test: *** $p < 0.001$, ** $0.001 < p < 0.01$ and * $0.001 < p < 0.05$. Median values are indicated as horizontal bars.

12,570) and 8,033 (range, 4,718–8,205), respectively. Lowest numbers of CD46 antigen levels were detected on CD8⁺T-cells from CB, PB and LP, i.e. 2,851 (range, 1,506–4,604), 2,965 (range, 2,451–4,343) and 2,442 (range, 2,079–3,053), respectively. Expression of CD46 on CD13⁺ myeloid cells was similar to CD3⁺ and CD4⁺ lymphocytes. CD46 antigen expression on CD34⁺ and CD38⁺ precursors and CD33⁺ cells is given in figure 2. Depending on the cell source or on the application of MACS separation, CD46 levels varied considerably. CD34⁺, CD38⁺ and CD33⁺ cells from CB expressed significantly more CD46 than corresponding cells from LP or from CB after MACS separation. The median number of CD46 molecules per CD34⁺ cell in CB, LP and CB after MACS separation was 6,232 (range, 5,219–7,956), 1,715 (range, 1,494–2,822) and 2,074 (range, 1,418–3,621), the number per CD38⁺ cell was 7,141 (range, 4,975–10,730), 2,195 (range, 1395–4058) and 2,169 (range, 1,945–3,665) and the number per CD33⁺ cell was 4,828 (range, 3,332–8455), 2,760 (range, 1,128–4,392) and 2,913 (range, 1,552–4,133), respectively. In addition, the T lymphoid cell line CRF-HSB-2 and the myeloid KG-1 cell line expressed 29,245 and 38,141 CD46 molecules per cell.

Our experiments show that mature lymphocytes and myeloid cells, as well as hematopoietic progenitor cells, express CD46. B-lymphocytes express higher levels of CD46 than T-lymphocytes; CD8⁺ T-lymphocytes exhibit less CD46 than CD4⁺ lymphocytes. Despite the discovery of HHV-6 as a B-lymphotropic virus, this does not correlate with the common view in the literature that T-lymphocytes would be *in vivo* and *in vitro* the preferred cells for HHV-6 infection [14]. Santoro *et al.* [15] suggested the existence of additional cellular factors, possibly co-receptors that are crucial for HHV-6 infection. However CD34⁺ and CD38⁺ hematopoietic progenitor cells from untreated CB express CD46 levels comparable to CD4⁺ cells. Thus, there is evidence that CD34⁺ progenitor cells as well as mature leukocytes carry HHV-6 receptors and thereby fulfil the basic requirement for susceptibility to HHV-6 infection.

The CD46 expression level on progenitor cells decreases to approximately one third on CD34⁺ and CD38⁺ cells from patients after G-CSF induced stem cell mobilisation and leukapheresis. Similarly, CD46 expression is reduced after immune affinity selection of CD34⁺ cells by MACS separation. We cannot explain why CD34⁺ cells in LP and CB after MACS separation bear lower amounts of CD46




Figure 2
 Detection of CD46 molecules on cell membranes of CD34⁺, CD38⁺, CD33⁺ hematopoietic progenitor cells from cord blood (CB) [squares], peripheral blood of adult donors (PB) [circles], leukapheresis products (LP) from G-CSF mobilised precursor cells [triangles] and MACS sorted CB CD34⁺ cells [diamonds]. Statistical analysis was performed by unpaired t-test: *** p < 0.001 and ** 0.001 < p < 0.01. Median values are indicated as horizontal bars.

than their native, non-manipulated counterparts. We cannot exclude that either *in vitro* manipulations lead to antigen down-regulation, antigen loss, sterical hindrance of antigen recognition, conformational change or that *in vivo* G-CSF treatment leads to a reduction of CD46 expression on the cell membrane. Seya *et al.* [16] showed a decrease of CD46 expression on leukaemia cell lines by *in vitro* G-CSF treatment.

Summing up, our results show significant expression of CD46 on various types of blood leukocytes including hematopoietic progenitor cells. Consequently these cells are fulfilling a requirement for HHV-6 infection. However, the level of expression appears not to be the only criterion for susceptibility to HHV-6.

Authors' contributions

ST contributed to the sample collection, performed FACS measurements, analysed the results and devised the manuscript.

AR contributed to study design and assisted the experiments as well as data analysis.

AN contributed to study design and mainly revised the manuscript.

WS composed the initial conception and contributed to data interpretation and manuscript revision.

All authors read and approved the final manuscript.

Acknowledgements

We gratefully acknowledge the technical assistance of Delia Barz.

This work was supported by a grant from Deutsche Krebshilfe (10-1362-Si I).

References

1. Salahuddin SZ, Ablashi DV, Markham PD, Josephs SF, Sturzenegger S, Kaplan M, Halligan G, Biberfeld P, Wong-Staal F, Kramarsky B, : **Isolation of a new virus, HBLV, in patients with lymphoproliferative disorders.** *Science* 1986, **234**:596-601.
2. Campadelli-Fiume G: **Virus receptor arrays, CD46 and human herpesvirus 6.** *Trends Microbiol* 2000, **8**:436-438.
3. Radonic A, Oswald O, Thulke S, Brockhaus N, Nitsche A, Siegert W, Schetelig J: **Infections with human herpesvirus 6 variant B delay platelet engraftment after allogeneic haematopoietic stem cell transplantation.** *Br J Haematol* 2005, **131**:480-482.
4. Andre-Garnier E, Milpied N, Boutolleau D, Saiagh S, Billaudel S, Imbert-Marcille BM: **Reactivation of human herpesvirus 6 during ex vivo expansion of circulating CD34+ haematopoietic stem cells.** *J Gen Virol* 2004, **85**:3333-3336.
5. Luppi M, Barozzi P, Morris C, Maiorana A, Garber R, Bonacorsi G, Donelli A, Marasca R, Tabilio A, Torelli G: **Human herpesvirus 6 latently infects early bone marrow progenitors in vivo.** *J Virol* 1999, **73**:754-759.
6. Nitsche A, Fleischmann J, Klima KM, Radonic A, Thulke S, Siegert W: **Inhibition of cord blood cell expansion by human herpesvirus 6 in vitro.** *Stem Cells Dev* 2004, **13**:197-203.
7. Santoro F, Kennedy PE, Locatelli G, Malnati MS, Berger EA, Lusso P: **CD46 is a cellular receptor for human herpesvirus 6.** *Cell* 1999, **99**:817-827.
8. Mori Y, Yang X, Akkapaiboon P, Okuno T, Yamanishi K: **Human Herpesvirus 6 Variant A Glycoprotein H-Glycoprotein L-Glycoprotein Q Complex Associates with Human CD46.** *J Virol* 2003, **77**:4992-4999.
9. Gaggar A, Shayakhmetov DM, Lieber A: **CD46 is a cellular receptor for group B adenoviruses.** *Nat Med* 2003, **9**:1408-1412.
10. Okada N, Liszewski MK, Atkinson JP, Caparon M: **Membrane cofactor protein (CD46) is a keratinocyte receptor for the M protein of the group A streptococcus.** *Proc Natl Acad Sci U S A* 1995, **92**:2489-2493.
11. Kallstrom H, Liszewski MK, Atkinson JP, Jonsson AB: **Membrane cofactor protein (MCP or CD46) is a cellular pilus receptor for pathogenic Neisseria.** *Mol Microbiol* 1997, **25**:639-647.
12. Russell S: **CD46: a complement regulator and pathogen receptor that mediates links between innate and acquired immune function.** *Tissue Antigens* 2004, **64**:111-118.
13. Greenstone HL, Santoro F, Lusso P, Berger EA: **Human herpesvirus 6 and measles virus employ distinct CD46 domains for receptor function.** *J Biol Chem* 2002.
14. Lusso P: **Human herpesvirus 6 (HHV-6).** *Antiviral Res* 1996, **31**:1-21.
15. Seya T, Hara T, Matsumoto M, Akedo H: **Quantitative analysis of membrane cofactor protein (MCP) of complement. High expression of MCP on human leukemia cell lines, which is down-regulated during cell differentiation.** *J Immunol* 1990, **145**:238-245.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

