

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Correspondence

Management of furcation-involved molar using Er:YAG laser-assisted bone regenerative therapy: A case report



Periodontitis is a multifactorial chronic disease. Furcation involvement (FI) is one of the most detrimental factors affecting tooth survival rate over time.¹ Recently, the dental application of Er:YAG lasers has increased,² especially in periodontal therapy.^{3,4} The Er:YAG laser effectively ablates soft and hard tissues and calculus on root and implant surfaces and effectively debrides bony defects in restricted areas, including furcation defects. Blood clots that form in the superficial areas of grafted bone when Er:YAG laser irradiation is used without water spray are convenient for bone regenerative procedures. Er:YAG laser-assisted bone regenerative therapy (Er-LBRT) is effective.⁵ This case report identified a potential method for degree III FI therapy for patients who refuse extraction.

A 52-year-old male complained of pain at tooth #36 during biting. The pocket depth was 12 mm buccally and lingually with degree III FI, bleeding on probing (BOP), and mobility I. The pocket depth was 3 mm at the mesial and distal sites buccally and lingually (Fig. 1A). Extraction was recommended due to the unfavorable prognosis of the tooth. As the patient desired to save the tooth, periodontal regenerative therapy was offered. Endodontic treatment and scaling and root planing (SRP) were initially performed, followed by a re-evaluation and surgery three months later. Local debridement at the furcation bone defect was achieved using Er:YAG laser (Erwin AdvErL™, J. Morita Mfg. Co., Osaka, Japan, Tip: PSM 600T) at 20 Hz and 50 mJ/pulse (Panel setting: 70 mJ/pulse) with water spray in near contact irradiation mode (Fig. 1B) and a mini curette. No

thermal damage was observed on the root surface. A 9-mm vertical bone defect was observed at the furcation area (Fig. 1C and D). Human freeze-dried bone allograft (FDBA, OraGRAFT®, LifeNet Health) and collected blood were mixed and grafted into the bony defect at the crest of the furcation (Fig. 1E), then the blood clot forming procedure was performed using Er:YAG laser irradiation without water spray in non-contact mode (Fig. 1F). The flaps were sutured (Fig. 1G). The soft tissue showed favorable wound healing after two weeks (Fig. 1H). Three years postoperatively, the soft tissue had healed favorably and the pocket depth was reduced to 4 mm at the buccal site and 3 mm at the lingual site without BOP or mobility (Fig. 1I). The periapical radiograph showed severe bone loss to the mesial root (Fig. 1J) preoperatively that was improved at three months (Fig. 1K). Dense, bone-like tissue was observed after the regenerative treatment. The original bone defects were successfully repaired, though some defects remained up to three years postoperatively (Fig. 1M, N, and O). The patient provided informed consent and the study was approved by the appropriate review board.

A molar with degree III FI is an advanced condition with an unfavorable prognosis. Er:YAG laser is a useful and effective treatment for granulation tissue and furcation root surface debridement within a restricted area. The Er-LBRT procedure simplifies the clinical procedure and can be used to treat molars with degree III FI if the interproximal bone level is acceptable for a trial of periodontal regenerative therapy.

<https://doi.org/10.1016/j.jds.2022.05.024>

1991-7902/© 2022 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1 Degree III furcation defect treated with Er:YAG laser-assisted bone regenerative therapy. (A) Baseline measurements. (B) A full-thickness flap is raised at the furcation bone defect at tooth #36, and the granulation tissue is debrided using Er:YAG laser and a mini curette. (C, D) A furcation defect is observed. The vertical bony defect is 9 mm following endodontic treatment and SRP initial therapy. (E) After debridement, bone material is grafted into the bony defect to the crest. (F) A blood clot forming procedure was performed using Er:YAG laser irradiation without water spray under non-contact mode. (G) The flaps are sutured. (H) At two weeks postoperatively, the soft tissue healing is favorable. (I) At three years postoperatively, the results are favorable. (J) The periapical radiograph obtained before treatment is shown. (K) A periapical radiograph obtained three months after the endodontic treatment and SRP is shown. (L) A dense, bone-like tissue is observed at tooth #36 furcation bone defect after surgery. (M, N, O) Radiographs obtained one, two, and three years after surgery show that the original bone defect at tooth #36 is successfully repaired, though some defect remains.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article.

References

1. Hamp SE, Nyman S, Lindhe J. Periodontal treatment of multirrooted teeth. Results after 5 years. *J Clin Periodontol* 1975;2:126–35.
2. Ng MY, Lin T, Yu CC, Chen CC. Er:YAG laser-assisted flapless esthetic crown lengthening procedure: a case report. *J Dent Sci* 2022;17:622–3.
3. Aoki A, Mizutani K, Mikami R, et al. Residual periodontal pocket treatment with Er:YAG laser-assisted comprehensive periodontal pocket therapy: a retrospective study. *Clin Oral Invest* 2022;26:761–71.
4. Lin T, Taniguchi Y, Aoki A, Chen CC. The "Er:YAG laser-assisted periimplantitis total therapy (Er:LPTT)"-a novel procedure to perform periimplantitis treatment with bone regeneration therapy. *J Dent Sci* 2021;16:1302–4.
5. Taniguchi Y, Aoki A, Sakai K, Mizutani K, Meinzer W, Izumi Y. A novel surgical procedure for Er:YAG laser-assisted periodontal regenerative therapy: case series. *Int J Periodontics Restor Dent* 2016;36:507–15.

Taichen Lin*
School of Dentistry, Chung Shan Medical University,
Taichung, Taiwan
Department of Dentistry, Chung Shan Medical University
Hospital, Taichung, Taiwan

Department of Periodontology, Graduate School of Medical
and Dental Science, Tokyo Medical and Dental University,
Japan

Yoichi Taniguchi
Department of Periodontology, Graduate School of Medical
and Dental Science, Tokyo Medical and Dental University,
Japan
Taniguchi Dental Clinic, Sapporo, Hokkaido, Japan

Akira Aoki
Department of Periodontology, Graduate School of Medical
and Dental Science, Tokyo Medical and Dental University,
Japan

Chun-Cheng Chen
School of Dentistry, Chung Shan Medical University,
Taichung, Taiwan
Department of Dentistry, Chung Shan Medical University
Hospital, Taichung, Taiwan

*Corresponding author. School of Dentistry, Chung Shan
Medical University, No.110, Sec.1, Jianguo N. Rd., Tai-
chung, 40201, Taiwan.
E-mail address: linperi@csmu.edu.tw (T. Lin)

Received 30 April 2022
Final revision received 28 May 2022
Available online 18 June 2022