

Effects of spaceflight on cancellous and cortical bone in proximal femur in growing rats[☆]

Amanda Gamboa^a, Adam J. Branscum^b, Dawn A. Olson^a, Lara H. Sattgast^a,
Urszula T. Iwaniec^{a,c}, Russell T. Turner^{a,c,*}

^a Skeletal Biology Laboratory, School of Biological and Population Health Sciences, Oregon State University, Corvallis, OR 97331, USA

^b Biostatistics Program, School of Biological and Population Health Sciences, Oregon State University, Corvallis, OR 97331, USA

^c Center for Healthy Aging Research, Oregon State University, Corvallis, OR 97331, USA

ARTICLE INFO

Keywords:

Spaceflight
Proximal femur
Bone microarchitecture
X-ray microtomography/micro-computed tomography
Bone volume/tissue volume
Rat

ABSTRACT

Mechanical loading of the skeleton during normal weight bearing plays an important role in bone accrual and turnover balance. We recently evaluated bone microarchitecture in the femoral head in 5.6-week-old male Sprague Dawley rats subjected to a 4-day spaceflight aboard STS-41. Compared to weight bearing ground controls, cancellous bone volume fraction was dramatically lower in animals subjected to microgravity. The effects of spaceflight on the rat skeleton are potentially influenced by factors such as age, duration of flight, strain and sex. To test the generalizability of our initial observation, we evaluated archived proximal femora from two additional spaceflight missions: a 10-day mission (STS-57) with 7.5-week-old male Fisher 344 rats, and a 14-day mission (STS-62) with 12-week-old ovariectomized (ovx) female Fisher 344 rats. Cancellous microarchitecture and cortical thickness were assessed using x-ray microtomography/microcomputed tomography. In male rats, cancellous bone volume fraction (bone volume/tissue volume) was lower in flight animals compared to flight controls, but differences were not significant compared to baseline. In ovx female rats, cancellous bone volume fraction was lower in flight animals compared to flight controls and baseline, indicating net bone loss. Cortical thickness did not differ among groups in either experiment. In summary, findings from three separate studies support the conclusion that spaceflight results in cancellous osteopenia in femoral head of growing rats.

1. Introduction

Life on Earth evolved in the presence of a nearly uniform gravitational field (Turner, 2000). Not surprisingly, the near absence of ‘weight’ during orbital spaceflight results in profound adaptive responses in multiple organ systems, including the skeleton. Abnormalities in bone and mineral homeostasis in astronauts during spaceflight result in net bone loss (Sibonga, 2013). Although it is unclear whether the changes in bone metabolism are self-limiting, evidence to date in animals and humans indicates that long-duration space missions will be detrimental to bone.

Animal studies, done primarily in rodents, are important for modeling the effects of spaceflight at all stages of skeletal maturation. Studies to date have investigated the effects of spaceflight on bone accrual and turnover balance at a variety of weight bearing, load

bearing, and unloaded skeletal sites in growing rodents (Bikle and Halloran, 1999; Turner, 2000). However, the majority of spaceflight experiments using rodents, primarily rats, were performed prior to wide spread availability of x-ray microtomography/microcomputed tomography (μ CT) (Keune et al., 2015). Consequently, limited information is available regarding the effects of spaceflight on bone microarchitecture.

The proximal tibia and distal femur are commonly evaluated following spaceflight. In contrast, the effects of spaceflight on proximal femur is much less commonly reported. Bone changes can occur at many skeletal sites but the greatest loss in astronauts is detected in the lower body and, in particular, the proximal femur (Keyak et al., 2009; Lang et al., 2006; Sibonga et al., 2007). In one of the few studies investigating multiple skeletal compartments, we analyzed gene expression in distal epiphysis, distal metaphysis, diaphysis, and proximal region of the femur following a 14-day spaceflight (STS-58) in growing male rats.

[☆] Funding: This work was supported by NASA (80NSSC19K0430 and NNX15AL15G).

* Corresponding author at: Skeletal Biology Laboratory, School of Biological and Population Health Sciences, 107B Milam Hall, Oregon State University, Corvallis, OR 97331, USA.

E-mail address: russell.turner@oregonstate.edu (R.T. Turner).

<https://doi.org/10.1016/j.bonr.2021.100755>

Received 16 August 2020; Received in revised form 1 February 2021; Accepted 10 February 2021

Available online 14 February 2021

2352-1872/© 2021 The Authors.

Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Notably, we found that the proximal region of the femur exhibited the largest reduction in mRNA levels for bone matrix proteins (Evans et al., 1998). Based on this initial observation, we evaluated bone architecture (using μ CT) in the proximal region of the femoral head of archived bone specimens from rapidly growing male Sprague Dawley rats flown aboard STS-41 (Turner et al., 2019), and observed strikingly lower cancellous bone volume fraction in the flight animals. The cancellous osteopenia was a surprise because of the short duration (4 days) of the spaceflight mission. To determine the generalizability of this finding across sex and strain, we assessed bone microarchitecture in the femoral head in archived bone specimens from two additional spaceflight experiments, STS-57 and STS-62. STS-57 was a 10-day mission performed using growing male Fisher 344 rats and STS-62 was a 14-day mission using growing ovariectomized (ovx) female Fisher 344 rats.

2. Methods

STS-57 was a Shuttle-SPACEHAB mission of Space Shuttle Endeavour. The mission launched June 21, 1993 from Kennedy Space Center, Florida. 7.5-week-old male Fisher 344 rats were flown on the flight. The experimental protocol was approved by the NASA Animal Care and Use Committee and flight details have been published (West-erlind and Turner, 1995). In brief, the rats were randomized into one of three groups: baseline control (baseline, $n = 6$), ground-based flight control (flight control, $n = 12$), or spaceflight (flight, $n = 12$). The baseline group was euthanized on day of launch. The flight animals were flown for 10 days. Flight control rats were housed in Animal Enclosure Modules (AEMs) (Moyer et al., 2016) identical to those housing the flight animals and temperature, humidity and light levels adjusted to values in flight AEMs. All animals were provided with NASA Nutrient-upgraded Rodent Foodbar (NuRFB) (Sun et al., 2010; Tou et al., 2003) and water ad libitum. Flight animals were euthanized 5–8 h after landing by decapitation and femora were removed, fixed overnight in 10% formalin, and stored in 70% ethanol for evaluation.

STS-62 was a Space Shuttle program mission flown aboard Space Shuttle Columbia that launched March 4, 1994. 12-week-old ovx female Fisher 344 rats were used in the study. The experimental protocol was approved by the NASA Animal Care and Use Committee and flight details have been published (Cavolina et al., 1997). In brief, the female rats were ovx 14 days prior to launch and randomized into one of three groups: baseline control (baseline, $n = 6$), ground-based flight control (flight control, $n = 12$), or spaceflight (flight, $n = 12$). The baseline group was euthanized on day of launch. The flight animals were housed in AEMs and flown for 14 days. Flight control rats were housed in AEMs and temperature, humidity and light levels were adjusted to values in flight AEMs. Flight and flight control rats were provided with food (AEM NuRFB) and water ad libitum. Flight animals were euthanized 4–6 h after landing. Asynchronous groups of ovary-intact vivarium-housed rats on a standard rat chow diet were euthanized to provide age-matched reference values for baseline and flight. The asynchronous component of the study was approved by the Animal Use and Care Committee at the Mayo Clinic (Rochester, MN) where the study was performed. The purpose of the ovary-intact controls was to evaluate normal age-related changes in bone over the 14-day flight period and to verify that ovx resulted in cancellous osteopenia. All rats were euthanized by decapitation and femora were removed, fixed overnight in 10% formalin, and stored in 70% ethanol for evaluation.

2.1. Micro-computed tomography

Bone volume and microarchitecture were assessed using a Scanco μ CT scanner (Scanco Medical AG, Basserdorf, Switzerland) at a voxel size of $12 \times 12 \times 12$ microns, 55 kVp X-ray voltage, 145 μ A intensity, and 200 ms integration time. Filtering parameters sigma and support were set to 0.8 and 1, respectively. Bone segmentation was conducted at a threshold value of 245 (gray scale, 0–1000) determined empirically. In

both studies, a sample of 20 slices (240 μ m) of cancellous bone was assessed in the proximal half of the femoral head as previously described (Turner et al., 2019). Automated irregular manual contouring was used to delineate cancellous from cortical bone. Direct cancellous bone measurements included cancellous bone volume fraction (bone volume/tissue volume, %), connectivity density (mm^{-3}), trabecular thickness (μ m), trabecular number (mm^{-1}), and trabecular separation (μ m). In addition, 62 slices (744 μ m) of cortical bone were assessed 690 slices (8.3 mm) distal to the top of femoral head and cortical thickness (μ m) measured.

2.2. Statistical analysis

Mean comparisons of quantitative variables (e.g., bone volume fraction) for the spaceflight, flight control, and baseline groups were made using separate one-way analysis of variance models for data from male and female rats. The mean structure of both models was specified as $\alpha \cdot \text{Spaceflight} + \beta \cdot \text{Flight Control} + \gamma \cdot \text{Baseline}$. The linear model conditions of normality and equal variance were assessed numerically using Anderson-Darling and Levene's tests, and graphically using box-plots, dotplots, normal quantile plots, and residual plots. Dunnett's procedure determined adjusted P-values for comparing mean outcomes of the spaceflight group to both the baseline and flight control groups. All data are reported as mean \pm SE with individual values presented. Differences are considered significant at $p \leq 0.05$. Data analysis was performed using R version 3.4.3.

3. Results

The effects of spaceflight on cancellous bone microarchitecture in the femoral head and cortical thickness in the proximal femur diaphysis in 7.5-week-old Fisher 344 male rats flown on STS-57 for 10 days are shown in Fig. 1. Flight rats had lower cancellous bone volume fraction compared to flight control rats (A). Flight rats also had lower connectivity density (B) and higher trabecular separation (E) compared to baseline rats. Significant differences among groups were not detected for any of the remaining cancellous endpoints evaluated. Significant differences in cortical thickness were likewise not detected among groups (F). Representative μ CT images of cancellous bone in the femur head from the 3 treatment groups are shown in Fig. 1G.

The effects of spaceflight on cancellous bone microarchitecture in the femoral head and cortical thickness in the proximal femur diaphysis in 12-week-old ovx female rats flown on STS-62 for 14 days are shown in Fig. 2. Age-matched (baseline and flight duration) asynchronous ovary-intact controls are shown as reference groups. Flight rats had lower cancellous bone volume fraction (A) and trabecular thickness (C), and higher trabecular separation (E) compared to flight control rats. Flight rats also had lower cancellous bone volume fraction (A) and connectivity density (B) and higher trabecular separation (E) compared to baseline rats. Significant differences in cortical thickness were not detected with treatment (F). Representative μ CT images of cancellous bone in the femur head from the treatment groups are shown in Fig. 2G.

The comparative effects of the 14-day spaceflight in 12-week-old female ovx rats flown on STS-62 (current and published studies) and in 16-week-old female C57BL6/J mice subjected to 15 days of spaceflight abroad STS-131 (Blaber et al., 2014) on bone microarchitecture at representative sites are shown in Table 1. Cancellous osteopenia was detected in femur (head, metaphysis and epiphysis) and lumbar vertebra (body), but not humerus (epiphysis) in rats and in the femur head in mice. Cortical thickness was unchanged in femur (rats and mice) and humerus (rats).

4. Discussion

Rodents—historically rats and more recently mice—are the primary animal model for investigating the effects and underlying mechanisms of

STS-57: Fisher 344 Male Rats, 10-day Flight




Fig. 1. Effects of a 10-day spaceflight aboard STS-57 in growing male Fisher 344 rats on cancellous bone volume fraction (A), connectivity density (B), trabecular thickness (C), trabecular number (D), and trabecular separation (E) in the femoral head, and on cortical thickness (F) in the proximal femur diaphysis. Representative three-dimensional images of cancellous bone from animals in each treatment group are shown in panel G. ^aDifferent from baseline, $P < 0.05$, ^bDifferent from flight control, $P < 0.05$.

action of spaceflight on the skeleton. Here we report data showing that spaceflight results in cancellous osteopenia in the proximal femoral head in rapidly growing male Fisher 344 rats and slowly growing ovx female Fisher 344 rats. The observed osteopenia in the femoral head following spaceflight supports the results of a 4-day study performed in very rapidly growing male Sprague-Dawley rats (STS-41). Spaceflight studies performed in C57BL/6 J mice (Blaber et al., 2014) suggest that this response may not be limited to rats.

The dramatically lower cancellous bone volume fraction noted in the femoral head of rats flown aboard STS-41 was a surprise (Turner et al., 2019) because histomorphometric evaluation and gene expression analyses failed to detect spaceflight-induced changes in other regions of the skeleton following this 4-day mission (Backup et al., 1994; Turner,

1995; Turner et al., 1995). Since the rats were very young and rapidly growing, it is likely that the lower cancellous bone volume fraction in the femoral head of the flight animals was due to reduced bone accrual. However, we cannot verify this because baseline controls were not available for analysis.

We analyzed femoral heads from rats flown aboard STS-57 and STS-62 to determine whether the results obtained for STS-41 are generalizable across sex and strain. Consistent with the original observation in male Sprague Dawley rats flown aboard STS-41, spaceflight resulted in osteopenia in the femoral head in male Fisher 344 rats flown aboard STS-57 and in ovx female Fisher 344 rats flown aboard STS-62. However, the magnitude of response appeared to be smaller in the latter two studies. Compared to ground controls, cancellous bone volume fraction

STS-62: Fisher 344 Ovariectomized Rats, 14-day Flight




Fig. 2. Effects of a 14-day spaceflight aboard STS-62 in growing ovariectomized (ovx) female Fisher 344 rats on bone volume fraction (A), connectivity density (B), trabecular thickness (C), trabecular number (D), and trabecular separation (E) in the femoral head, and on cortical thickness (F) in the proximal femur diaphysis. Representative three-dimensional images of cancellous bone from animals in each treatment group are shown in panel G. Asynchronous groups of age-matched ovary-intact rats (Intact) are shown as reference values for baseline and flight animals. ^aDifferent from baseline, $P < 0.05$, ^bDifferent from flight control, $P < 0.05$.

in flight rats was -46% in STS-41, -11% in STS-57 and -16% in STS-62. To address the possibility of error in the initial analysis, a different observer blinded to treatment re-measured the μ CT scans of the femoral head in rats flown aboard STS-41. The comparable results of the 2nd analysis excludes measurement error but does not provide additional insight. Based on absence of reduction in longitudinal bone growth, absence of reduction in mRNA levels for bone matrix proteins, and absence of histomorphometric evidence for decreased bone formation at multiple skeletal sites assessed, the most likely explanation for the dramatic osteopenia in femoral head in rats from STS-41 is increased bone resorption (Backup et al., 1994; Turner, 1995; Turner et al., 1995). It remains unclear, however, whether the more modest osteopenia in rats flown aboard STS-57 and STS-62 compared to their respective flight controls reflects differences due to strain, age, duration of flight, or other factors. However, it is notable that BV/TV in ovx female rats flown aboard STS-62 was lower than baseline controls, indicating net bone loss contributed to the osteopenia.

There is evidence for location-specific (e.g., appendicular versus axial) and compartment-specific (e.g., cancellous versus cortical) effects of microgravity on bone microarchitecture (Keune et al., 2015). Mechanistic inference is challenging because of the limited number of studies and variation in experimental design and endpoints used to evaluate the bone response to spaceflight (Turner, 2000). Variables in design include age at launch, strain, duration of flight, housing conditions during flight, and interval of time between reloading and necropsy (Morey-Holton et al., 2000; Vico et al., 1993; Wronski et al., 1987; Zerath et al., 2000). In general, investigators have had limited access to tissues (often a single bone). As for endpoints, histomorphometric analysis was the principle method available for evaluation of bone architecture during the interval when most spaceflight studies using rats as models were performed (1970s–1990s) (Keune et al., 2015). In this regard, we have been very fortunate in being able to analyze archived femur, humerus, lumbar vertebra and calvarium from ovx female Fisher 344 rats flown on STS-62 (Keune et al., 2015). Additionally, we measured dynamic and static

Table 1

Effects of spaceflight on cancellous bone volume fraction (femur, humerus, lumbar vertebra) and cortical thickness (femur, humerus) in 12-week-old ovariectomized female Fisher 344 rats flown aboard STS-62 for 14 days and on cancellous bone volume fraction (femur) and cortical thickness (femur) in 16-week-old female C57BL/6J mice flown aboard STS-131 for 15 days.

	Flight control	Flight	% difference	Reference
Ovariectomized female Fisher 344 rats				
Cancellous bone volume fraction (%)				
Femur				
Head	43.9 ± 0.7	36.9 ± 0.9	16.0*	Current study
Metaphysis	6.8 ± 0.3	4.9 ± 0.3	27.9*	Keune et al. 2016
Epiphysis	28.2 ± 0.2	24.1 ± 0.4	14.5*	Keune et al. 2016
Humerus				
Epiphysis	32.6 ± 0.5	32.6 ± 0.3	0.0	Keune et al. 2016
Lumbar vertebra				
Body	20.9 ± 0.4	16.9 ± 0.8	19.1*	Keune et al. 2016
Cortical thickness (μm)				
Femur				
Proximal diaphysis	489 ± 7	482 ± 6	1.4	Current study
Mid-diaphysis	492 ± 4	478 ± 4	2.8	Keune et al. 2016
Humerus				
Mid-diaphysis	486 ± 4	477 ± 4	1.9	Keune et al. 2016
Female C57BL/6J mice				
Cancellous bone volume fraction (%)				
Femur				
Head	47.0 ± 0.7	38.8 ± 1.3	17.5*	Blaber et al. 2014
Cortical thickness (μm)				
Femur				
Neck	140 ± 1	138 ± 1	1.0	Blaber et al. 2014

* Different from flight control, $P < 0.05$.

histomorphometry in tibia and lumbar vertebra in rats from this flight (Cavolina et al., 1997; Keune et al., 2015; Keune et al., 2016; Westerlind et al., 1997). This wealth of material allows a more complete description of site-specific effects of spaceflight on the skeleton. Compared to ground-based flight controls, flight animals had lower cancellous bone volume fraction in femur at all sites evaluated (head, distal metaphysis and distal epiphysis) but showed no change in cortical thickness (proximal femur and mid-diaphysis). Additionally, the effects of spaceflight were bone-specific, with cancellous bone loss being detected in femur and lumbar vertebra but not in humerus. Although fewer skeletal sites were reported, differential effects of a 15-day spaceflight on cancellous bone in the femur head (lower BV/TV in flight animals) and cortical bone thickness in the femur neck (no change) were also noted in C57BL/6 J mice (Blaber et al., 2014).

While suitable for assessing bone microarchitecture, μ CT provides limited insight into the underlying cellular and molecular mechanisms responsible for the site-specific alterations in bone occurring during spaceflight. Histomorphometric analysis of bones of rats flown on STS-62 suggests differential responses at cortical and cancellous bone sites. Specifically, spaceflight resulted in decreased periosteal bone formation in weight-bearing bone (Cavolina et al., 1997), likely contributing to decreased bone mineral content (Keune et al., 2015). Reduced bone formation at the periosteum was associated with lower mRNA levels for bone matrix proteins (Cavolina et al., 1997). Decreased bone formation was also commonly detected in longer duration flights (>7 days) in male rats (Backup et al., 1994; Durnova et al., 1996; Morey and Baylink, 1978; Turner, 2000; Vico et al., 1993; Wronski and Morey, 1983). In contrast, the effect of spaceflight on bone formation at cancellous sites is

less clear (Jee et al., 1983; Wronski et al., 1987, 1998; Zerath et al., 2000). The most detailed analyses to date were performed on rats flown aboard STS-62. In these ovx female animals, cancellous bone loss in tibia (Cavolina et al., 1997) and lumbar vertebra (Keune et al., 2015) in excess of that attributable to ovx resulted from increased bone resorption, as ascertained by fluorochrome label disappearance, with no change in bone formation, as ascertained by static and dynamic bone histomorphometry. Finite element modeling of distal femur in these rats, as well as a theoretical model based on fuzzy decision, suggest that mechanical strain modulates the balance between bone formation and bone resorption. Specifically, the overall rate of bone turnover was influenced by endocrine status but bone loss occurred preferentially at sites experiencing low mechanical strain energies (Luo et al., 2000; Westerlind et al., 1997). At the molecular level, expression levels of two cytokines known to increase bone resorption, interleukin-1 β and interferon gamma, were elevated in proximal tibial metaphysis (Zhang and Turner, 1998).

In recent years, the mouse has become the exclusive rodent model flown on spaceflight missions. However, there remains a paucity of publications detailing the skeletal response of mice to microgravity (Blaber et al., 2014; Gerbaix et al., 2017; Smith et al., 2020; Tavella et al., 2012). Smith et al. reported osteopenia in the distal femur of growing female BALB/c mice following spaceflight (Smith et al., 2020). As mentioned, Blaber et al. reported changes in BV/TV in femoral head of 16-week-old female C57BL/6 J mice following a 15-day spaceflight (Blaber et al., 2014) that were similar in magnitude to changes observed in ovx female Fisher 344 rats following a 14 day spaceflight. In the ovx female rats, cancellous osteopenia in the flight animals was primarily due to lower trabecular thickness, whereas in ovary-intact mice both trabecular number and thickness were lower in flight animals compared to ground controls.

The proximal femur is clinically relevant but not commonly evaluated in animal models. Low trauma fractures are common in the femoral neck of individuals with osteoporosis (Tsuda, 2017). Fractures occur at several locations in the proximal femur, including intertrochanteric region, transcervical neck, subcapital neck, and subtrochanteric region. Fractures occurring through the femoral head, the site investigated in this study, are more typically associated with trauma and are particularly difficult to manage (Scolaro et al., 2017). That being said, traumatic injury would be the likely cause of fracture during a spaceflight mission and a microgravity-induced reduction in bone volume fraction at this site would likely reduce the magnitude of trauma required for fracture. This conclusion is supported by studies demonstrating that spaceflight and simulated spaceflight result in lower mechanical competence (Patterson-Buckendahl et al., 1987; Shirazi-Fard et al., 2013; Spengler et al., 1983; Zernicke et al., 1990).

In summary, spaceflight ranging from 4 to 14 days in duration in two strains of growing male rats (Sprague Dawley and Fisher 344) and in growing ovx female Fisher 344 rats resulted in osteopenia in the femoral head. It is noteworthy that significant bone loss during long duration spaceflight in astronauts occurs in the proximal femur. Thus, future studies performed in skeletally-mature animals should carefully evaluate this clinically relevant site.

Declaration of interest

The authors have no conflict of interest.

CRediT authorship contribution statement

Conceptualization: RT
Data Collection: AG, DO, LS
Data analysis: AB
Drafting manuscript: AG
Revising manuscript content: RT, AB, DO, LS, UI, and AG
Approving final version: RT, AB, DO, LS, UI, and AG

RT takes responsibility for the integrity of the data.

References

- Backup, P., Westerlind, K., Harris, S., Spelsberg, T., Kline, B., Turner, R., 1994. Spaceflight results in reduced mRNA levels for tissue-specific proteins in the musculoskeletal system. *Am. J. Phys.* 266, E567–E573.
- Bikle, D.D., Halloran, B.P., 1999. The response of bone to unloading. *J. Bone Miner. Metab.* 17, 233–244.
- Blaber, E.A., Dvorochkin, N., Torres, M.L., Yousuf, R., Burns, B.P., Globus, R.K., Almeida, E.A., 2014. Mechanical unloading of bone in microgravity reduces mesenchymal and hematopoietic stem cell-mediated tissue regeneration. *Stem Cell Res.* 13, 181–201.
- Cavolina, J.M., Evans, G.L., Harris, S.A., Zhang, M., Westerlind, K.C., Turner, R.T., 1997. The effects of orbital spaceflight on bone histomorphometry and messenger ribonucleic acid levels for bone matrix proteins and skeletal signaling peptides in ovariectomized growing rats. *Endocrinology* 138, 1567–1576.
- Durnova, G.N., Kaplanskii, A.S., Morey-Holton, E.R., Vorobeva, V.N., 1996. Investigation of tibial bones of the rats exposed on board “Spacelab-2”: histomorphometric analysis. *Aviakosm. Ekolog. Med.* 30, 21–26.
- Evans, G.L., Morey-Holton, E., Turner, R.T., 1998. Spaceflight has compartment- and gene-specific effects on mRNA levels for bone matrix proteins in rat femur. *J. Appl. Physiol.* 84, 2132–2137.
- Gerbaix, M., Gnyubkin, V., Farlay, D., Olivier, C., Ammann, P., Courbon, G., Laroche, N., Gentil, R., Follet, H., Peyrin, F., et al., 2017. One-month spaceflight compromises the bone microstructure, tissue-level mechanical properties, osteocyte survival and lacunae volume in mature mice skeletons. *Sci. Rep.* 7, 2659.
- Jee, W.S., Wronski, T.J., Morey, E.R., Kimmel, D.B., 1983. Effects of spaceflight on trabecular bone in rats. *Am. J. Phys.* 244, R310–R314.
- Keune, J.A., Branscum, A.J., Iwaniec, U.T., Turner, R.T., 2015. Effects of spaceflight on bone microarchitecture in the axial and appendicular skeleton in growing ovariectomized rats. *Sci. Rep.* 5, 18671.
- Keune, J.A., Philbrick, K.A., Branscum, A.J., Iwaniec, U.T., Turner, R.T., 2016. Spaceflight-induced vertebral bone loss in ovariectomized rats is associated with increased bone marrow adiposity and no change in bone formation. *NPJ Microgravity* 2, 16016.
- Keyak, J.H., Koyama, A.K., LeBlanc, A., Lu, Y., Lang, T.F., 2009. Reduction in proximal femoral strength due to long-duration spaceflight. *Bone* 44, 449–453.
- Lang, T.F., Leblanc, A.D., Evans, H.J., Lu, Y., 2006. Adaptation of the proximal femur to skeletal reloading after long-duration spaceflight. *J. Bone Miner. Res.* 21, 1224–1230.
- Luo, Z.P., Zhang, L., Turner, R.T., An, K.N., 2000. Effects of mechanical stress/strain and estrogen on cancellous bone structure predicted by fuzzy decision. *IEEE Trans. Biomed. Eng.* 47, 344–351.
- Morey, E.R., Baylink, D.J., 1978. Inhibition of bone formation during space flight. *Science* 201, 1138–1141.
- Morey-Holton, E.R., Halloran, B.P., Garetto, L.P., Doty, S.B., 2000. Animal housing influences the response of bone to spaceflight in juvenile rats. *J. Appl. Physiol.* 88, 1303–1309.
- Moyer, E.L., Dumars, P.M., Sun, G.S., Martin, K.J., Heathcote, D.G., Boyle, R.D., Skidmore, M.G., 2016. Evaluation of rodent spaceflight in the NASA animal enclosure module for an extended operational period (up to 35 days). *NPJ Microgravity* 2, 16002.
- Patterson-Buckendahl, P., Arnaud, S.B., Mechanic, G.L., Martin, R.B., Grindeland, R.E., Cann, C.E., 1987. Fragility and composition of growing rat bone after one week in spaceflight. *Am. J. Phys.* 252, R240–R246.
- Scolaro, J.A., Marecek, G., Firoozabadi, R., Krieg, J.C., Routt, M.L.C., 2017. Management and radiographic outcomes of femoral head fractures. *J. Orthop. Traumatol.* 18, 235–241.
- Shirazi-Fard, Y., Anthony, R.A., Kwaczala, A.T., Judex, S., Bloomfield, S.A., Hogan, H.A., 2013. Previous exposure to simulated microgravity does not exacerbate bone loss during subsequent exposure in the proximal tibia of adult rats. *Bone* 56, 461–473.
- Sibonga, J.D., 2013. Spaceflight-induced bone loss: is there an osteoporosis risk? *Curr Osteoporos Rep* 11, 92–98.
- Sibonga, J.D., Evans, H.J., Sung, H.G., Spector, E.R., Lang, T.F., Oganov, V.S., Bakulin, A. V., Shackelford, L.C., LeBlanc, A.D., 2007. Recovery of spaceflight-induced bone loss: bone mineral density after long-duration missions as fitted with an exponential function. *Bone* 41, 973–978.
- Smith, R.C., Cramer, M.S., Mitchell, P.J., Lucchesi, J., Ortega, A.M., Livingston, E.W., Ballard, D., Zhang, L., Hanson, J., Barton, K., et al., 2020. Inhibition of myostatin prevents microgravity-induced loss of skeletal muscle mass and strength. *PLoS One* 15, e0230818.
- Spengler, D.M., Morey, E.R., Carter, D.R., Turner, R.T., Baylink, D.J., 1983. Effects of spaceflight on structural and material strength of growing bone. *Proc. Soc. Exp. Biol. Med.* 174, 224–228.
- Sun, G.S., Tou, J.C., Liittschwager, K., Herrera, A.M., Hill, E.L., Girten, B., Reiss-Bubenheim, D., Vasques, M., 2010. Evaluation of the nutrient-upgraded rodent food bar for rodent spaceflight experiments. *Nutrition* 26, 1163–1169.
- Tavella, S., Ruggiu, A., Giuliani, A., Brun, F., Canciani, B., Manescu, A., Marozzi, K., Cilli, M., Costa, D., Liu, Y., et al., 2012. Bone turnover in wild type and pleiotrophin-transgenic mice housed for three months in the international Space Station (ISS). *PLoS One* 7, e33179.
- Tou, J., Grindeland, R., Barrett, J., Dalton, B., Mandel, A., Wade, C., 2003. Evaluation of NASA Foodbars as a standard diet for use in short-term rodent space flight studies. *Nutrition* 19, 947–954.
- Tsuda, T., 2017. Epidemiology of fragility fractures and fall prevention in the elderly: a systematic review of the literature. *Curr Orthop Pract* 28, 580–585.
- Turner, R.T., 1995. Effects of short-term spaceflight and recombinant human growth hormone (rhGH) on bone growth in young rats. *Aviat. Space Environ. Med.* 66, 763–769.
- Turner, R.T., Evans, G.L., Wakley, G.K., 1995. Spaceflight results in depressed cancellous bone formation in rat humeri. *Aviat. Space Environ. Med.* 66, 770–774.
- Turner, R.T., 2000. Invited review: what do we know about the effects of spaceflight on bone? *J. Appl. Physiol.* 89, 840–847.
- Turner, R.T., Deyle Jr., R.T., Branscum, A.J., Iwaniec, U.T., 2019. Effects of a four-day spaceflight and recombinant human growth hormone on cancellous bone microarchitecture in femoral head of rapidly growing male mice. *Matters Select* 5.
- Vico, L., Bourrin, S., Genty, C., Palle, S., Alexandre, C., 1993. Histomorphometric analyses of cancellous bone from COSMOS 2044 rats. *J. Appl. Physiol.* 75, 2203–2208.
- Westerlind, K.C., Turner, R.T., 1995. The skeletal effects of spaceflight in growing rats: tissue-specific alterations in mRNA levels for TGF-beta. *J. Bone Miner. Res.* 10, 843–848.
- Westerlind, K.C., Wronski, T.J., Ritman, E.L., Luo, Z.P., An, K.N., Bell, N.H., Turner, R.T., 1997. Estrogen regulates the rate of bone turnover but bone balance in ovariectomized rats is modulated by prevailing mechanical strain. *Proc. Natl. Acad. Sci. U. S. A.* 94, 4199–4204.
- Wronski, T.J., Morey, E.R., 1983. Effect of spaceflight on periosteal bone formation in rats. *Am. J. Phys.* 244, R305–R309.
- Wronski, T.J., Morey-Holton, E.R., Doty, S.B., Maese, A.C., Walsh, C.C., 1987. Histomorphometric analysis of rat skeleton following spaceflight. *Am. J. Phys.* 252, R252–R255.
- Wronski, T.J., Li, M., Shen, Y., Miller, S.C., Bowman, B.M., Kostenuik, P., Halloran, B.P., 1998. Lack of effect of spaceflight on bone mass and bone formation in group-housed rats. *J. Appl. Physiol.* 85, 279–285.
- Zerath, E., Holy, X., Roberts, S.G., Andre, C., Renault, S., Hott, M., Marie, P.J., 2000. Spaceflight inhibits bone formation independent of corticosteroid status in growing rats. *J. Bone Miner. Res.* 15, 1310–1320.
- Zernicke, R.F., Vailas, A.C., Grindeland, R.E., Kaplansky, A., Salem, G.J., Martinez, D.A., 1990. Spaceflight effects on biomechanical and biochemical properties of rat vertebrae. *Am. J. Phys.* 258, R1327–R1332.
- Zhang, M., Turner, R.T., 1998. The effects of spaceflight on mRNA levels for cytokines in proximal tibia of ovariectomized rats. *Aviat. Space Environ. Med.* 69, 626–629.