

Adjustment for time-invariant and time-varying confounders in ‘unexplained residuals’ models for longitudinal data within a causal framework and associated challenges

KF Arnold,^{1,2}  GTH Ellison,^{1,2} SC Gadd,² J Textor,³  PWG Tennant,^{1,4}
A Heppenstall^{1,5} and MS Gilthorpe^{1,2} 

Statistical Methods in Medical Research
2019, Vol. 28(5) 1347–1364

© The Author(s) 2018



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0962280218756158

journals.sagepub.com/home/smm



Abstract

‘Unexplained residuals’ models have been used within lifecourse epidemiology to model an exposure measured longitudinally at several time points in relation to a distal outcome. It has been claimed that these models have several advantages, including: the ability to estimate multiple total causal effects in a single model, and additional insight into the effect on the outcome of greater-than-expected increases in the exposure compared to traditional regression methods. We evaluate these properties and prove mathematically how adjustment for confounding variables must be made within this modelling framework. Importantly, we explicitly place unexplained residual models in a causal framework using directed acyclic graphs. This allows for theoretical justification of appropriate confounder adjustment and provides a framework for extending our results to more complex scenarios than those examined in this paper. We also discuss several interpretational issues relating to unexplained residual models within a causal framework. We argue that unexplained residual models offer no additional insights compared to traditional regression methods, and, in fact, are more challenging to implement; moreover, they artificially reduce estimated standard errors. Consequently, we conclude that unexplained residual models, if used, must be implemented with great care.

Keywords

Unexplained residuals model, conditional regression model, conditional analysis, conditional growth, conditional weight, conditional size, directed acyclic graph, causal inference, lifecourse epidemiology

1 Background

Within the field of lifecourse epidemiology, there is substantial interest in modelling the relationship between an exposure x measured longitudinally at several time points (i.e. x_1, x_2, \dots, x_k) and a subsequent outcome y measured once later in life (hereafter referred to as a distal outcome); such a relationship can be helpfully summarised in Figure 1(a) in the form of a directed acyclic graph (DAG).¹ DAGs are pictorial representations of hypothesised causal relationships between variables in which: variables (nodes) are connected via unidirectional arrows (directed edges), which represent direct causal relationships; and no directed loops (i.e. circular paths) between variables are permitted. Nodes may be either: endogenous, having at least one causally preceding variable represented in the graph; or exogenous, having none.² All unexplained causes of the endogenous nodes x_2, \dots, x_k, y in Figure 1(a) are represented by the variables $e_{x_2}, \dots, e_{x_k}, e_y$, respectively. While there are many

¹Leeds Institute for Data Analytics, University of Leeds, Leeds, UK

²School of Medicine, University of Leeds, Leeds, UK

³Tumor Immunology Lab, Radboud University Medical Centre, Nijmegen, The Netherlands

⁴School of Healthcare, University of Leeds, Leeds, UK

⁵School of Geography, University of Leeds, Leeds, UK

Corresponding author:

KF Arnold, Leeds Institute for Data Analytics, University of Leeds, Worsley Building Level 11, Leeds LS2 9NL, UK.

Email: K.F.Arnold@leeds.ac.uk




Figure 1. (a) Nonparametric causal diagram (DAG) representing the hypothesised data-generating process for k longitudinal measurements of exposure x (i.e. x_1, x_2, \dots, x_k) and one distal outcome y . The terms e_{x_2}, \dots, e_{x_k} and e_y represent all unexplained causes of x_2, \dots, x_k and y , respectively, and are included to explicitly reflect uncertainty in all endogenous nodes (whether modelled or not). (b) Path diagrams depicting the k standard regression models that would be constructed to estimate the total causal effect of each of x_1, x_2, \dots, x_k on y (i.e. equation (5)). For each model, only the final coefficient may be interpreted as a total causal effect; all other coefficients are greyed to illustrate that no such interpretation should be made for them. (c) Path diagrams depicting the UR model, consisting of $k - 1$ preparation regressions (i.e. equation (6)) and a final composite regression model (i.e. equation (7), with $i = k$).

useful applications for DAGs in epidemiologic research, perhaps the most beneficial is their ability to identify suitable sets of covariates for removing bias due to confounding between an exposure and outcome,^{3,4} which occurs whenever both variables share one or more common causes. For this reason, DAGs are increasingly being used in epidemiology, as they provide a framework for estimating the total causal effect of an exposure on an outcome.⁴

Using a causal framework to (correctly) model the scenario in Figure 1(a) may also have additional utility in identifying and quantifying important periods of change in the exposure that are causally related to the outcome. However, one challenge to such applications is that successive measurements of an exposure over time may be highly correlated with one another and therefore likely to suffer collinearity when analysed in relation to a distal outcome. Consequently, there has been extensive debate regarding the best way to model these types of longitudinally measured variables; a recent review⁵ of analytical and modelling techniques has identified a range of different approaches, including z-score plots, regression with change scores, multilevel and latent growth curve models, and growth mixture models. Nonetheless, one of the most straightforward methods in use is a series of standard multivariable regression models.

1.1 Standard regression method

When using this approach, each longitudinal measurement of the exposure variable is treated as a separate entity that is subject to confounding by all previous measurements of that variable – the total number of models needed therefore being equal to the total number of time points at which the exposure has been measured.

As an example, the simplest scenario would involve just two measurements of the exposure x (i.e. x_1 and x_2 , measured at time points 1 and 2, respectively), and a distal outcome, y , where all variables are continuous in nature. Here, two standard regression models (denoted $\hat{y}_S^{(i)}$, for $i = 1, 2$) would need to be constructed to estimate the total causal effect of each distinct measurement of x on y , i.e.

$$\hat{y}_S^{(1)} = \hat{\alpha}_0^{(1)} + \hat{\alpha}_{x_1}^{(1)}x_1 \tag{1}$$

$$\hat{y}_S^{(2)} = \hat{\alpha}_0^{(2)} + \hat{\alpha}_{x_1}^{(2)}x_1 + \hat{\alpha}_{x_2}^{(2)}x_2. \quad (2)$$

Importantly, to estimate the total causal effect of x_1 on y in equation (1), adjustment for x_2 is inappropriate, as it lies on the causal path between x_1 and y (i.e. x_2 is a mediator); in fact, adjustment for x_2 might invoke bias in the causal interpretation due to a phenomenon known as the ‘reversal paradox’.^{5–7} In contrast, to estimate the total causal effect of x_2 on y in equation (2), adjustment for x_1 is appropriate, since it confounds the desired relationship (i.e. x_1 causally precedes both x_2 and y , potentially creating a spurious relationship between them). For this reason, in either model, it is *only* possible to interpret the coefficient of the last/most recent measurement of x (the exposure) as a total causal effect,¹ which encompasses all direct and indirect causal pathways between an exposure and outcome. No such interpretation is possible (nor should it be attempted) for the coefficient of the earlier measurement of x in equation (2), as it operates purely as a confounder.

1.2 Unexplained residuals method

To circumvent the need for multiple models, Keijzer-Veen⁸ has suggested an alternative approach that would combine the information contained within each of the two separate models (equations (1) and (2)) into a single composite regression model using ‘unexplained residuals’. As originally proposed,⁹ such a model allows the researcher to quantify the total effects of both the initial measurement of x (i.e. x_1) and subsequent change in x on the outcome y . The proposed approach contains two steps but is straightforward in principle.

First, the most recent measurement of x (i.e. x_2) is regressed on the earlier measurement of x (i.e. x_1):

$$x_2 = \hat{\gamma}_0^{(2)} + \hat{\gamma}_{x_1}^{(2)}x_1 + e_{x_2}. \quad (3)$$

This produces a measure of each observation’s ‘expected’ value of x_2 as predicted by its value of x_1 . The difference between the expected value of x_2 (i.e. $\hat{\gamma}_0^{(2)} + \hat{\gamma}_{x_1}^{(2)}x_1$) and the observed value of x_2 amounts to the residual term e_{x_2} . Put another way, e_{x_2} represents the part of x_2 ‘unexplained’ by x_1 .

Second, y is regressed on both the initial exposure x_1 and subsequent residual term e_{x_2} :

$$\hat{y}_{UR}^{(2)} = \hat{\lambda}_0^{(2)} + \hat{\lambda}_{x_1}^{(2)}x_1 + \hat{\lambda}_{e_{x_2}}^{(2)}e_{x_2}. \quad (4)$$

According to Keijzer-Veen et al.,⁹ the key advantages of conducting regression using the composite ‘unexplained residuals’ (UR) model (4) are that:

- (1) The UR model produces the same estimated outcome values as the standard regression model in equation (2) (i.e. $\hat{y}_S^{(2)} = \hat{y}_{UR}^{(2)}$);
- (2) The estimated total effect sizes (coefficient values) produced by individual standard regression models (equations (1) and (2)) are equal to those estimated within the UR model (i.e. $\hat{\alpha}_{x_1}^{(1)} = \hat{\lambda}_{x_1}^{(2)}$ and $\hat{\alpha}_{x_2}^{(2)} = \hat{\lambda}_{e_{x_2}}^{(2)}$); thus, multiple coefficients in a single model may be interpreted;
- (3) The UR model provides additional insight (via the coefficient $\hat{\lambda}_{e_{x_2}}^{(2)}$ in equation (4)) into the effect of x increasing *more than expected* upon y ; and
- (4) The initial exposure x_1 and subsequent residual term e_{x_2} are mathematically independent (i.e. orthogonal).

Succinctly, the two models $\hat{y}_S^{(2)}$ and $\hat{y}_{UR}^{(2)}$ are algebraically equivalent, but $\hat{y}_{UR}^{(2)}$ makes interpretation of the separate influence of the initial measurement of the exposure x (i.e. x_1) and subsequent changes in x more straightforward than do (multiple) standard regression models $\hat{y}_S^{(1)}$ and $\hat{y}_S^{(2)}$.

Within the epidemiological literature, UR models have been used under a number of different names. In addition to ‘regression with unexplained residuals’ (as first proposed by Keijzer-Veen et al.^{9–11}), other studies have referred to: ‘unexplained residual regression’¹²; ‘method of unexplained residuals’¹³; ‘conditional linear regression’¹²; ‘conditional (regression) models’^{5,14}; ‘regression with conditional growth measures’¹⁴; ‘conditional growth models’^{15–18}; ‘conditional weight models’¹⁹; and ‘conditional (regression) analysis’.^{20–24} The terms ‘conditional growth’ and ‘conditional size’ – and additional variations thereof – are also commonly used to refer to the difference between observed and expected size measurements.^{5,15,18,25–39} To avoid further confusion, the residual term representing the difference between the observed and expected values of an exposure produced in the manner proposed by Keijzer-Veen et al. (as in equation (3)) will be henceforth referred to as the ‘*unexplained*

residuals (UR) term, and the models themselves (as in equation (4)) will be referred to as ‘*unexplained residuals (UR) models*’.

Despite the numerous names given to these models, the process remains essentially the same as that first proposed. Indeed, several authors have extended the original model to examine scenarios involving several measurements of an exposure x (i.e. x_1, x_2, \dots, x_k); UR models in these extended applications thus include several UR terms.^{5,12,13,16–41} In general, each UR term e_{xi} is derived from the regression of each measured value x_i on all previous measurements x_1, x_2, \dots, x_{i-1} , for $2 \leq i \leq k$,^{12,16,18–22,24,25,27,29,31–34,36,39,40} though some researchers have deviated from this procedure^{13,26,35,37,41}; the outcome y is then regressed on x_1 and all subsequent UR terms $e_{x2}, e_{x3}, \dots, e_{xk}$.

Many researchers have further extended the original UR models by adjusting for additional confounding variables (i.e. over and above the confounding of prior measurements of the exposure), though there is, as yet, little consensus as to whether or how such adjustments should be performed. For example, Horta et al.¹⁶ made no adjustments for potential confounders when deriving their UR terms, but did make adjustments within their composite UR model. In contrast, Gandhi et al.¹⁸ adjusted for just one potential confounder (gender) when creating their UR terms, but also made further adjustments to the composite UR model (for gender and other variables). Adair et al.²⁵ created their UR terms using site- and sex-stratified linear regressions that were also adjusted for age, and made further adjustments for age, sex, and study site in their subsequent composite UR models. Indeed, there are many other examples of different approaches to confounder adjustment, but none of these have been adequately and explicitly justified by the researchers concerned, even though it appears that they did so in order to make causal inferences.

2 Research aims

The potential impact of using alternative approaches to adjust for confounding when constructing and using UR terms has yet to be fully evaluated. Indeed, Keijzer-Veen et al.⁹ did not address confounding variables in their original paper, and there has been little to no discussion or analysis of this issue by subsequent authors using this approach. It therefore remains unclear whether UR models that include confounders offer the same purported benefits as those lacking (or ignoring) confounders, and there is no clear indication of how potential confounders should be treated by analyses using these models. This is an issue of particular relevance to researchers seeking to infer causality from individual coefficient estimates, since inappropriate adjustment for covariates (which includes both the failure to adjust for genuine confounders and the adjustment for mediators mistaken for confounders) can lead to biased causal inferences. For this reason, UR models are likely to have limited practical utility unless they are able to accommodate confounding variables appropriately. The fact that UR models have not been developed or analysed within a causal framework also creates uncertainty about their utility for making causal inferences.

Therefore, the aims of the present study were to: (1) confirm that the approach proposed by Keijzer-Veen et al. may be extended to a scenario involving k longitudinal measurements of an exposure x in the absence of any additional confounding; (2) determine whether it is possible (and if so, how might it be possible) to adjust for additional confounders within the UR modelling framework; (3) evaluate the benefits of UR models claimed by Keijzer-Veen et al.; and (4) offer recommendations for future use of UR models. The present study examines two very different types of potential confounders: time-invariant (which require/provide measurements taken at a single time point and remain constant across the lifecourse, e.g. sex); and time-varying (for which measurements are collected at multiple time points across the lifecourse – usually concurrent to measurements of the exposure – because the value of the variable may change, e.g. socioeconomic position).

These aims are summarised in the DAGs presented in Figures 1(a), 2(a), and 3(a), which depict three general scenarios drawn from lifecourse epidemiology, each of which will be examined in the analyses that follow. Each DAG relates k longitudinally measured exposure variables x_1, x_2, \dots, x_k (i.e. x measured at time points $1, 2, \dots, k$) to a distal outcome y (measured at some point either concurrent to or following k) under three very different circumstances: (1a) in the absence of any additional confounders; (2a) in the presence of an additional time-invariant confounder m ; and (3a) in the presence of an additional time-varying confounder m_1, m_2, \dots, m_k . All DAGs are drawn forwardly saturated (i.e. where each node may causally affect all future nodes), and all unexplained causes of endogenous nodes are represented by the variable e and depicted as independent (i.e. we assume no unobserved confounding). The explicit inclusion of these three DAGs in Figures 1(a), 2(a), and 3(a) is intended not only to visually illustrate each of the scenarios that will be examined, but also, importantly, to situate the analyses that follow within a causal framework.




Figure 2. (a) Nonparametric causal diagram (DAG) representing the hypothesised data-generating process for k longitudinal measurements of exposure x (i.e. x_1, x_2, \dots, x_k), one distal outcome y , and one time-invariant confounder m . The terms $e_m, e_{x_1}, \dots, e_{x_k}$ and e_y represent all unexplained causes of m, x_1, \dots, x_k , and y , respectively, and are included to explicitly reflect uncertainty in all endogenous nodes (whether modelled or not). (b) Path diagrams depicting the k standard regression models that would be constructed to estimate the total causal effect of each of x_1, x_2, \dots, x_k on y (i.e. equation (9)). For each model, only the final coefficient may be interpreted as a total causal effect; all other coefficients are greyed to illustrate that no such interpretation should be made for them. (c) Path diagrams depicting the UR model, consisting of $k - 1$ preparation regressions (i.e. equation (10)) and a final composite regression model (i.e. equation (11), with $i = k$).

Sections 3 through 9, which follow, provide: the three key properties of UR models that will be evaluated for the scenarios in Figures 1(a), 2(a), and 3(a) (§3); DAG-based and mathematical examinations of the UR models for the scenarios given in Figure 1(a) (§4), 2(a) (§5), and 3(a) (§6); a discussion of several interpretational issues that arise for UR models when placed within a causal framework, including an evaluation of the claim that UR models provide greater insight than standard regression methods (§7); an argument outlining how UR models produce artificially reduced standard errors (SEs) and how this might be corrected (§8); and recommendations for future use and interpretation of UR models, particularly as these relate to the inclusion of confounders (§9).

3 Key properties of UR models

In the following sections, we evaluate the mathematical properties of the original UR models after extending them to include k measurements of a continuous exposure x : in the absence of any additional confounding (§4); in the presence of a single additional time-invariant confounder m (§5); and in the presence of a single additional time-varying confounder with sequential values m_1, m_2, \dots, m_k (§6). These properties are:

- **Property (i):** The outcome values predicted by the final standard regression model (for the final measurement of the exposure variable, x_k) are equal to those predicted by the composite UR model.
- **Property (ii):** The estimated coefficient for x_1 in the initial standard regression model (for the first measurement of the exposure variable, x_1) is equal to the estimated coefficient for x_1 in the composite UR model.
- **Property (iii):** The estimated coefficient for each x_i in its individual standard regression model (i.e. for designated exposure x_i) is equal to the estimated coefficient for the corresponding UR term e_{x_i} in the composite UR model.

From a causal inference perspective, only Properties (ii) and (iii) are meaningful, since the focus is on individual coefficient estimates as opposed to predicted outcomes. Nevertheless, we evaluate all three properties in Sections 4 through 6, and leave discussion of interpretational issues until later in the paper (§8).




Figure 3. (a) Nonparametric causal diagram (DAG) representing the hypothesised data-generating process for k longitudinal measurements of exposure x (i.e. x_1, x_2, \dots, x_k), one distal outcome y , and k longitudinal measurements of one time-varying confounder m_1, m_2, \dots, m_k . The terms $e_{m_2}, \dots, e_{m_k}, e_{x_1}, \dots, e_{x_k}$ and e_y represent all unexplained causes of $m_2, \dots, m_k, x_1, \dots, x_k$, and y , respectively, and are included to explicitly reflect uncertainty in all endogenous nodes (whether modelled or not). (b) Path diagrams depicting the k standard regression models that would be constructed to estimate the total causal effect of each of x_1, x_2, \dots, x_k on y (i.e. equation (12)). For each model, only the final coefficient may be interpreted as a total causal effect; all other coefficients are greyed to illustrate that no such interpretation should be made for them. (c) Path diagrams depicting the UR model, consisting of $2(k - 1)$ preparation regressions (i.e. equations (13) and (14)) and a final composite regression model (i.e. equation (15), with $i = k$).

4 UR models: No confounders (Figure 1(a))

Before considering any additional confounding variables, we first consider the straightforward scenario depicted in Figure 1(a). We provide: definitions of the standard regression models, UR terms, and UR models (§4.1); an analysis of UR models within a causal framework (§4.2); and arguments for why Properties (i)–(iii) are upheld (§4.3).

4.1 Definitions

We define the ordinary least-squares (OLS) regression model $\hat{y}_S^{(i)}$ for estimating the total causal effect of each measurement of the exposure variable x_i (for $1 \leq i \leq k$) on y as:

$$\hat{y}_S^{(i)} = \hat{\alpha}_0^{(i)} + \hat{\alpha}_{x_1}^{(i)}x_1 + \hat{\alpha}_{x_2}^{(i)}x_2 + \dots + \hat{\alpha}_{x_i}^{(i)}x_i \tag{5}$$

A visual depiction of equation (5) is given in Figure 1(b). Because the relationship between each x_i and y is confounded by all previous measurements of x (i.e. x_1, \dots, x_{i-1}), these covariates must be adjusted for. However, as discussed in Section 1, only the coefficient of the last/most recent measurement of x (i.e. $\hat{\alpha}_{x_i}^{(i)}$) may be interpreted as a total causal effect.

To create UR terms according to the process established by Keijzer-Veen et al.,⁹ each measurement of the exposure x_i is regressed on all previous measurements of x (for $2 \leq i \leq k$):

$$x_i = \hat{\gamma}_0^{(i)} + \hat{\gamma}_{x_1}^{(i)}x_1 + \hat{\gamma}_{x_2}^{(i)}x_2 + \dots + \hat{\gamma}_{x_{i-1}}^{(i)}x_{i-1} + e_{xi} \tag{6}$$

The UR term e_{xi} thus represents the difference between the actual value of x_i and the value of x_i as predicted by all previous measurements of x .

Lastly, we define the UR model $\hat{y}_{UR}^{(i)}$ (for $1 \leq i \leq k$), which represents the outcome y as function of the initial value of the exposure x_1 and subsequent ‘unexplained’ increases e_{x2}, \dots, e_{xi} :

$$\hat{y}_{UR}^{(i)} = \hat{\lambda}_0^{(i)} + \hat{\lambda}_{x1}^{(i)}x_1 + \hat{\lambda}_{ex2}^{(i)}e_{x2} + \dots + \hat{\lambda}_{exi}^{(i)}e_{xi} \tag{7}$$

The composite UR model $\hat{y}_{UR}^{(k)}$ thus represents the outcome y as function of the initial value of the exposure x_1 and all subsequent ‘unexplained’ increases e_{x2}, \dots, e_{xk} . The UR modelling process is summarised in Figure 1(c), depicting $k - 1$ regressions of x_i on x_1, \dots, x_{i-1} (equation (6)) and one composite UR regression model (equation (7), with $i = k$).

4.2 A causal framework

Within the causal framework provided by Figure 1(a), the unique properties of UR models can be visualised. If we were naively to model x_1, x_2, \dots, x_k simultaneously, only the coefficient of the final measurement x_k could be interpreted as a total causal effect on y ; the coefficients of x_1, \dots, x_{k-1} would represent only the direct effects of each measurement on y , because all future measurements would fully mediate the respective relationship and all backdoor paths¹ would be blocked by preceding measurements. However, by modelling $x_1, e_{x2}, \dots, e_{xk}$ (as in a UR model), we encounter no mediation problems due to the fact that, by construction, the UR terms remain wholly independent of the other terms in the model. In fact, by placing the UR model in a causal framework, we are able to see that the UR terms e_{x2}, \dots, e_{xk} are essentially instrumental variables (IVs)⁴² for x_2, \dots, x_k , respectively, which have been produced by the modelling process (Note: The process has similarities with the two-stage least squares regression method,⁴³ a form of instrumental variable analysis commonly encountered in economics research).

All techniques based on linear regression, including UR models, assume that the causal relationships between variables are linear functions. If that is the case, we may parameterise a DAG (as in Figure 1(a)) by assigning a single coefficient to every arrow and assuming all variables to have a variance of one. The method of path coefficients⁴⁴ then allows us to determine the ‘true’ total causal effects in the data generating process. Take x_2 as an example, where $k = 3$. The total effect of x_2 on y encompasses the direct effect from $x_2 \rightarrow y$ and all indirect effects (of which there is only one in this scenario): $x_2 \rightarrow x_3 \rightarrow y$. We introduce the notation p_{ba} to represent the coefficient of the arrow $a \rightarrow b$. Table 1 gives the total effects of x_2 on y and of e_{x2} on y , with both total effects decomposed into their respective direct and indirect effects. From Table 1, we see that the total effect of x_2 on y is equal to the total effect of e_{x2} on y ; this is because there are no direct paths between e_{x2} and y , and all indirect paths pass through x_2 (with $p_{x_2e_{x2}}$ being equal to one, as in Figure 1(c)).

4.3 Covariate orthogonality and Properties (i)–(iii)

In addition to the graph-based approach in the preceding section, we are able to prove mathematically that Properties (i)–(iii) are upheld for the scenario given in Figure 1(a). In summary, these properties are:

- **Property (i):** $\hat{y}_S^{(k)} = \hat{y}_{UR}^{(k)}$
- **Property (ii):** $\hat{\alpha}_{x1}^{(1)} = \hat{\lambda}_{x1}^{(k)}$
- **Property (iii):** $\hat{\alpha}_{xi}^{(i)} = \hat{\lambda}_{exi}^{(k)}$

Equations (5) to (7) are summarised in Table 2; the standard regression models $\hat{y}_S^{(i)}$ (for $1 \leq i \leq k$) and composite UR model $\hat{y}_{UR}^{(k)}$ (in which the UR terms have been produced via the regression of each measurement of x on all previous measurements, as in equation (5)) contained therein are guaranteed to satisfy Properties (i)–(iii). These properties of UR models rely crucially on all UR terms e_{x2}, \dots, e_{xk} being orthogonal to all other covariates in the composite UR model $\hat{y}_{UR}^{(k)}$.

We illustrate this property, and explain how it is exploited to ensure Properties (i)–(iii) are upheld. Formal proofs are provided in online supplementary Appendix 1.

Table 1. Total effect of x_2 on y estimated by a standard regression model compared to total effect of e_{x2} on y estimated by an equivalent UR model (Figure 1(a), with $k = 3$).

Exposure	Path	Effect size	Total effect
x_2	Direct:	$x_2 \rightarrow y$	p_{yx_2}
	Indirect:	$x_2 \rightarrow x_3 \rightarrow y$	$p_{x_3x_2} \cdot p_{yx_3}$
e_{x2}	Direct:	n/a	$p_{yx_2} + p_{x_3x_2} \cdot p_{yx_3}$
	Indirect:	$e_{x2} \rightarrow x_2 \rightarrow y$	$p_{x_2e_{x2}} \cdot p_{yx_2}$
		$e_{x2} \rightarrow x_2 \rightarrow x_3 \rightarrow y$	$p_{x_2e_{x2}} \cdot p_{x_3x_2} \cdot p_{yx_3}$

Table 2. For the scenario depicted in Figure 1(a), the standard regression model $\hat{y}_S^{(i)}$ necessary for estimating the total causal effect of each exposure x_i on y , and the corresponding UR model $\hat{y}_{UR}^{(i)}$, for $1 \leq i \leq k$.

	Standard regression model $\hat{y}_S^{(i)}$	UR model $\hat{y}_{UR}^{(i)}$
$i = 1:$	$\hat{\alpha}_0^{(1)} + \hat{\alpha}_{x_1}^{(1)}x_1$	$\hat{\lambda}_0^{(1)} + \hat{\lambda}_{x_1}^{(1)}x_1$
$i = 2:$	$\hat{\alpha}_0^{(2)} + \hat{\alpha}_{x_1}^{(2)}x_1 + \hat{\alpha}_{x_2}^{(2)}x_2$	$\hat{\lambda}_0^{(2)} + \hat{\lambda}_{x_1}^{(2)}x_1 + \hat{\lambda}_{e_{x2}}^{(2)}e_{x2}$
\vdots	\vdots	\vdots
$i = k:$	$\hat{\alpha}_0^{(k)} + \hat{\alpha}_{x_1}^{(k)}x_1 + \hat{\alpha}_{x_2}^{(k)}x_2 + \dots + \hat{\alpha}_{x_k}^{(k)}x_k$	$\hat{\lambda}_0^{(k)} + \hat{\lambda}_{x_1}^{(k)}x_1 + \hat{\lambda}_{e_{x2}}^{(k)}e_{x2} + \dots + \hat{\lambda}_{e_{xk}}^{(k)}e_{xk}$

In Table 2, note that each regression model (for both the standard and UR methods) contains one more covariate than the model preceding it. In the column of standard regression models, each row contains an additional x_i term; in the column of UR models, each row contains an additional e_{x_i} term.

Typically, the inclusion of an additional covariate in a regression model changes the coefficient(s) estimated for other covariates because their covariance would be nonzero. For example, the addition of x_2 in $\hat{y}_S^{(2)}$ will undoubtedly change the estimated coefficient for x_1 in $\hat{y}_S^{(2)}$ compared to $\hat{y}_S^{(1)}$, because x_1 and x_2 are two measurements of the same variable and thus will have a nonzero covariance (i.e. correlation $\neq 0$). This nonzero covariance is what is exploited by adjustment for confounders – if two covariates did not covary, then adjustment would not be necessary in the first place.

However, a UR model upholds Properties (ii) and (iii) specifically because its covariates do not covary. The addition of e_{x2} in $\hat{y}_{UR}^{(2)}$ does not change the estimated coefficient for x_1 in $\hat{y}_{UR}^{(2)}$ compared to $\hat{y}_{UR}^{(1)}$ because x_1 and e_{x2} are orthogonal (i.e. correlation = 0). This orthogonality is ensured as an artefact of OLS regression; because e_{x2} represents the residual term from the regression of x_2 on x_1 by definition (equation (6)), it is guaranteed to be orthogonal to x_1 .

In fact, it can easily be shown that all UR terms e_{x2}, \dots, e_{xk} are orthogonal to one another by construction. For any UR term e_{x_i} , it holds that e_{x_i} is orthogonal to x_1, \dots, x_{i-1} . Because preceding UR terms $e_{x2}, \dots, e_{x_{i-1}}$ are themselves linear combinations of x_1, \dots, x_{i-1} (equation (6)), it follows that e_{x_i} is orthogonal to $e_{x2}, \dots, e_{x_{i-1}}$, for $2 \leq i \leq k$. Using this information, we can easily conclude that the addition of subsequent UR terms in the set of UR models in Table 2 will leave the coefficients of all other covariates unchanged. Thus, it only remains to be shown that the estimated coefficients for x_1 and the UR terms e_{x2}, \dots, e_{xk} are themselves equivalent to the coefficients for x_1, x_2, \dots, x_k as estimated in their individual standard regression models, respectively.

Property (i):

First, it must be noted that each UR model is nothing more than a reparameterisation of the corresponding standard regression model (i.e. $\hat{y}_S^{(i)} = \hat{y}_{UR}^{(i)}$ for each row in Table 2). Each standard regression model $\hat{y}_S^{(i)}$ represents y as a function of x_1, \dots, x_i . In contrast, each UR model $\hat{y}_{UR}^{(i)}$ represents y as a function of $x_1, e_{x2}, \dots, e_{x_i}$. However, e_{x_i} is itself a function of x_1, \dots, x_i (equation (5)), and thus it follows that the UR model $\hat{y}_{UR}^{(i)}$ itself is also a function of x_1, \dots, x_i . Because $\hat{y}_S^{(i)}$ and $\hat{y}_{UR}^{(i)}$ are both functions of the same covariates, it follows that $\hat{y}_S^{(k)} = \hat{y}_{UR}^{(k)}$, thereby satisfying Property (i).

Property (ii):

It is trivially true that the coefficients estimated for x_1 in the first standard regression model $\hat{y}_S^{(1)}$ and corresponding UR model $\hat{y}_{UR}^{(1)}$ will be equal (i.e. $\hat{\alpha}_{x_1}^{(1)} = \hat{\lambda}_{x_1}^{(1)}$) because the models are themselves equivalent. All subsequent UR

terms e_{x_2}, \dots, e_{x_k} are orthogonal to x_1 and to one another; therefore, it follows that the estimated coefficient of x_1 will be equivalent for all UR models in Table 1 (i.e. $\hat{\lambda}_{x_1}^{(1)} = \hat{\lambda}_{x_1}^{(2)} = \dots = \hat{\lambda}_{x_1}^{(k)}$). This ensures that the coefficient of x_1 in $\hat{y}_S^{(1)}$ (which represents the total effect of x_1 on y) will be unchanged in the composite UR model $\hat{y}_{UR}^{(k)}$ (i.e. $\hat{\alpha}_{x_1}^{(1)} = \hat{\lambda}_{x_1}^{(k)}$).

Property (iii):

Lastly, we can show that the coefficient for e_{x_i} (i.e. $\hat{\lambda}_{e_{x_i}}^{(i)}$) in a UR model is equal to the estimated total effect of x_i (i.e. $\hat{\alpha}_{x_i}^{(i)}$) in the corresponding standard regression model. To this end, we consider the following standard regression and corresponding UR models, respectively:

$$\begin{aligned} \hat{y}_S^{(i)} &= \hat{\alpha}_0^{(i)} + \hat{\alpha}_{x_1}^{(i)}x_1 + \hat{\alpha}_{x_2}^{(i)}x_2 + \dots + \hat{\alpha}_{x_i}^{(i)}x_i \\ \hat{y}_{UR}^{(i)} &= \hat{\lambda}_0^{(i)} + \hat{\lambda}_{x_1}^{(i)}x_1 + \hat{\lambda}_{e_{x_2}}^{(i)}e_{x_2} + \dots + \hat{\lambda}_{e_{x_i}}^{(i)}e_{x_i} \end{aligned}$$

We may set these two equations equal to one another (due to Property (i)), substitute the expansions for e_{x_2}, \dots, e_{x_i} (equation (5)) into the UR model and rearrange, thereby producing:

$$\begin{aligned} \hat{\alpha}_0^{(i)} + \hat{\alpha}_{x_1}^{(i)}x_1 + \hat{\alpha}_{x_2}^{(i)}x_2 + \dots + \hat{\alpha}_{x_i}^{(i)}x_i &= \hat{\lambda}_0^{(i)} + \hat{\lambda}_{x_1}^{(i)}x_1 + \hat{\lambda}_{e_{x_2}}^{(i)}e_{x_2} + \dots + \hat{\lambda}_{e_{x_i}}^{(i)}e_{x_i} \\ &= \hat{\lambda}_0^{(i)} + \hat{\lambda}_{x_1}^{(i)}x_1 + \hat{\lambda}_{e_{x_2}}^{(i)}[-\hat{\gamma}_0^{(2)} - \hat{\gamma}_{x_1}^{(2)}x_1 + x_2] + \dots + \hat{\lambda}_{e_{x_i}}^{(i)}[-\hat{\gamma}_0^{(i)} - \hat{\gamma}_{x_1}^{(i)}x_1 \\ &\quad - \hat{\gamma}_{x_2}^{(i)}x_2 - \dots - \hat{\gamma}_{x_{i-1}}^{(i)}x_{i-1} + x_i] \\ &= \left[\hat{\lambda}_0^{(i)} - \hat{\lambda}_{e_{x_2}}^{(i)}\hat{\gamma}_0^{(2)} - \dots - \hat{\lambda}_{e_{x_i}}^{(i)}\hat{\gamma}_0^{(i)} \right] + \left[\hat{\lambda}_{x_1}^{(i)} - \hat{\lambda}_{e_{x_2}}^{(i)}\hat{\gamma}_{x_1}^{(2)} - \dots - \hat{\lambda}_{e_{x_i}}^{(i)}\hat{\gamma}_{x_1}^{(i)} \right]x_1 \\ &\quad + \left[\hat{\lambda}_{e_{x_2}}^{(i)} - \hat{\lambda}_{e_{x_3}}^{(i)}\hat{\gamma}_{x_2}^{(3)} - \dots - \hat{\lambda}_{e_{x_i}}^{(i)}\hat{\gamma}_{x_2}^{(i)} \right]x_2 + \dots + \left[\hat{\lambda}_{e_{x_i}}^{(i)} \right]x_i \end{aligned} \tag{8}$$

From equation (8) above, it becomes clear that the coefficients for x_i in $\hat{y}_S^{(i)}$ and e_{x_i} in $\hat{y}_{UR}^{(i)}$ are equal (i.e. $\hat{\alpha}_{x_i}^{(i)} = \hat{\lambda}_{e_{x_i}}^{(i)}$). Again, we invoke the property of orthogonality to conclude that the estimated coefficient for e_{x_i} will be equivalent for all UR models in Table 2 (i.e. $\hat{\lambda}_{e_{x_i}}^{(1)} = \hat{\lambda}_{e_{x_i}}^{(2)} = \dots = \hat{\lambda}_{e_{x_i}}^{(k)}$). This ensures that the coefficient of e_{x_i} in $\hat{y}_S^{(i)}$ (which represents the total effect of x_i on y) will be unchanged in the composite UR model $\hat{y}_{UR}^{(k)}$ (i.e. $\hat{\alpha}_{x_i}^{(i)} = \hat{\lambda}_{e_{x_i}}^{(k)}$).

5 UR models: Time-invariant confounder (Figure 2(a))

We next consider the scenario in Figure 2(a), in which a time-invariant covariate m confounds the relationship between x_1, x_2, \dots, x_k and y . This section is structured similarly to the preceding one. We provide: definitions of the standard regression models, UR terms, and UR models, all adjusted for the confounder m based upon the DAG in Figure 2(a) (§5.1); an analysis of UR models within a causal framework (§5.2); arguments for why Properties (i)–(iii) are upheld when the defined adjustments for m have been made (§5.3); and a discussion regarding the implications of insufficient adjustment for m (§5.4).

5.1 Definitions (with correct adjustment for m)

Using the DAG in Figure 2(a) as guidance, we extend the original definitions of the standard regression models, UR terms, and UR models (equations (5) to (7), respectively) to properly account for the confounding effect of m , a time-invariant covariate.

We define the OLS regression model $\hat{y}_S^{(i)}$ for estimating the total causal effect of each measurement of the exposure variable x_i (for $1 \leq i \leq k$) on y as:

$$\hat{y}_S^{(i)} = \hat{\alpha}_0^{(i)} + \hat{\alpha}_m^{(i)}m + \hat{\alpha}_{x_1}^{(i)}x_1 + \hat{\alpha}_{x_2}^{(i)}x_2 + \dots + \hat{\alpha}_{x_i}^{(i)}x_i \tag{9}$$

Because the relationship between each x_i and y is confounded by all previous measurements of x (i.e. x_1, \dots, x_{i-1}) and m , these covariates must be adjusted for to obtain an inferentially unbiased estimate of the total causal effect of each measurement of the exposure. As previously, only the coefficient of the last/most recent measurement of x (i.e. $\hat{\alpha}_{x_i}^{(i)}$) may be interpreted as a total causal effect.

We further extend the process of Keijzer-Veen et al.⁹ to create UR terms for this scenario. As is evident, the relationship between each measurement of the exposure variable x_i and all previous measurements x_1, \dots, x_{i-1} is confounded by m (for $2 \leq i \leq k$); thus, adjustment for m is necessary:

$$x_i = \hat{\gamma}_0^{(i)} + \hat{\gamma}_m^{(i)}m + \hat{\gamma}_{x_1}^{(i)}x_1 + \hat{\gamma}_{x_2}^{(i)}x_2 + \dots + \hat{\gamma}_{x_{(i-1)}}^{(i)}x_{i-1} + e_{x_i} \tag{10}$$

Therefore, the UR term e_{x_i} represents the difference between the actual value of x_i and the value of x_i as predicted by all previous measurements x_1, \dots, x_{i-1} , adjusted for the confounding effect of m .

Finally, we define the UR model $\hat{y}_{UR}^{(i)}$ (for $1 \leq i \leq k$); this model must be also be adjusted for m , since m confounds the relationship between x_1 and y :

$$\hat{y}_{UR}^{(i)} = \hat{\lambda}_0^{(i)} + \hat{\lambda}_m^{(i)}m + \hat{\lambda}_{x_1}^{(i)}x_1 + \hat{\lambda}_{e_{x_2}}^{(i)}e_{x_2} + \dots + \hat{\lambda}_{e_{x_i}}^{(i)}e_{x_i} \tag{11}$$

The composite UR model $\hat{y}_{UR}^{(k)}$ thus represents the outcome y as function of the initial value of the exposure x_1 , all subsequent ‘unexplained’ increases e_{x_2}, \dots, e_{x_k} , and the time-invariant confounder m .

As in the preceding section, visual depictions of the previous equations are provided, with Figure 2(b) corresponding to equation (8) and Figure 2(c) corresponding to equation (8) and equation (9) (with $i = k$).

5.2 A causal framework

We may easily extend the reasoning from the previous scenario (§4.2) to explain why the UR model (equation (11)) satisfies Properties (i)–(iii) before resorting to mathematics, by considering the diagram in Figure 2(a) as a path diagram. A regression model containing all of m, x_1, x_2, \dots, x_k (as in equation (9)) would only allow for the interpretation of the coefficient of x_k as a total causal effect on y ; the coefficients of x_1, \dots, x_{k-1} would represent only the direct effects of each measurement on y , because all future measurements would mediate the respective relationship and all backdoor paths would be blocked by preceding measurements (including m). Within the UR model, the independence of all UR terms e_{x_2}, \dots, e_{x_k} ensures no mediating paths are blocked, and the only backdoor path between x_1 and y is blocked by m .

5.3 Covariate orthogonality and Properties (i)–(iii)

In addition to the graph-based approach in the preceding section (§5.2), we are able to illustrate mathematically that adjustment for m both when generating each UR term e_{x_i} (equation (10)) and in the composite UR model (Eq.11) will result in Properties (i)–(iii) being satisfied. Note that the scenario depicted in Figure 2(a) is nearly indistinguishable, both visually and mathematically, from the scenario in Figure 1(a). The confounder m (which affects y and all measurements of x) could be reimagined as variable x_0 ; viewed in this way, the need for its adjustment becomes clear and the proofs from the previous section apply with only minor notational adjustments. Even though a distinction must be drawn between exposure variables and confounding variables within a causal framework, OLS regression treats both equivalently (i.e. as ‘independent variables’). Therefore, we give a brief outline only of how the adjustments deemed necessary by the causal diagram in Figure 2(a) will result in Properties (i)–(iii) being upheld and attach the formal mathematical proofs in online supplementary Appendix 2.

Equations (9) to (11), which are summarised in Table 3, are guaranteed satisfy Properties (i)–(iii). As in the previous scenario (§4.3), each regression model (for both the standard and UR methods) in Table 3 contains one more covariate than the model preceding it – an additional x_i term in the column of standard regression models,

Table 3. For the scenario depicted in Figure 2(a), the standard regression model $\hat{y}_S^{(i)}$ necessary for estimating the total causal effect of each exposure x_i on y , and the corresponding UR model $\hat{y}_{UR}^{(i)}$, for $1 \leq i \leq k$.

	Standard regression model $\hat{y}_S^{(i)}$	UR model $\hat{y}_{UR}^{(i)}$
$i = 1$:	$\hat{\alpha}_0^{(1)} + \hat{\alpha}_m^{(1)}m + \hat{\alpha}_{x_1}^{(1)}x_1$	$\hat{\lambda}_0^{(1)} + \hat{\lambda}_m^{(1)}m + \hat{\lambda}_{x_1}^{(1)}x_1$
$i = 2$:	$\hat{\alpha}_0^{(2)} + \hat{\alpha}_m^{(2)}m + \hat{\alpha}_{x_1}^{(2)}x_1 + \hat{\alpha}_{x_2}^{(2)}x_2$	$\hat{\lambda}_0^{(2)} + \hat{\lambda}_m^{(2)}m + \hat{\lambda}_{x_1}^{(2)}x_1 + \hat{\lambda}_{e_{x_2}}^{(2)}e_{x_2}$
\vdots	\vdots	\vdots
$i = k$:	$\hat{\alpha}_0^{(k)} + \hat{\alpha}_m^{(k)}m + \hat{\alpha}_{x_1}^{(k)}x_1 + \hat{\alpha}_{x_2}^{(k)}x_2 + \dots + \hat{\alpha}_{x_k}^{(k)}x_k$	$\hat{\lambda}_0^{(k)} + \hat{\lambda}_m^{(k)}m + \hat{\lambda}_{x_1}^{(k)}x_1 + \hat{\lambda}_{e_{x_2}}^{(k)}e_{x_2} + \dots + \hat{\lambda}_{e_{x_k}}^{(k)}e_{x_k}$

and an additional e_{xi} term in the column of UR models. Proofs for the previous scenario relied on the property of each UR term being orthogonal to all preceding terms in the model. Adjustment for m when generating each UR term e_{xi} (equation (10)) guarantees that this property will be upheld, because it ensures that e_{xi} is orthogonal to m in addition to $e_{x1}, \dots, e_{x(i-1)}$; this cannot be guaranteed without explicit adjustment for m . Furthermore, adjustment for m in each UR model in Table 3 ensures that $\hat{y}_S^{(i)} = \hat{y}_{UR}^{(i)}$ for each row in Table 3.

5.4 Incorrect adjustment for m

We have used the causal diagram in Figure 2(a) to argue for the necessity of adjusting for a time-invariant confounder m during both stages of the UR modelling process, and have demonstrated how such adjustments will produce a composite UR model that satisfies Properties (i)–(iii), as Keijzer-Veen et al. intended. We now consider the implications of insufficient adjustment.

Without adjustment for m when generating each UR term e_{xi} , the coefficients of x_1, \dots, x_{i-1} (i.e. $\hat{\gamma}_{xi}^{(j)}$, for $1 \leq i \leq k - 1$ and $1 \leq j \leq k$) and the UR term will absorb the effect of the omitted variable m on x_i , thereby biasing the total effect of e_{xi} estimated within the UR model (so-called ‘omitted variable bias’). Further, it is evident that m confounds the relationship between x_1 and y , so that failure to adjust for m in the composite UR model will produce different predicted outcome values and bias the estimated coefficient of x_1 .

6 UR models: Time-varying confounder (Figure 3(a))

Finally, we consider the scenario in Figure 3(a), in which a time-varying covariate m_1, m_2, \dots, m_k confounds the relationship between x_1, x_2, \dots, x_k and y .

In this section, we again provide: definitions of the standard regression models, UR terms, and UR models, all adjusted for the confounder m_1, m_2, \dots, m_k based upon the DAG in Figure 3(a) (§6.1); an analysis of UR models within a causal framework (§6.2); arguments for why Properties (i)–(iii) are upheld when the defined adjustments for m_1, m_2, \dots, m_k have been made (§6.3); and a discussion regarding the implications of insufficient adjustment for m_1, m_2, \dots, m_k (§6.4).

6.1 Definitions (with correct adjustment for m_1, m_2, \dots, m_k)

Using the DAG in Figure 3(a), we extend the original definitions of the standard regression models, UR terms, and UR models (equations (5) to (7), respectively) to properly account for the confounding effect of m_1, m_2, \dots, m_k , a time-varying covariate.

We define the OLS regression model $\hat{y}_S^{(i)}$ for estimating the total causal effect of each measurement of the exposure variable x_i (for $1 \leq i \leq k$) on y as:

$$\hat{y}_S^{(i)} = \hat{\alpha}_0^{(i)} + \hat{\alpha}_{m_1}^{(i)} m_1 + \hat{\alpha}_{x_1}^{(i)} x_1 + \dots + \hat{\alpha}_{m_i}^{(i)} m_i + \hat{\alpha}_{x_i}^{(i)} x_i \tag{12}$$

The relationship between each x_i and y is not only confounded by all previous values of the exposure x_1, \dots, x_{i-1} but also by the current measurement and all previous measurements of the confounder m_1, \dots, m_i . Therefore, adjustment for $m_1, \dots, m_i, x_1, \dots, x_{i-1}$ is necessary to obtain an inferentially unbiased estimate of the total causal effect of each measurement of the exposure. We reiterate that only the coefficient of the last/most recent measurement of x (i.e. $\hat{\alpha}_{x_i}^{(i)}$) may be interpreted as a total causal effect.

Extending the process of Keijzer-Veen et al.⁹ to create UR terms for each measurement of the exposure x_i in this scenario necessitates adjustment for the current measurement and all previous measurements of the confounder m_1, m_2, \dots, m_i (for $2 \leq i \leq k$), since these variables confound the relationship between each measurement of the exposure variable x_i and all previous measurements x_1, \dots, x_{i-1} , i.e.:

$$x_i = \hat{\gamma}_0^{(i)} + \hat{\gamma}_{m_1}^{(i)} m_1 + \hat{\gamma}_{x_1}^{(i)} x_1 + \dots + \hat{\gamma}_{m_{(i-1)}}^{(i)} m_{i-1} + \hat{\gamma}_{x_{(i-1)}}^{(i)} x_{i-1} + \hat{\gamma}_{m_i}^{(i)} m_i + e_{xi} \tag{13}$$

In this way, e_{xi} represents the difference between the observed value of x_i and the value of x_i as predicted by all previous measurements x_1, \dots, x_{i-1} , adjusted for the confounding effects of m_1, m_2, \dots, m_i .

As we have demonstrated previously (§4.3, §5.3), UR models rely upon the orthogonality of terms in the composite UR model. This necessitates the creation of UR terms e_{mi} for each measurement of the time-varying confounding variable m_i (for $2 \leq i \leq k$) in a similar manner to that of the UR terms e_{xi} (equation (13)). Each e_{mi} is

derived from the OLS regression of m_i on all previous values of the confounder m_1, \dots, m_{i-1} , as well as all previous values of the exposure x_1, x_2, \dots, x_{i-1} which confound this relationship:

$$m_i = \hat{\eta}_0^{(i)} + \hat{\eta}_{m_1}^{(i)} m_1 + \hat{\eta}_{x_1}^{(i)} x_1 + \dots + \hat{\eta}_{m_{(i-1)}}^{(i)} m_{i-1} + \hat{\eta}_{x_{(i-1)}}^{(i)} x_{i-1} + e_{mi} \tag{14}$$

Thus, e_{mi} has a similar interpretation to the original UR term e_{xi} , in that it represents the part of m_i unexplained by all previous values m_1, \dots, m_{i-1} , adjusted for the confounding effects of x_1, \dots, x_{i-1} .

Lastly, we define the UR model $\hat{y}_{UR}^{(i)}$ (for $1 \leq i \leq k$) as a function of the initial value of the confounder m_1 and its subsequent ‘unexplained’ increases e_{m2}, \dots, e_{mi} , and the initial value of the exposure x_1 and its subsequent ‘unexplained’ increases e_{x2}, \dots, e_{xi} :

$$\hat{y}_{UR}^{(i)} = \hat{\lambda}_0^{(i)} + \hat{\lambda}_{m_1}^{(i)} m_1 + \hat{\lambda}_{x_1}^{(i)} x_1 + \hat{\lambda}_{em2}^{(i)} e_{m2} + \hat{\lambda}_{ex2}^{(i)} e_{x2} + \dots + \hat{\lambda}_{emi}^{(i)} e_{mi} + \hat{\lambda}_{exi}^{(i)} e_{xi} \tag{15}$$

As previously, visual depictions of these equations are provided. Figure 3(b) corresponds to the standard regression models given by equation (12); Figure 3(c) corresponds to the $k - 1$ regressions of x_i on all preceding measurements of x and m (equation (13)), the $k - 1$ regressions of m_i on all preceding measurements of x and m (equation (14)), and one composite UR regression model (equation (15), with $i = k$).

6.2 A causal framework

The similarities amongst the three causal scenarios depicted in Figures 1(a), 2(a), and 3(a) are evident, and shed light on how the reasoning from the previous scenarios (§4.2 and §5.2) can be extended to demonstrate why the UR model in equation (15) satisfies Properties (i)–(iii). In a regression model containing all of $m_1, \dots, m_k, x_1, \dots, x_k$ (as in equation (12), with $i = k$), only the coefficient of x_k could be interpreted as a total causal effect on y ; the coefficients of x_1, \dots, x_{k-1} may only be interpreted as the direct effects of each measurement of the exposure on y , because all future measurements of both x and m would fully mediate the respective relationship and all preceding measurements of x and m would block all backdoor paths. Within the UR model, however, the independence of all UR terms for both the exposure (i.e. e_{x2}, \dots, e_{xk}) and confounder (i.e. e_{m2}, \dots, e_{mk}) ensures no mediating paths are blocked, and the only backdoor path between x_1 and y is blocked by m_1 .

6.3 Covariate orthogonality and Properties (i)–(iii)

In addition to the graph-based approach in the preceding section (§6.2), we can illustrate mathematically that the standard regression models $\hat{y}_S^{(i)}$ (equation (12)), UR terms for measurements of the exposure (equation (13)) and confounder (equation (14)), and composite UR model $\hat{y}_{UR}^{(k)}$ (equation (15), with $i = k$) satisfy Properties (i)–(iii). Although seemingly more complex, the scenario depicted in Figure 3(a) also has very little to distinguish it from the scenarios in Figures 1(a) and 2(a). The confounder m_1 , being the only exogenous node on the graph, could be imagined as variable x_0 , with all nodes subsequent to x_1 having an associated UR term. Viewed as such, the necessity of adjusting for m_1 and creating UR terms for both the exposure and the time-varying confounder becomes apparent, as the causal diagram in Figure 3(a) is equivalent to that of Figure 2(a) with minor notational adjustments. Therefore, we provide only a brief outline of how the adjustments deemed necessary by the causal diagrams in Figure 3(a) will result in Properties (i)–(iii) being upheld; formal mathematical proofs are provided in online supplementary Appendix 3.

Equations (12) to (15) are summarised in Table 4 and are guaranteed to satisfy Properties (i)–(iii). In contrast to previous scenarios (§4.3 and §5.3), each regression model (for both the standard and UR models) contains *two* more covariates than the model preceding it. In the column of standard regression models, each row contains an additional x_i and m_i term; in the column of UR models, each row contains an additional e_{xi} and e_{mi} term. Thus, for Properties (i)–(iii) to be upheld in in each UR model $\hat{y}_{UR}^{(i)}$, these two additional terms must be orthogonal to one another and to all preceding terms.

Proving this is relatively straightforward. For any UR term e_{mi} for the confounder, it holds that e_{mi} is orthogonal to $m_1, \dots, m_{i-1}, x_1, \dots, x_{i-1}$ by construction (equation (14)). Because preceding UR terms $e_{x2}, \dots, e_{x(i-1)}$ (equation (13)) and $e_{m2}, \dots, e_{m(i-1)}$ (equation (14)) may be expressed as linear combinations of $m_1, \dots, m_i, x_1, \dots, x_{i-1}$, it follows that e_{mi} is orthogonal to $e_{m2}, \dots, e_{m(i-1)}, e_{x2}, \dots, e_{x(i-1)}$. Furthermore, for any UR term e_{xi} for the exposure, it holds that e_{xi} is orthogonal to $m_1, \dots, m_i, x_1, \dots, x_{i-1}$ by construction (equation (13)). Because preceding UR terms $e_{x2}, \dots, e_{x(i-1)}$ (equation (13)) and e_{m2}, \dots, e_{mi} (equation (14)) may be expressed

Table 4. For the scenario depicted in Figure 3(a), the standard regression model $\hat{y}_S^{(i)}$ necessary for estimating the total causal effect of each exposure x_i on y , and the corresponding UR model $\hat{y}_{UR}^{(i)}$, for $1 \leq i \leq k$.

	Standard regression model $\hat{y}_S^{(i)}$	UR model $\hat{y}_{UR}^{(i)}$
$i = 1$:	$\hat{\alpha}_0^{(1)} + \hat{\alpha}_{m_1}^{(1)}m_1 + \hat{\alpha}_{x_1}^{(1)}x_1$	$\hat{\lambda}_0^{(1)} + \hat{\lambda}_{m_1}^{(1)}m_1 + \hat{\lambda}_{x_1}^{(1)}x_1$
$i = 2$:	$\hat{\alpha}_0^{(2)} + \hat{\alpha}_{m_1}^{(2)}m_1 + \hat{\alpha}_{x_1}^{(2)}x_1 + \hat{\gamma}_{m_2}^{(2)}m_2 + \hat{\alpha}_{x_2}^{(2)}x_2$	$\hat{\lambda}_0^{(2)} + \hat{\lambda}_{m_1}^{(2)}m_1 + \hat{\lambda}_{x_1}^{(2)}x_1 + \hat{\lambda}_{em_2}^{(2)}e_{m_2} + \hat{\lambda}_{ex_2}^{(2)}e_{x_2}$
\vdots	\vdots	\vdots
$i = k$:	$\hat{\alpha}_0^{(k)} + \hat{\alpha}_{m_1}^{(k)}m_1 + \hat{\alpha}_{x_1}^{(k)}x_1 + \hat{\alpha}_{m_2}^{(k)}m_2 + \hat{\alpha}_{x_2}^{(k)}x_2$ $+ \dots + \hat{\alpha}_{m_k}^{(k)}m_k + \hat{\alpha}_{x_k}^{(k)}x_k$	$\hat{\lambda}_0^{(k)} + \hat{\lambda}_{m_1}^{(k)}m_1 + \hat{\lambda}_{x_1}^{(k)}x_1 + \hat{\lambda}_{em_2}^{(k)}e_{m_2} + \hat{\lambda}_{ex_2}^{(k)}e_{x_2}$ $+ \dots + \hat{\lambda}_{emk}^{(k)}e_{mk} + \hat{\lambda}_{exk}^{(k)}e_{xk}$

as linear combinations of $m_1, \dots, m_i, x_1, \dots, x_{i-1}$, it follows that e_{x_i} is orthogonal to $e_{m_2}, \dots, e_{m_i}, e_{x_2}, \dots, e_{x_{i-1}}$. Thus, we are able to conclude that e_{m_i} and e_{x_i} are orthogonal to one another and to all preceding terms in for any UR model $\hat{y}_{UR}^{(i)}$; adjustment for all causally preceding measurements of both m and x when generating UR terms for both the confounder and the exposure ensures this orthogonality.

6.4 Incorrect adjustment for m_1, m_2, \dots, m_k

The DAG in Figure 3(a) demonstrates the necessity of adjusting for a time-varying confounder m_1, m_2, \dots, m_k in the manner described in Section 6.1, and we have demonstrated how such adjustments will produce a composite UR model that satisfies Properties (i)–(iii). The implications of incorrect adjustment for a time-varying confounder m_1, m_2, \dots, m_k in a UR model are similar to those of incorrect adjustment for a time-invariant confounder m , which were previously outlined in Section 5.4. Without adjustment for any of m_1, \dots, m_i when constructing each UR term for the exposure e_{x_i} , the coefficients of x_1, \dots, x_{i-1} (i.e. $\hat{\gamma}_{x_i}^{(j)}$, for $1 \leq i \leq (k-1)$ and $1 \leq j \leq k$) and the UR term will absorb the effect of each omitted variable on x_i ; this will result in the coefficient estimated for each e_{x_i} in the composite UR model being unequal to the total effect of x_i in its corresponding standard regression model.

The requirement of orthogonal covariates within the composite UR model also sheds light on the necessity for generating UR terms $e_{m_2}, e_{m_3}, \dots, e_{m_k}$ for measurements of a time-varying confounder, if present. We might easily imagine a scenario in which we considered only the original covariates m_1, m_2, \dots, m_k in the UR model. In such a scenario, the terms would remain correlated with each other and with x_1 ; therefore, the inclusion of subsequent m terms in the UR model would necessarily change the coefficient estimates for x_1 and all other covariates.

7 UR model interpretation

Having demonstrated that confounder adjustment within UR models is possible, we consider the claim⁹ that UR models offer additional insight via the coefficients for each UR term e_{x_i} (e.g. $\hat{\lambda}_{ex_i}^{(k)}$ in equation (7), for $2 \leq i \leq k$) into the effect of x_i increasing *more than expected* upon y .

Consider again the simple example with two longitudinal measurements of a continuous exposure x (i.e. x_1 and x_2), outcome y , and no additional confounders (i.e. Figure 1(a), with $k = 2$); the standard regression model (with x_2 as the specified exposure variable) and ‘equivalent’ UR model are given below, respectively:

$$\begin{aligned}\hat{y}_S^{(2)} &= \hat{\alpha}_0^{(2)} + \hat{\alpha}_{x_1}^{(2)}x_1 + \hat{\alpha}_{x_2}^{(2)}x_2 \\ \hat{y}_{UR}^{(2)} &= \hat{\lambda}_0^{(2)} + \hat{\lambda}_{x_1}^{(2)}x_1 + \hat{\lambda}_{ex_2}^{(2)}e_{x_2}\end{aligned}$$

It has been shown (§4.3) that $\hat{\alpha}_{x_2}^{(2)}$ and $\hat{\lambda}_{ex_2}^{(2)}$ are equal, yet $\hat{\alpha}_{x_2}^{(2)}$ is interpreted as the total effect of a one-unit increase in x_2 on y , whereas $\hat{\lambda}_{ex_2}^{(2)}$ is (supposedly) interpreted as the total effect of a one-unit *higher than expected* increase in x_2 on y . If these two variables truly are distinct, their regression coefficients should likewise be distinct. This issue has also been addressed by Tu and Gilthorpe,¹¹ who have argued that the two coefficients are equivalent because adjustment for x_1 in $\hat{y}_S^{(2)}$ amounts to testing the relation between y and the part of x_2 *unexplained by* x_1 (i.e. the unexplained residual). In fact, the two coefficients are equal simply because they mean the same thing. The UR model does not, therefore, offer any additional insight into the effect of higher than expected change in x on the outcome.¹⁵

We also raise a more philosophical point, which speaks to the need for any model to reflect accurately the underlying data-generation process of a given scenario. As an artefact of OLS regression, the UR terms will always

be mathematically independent of the value of the initial measurement of the exposure and all subsequent measurements. This is unlikely to be an accurate representation of real-world exposure variables. Many of these, such as body size, exhibit a consistent, cumulative presence that is only manifest at the discrete time points at which it is measured; these measurements are thus distinct only as a result of the discretisation of time within the measurement processes adopted. Moreover, in auxological studies, the phenomenon of so-called compensatory (or ‘catch up’) growth has been well documented, with accelerated growth being observed in individuals who begin with a low value of some measure, e.g. birthweight.^{45,46} Therefore, however convenient and mathematically sound it may be to model data in a way that implies complete statistical independence amongst an exposure variable’s initial value and its subsequent measurements, this assumption is likely to be implausible and unrealistic for most biological and social variables of interest to epidemiologists. This is a weakness shared by all conditional approaches (of which UR models are one), which has led several authors⁴⁷ to recommend that the results be considered alongside those produced by other methods, rather than in isolation.

8 Standard error reduction

Finally, we address an important consequence of the use of UR models; namely, that they underestimate the standard errors (SEs) of estimated coefficients, thereby resulting in artificial precision of estimated effect sizes. Although focus on *statistical* significance by way of p -values and confidence intervals is not in and of itself justifiable within a causal framework (as focus is effect size and likely *functional* significance, e.g. the absolute risk posed or the potential for substantive intervention), we consider it an important issue to address as a matter of clarity for researchers seeking to use UR models.

To demonstrate, we have simulated 1000 non-overlapping random samples of 1000 observations from a multivariate normal distribution based upon the DAG in Figure 1(a) with $k = 2$, using the ‘dagitty’ package (v. 0.2–2)^{4,48} in R (v. 3.3.2).⁴⁹ Each sample was used to create: (1) the two standard regression models necessary for estimating the total causal effect of each of x_1, x_2 on y (equation (5)); (2) the UR term e_{x_2} , derived by regressing x_2 on x_1 (equation (6)); and (3) the composite UR model in which y is regressed on x_1 and e_{x_2} (equation (7)). For each standard regression model $\hat{y}_S^{(i)}$ (for $i = 1, 2$), the reported SE of the regression coefficient for exposure x_i is stored. For each composite UR model $\hat{y}_{UR}^{(2)}$, the SE of the regression coefficient for each of x_1, e_{x_2} is stored in two forms: (1) as reported in the UR model summary output; and (2) as estimated by bootstrapping 1000 samples and calculating the standard deviation of the distribution of estimated coefficients. Additional details relating to this simulation – including parameters and code – are located in online supplementary Appendix 4. (Note: The specific correlation structure and parameter values used to simulate the data are unimportant for the purposes of this demonstration).

By definition, the SE of an estimated regression coefficient is a point estimate of the standard deviation of an (infinitely) large sampling distribution of estimated regression coefficients. We have shown that standard regression and UR models elicit identical point estimates of the total causal effects of each measure of the longitudinal exposure (§4); from this, it follows that the associated SEs should themselves be equal.

Violin plots of the SEs estimated for each coefficient representing a total causal effect across the 1000 simulations are displayed in Figure 4 for each method considered. As is evident, the *reported* SEs within the UR models are reduced in comparison to those within the first standard regression models (for designated exposure x_1) and equal to those within the final standard regression models (for designated exposure x_2). This demonstrates an apparent paradox: the coefficient values are equivalent, yet the associated SEs are unequal.

We argue that the apparent reduction in SEs achieved by using UR models is purely artefactual and arises from the explicit conditioning on future measurements of x within a UR model. In the standard regression analysis, the only information within the data that is used to inform SE estimation lies in the past (i.e. past measures of the exposure plus any confounders). In contrast, the UR modelling process generates (orthogonal) residuals for the entire exposure period and combines these into a single model, thereby using information within the data that is from both the past *and the future*. If we possessed data pertaining to any true independent causes of future measurements of the exposure, such a method would indeed be valid; however, the UR terms are simply estimated using prior measurements of the exposure. Moreover, due to the fact that they are estimates, the UR terms themselves contain additional variation that is not accommodated by traditional regression methods which assume covariates are measured without error. Consequently, the SEs of estimated causal effect derived from UR models are artefactually reduced and should not be inferred as robust. Indeed, when the SEs within the UR models are estimated via bootstrapping, they are similar to those within the standard regression models.

Comparing the two plots in Figure 4 offers clarity to this argument: (a) displays *differing* distributions of the reported SEs for the coefficient estimates of x_1 (where conditioning on the future information given by x_2 reduces




Figure 4. Violin plots comparing the standard errors associated with equivalent coefficients estimated in standard regression vs. UR models, for data simulated based upon the scenario depicted in Figure 1(a) (with $k = 2$). Horizontal bars within each distribution represent the mean ± 1 standard deviation.

the standard error in the UR model); whereas (b) displays the *same* distribution of the reported SEs for the coefficient estimates of x_2 and e_{x_2} (where the standard regression model correctly exploits all prior information given by x_1 , as does the UR model). Although the magnitude of bias in estimated SEs is small in this simulated example, it will always be present due to the way in which UR models are constructed. Quantifying the magnitude of this bias is not trivial and is beyond the scope of the present study.

9 Conclusion

The mathematical appraisal of UR models that we have undertaken confirms that the method proposed by Keijzer-Veen et al.⁹ is capable of accommodating more than two longitudinal measurements of an exposure

variable and demonstrates how adjustment for confounding variables should be made in this framework to uphold the property that the coefficients for the terms $x_1, e_{x2}, \dots, e_{xk}$ estimated within a UR model are equal to the total effects for x_1, x_2, \dots, x_k estimated by their respective standard regression models. This result will only be guaranteed to hold when adjustment for *all* confounding variables has been made at both stages in the UR modelling process (i.e. when generating UR terms for subsequent measurements of the exposure and in the composite UR model). From a statistical perspective, adjustment for all preceding variables (including confounders) ensures orthogonality amongst the covariates in a composite UR model. Therefore, when the potential confounder is time-varying, it is also necessary to generate UR terms for subsequent measurements of the confounder itself and include these in the final composite models used.

As our proofs only consider one confounding variable, the causal framework provided by DAGs should aid future researchers who wish to extend robustly UR models to situations involving multiple, possibly causally linked, time-invariant and time-varying confounders. Such a DAG will be useful in identifying confounders and establishing the temporal ordering of variables, thereby ensuring that all preceding variables are adjusted for when generating the necessary UR terms.

Although UR models can accommodate multiple measurements of an exposure variable in addition to confounding variables, we have concerns about their practical implementation. Although only one UR model need ultimately be presented, the necessity of generating orthogonal covariates for that UR model requires that many models be created; this has the potential to be quite substantial when multiple confounders are considered. For an exposure x measured at k points in time, the standard regression approach necessitates k separate models for estimating the total causal effect of each measurement on the outcome *regardless of the number of confounders*. In the case of one time-invariant confounder (§5), k models are also created ($k - 1$ models to generate all UR terms and 1 composite UR model); for a time-varying confounder (§6), $2k - 1$ models are created (i.e. $2k - 2$ models to generate all UR terms and 1 composite UR model). The total number of models created by the UR process will always be either equal to or greater than the total number of models created by the standard regression process. If such a process offered real gains in insight into the scenario under consideration, it may indeed be worth it; however, UR models offer no additional insight compared to standard regression methods. Moreover, the inclusion of multiple covariates that are explicitly conditional on one another within the same model also results in artificially reduced standard error estimates, the extent of which has yet to be fully evaluated; the issue can be avoided by bootstrapping, but such a solution may be computationally intensive and require more programming skills than those necessary for implementing the built-in regression functionalities in statistical software packages. Previous research that has utilised UR models without undertaking sufficient adjustment for confounders and correcting SEs via bootstrapping should not be considered robust.

We therefore have strong reservations about the use and implementation of UR models within lifecourse epidemiology, and suggest that researchers considering using them should instead rely on standard regression methods, which produce the same results but are much less likely to be mis-specified and misleading. However, for researchers wishing to use these models, the hypothesised DAG or causal diagram should be presented so that any readers and/or reviewers can confirm that sufficient adjustment for confounders has been undertaken; moreover, SEs should be estimated via bootstrapping and not simply reported as in the model output, as these have the potential to be misleading. We support the recommendation of previous authors⁴⁷ that additional analytical approaches should be considered alongside conditional approaches (e.g. UR models) in order to achieve robust causal conclusions. For example, multilevel, latent growth curve, and growth mixture models may be used to estimate the effects of growth across the lifecourse on a distal outcome, and are more flexible than standard regression methods.⁵ Moreover, the three G-methods^{50,51} are explicitly grounded in a causal framework and allow for the simultaneous consideration of multiple measurements of a longitudinally measured exposure, as well as time-varying confounding; these methods provide exciting avenues of research for lifecourse epidemiologists.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.


Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: KFA and SCG were supported by the Economic and Social Research Council [grant numbers ES/J500215/1 and ES/P000746/1, respectively]. GTHE, PWGT, AH, and MSG were supported by the Higher Education Funding Council for England.

Orcid ID

KF Arnold  <http://orcid.org/0000-0002-0911-5029>

J Textor  <http://orcid.org/0000-0002-0459-9458>

MS Gilthorpe  <http://orcid.org/0000-0001-8783-7695>

Supplemental material

Supplemental material for this article is available online.

References

1. Pearl J, Glymour M and Jewell NP. *Causal inference in statistics: a primer*, 1st ed. Chichester: John Wiley & Sons Ltd, 2016.
2. Kline RB. *Principles and practice of structural equation modelling*, 4th ed. New York: The Guilford Press, 2016.
3. Textor J, Hardt J and Knüppel S. DAGitty: a graphical tool for analyzing causal diagrams. *Epidemiology* 2011; **22**: 745.
4. Textor J, van der Zander B, Gilthorpe MS, et al. Robust causal inference using directed acyclic graphs: the R package ‘dagitty’. *Int J Epidemiol* 2017; **15**: 15.
5. Tu YK, Tilling K, Sterne JAC and Gilthorpe MS. A critical evaluation of statistical approaches to examining the role of growth trajectories in the developmental origins of health and disease. *Int J Epidemiol* 2013; **42**: 1327–1339.
6. Tu YK, West R, Ellison GTH, et al. Why evidence for the fetal origins of adult disease might be a statistical artifact: the “reversal paradox” for the relation between birth weight and blood pressure in later life. *Am J Epidemiol* 2005; **161**: 27–32.
7. Tu YK, Gilthorpe MS and Ellison GTH. What is the effect of adjusting for more than one measure of current body size on the relation between birthweight and blood pressure? *J Hum Hypertens* 2006; **20**: 646–657.
8. Keijzer-Veen MG. Response to Tu and Gilthorpe: preventing misinterpretation of coefficients in analysis of fetal origins of adult disease. *J Clin Epidemiol* 2007; **60**: 319–320.
9. Keijzer-Veen MG, Euser AM, Van Montfoort N, et al. A regression model with unexplained residuals was preferred in the analysis of the fetal origins of adult diseases hypothesis. *J Clin Epidemiol* 2005; **58**: 1320–1324.
10. Cournil A, Coly AN, Diallo A, et al. Enhanced post-natal growth is associated with elevated blood pressure in young Senegalese adults. *Int J Epidemiol* 2009; **38**: 1401–1410.
11. Tu YK and Gilthorpe MS. Unexplained residuals models are not solutions to statistical modeling of the fetal origins hypothesis. *J Clin Epidemiol* 2007; **60**: 318–319.
12. Chioloro A, Paradis G, Madeleine G, et al. Birth weight, weight change, and blood pressure during childhood and adolescence: a school-based multiple cohort study. *J Hypertens* 2011; **29**: 1871–1879.
13. Grijalva-Eternod CS, Wells JC, Girma T, et al. Midupper arm circumference and weight-for-length z scores have different associations with body composition: evidence from a cohort of Ethiopian infants. *Am J Clin Nutr* 2015; **102**: 593–599.
14. Johnson W. Analytical strategies in human growth research. *Am J Hum Biol* 2015; **27**: 69–83.
15. Wills AK, Strand BH, Glavin K, et al. Regression models for linking patterns of growth to a later outcome: infant growth and childhood overweight. *BMC Med Res Methodol* 2016; **16**: 9.
16. Horta BL, Gigante DP, Osmond C, et al. Intergenerational effect of weight gain in childhood on offspring birthweight. *Int J Epidemiol* 2009; **38**: 724–732.
17. Richter LM, Victora CG, Hallal PC, et al. Cohort profile: the consortium of health-orientated research in transitioning societies. *Int J Epidemiol* 2012; **41**: 621–626.
18. Gandhi M, Ashorn P, Maleta K, et al. Height gain during early childhood is an important predictor of schooling and mathematics ability outcomes. *Acta Paediatr* 2011; **100**: 1113–1118.
19. Gonzalez DA, Nazmi A and Victora CG. Growth from birth to adulthood and abdominal obesity in a Brazilian birth cohort. *Int J Obesity* 2010; **34**: 195–202.
20. Yesil GD, Gishti O, Felix JF, et al. Influence of maternal gestational hypertensive disorders on microvasculature in school-age children. *Am J Epidemiol* 2016; **184**: 605–615.
21. Toemen L, Gishti O, Van Osch-Gevers L, et al. Maternal obesity, gestational weight gain and childhood cardiac outcomes: role of childhood body mass index. *Int J Obesity* 2016; **40**: 1070–1078.
22. Toemen L, de Jonge LL, Gishti O, et al. Longitudinal growth during fetal life and infancy and cardiovascular outcomes at school-age. *J Hypertens* 2016; **34**: 1396–1406.
23. Sonnenschein-van der Voort AMM, Gaillard R, de Jongste JC, et al. Foetal and infant growth patterns, airway resistance and school-age asthma. *Respirology* 2016; **21**: 674–682.
24. Lira PIC, Eickmann SH, Lima MC, et al. Early head growth: relation with IQ at 8 years and determinants in term infants of low and appropriate birthweight. *Develop Med Child Neurol* 2010; **52**: 40–46.
25. Adair LS, Martorell R, Stein AD, et al. Size at birth, weight gain in infancy and childhood, and adult blood pressure in 5 low- and middle-income-country cohorts: when does weight gain matter? *Am J Clin Nutr* 2009; **89**: 1383–1392.

26. Hardy R, Ghosh AK, Deanfield J, et al. Birthweight, childhood growth and left ventricular structure at age 60-64 years in a British birth cohort study. *Int J Epidemiol* 2016; **13**: 13.
27. Martorell R, Horta BL, Adair LS, et al. Weight gain in the first two years of life is an important predictor of schooling outcomes in pooled analyses from five birth cohorts from low- and middle-income countries. *J Nutr* 2010; **140**: 348–354.
28. Osmond C, Kajantie E, Forsen TJ, et al. Infant growth and stroke in adult life: the Helsinki birth cohort study. *Stroke* 2007; **38**: 264–270.
29. Antonisamy B, Vasani SK, Geethanjali FS, et al. Weight gain and height growth during infancy, childhood, and adolescence as predictors of adult cardiovascular risk. *J Pediatr* 2017; **180**: 53–61.e3.
30. Kagura J, Adair LS, Munthali RJ, et al. Association between early life growth and blood pressure trajectories in black South African children. *Hypertension* 2016; **68**: 1123–1131.
31. Norris S, Osmond C, Gigante D, et al. Size at birth, weight gain in infancy and childhood, and adult diabetes risk in five low- or middle-income country birth cohorts. *Diab Care* 2012; **35**: 72–79.
32. Stein AD, Barros FC, Bhargava SK, et al. Birth status, child growth, and adult outcomes in low- and middle-income countries. *J Pediatr* 2013; **163**: 1740–1746.e4.
33. De Beer M, Vrijkotte TGM, Fall CHD, et al. Associations of infant feeding and timing of linear growth and relative weight gain during early life with childhood body composition. *Int J Obesity* 2015; **39**: 586–592.
34. Menezes AMB, Hallal PC, Dumith SC, et al. Adolescent blood pressure, body mass index and skin folds: sorting out the effects of early weight and length gains. *J Epidemiol Community Health* 2012; **66**: 149–154.
35. Wells JCK, Hallal PC, Reichert FF, et al. Associations of birth order with early growth and adolescent height, body composition, and blood pressure: prospective birth cohort from Brazil. *Am J Epidemiol* 2011; **174**: 1028–1035.
36. Adair LS, Fall CHD, Osmond C, et al. Associations of linear growth and relative weight gain during early life with adult health and human capital in countries of low and middle income: findings from five birth cohort studies. *Lancet* 2013; **382**: 525–534.
37. Grijalva-Eternod CS, Lawlor DA and Wells JCK. Testing a capacity-load model for hypertension: disentangling early and late growth effects on childhood blood pressure in a prospective birth cohort. *PLoS One* 2013; **8**: e56078.
38. Fink G and Rockers PC. Childhood growth, schooling, and cognitive development: further evidence from the Young Lives study. *Am J Clin Nutr* 2014; **100**: 182–188.
39. Araujo De Franca GV, Lucia Rolfe ED, Horta BL, et al. Associations of birth weight, linear growth and relative weight gain throughout life with abdominal fat depots in adulthood: the 1982 Pelotas (Brazil) birth cohort study. *Int J Obesity* 2016; **40**: 14–21.
40. Ghosh AK, Hughes AD, Francis D, et al. Midlife blood pressure predicts future diastolic dysfunction independently of blood pressure. *Heart* 2016; **102**: 1380–1387.
41. Sammallahti S, Pyhala R, Lahti M, et al. Infant growth after preterm birth and neurocognitive abilities in young adulthood. *J Pediatr* 2014; **165**: 1109–1115.e3.
42. Hernan MA and Robins JM. Instruments for causal inference: an epidemiologist's dream? *Epidemiology* 2006; **17**: 360–372.
43. Angrist JD and Imbens GW. Two-stage least squares estimation of average causal effects in models with variable treatment intensity. *J Am Stat Assoc* 1995; **90**: 431–442.
44. Wright S. The method of path coefficients. *Ann Math Stat* 1934; **5**: 161–215.
45. Hack M, Weissman B and Borawski-Clark E. Catch-up growth during childhood among very low-birth-weight children. *Arch Pediatr Adolesc Med* 1996; **150**: 1122–1129.
46. Ong KKL, Ahmed ML, Dunger DB, et al. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *Br Med J* 2000; **320**: 967–971.
47. De Stavola BL, Nitsch D, dos Santos Silva I, et al. Statistical issues in life course epidemiology. *Am J Epidemiol* 2005; **163**: 84–96.
48. Textor J and van der Zander B. Dagitty: graphical analysis of structural causal models. *R package version 0.2-2*. 2016.
49. Team RC. *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing, 2013.
50. Robins JM. A new approach to causal inference in mortality studies with a sustained exposure period – application to control of the healthy worker survivor effect. *Math Modell* 1986; **7**: 1393–1512.
51. Robins JM and Hernan MA. Estimation of the causal effects of time-varying exposures. In: Fitzmaurice G, Davidian M and Verbeke G (eds) *Longitudinal data analysis*. Boca Raton: Chapman & Hall/CRC, 2009, pp.553–599.