

POSTER PRESENTATION

Open Access

Critical slowing in a Hodgkin-Huxley neuron near spiking threshold

Alex Bukoski^{1*}, D Alistair Steyn-Ross², Moira L Steyn-Ross²

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012
Decatur, GA, USA. 21-26 July 2012

As noted by Freeman [1], a quiescent neuron approaching spiking threshold exhibits a nonlinearly increasing sensitivity to stimulus. This growth of subthreshold susceptibility can be quantified by applying a linear multivariate Ornstein-Uhlenbeck analysis to the neuron equations, and this has been verified recently [2] for a reduced two-variable spiking model due to Wilson [3].

Here we generalize this stochastic analysis to the classical four-variable conductance-based Hodgkin-Huxley neuron with type-I excitability [4], perturbed by independent white noises entering the drive current and gating variables. We demonstrate critical slowing down—growth in amplitude simultaneous with decay in frequency of soma voltage fluctuations—as the neuron




Figure 1 Subthreshold response to white-noise perturbation as a function of I_{dc} stimulus current. Solid black curves show theoretical ± 3 standard deviation limits for voltage excursions δV away from equilibrium and each vertical gray trace shows actual maximal excursions recorded in 200 ms stochastic simulation runs of the four-variable Hodgkin-Huxley equations at each of 2000 settings for stimulus current ranging from -0.05 to +0.05 nA.

* Correspondence: bukoskia@missouri.edu

¹College of Veterinary Medicine, University of Missouri, Columbia, MO 65203, USA

Full list of author information is available at the end of the article

approaches firing threshold. We show that this behavior is a direct result of the interaction between the model's eigenvalue structure and the noisy environment in which a biological neuron is presumed to function. Stochastic calculus results applied to this four-variable system predict fractional power-law scaling in the divergences for both voltage fluctuations (see Fig. 1) and correlation time as the critical point of saddle-node annihilation is closely approached. Such divergences are expected to be universal characteristics for all type-I neuron models. If these critical fluctuations are communicated to neighboring neurons via ubiquitous electrical gap junctions, then subthreshold neuronal dynamics may play an important role in overall cortical dynamics.

Author details

¹College of Veterinary Medicine, University of Missouri, Columbia, MO 65203, USA. ²School of Engineering, University of Waikato, Hamilton 3240, New Zealand.

Published: 16 July 2012

References

1. Freeman WJ: **Neurodynamics, volume transmission, and self-organization in brain dynamics.** *J Integ Neurosci* 2005, **4**(4):407-421.
2. Steyn-Ross DA, Steyn-Ross ML, Wilson MT, Sleigh JW: **White-noise susceptibility and critical slowing in neurons near spiking threshold.** *Phys Rev E* 2006, **74**:051920.
3. Wilson HR: **Spikes, Decisions, and Actions: The Dynamical Foundations of Neuroscience.** New York: Oxford University Press; 1999.
4. Nowotny T, Rabinovich MI: **Dynamical origin of independent spiking and bursting activity in neural microcircuits.** *Phys Rev Lett* 2007, **98**:128106.

doi:10.1186/1471-2202-13-S1-P34

Cite this article as: Bukoski et al.: Critical slowing in a Hodgkin-Huxley neuron near spiking threshold. *BMC Neuroscience* 2012 **13**(Suppl 1):P34.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

