

RESEARCH

Open Access



Excellent outcomes with Oxford Uni-compartmental knee arthroplasty in anteromedial osteoarthritis patients (≤ 60 years) at mid-term follow-up

Zhen Li¹, Zhenyue Chen², Jinqiang Wei¹, Xianzhong Zeng¹, He Sun³, Zehui Li^{3*} and Xuewei Cao^{3*}

Abstract

Background: The use of Oxford uni-compartmental knee arthroplasty (UKA) has rapidly increased worldwide, however, the relevance of younger patients for postoperative function after Oxford UKA remains unclear. The main purpose of our study is to clarify the effectiveness of Oxford UKA in the younger Chinese patients with anteromedial osteoarthritis (AMOA).

Methods: We retrospectively enrolled 252 consecutive patients who underwent Oxford UKA for AMOA with a minimum follow-up of 5 years between March 2013 and December 2016. The patients were divided into the younger (≤ 60 years) and elderly (> 60 years) age groups. The demographic data and surgery variables were recorded and compared. Patient satisfaction grade, range of motion (ROM), Oxford knee score (OKS), Hospital for Special Surgery (HSS) score, Western Ontario and McMaster (WOMAC) Universities Osteoarthritis Index score and postoperative complications were recorded. The 5-year survival of the implants were also compared with TKA revision as the endpoint.

Results: A total of 252 consecutive patients were recruited, including 96 aged 60 years or less and 156 aged over 60 years. The mean follow-up duration in the younger and elderly groups were 73.6 months (SD, standard deviation, 4.1) and 74.7 months (SD 6.2) respectively. Patient satisfaction rate was high in both groups ($P = 0.805$). Furthermore, no significant differences were observed in postoperative ROM ($P = 0.299$), OKS ($P = 0.117$), HSS ($P = 0.357$) and WOMAC scores ($P = 0.151$) between the younger and elderly groups ($P > 0.05$). However, the incidence of joint stiffness ($P = 0.033$) and delayed wound dehiscence ($P = 0.026$) were significantly different between both groups. Five-year implant survival without revision were also similar in both groups (96.9% vs 97.4%, $P = 0.871$), and that for the entire cohort was 97.2% (95% CI 95.4–99.6).

Conclusion: Oxford UKA for AMOA demonstrated favorable results in younger patients aged ≤ 60 years at a minimum 5-year follow-up in terms of patient satisfaction, functional outcomes, implant survival and postoperative complications. Therefore, younger patients might not be considered as an absolute contraindication to Oxford UKA.

Keywords: Oxford Uni-compartmental knee Arthroplasty, 60 years, Implant survivorship, Mid-term follow-up

Introduction

Oxford uni-compartmental knee arthroplasty (UKA) is a minimally invasive surgery that replaces the surface of the knee joint, and is an ideal treatment for isolated uni-compartmental osteoarthritis (OA). It is increasingly

*Correspondence: cysycaocao@126.com; caoxuewei2021@163.com

³ Department of Orthopaedic Surgery, Guangdong Provincial Hospital of Traditional Chinese Medicine, 111 Dade Road, Yuexiu District, Guangzhou 510120, Guangdong, China

Full list of author information is available at the end of the article



replacing total knee arthroplasty (TKA) on account of less blood loss, shorter operation time, faster postoperative recovery, fewer complications, knee proprioception and bone stock preservation [1–3]. The indications for UKA have also been expanded in recent years [3–6], and the high survival rates of the implants have been demonstrated [7–10].

Based on their experience with fixed-bearing devices, Kozinn and Scott [11] proposed age greater than 60 years, weight less than 180 pounds (82 kg), high activity requirement, patellofemoral joint or lateral compartment degeneration, cartilage calcium deposition, preoperative flexion angle $>90^\circ$, flexion contracture $<5^\circ$, and angular knee deformity $<15^\circ$ as the indications for Oxford UKA. On the other hand, Goodfellow et al. [12, 13] established the indications for Oxford UKA based on the anteromedial osteoarthritis (AMOA) and spontaneous osteonecrosis of the knee (SONK). The ideal indications for AMOA patients are medial compartment OA with bone-on-bone wear, full thickness cartilage in the lateral compartment, and functionally normal anterior cruciate ligament, posterior cruciate ligament and medial collateral ligament. However, studies increasingly show that Oxford UKA can also be performed for younger patients without these classic indications [14–19].

In Caucasian populations, younger patients have demonstrated excellent functional outcomes after Oxford UKA [14, 20]. However, the functional outcomes and implant survival are ambiguous for younger Asian patients with completely different lifestyles and physical activities [20–22]. The aim of this study was to assess the mid-term patient satisfaction, functional outcomes, implant survival and postoperative complications of Oxford UKA in Asian patients aged ≤ 60 years, and to determine the ideal indications for this procedure.

Materials and methods

Participants

A total of 252 consecutive patients with AMOA that underwent medial Oxford UKA surgery were retrospectively recruited between March 2013 and December 2016. Follow-up period referred to the time from the end of surgery to June 2021. All patients had been operated on by one surgical team. The patients were divided into the younger (≤ 60 years) and elderly (> 60 years) age groups. The inclusion criteria were as follows: (1) primary severe AMOA with at least K-L grade three, (2) functionally intact ACL (Anterior Cruciate Ligament) and MCL (Medial Collateral Ligament), (3) correctable varus deformity (varus $< 15^\circ$), (4) flexion deformity $< 15^\circ$ and an active range of motion (at least $0-100^\circ$), (5) intact lateral compartment with no evidence of OA, and (6) minimum 5 years of follow-up and availability of

complete follow-up data. Patients with (1) bone groove-like changes on the lateral side of the patellofemoral joint (Qutebridge grade IV), (2) prior surgery on the same knee, and (3) rheumatoid arthritis or fixed varus deformity were excluded. Obesity, age, high activity or patellofemoral joint degeneration are not considered absolute contraindications. This study was approved by the institutional review committee of our hospital and informed consent was also obtained from all patients.

Surgical procedure and perioperative management

Primary Oxford UKAs were performed using the Oxford Uni-compartmental Phase III mobile-bearing prosthesis (Biomet Merck Ltd., South Wales, UK) by the same group of surgeons. After administering general or spinal anesthesia, the patient was laid in a supine position, and the right lower limb was routinely disinfected. A tourniquet was applied to the proximal thigh on the operative side. Pump up the tourniquet before cutting the skin, and loosen the tourniquet before suturing the surgical incision to stop bleeding. The operative time refers to the time from cutting the skin to closing the wound, including two parts, prosthesis and wound suture. It takes about half an hour to install the prosthesis and half hour to close the wound. A tourniquet is only needed when installing a prosthesis, so the tourniquet time was far shorter than the operative time.

The knee joint was flexed to 90° , and an incision was made from the medial edge of the patella to 3 cm below the joint line. The surgery was performed as per the specifications of the Oxford Microplasty instrumentation. All patients received an analgesia cocktail formulated in our hospital, along with postoperative intravenous medication, acupuncture analgesia etc. The patients were instructed to walk within 2–6 h post-operation, and were given routine anticoagulant and antibiotics the following day. Rehabilitation treatment was started on the first day after surgery with walker-assisted movement, and full weight-bearing exercises were started 3 days later.

Outcome measures

Patients were instructed to go to the outpatient clinic two and 4 weeks after discharge, and then 3, 6, 12 months and once a year postoperatively. Patient records were reviewed and the following data was collected: age, sex, operation side, body mass index (BMI), American society of anesthesiologist (ASA) classification, osteoarthritis (OA) grade and follow-up duration. The preoperative and postoperative values were compared. In addition, the anesthesia type, hospital stay duration, blood loss volume, tourniquet time and operative time were also recorded. At the final follow-up, patient satisfaction was documented as disappointed, dissatisfied, satisfied or very

satisfied. Preoperative and postoperative range of motion (ROM), the Oxford knee score (OKS), hospital for special surgery (HSS) knee score and Western Ontario and McMaster (WOMAC) University osteoarthritis index score were used to evaluate the function of the knee joint at final follow-up. Postoperative complications such as patellar ligament injury, deep vein thrombosis, joint stiffness, wound infection, delayed wound dehiscence, radiographic lucency and revision rate were recorded and compared. The details of knee revision for TKA and the 5-year survivorship of Oxford UKA implants were also compared.

Postoperative radiographs of both groups were reviewed to determine the differences in implants. In the follow-up period, we analyzed the x-ray of each patient to find signs of complications, such as prosthesis position, size and dislocation, pathological radiolucent lines or lateral compartment OA progression. It is worth noting that the anterior radiograph should directly face the tibial prosthesis, and the lateral radiograph should directly face the femoral prosthesis. Only in this way can we correctly compare the postoperative complications between the two groups from the x-ray.

Statistical analysis

All statistical analysis was performed using GraphPad Prism 7.0 (GraphPad Software, San Diego, USA) and SPSS 23.0 (SPSS Inc., Chicago, IL, USA). Continuous variables were reported as mean \pm standard deviation (SD). The inter-group differences of variables conforming

to normal distribution were compared by independent sample *t*-tests. Categorical variables (demographic data, anesthesia type, satisfaction grade and postoperative complications) were described using frequency distributions and percentages, and compared by Chi-square test or Fisher's exact test. Log-rank test was used to compare the Kaplan-Meier survival curves with revision to total knee arthroplasty (TKA) as the end point. All tests were two-sided and *P*-value < 0.05 was considered statistically significant.

Results

Two hundred and fifty-two consecutive patients with a minimum 5-year follow-up were enrolled in this study, of which 96 were aged 60 years or younger, and 156 were over 60 years. The mean follow-up in the younger and elderly groups was 73.6 months (SD4.1) and 74.7 months (SD6.2) respectively. The demographic data of both groups are summarized in Table 1 and the clinical variables are listed in Table 2. There were no significant differences between the two age groups in terms of anesthesia type, hospital stay volume, blood loss volume, tourniquet time or operative time ($P < 0.05$). In addition, patients in both groups reported high satisfaction ($P = 0.805$), and the overall patient satisfaction rate was almost 93% (Fig. 1). Oxford UKA significantly improved the functional outcomes at follow-up compared to preoperative scores, and the postoperative ROM ($P = 0.299$), OKS ($P = 0.117$), HSS ($P = 0.357$) and WOMAC scores ($P = 0.151$) were similar in both groups

Table 1 Patients' demographic data between the two groups


Variables	Younger group (≤ 60 years)	Elderly group (> 60 years)	F value/ χ^2	P value
Knees	96	156	–	–
Age (years)	55.6 \pm 4.0	69.0 \pm 5.9	–19.70	< 0.001
Sex			0.231	0.631
female	68 (70.8%)	106 (67.9%)		
male	28 (29.2%)	50 (32.1%)		
Operation Side			0.632	0.427
right	56 (58.3%)	83 (53.2%)		
left	40 (41.7%)	73 (46.8%)		
BMI (Kg/m ²)	26.9 \pm 1.6	27.2 \pm 2.7	–1.413	0.159
ASA classification			0.718	0.397
1 to 2	81 (84.4%)	125 (80.1%)		
3 to 4	15 (15.6%)	31 (19.9%)		
OA grade in MC (KL)			1.462	0.227
III	36 (37.5%)	47 (30.1%)		
IV	60 (62.5%)	109 (69.9%)		
Follow-up (months)	73.6 \pm 4.1	74.7 \pm 6.2	–1.521	0.130

Values are expressed as mean \pm SD (standard deviation) or number (percentage); BMI body mass index, ASA American society of anesthesiologist, OA osteoarthritis, MC medial compartment, KL Kellgren-Lawrence score

Table 2 Comparisons of preoperative and postoperative variables between the two groups

Variables	Younger group(n = 96)	Elderly group(n = 156)	F value/ χ^2	P value
Anaesthesia type			0.321	0.571
general	76(79.2%)	128(82.1%)		
spinal	20(20.8%)	28(17.9%)		
Hospital stay(d)	9.7 ± 1.9	10.2 ± 2.2	-1.73	0.086
Blood loss volume(ml)	82.6 ± 26.4	81.7 ± 23.3	0.258	0.796
Tourniquet time(min)	32.4 ± 3.0	33.1 ± 4.1	-1.569	0.118
Operative time(min)	82.0 ± 5.1	82.8 ± 4.7	-1.374	0.171

Values are expressed as mean ± SD or number (percentage)



($P > 0.05$) (Fig. 2). For the elderly patients, the OKS, HSS and WOMAC scores improved by an average of 16.1, 26.3 and 19.8 points relative to preoperative values, and the mean improvement in these scores among younger patients were 16.3, 26.2 and 20.0 respectively (Table 3).

There were no significant differences in the incidence rates of postoperative complications such as patellar ligament injury (1% vs 0%, $P = 0.381$), deep vein thrombosis (2.1% vs 1.9%, $P = 1.000$), wound infection (1% vs 1.3%, $P = 1.000$), radiographic lucency (4.2% vs 1.9%, $P = 0.432$) and revision rate (4.2% vs 3.8%, $P = 1.000$) between the younger and elderly groups (Table 4). However, the significant differences in terms of joint stiffness ($P = 0.033$) and delayed wound dehiscence ($P = 0.026$) were found in both groups (Table 4). As shown in Table 5, TKA was performed for 10 knees due to progression of lateral OA (5 cases, 50%), persistent unexplained pain (3 cases, 30%) or aseptic loosening (2 cases, 20%). Furthermore, seven knees required posterior-stabilized primary TKA implants, three knees required tibial stem and medial tibial augmentation was performed for one knee. Regardless of the reason, the mean time to TKA [47 (20–73) months vs 49.2 (21–74) months; ($P = 0.878$)] and 5-year survival rate of the implant [96.9% (95% CI 95.1–99) vs 97.4% (95% CI 95.6–99.4); ($P = 0.871$)] were similar in the younger and elderly age groups (Fig. 3). Furthermore, the 5-year implant survival rate was 97.2% (95% CI




Table 3 Comparisons of patient satisfaction and functional outcomes scores between the two groups

Parameter	Younger group (n = 96)	Elderly group (n = 156)	Mean difference	95% CI of mean diff	F value/ χ^2	P value
Satisfaction grade					0.061	0.805
Satisfied-very satisfied	90(93.8%)	145(92.9%)				
Disappointed-dissatisfied	6(6.2%)	11(7.1%)				
ROM						
Preoperative	110.1 ± 3.4	109.2 ± 4.1	0.85	−0.12 to 1.83	1.720	0.087
Postoperative	122.2 ± 3.8	121.7 ± 3.1	0.48	−0.43 to 1.39	1.043	0.299
OXS						
Preoperative	24.3 ± 2.9	23.8 ± 3.7	0.44	−0.38 to 1.27	1.056	0.292
Postoperative	40.6 ± 3.3	39.9 ± 4.0	0.72	−0.18 to 1.62	1.571	0.117
HSS						
Preoperative	60.3 ± 3.7	59.7 ± 3.8	0.63	−0.33 to 1.58	1.296	0.196
Postoperative	86.5 ± 4.3	86.0 ± 3.5	0.48	−0.55 to 1.51	0.923	0.357
WOMAC						
Preoperative	45.5 ± 3.5	46.2 ± 4.3	−0.61	−1.64 to 0.42	−1.173	0.242
Postoperative	25.5 ± 4.8	26.4 ± 3.7	−0.83	−1.96 to 0.31	−1.443	0.151

Values are expressed as mean ± SD or number (percentage); Diff difference, OXS Oxford knee score, HSS hospital for surgery score, ROM range of motion

Table 4 Postoperative complications after Oxford UKA

Variables	Younger group(n = 96)	Elderly group(n = 156)	P value
Patellar ligament injury	1(1.0%)	0(0%)	0.381
Deep vein thrombosis	2(2.1%)	3(1.9%)	1.000
Joint stiffness	1(1.0%)	11(7.1%)	0.033
Wound infection	1(1.0%)	2(1.3%)	1.000
Delayed wound dehiscence	0(0%)	8(5.1%)	0.026
Radiographic lucency	4(4.2%)	3(1.9%)	0.432
Revision for any reason	4(4.2%)	6(3.8%)	1.000

Values are all expressed as number (percentage); UKA unicompartmental knee arthroplasty

95.4–99.6) for the entire cohort (Fig. 4). Comparison of the radiographs of a 54-years-old patient and 69-years-old patient indicated similar clinical outcomes at the 5-year follow-up (Fig. 5).

Discussion

The most significant finding of our study was that patient satisfaction, functional outcomes, implant survival and postoperative complications were similar in the younger and elderly OA patients at a minimum 5-year follow-up after Oxford UKA. Therefore, younger patients might not be considered as an absolute contraindication to Oxford UKA.

Table 5 Details of the ten Knees Revised to TKA

Cases	Time to Revision(m)	age	Reason for Revision	TKA
1	20	58	Persistent unexplained pain	Primary PS TKA,no stems or augments
2	21	67	Persistent unexplained pain	Primary PS TKA,no stems or augments
3	35	75	Progression of lateral OA	Primary PS TKA,no stems or augments
4	45	59	Persistent unexplained pain	Primary PS TKA,no stems or augments
5	45	70	Progression of lateral OA	Primary PS TKA,no stems or augments
6	50	55	Aseptic loosening	PS TKA,tibial stem,no augments
7	55	67	Aseptic loosening	PS TKA,tibial stem,no augments
8	65	74	Progression of lateral OA	Primary PS TKA,no stems or augments
9	73	60	Progression of lateral OA	PS TKA,tibial stem and medial tibial augments
10	74	72	Progression of lateral OA	Primary PS TKA,no stems or augments

OA osteoarthritis, PS posterior stabilized, TKA total knee arthroplasty




Fig. 3 Kaplan-Meier survival curves with 95% confidence intervals in the younger and older patients with revision to TKA as the end point




Fig. 4 Kaplan-Meier survival curves 95% confidence intervals for all 252 consecutive patients with revision to TKA as the end point

Patient satisfaction

The overall patient satisfaction rate in our cohort was 93%, and was similar in both age groups. Several studies have reported high satisfaction in younger patients after UKA [15, 17]. Dalury et al. [16] reported satisfactory results for patients aged 46–59 years at mid-term follow-up. Mannan et al. [17] found that 97% of the patients aged 60 years or less were pleased or very pleased after UKA at minimum 10-year follow-up.

Functional outcomes

Initial studies have shown that ideal candidates for UKA are older than 60 years of age [11]. However, we observed significant improvement and satisfactory functional outcomes in both age groups. Mannan et al. [17] reviewed a consecutive series of 91 medial UKAs performed in patients aged 60 years or younger at minimum 10-year follow-up, and demonstrated good-to-excellent

long-term patient outcomes with 92.9 and 87.8% implant survival at 10 years and 15 years respectively. Lee et al. [18] also reported similar post-UKA outcome scores and implant survival in patients younger or older than 55 years at minimum 10 years follow-up. Hamilton et al. [23] evaluated 1000 Oxford UKAs and did not find any significant differences in the functional outcomes of patients aged younger and older than 60 years of age at 10-year follow-up. Walker et al. [19] reviewed 118 knees following medial mobile bearing UKA in patients aged 60 or younger and showed that the patients were able to resume normal physical activities, and nearly two-thirds could engage in high-level activities. Thompson et al. [5] reported that patients younger than 60 years appeared to have better results at 2-year follow-up, and Kleebblad et al. [24] showed that patients younger than 65 years were satisfied with the outcomes of UKA and TKA. Therefore, younger age is not a contraindication for UKA.

Given that squatting and sitting on the floor with high flexion angles are common daily activities in the Asian population, their post-operative functional requirements are greater compared to that of Western populations. Yoshida et al. [25] found that the 10-year implant survival rate in 1279 Japanese patients who underwent Oxford UKAs was 95%. Lim et al. [26] likewise reported a 94% 10-year implant survival rate in Korean patients. Xue et al. [27] assessed 708 consecutive medial Oxford UKAs in Chinese patients and found that the 5-year cumulative survival rate was 98.8% and the 10-year survival rate was 94.3%. The clinical outcomes of the patients in our cohort were similar to that of Western patients, and satisfactory ROM was achieved for daily activities.

Implant survival

A recent review of twenty-six studies on 42,791 knees reported 95.3 and 91.3% 5- and 10-year pooled survival rates respectively after medial UKA [7]. Mohammad et al. [8] reported a 10-year survival of 93% and 15-year survival of 89% for 15 studies involving 8658 knees. Walker et al. [10] reported that medial Oxford UKA ensured a survival rate of 92.4% at 10 years and 88.6% at 15 years with an excellent functional outcome. Xue et al. [27] also reported 98 and 94.3% 5- and 10-year cumulative survival rates in Chinese patients. Several studies have also confirmed that patient age is not a determinant of implant survival, which is consistent with the 96.9 and 97.4% survival rates observed for the younger and elderly patients respectively at the 5-year follow-up. The Oxford mobile-bearing implant was used for all patients, indicating that it is a suitable choice for UKA.

The survival rate of UKA implant is usually evaluated with TKA revision as the endpoint. The major reasons for the failure of Oxford UKA are weight-bearing




Fig. 5 Radiographs of (A–D) a 54-year-old female patient and (E–H) a 69-year-old male patient with isolated compartment OA who underwent Oxford UKA and showed excellent outcomes at 5-year follow-up

dislocation, aseptic loosening, lateral compartment osteoarthritis progression, and pain with no identifiable etiology. Xue et al. [27] reported that the most common reason of revision in Chinese patients were dislocation and osteoarthritis of the lateral compartment. Liddle et al. [28] reported that aseptic loosening was the most common reason for revision after UKA, and progression of arthritis and weight-bearing dislocation were almost exclusive to UKA. Other studies have implicated lateral compartment osteoarthritis progression and pain as the most frequent causes of revision [8, 9]. van der List et al. [29] found that aseptic loosening lead early failure of mobile-bearing UKA, whereas OA progression lead to the failure of fixed-bearing UKA in later years. Crawford et al. [6] also reported aseptic loosening as the most common early indication for revision of UKA, and arthritic progression as the most common indication in middle and late stages. In our study, progression of lateral OA, persistent unexplained pain and aseptic loosening were the most frequent causes of

implant failure and revision.

Ideal indications

The outcome of surgery and revision rate is directly determined by the surgeon's experience and proficiency [30, 31]. In a review of 37,131 UKAs conducted in England and Wales, Liddle et al. [32] found that surgical caseload was a determining factor of implant survival, and the revision rate between surgeons with the lowest

and highest-caseload had a four-fold difference. Kozinn and Scott [11] recommended strict inclusion criteria for UKA, whereas Goodfellow et al. [12, 13] recommended Oxford UKA for AMOA and SONK. In fact, with the increasing range of applications and excellent functional outcomes, the traditional indications of UKA have been expanded in recent years [3–6]. Thompson et al. [5] optimized the indications proposed by Kozinn and Scott based on the findings that younger patients had better clinical outcomes, and obese patients had a low revision rate at short or medium-term follow-up. In a large case series, Hamilton et al. [23] reported more indications based on the occurrence and development stage of the disease as proposed by Goodfellow et al. [12] instead of the contraindications set forth by Kozinn and Scott.

The aim of this study was to discuss the mid-term patient satisfaction, functional outcomes, implant survival and postoperative complications between younger and elderly groups. On the one hand, there were no obvious differences between the two groups in terms of patient satisfaction, functional outcomes and implant survival, which indicated that the indications for UKA have been expanded. There are many reasons for the expansion of UKA's traditional indications. Among them, the surgeon's experience and proficiency are crucial, and the development of prostheses and devices also play an important role. With the advancement of surgical technology, the surgical outcomes will be better, and the complications will be less than before. On the other

hand, there were obvious differences in terms of postoperative complications including joint stiffness and delayed wound dehiscence, which may be related to the different age attributes of the two groups. The younger age group is presumed to have higher physical activity and the elderly age group is presumed to have low immunity.

Conclusion

Oxford UKA for AMOA demonstrated favorable results in younger patients aged ≤ 60 years at a minimum 5-year follow-up. Patient satisfaction, functional outcomes, implant survivorship or postoperative complications were largely similar in both age groups. Therefore, younger patients might not be considered as an absolute contraindication to Oxford UKA.

Limitations

The sample size of the study was relatively small, and the younger patients accounted for only 38% of the total cases. Secondly, the relative mid-term follow-up could not fully explain the exact difference between the two groups. Thirdly, we did not consider that different age groups have different attributes, such as physical activity and immunity, which may bias the study results. Further studies on larger cohorts with samples randomization and long-term follow-up are required to improve the difference between age groups.

Abbreviations

UKA: Uni-compartmental knee arthroplasty; AMOA: Anteromedial osteoarthritis; SONK: Spontaneous osteonecrosis of the knee; ACL: Anterior Cruciate Ligament; MCL: Medial Collateral Ligament; ROM: Range of motion; OKS: Oxford knee score; HSS: Hospital for Special Surgery score; WOMAC: Western Ontario and McMaster Universities Osteoarthritis Index score; SD: Standard deviation; BMI: Body mass index; ASA: American society of anesthesiologist; OA: Osteoarthritis; MC: Medial compartment; KL: Kellgren-Lawrence score; PS: Posterior stabilized; TKA: Total knee arthroplasty.

Acknowledgements

We express deep thanks to Dr. Xuewei Cao and Zehui Li for their technical support.

Authors' contributions

Zhen Li and Zhenyue Chen conceived and designed the experiments. He Sun, Zehui Li and Xuewei Cao performed the surgeries. Jinqiang Wei and Xianzhong Zeng analyzed the data and prepared the figures. Zhen Li wrote the main manuscript text. All authors read and approved the final manuscript.

Funding

This study was supported by a grant from the Traditional Chinese Medicine Talent Training Project of Guangdong provincial Bureau of Traditional Chinese Medicine (NO.0103030908) and Guangdong Provincial Hospital of Traditional Chinese Medicine and the School of Biomedicine, Chinese University of Hong Kong School of Medicine, Basic Clinical Collaborative Innovation Project (NO. YN2018HK04).

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study. The consent and study plan was approved by the Ethics Committee of Guangdong Provincial Hospital of Traditional Chinese Medicine (BF2021-004).

Consent for publication

We obtained permission from the participants to publish their data. All participants gave written consent for publication.

Competing interests

The authors have no conflicts of interest to declare.

Author details

¹The Second Clinical Medical College of Guangzhou University of Chinese Medicine, Guangzhou 510405, Guangdong, China. ²The First Clinical Medical College of Guangzhou University of Chinese Medicine, Guangzhou 510405, Guangdong, China. ³Department of Orthopaedic Surgery, Guangdong Provincial Hospital of Traditional Chinese Medicine, 111 Dade Road, Yuexiu District, Guangzhou 510120, Guangdong, China.

Received: 13 August 2021 Accepted: 28 September 2021

Published online: 08 October 2021

References

- Uzun E, Misir A, Kizkapan TB, Ozcamdalli M, Gunay AE, Husrevoglu K. Mid-term functional, clinical, and radiological outcomes with factors affecting revision of mobile-bearing medial unicompartmental knee arthroplasty. *Knee*. 2020;27(2):527–34.
- Tu Y, Ma T, Wen T, Yang T, Xue L, Xue H. Does Unicompartmental knee replacement offer improved clinical advantages over Total knee replacement in the treatment of isolated lateral osteoarthritis? A matched cohort analysis from an independent center. *J Arthroplast*. 2020;35(8):2016–21.
- Murray DW, Parkinson RW. Usage of unicompartmental knee arthroplasty. *Bone Joint J*. 2018;100-B(4):432–5.
- Pandit H, Jenkins C, Gill HS, et al. Unnecessary contraindications for mobile-bearing unicompartmental knee replacement. *J Bone Joint Surg Br*. 2011;93(5):622–8.
- Thompson SA, Liabaud B, Nellans KW, Geller JA. Factors associated with poor outcomes following unicompartmental knee arthroplasty: redefining the "classic" indications for surgery. *J Arthroplast*. 2013;28(9):1561–4.
- Crawford DA, Berend KR, Thienpont E. Unicompartmental knee Arthroplasty: US and global perspectives. *Orthop Clin North Am*. 2020;51(2):147–59.
- Heaps BM, Blevins JL, Chiu YF, Konopka JF, Patel SP, McLawhorn AS. Improving estimates of annual survival rates for medial Unicompartmental knee Arthroplasty, a Meta-analysis. *J Arthroplast*. 2019;34(7):1538–45.
- Mohammad HR, Strickland L, Hamilton TW, Murray DW. Long-term outcomes of over 8,000 medial Oxford phase 3 Unicompartmental knees—a systematic review. *Acta Orthop*. 2018;89(1):101–7.
- Neufeld ME, Albers A, Greidanus NV, Garbus DS, Masri BA. A comparison of Mobile and fixed-bearing Unicompartmental knee Arthroplasty at a minimum 10-year follow-up. *J Arthroplast*. 2018;33(6):1713–8.
- Walker T, Hetto P, Bruckner T, et al. Minimally invasive Oxford unicompartmental knee arthroplasty ensures excellent functional outcome and high survivorship in the long term. *Knee Surg Sports Traumatol Arthrosc*. 2019;27(5):1658–64.
- Kozinn SC, Marx C, Scott RD. Unicompartmental knee arthroplasty. A 4.5–6-year follow-up study with a metal-backed tibial component. *J Arthroplast*. 1989;4(Suppl):S1–S10.
- Goodfellow JW, Kershaw CJ, Benson MK, O'Connor JJ. The Oxford knee for unicompartmental osteoarthritis. The first 103 cases. *J Bone Joint Surg Br*. 1988;70(5):692–701.

13. Murray DW, Goodfellow JW, O'Connor JJ. The Oxford medial unicompartmental arthroplasty: a ten-year survival study. *J Bone Joint Surg Br.* 1998;80(6):983–9.
14. Pennington DW, Swienckowski JJ, Lutes WB, Drake GN. Unicompartmental knee arthroplasty in patients sixty years of age or younger. *J Bone Joint Surg Am.* 2003;85(10):1968–73.
15. Biswas D, Van Thiel GS, Wetters NG, Pack BJ, Berger RA, Della Valle CJ. Medial unicompartmental knee arthroplasty in patients less than 55 years old: minimum of two years of follow-up. *J Arthroplast.* 2014;29(1):101–5.
16. Dalury D, Kelley TC, Adams MJ. Medial UKA: favorable mid-term results in middle-aged patients. *J Knee Surg.* 2013;26(2):133–7.
17. Mannan A, Pilling RWD, Mason K, Stirling P, Duffy D, London N. Excellent survival and outcomes with fixed-bearing medial UKA in young patients (≤ 60 years) at minimum 10-year follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(12):3865–70.
18. Lee M, Chen J, Shi Lu C, Lo NN, Yeo SJ. No differences in outcomes scores or survivorship of Unicompartmental knee Arthroplasty between patients younger or older than 55 years of age at minimum 10-year Follow-up. *Clin Orthop Relat Res.* 2019;477(6):1434–46.
19. Walker T, Streit J, Gotterbarm T, Bruckner T, Merle C, Streit MR. Sports, physical activity and patient-reported outcomes after medial Unicompartmental knee Arthroplasty in young patients. *J Arthroplast.* 2015;30(11):1911–6.
20. Kim SJ, Postigo R, Koo S, Kim JH. Causes of revision following Oxford phase 3 unicompartmental knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2014;22(8):1895–901.
21. Choy WS, Lee KW, Kim HY, Kim KJ, Chun YS, Yang DS. Mobile bearing medial unicompartmental knee arthroplasty in patients whose lifestyles involve high degrees of knee flexion: a 10-14year follow-up study. *Knee.* 2017;24(4):829–36.
22. Cheng J, Feng M, Cao G, Li Z, An S, Lu S. Patient outcomes in Antero-medial osteoarthritis patients over 80 years old undergoing Oxford Unicompartmental knee Arthroplasty in China. *BMC Musculoskelet Disord.* 2020;21(1):446 Published 2020 Jul 8.
23. Hamilton TW, Pandit HG, Jenkins C, Mellon SJ, Dodd CAF, Murray DW. Evidence-based indications for Mobile-bearing Unicompartmental knee Arthroplasty in a consecutive cohort of thousand knees. *J Arthroplast.* 2017;32(6):1779–85.
24. Kleefeld LJ, van der List JP, Zuiderbaan HA, Pearle AD. Larger range of motion and increased return to activity, but higher revision rates following unicompartmental versus total knee arthroplasty in patients under 65: a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(6):1811–22.
25. Yoshida K, Tada M, Yoshida H, Takei S, Fukuoka S, Nakamura H. Oxford phase 3 unicompartmental knee arthroplasty in Japan—clinical results in greater than one thousand cases over ten years. *J Arthroplast.* 2013;28(9 Suppl):168–71.
26. Lim HC, Bae JH, Song SH, Kim SJ. Oxford phase 3 unicompartmental knee replacement in Korean patients. *J Bone Joint Surg Br.* 2012;94(8):1071–6.
27. Xue H, Tu Y, Ma T, Wen T, Yang T, Cai M. Up to twelve year follow-up of the Oxford phase three unicompartmental knee replacement in China: seven hundred and eight knees from an independent Centre. *Int Orthop.* 2017;41(8):1571–7.
28. Liddle AD, Judge A, Pandit H, Murray DW. Adverse outcomes after total and unicompartmental knee replacement in 101,330 matched patients: a study of data from the National Joint Registry for England and Wales [published correction appears in *lancet.* 2015 Feb 28;385(9970):774]. *Lancet.* 2014;384(9952):1437–45.
29. van der List JP, Zuiderbaan HA, Pearle AD. Why do medial Unicompartmental knee Arthroplasties fail today? *J Arthroplast.* 2016;31(5):1016–21.
30. Rajasekhar C, Das S, Smith A. Unicompartmental knee arthroplasty. 2- to 12-year results in a community hospital. *J Bone Joint Surg Br.* 2004;86(7):983–5.
31. Robertsson O, Knutson K, Lewold S, Lidgren L. The routine of surgical management reduces failure after unicompartmental knee arthroplasty. *J Bone Joint Surg Br.* 2001;83(1):45–9.
32. Liddle AD, Pandit H, Judge A, Murray DW. Effect of surgical caseload on revision rate following Total and Unicompartmental knee replacement. *J Bone Joint Surg Am.* 2016;98(1):1–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

