

RESEARCH ARTICLE

Estradiol-induced immune suppression via prostaglandin E₂ during parturition in bovine leukemia virus-infected cattle

Yamato Sajiki¹, Satoru Konnai^{1,2*}, Tomohiro Okagawa², Naoya Maekawa², Shinya Goto¹, Junko Kohara³, Atsushi Nitana⁴, Hirofumi Takahashi⁴, Kentaro Kubota⁵, Hiroshi Takeda⁶, Shiro Murata^{1,2}, Kazuhiko Ohashi^{1,2}

1 Department of Disease Control, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Japan, **2** Department of Advanced Pharmaceutics, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Japan, **3** Animal Research Center, Agriculture Research Department, Hokkaido Research Organization, Shintoku, Japan, **4** Hokkaido Higashi Agricultural Mutual Aid Association, Nakashibetsu, Japan, **5** Rumoi Nambu Veterinary Clinic, Hokkaido Chuo Agricultural Mutual Aid Association, Tomamae, Japan, **6** Osyamanbe Veterinary Clinic, Minami Hokkaido Agricultural Mutual Aid Association, Osyamanbe, Japan

* konnai@vetmed.hokudai.ac.jp



OPEN ACCESS

Citation: Sajiki Y, Konnai S, Okagawa T, Maekawa N, Goto S, Kohara J, et al. (2022) Estradiol-induced immune suppression via prostaglandin E₂ during parturition in bovine leukemia virus-infected cattle. PLoS ONE 17(3): e0263660. <https://doi.org/10.1371/journal.pone.0263660>

Editor: Juan J. Loor, University of Illinois, UNITED STATES

Received: May 23, 2021

Accepted: January 21, 2022

Published: March 9, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0263660>

Copyright: © 2022 Sajiki et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Abstract

Immune suppression during pregnancy and parturition is considered a risk factor that is related to the progression of bovine chronic diseases, such as bovine leukosis, which is caused by bovine leukemia virus (BLV). Our previous studies have demonstrated that prostaglandin E₂ (PGE₂) suppresses BLV-specific Th1 responses and contributes to the disease progression during BLV infection. Although PGE₂ reportedly plays important roles in the induction of parturition, PGE₂ involvement in immune suppression during parturition is unknown. To investigate its involvement, we analyzed PGE₂ kinetics and Th1 responses in BLV-infected pregnant cattle. PGE₂ concentrations in sera were increased, whereas IFN-γ responses were decreased before delivery. PGE₂ is known to suppress Th1 immune responses in cattle. Thus, these data suggest that PGE₂ upregulation inhibits Th1 responses during parturition. We also found that estradiol was important for PGE₂ induction in pregnant cattle. *In vitro* analyses indicated that estradiol suppressed IFN-γ production, at least in part, via PGE₂/EP4 signaling. *In vivo* analyses showed that estradiol administration significantly influenced the induction of PGE₂ production and impaired Th1 responses. Our data suggest that estradiol-induced PGE₂ is involved in the suppression of Th1 responses during pregnancy and parturition in cattle, which could contribute to the progression of BLV infection.

Introduction

In dairy cattle, efficient milk production continues to require for pregnancy and 3 parturition each year. During pregnancy and parturition, maternal immune responses are suppressed to contribute to fetal survival, which is also known as maternal tolerance of the fetus [1, 2]. In general, Th1/Th2 imbalance and regulatory T cell (Treg) induction are known to be involved

Funding: This work was supported by JSPS KAKENHI grant number 19KK0172 [to S.K.], grants from the Project of the NARO, Bio-oriented Technology Research Advancement Institution (Research Program on Development of Innovative Technology 26058 BC [to S.K.] and Special Scheme Project on Regional Developing Strategy, Grant 16817557 [to S.K.]), Regulatory research projects for food safety, animal health and plant protection (JPJ008617.17935709) funded by the Ministry of Agriculture, Forestry and Fisheries of Japan and Clinical Research Promotion Fund 2021 by Hokkaido University Veterinary Teaching Hospital[to S.K.]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

in maternal tolerance [3, 4]. Sex hormones, such as estradiol and progesterone, play important roles in the inhibition of Th1 responses and induction of Tregs [5, 6]. Previous report has shown that sex hormones contribute to the suppression of immune responses in pregnant cattle [7]. Although immune suppression in pregnant animals is essential for the prevention of fetal injury by the maternal immune system, this suppression may affect the progression and onset of bovine chronic diseases, such as bovine leukosis. Bovine leukemia virus (BLV) is a member of the genus *Deltaretrovirus*, and causes bovine leukosis after a long latent period [8]. Most BLV-infected cattle remain subclinical and are referred to as aleukemic. However, approximately 30% of BLV-infected cattle develop persistent lymphocytosis, which is characterized by the nonmalignant polyclonal expansion of infected B cells in peripheral blood. Less than 5% of BLV-infected cattle develop bovine leukosis, which is called enzootic bovine leukosis (EBL) and is characterized by fatal lymphoma or lymphosarcoma [9]. The suppression of the immune system, particularly in a stressful situation such as parturition, is considered a risk factor for the development of bovine leukosis [10, 11]. Although parturition is an important event for progression to the clinical stages of BLV infection, the relationship between parturition and the progression of these diseases has not been fully elucidated.

Prostaglandin E₂ (PGE₂) is an inflammatory mediator that is metabolically derived from arachidonic acid by cyclooxygenase enzymes (COX-1 and COX-2) [12]. COX-1 is a constitutive enzyme that is expressed in many tissues including the kidneys, stomach, and vascular endothelium, and it is involved in a multitude of physiological processes [13]. Conversely, COX-2 is an inducible enzyme that is regulated by inflammatory cytokines, antigen-stimulation, and growth factors via the activation of nuclear factor-kappa B (NF-κB) [14]. PGE₂ inhibits the activities of immune cells such as T cells, dendritic cells, and natural killer cells via prostaglandin E receptor (EP) 2 and EP4, thereby contributing to immune evasion during chronic infections [15]. Our previous studies have demonstrated that PGE₂ inhibits Th1 responses in cattle, and is associated with the progression of BLV infection [16, 17]. In addition, PGE₂, as well as PGF_{2α}, is an important factor related to the induction of parturition in cattle [18]. However, little information is available regarding the immune dysfunctional effect of PGE₂ in periparturient period.

Here, to investigate whether PGE₂ is involved in the suppression of Th1 immune responses in the periparturient period, PGE₂ kinetics and Th1 cytokine production were analyzed by using BLV-infected pregnant cattle. Additionally, to examine whether PGE₂-induced suppression of Th1 response is mediated by estradiol, the effects of estradiol on PGE₂ production and Th1 responses was analyzed by using BLV-infected and uninfected cattle.

Materials and methods

Animals and blood collection

The blood samples of BLV-infected and BLV-uninfected cattle (adult female, Holstein breed), which were used in this study, were obtained from several farmers and veterinarians in Japan. BLV infection was confirmed by detection of the provirus using nested polymerase chain reaction (PCR) that targets the viral LTR and by detection of the anti-BLV antibody using a commercial ELISA kit (JNC, Tokyo, Japan), as described previously [19]. The blood samples of BLV-infected pregnant cattle (animals #1–#5, Table 1) were collected at two points before delivery, on the day of delivery (day 0), and on three points after delivery (days 1, 7, and 14). Blood collection on day 0 was performed before parturition. These animal experiments were approved by the Ethics Committee of the Faculty of Veterinary Medicine, Hokkaido University. Verbal informed consent was obtained from the owners for the participation of their animals in this study.

Table 1. BLV-infected cattle used in this study.

Cattle	#1	#2	#3	#4
Age	98 months old	41 months old	56 months old	51 months old
Breed	Holstein	Holstein	Holstein	Holstein
Sex	female	female	female	female
BLV infection	+	+	+	+
Pregnancy	+	+	+	+
Cattle	#5	#6	#7	
Age	52 months old	98 months old	75 months old	
Breed	Holstein	Holstein	Holstein	
Sex	female	female	female	
BLV infection	+	+	+	
Pregnancy	+	-	-	
Cattle	#8	#9	#10	
Age	85 months old	106 months old	152 months old	
Breed	Holstein	Holstein	Holstein	
Sex	female	female	female	
BLV infection	+	+	+	
Pregnancy	-	-	-	

<https://doi.org/10.1371/journal.pone.0263660.t001>

For estradiol administration to BLV-infected cattle, 5 BLV-infected cattle (animals #6–#10, [Table 1](#)) were kept in an animal facility at the Faculty of Veterinary Medicine, Hokkaido University (Sapporo, Hokkaido, Japan, animals #6 and #7), and a biosafety level I animal facility at the Animal Research Center, Agricultural Research Department, Hokkaido Research Organization (Shintoku, Hokkaido, Japan, animals #8–#10). These animal experiments were approved by the Ethics Committee of the Faculty of Veterinary Medicine, Hokkaido University, and the Ethics Committee of the Animal Research Center, Agricultural Research Department, Hokkaido Research Organization.

Blood samples were shipped immediately after collection and stored at 4°C until the experiment. After 24 hours from collection, blood samples were used for the following analyses.

Specimens of cattle with bovine leukosis

In this study, we collected clinical data of 68 cattle which were diagnosed with bovine leukosis in a specific area in Hokkaido, Japan from 2016 to 2018, and analyzed the time intervals between parturition and the onset of bovine leukosis. Animals with lymphoma were clinically diagnosed as onset. The diagnosis was carried out by clinical veterinarians in Livestock Hygiene Service Center, Meat Inspection Center or Agricultural Mutual Aid Association in Hokkaido, Japan.

Preparation of peripheral blood mononuclear cells (PBMCs)

Buffy coat fraction was separated from blood samples by centrifugation (1,700 × g, 15 min, 25°C, without break). Next, PBMCs were purified from the buffy coat fraction by density gradient centrifugation (1,200 × g, 20 min, 25°C, without break) on 60% Percoll (GE Healthcare, Little Chalfont, UK). Then, collected PBMCs were washed 3 times by centrifugation (770 × g, 10 min, 25°C) in phosphate buffered saline and filtered through a 40-µm cell strainer (BD Biosciences, San Jose, CA, USA). Then, PBMCs were stained with 0.4% Trypan Blue Stain

(Thermo Fisher Scientific, Waltham, MA, USA) and the number of the viable cells was counted using Countess II FL Automated Cell Counter (Thermo Fisher Scientific).

Whole-blood culture

Whole blood cells (500 μ L of blood) were incubated with 20 μ g/ml concanavalin A (Con A, Sigma-Aldrich, St. Louis, MO, USA) at 37°C under 5% CO₂ for 24 hours using 48-well plates (Corning Inc., Corning, NY, USA). Three culture wells were prepared in each sample. Collected culture supernatants were assayed for IFN- γ by Bovine IFN- γ ELISA Development Kit (Mabtech, Nacka Strand, Sweden) according to the manufacturer's instructions.

Real-time PCR

Total RNA of PBMCs was extracted using TRI Reagent (Molecular Research Center, Cincinnati, OH, USA) in accordance with the manufacturer's instructions. cDNA was synthesized from the total RNA by using PrimeScript Reverse Transcriptase (Takara Bio, Otsu, Japan) according to the manufacturer's instructions. To confirm the mRNA expression of COX2 in PBMCs, real-time PCR was performed in triplicate with the LightCycler 480 System II (Roche Diagnostic, Mannheim, Germany) using SYBR Premix DimerEraser (Takara Bio) following the manufacturer's instructions. β -actin (*ACTB*) was used as a reference gene. The relative expression levels were calculated by using the $\Delta\Delta$ Ct method, and the results were indicated as relative change to no treatment group. Primers were 5' -ACG TTT TCT CGT GAA GCC CT-3' and 5' -TCT ACC AGA AGG GCG GGA TA-3' for COX2, and 5' -TCT TCC AGC CTT CCT TCC TG-3' and 5' -ACC GTG TTG GCG TAG AGG TC-3' for *ACTB*.

Quantitation of PGE₂ and estradiol by ELISA

PGE₂ concentrations in sera were determined using Prostaglandin E₂ Express ELISA Kit (Cayman Chemical, Ann Arbor, MI, USA) in accordance with the previous report [16]. Estradiol concentrations in sera were determined using Estradiol ELISA Kit (Cayman Chemical) following the manufacturer's instructions. The measurement of both PGE₂ and estradiol was performed in triplicate.

PBMC culture assays

PBMCs were cultured in 200 μ L of RPMI 1640 medium (Sigma-Aldrich) containing 10% heat-inactivated fetal calf serum (Thermo Fisher Scientific), 100 U/ml penicillin (Thermo Fisher Scientific), 100 μ g/ml streptomycin (Thermo Fisher Scientific), and 2 mM L-glutamine (Thermo Fisher Scientific). 96-well Plates (Corning Inc.) were used for all of the PBMC cultures.

To evaluate the immunosuppressive functions of estradiol *in vitro*, PBMCs (1×10^6 cells/well) from uninfected and BLV-infected cattle ($n = 6$ each) were incubated with 2 μ g/ml of 17 β -Estradiol (Cayman Chemical) and optimal antigen stimulations as shown below. Dimethyl sulfoxide (DMSO, Nacalai Tesque Inc., Kyoto, Japan) was used as a vehicle of 17 β -Estradiol. PBMCs from BLV-uninfected cattle were cultured in the presence of 1 μ g/ml anti-CD3 monoclonal antibody (mAb) (MM1A; WSU Monoclonal Antibody Center, Pullman, WA, USA) and 1 μ g/ml anti-CD28 mAb (CC220; Bio-Rad, Hercules, CA, USA) for 24 hours. PBMCs from BLV-infected cattle were cultured in the presence of fetal lamb kidney (FLK)-BLV antigen (2% heat-inactivated culture supernatant of FLK-BLV cells) for 6 days. As described in a previous paper, FLK-BLV supernatant contained BLV antigens to activate BLV-

specific T cells in PBMC cultures [20]. After incubation, culture supernatants were collected and IFN- γ concentrations were quantitated by ELISA, as described above.

To examine the effects of estradiol on the gene expression of COX2, PBMCs (1×10^6 cells/well) from BLV-uninfected cattle ($n = 7$ or 8) were incubated with $2 \mu\text{g/ml}$ 17β -Estradiol for 24 hours. Then, cells were harvested and the expression of COX2 was quantitated by real-time PCR as described above. In addition, culture supernatants were collected after incubation, and then, PGE₂ concentrations were measured by ELISA as described above.

To assess the effects of the inhibition of PGE₂ receptors in the presence of estradiol, PBMCs (1×10^6 cells/well) from uninfected cattle ($n = 10$) were pretreated with each EP antagonist (EP1: SC-19220, EP2: AH 6809, EP3: L-798,106, EP4: ONO-AE3-208; Cayman Chemical) for 1 hour. After preincubation, PBMCs were washed, and then, cultured with $2 \mu\text{g/ml}$ of 17β -Estradiol in the presence of $1 \mu\text{g/ml}$ anti-CD3 mAb and $1 \mu\text{g/ml}$ anti-CD28 mAb for 24 hours. After cultivation, culture supernatants were collected, and IFN- γ concentrations were quantitated by ELISA as described above. In the cultures using PBMCs from uninfected cattle, two or three independent PBMC-cultures were performed, and pooled data from these experiments were analyzed by the statistical analysis.

To evaluate BLV-specific Th1 responses in BLV-infected pregnant cattle (animals #1–#5), PBMCs (4×10^5 cells/well) from BLV-infected cattle were incubated with $0.1 \mu\text{g/ml}$ BLV gp51 peptide mix [21] for 6 days. Three culture wells were prepared in each sample. After incubation, IFN- γ concentrations in culture supernatants were quantitated by ELISA, as described above.

Estradiol administration

To evaluate the effects of exogenous estradiol *in vivo*, estradiol administration was performed using BLV-infected cattle (animals #6–#10, Table 1). $2.5 \text{ mg}/100 \text{ kg}$ of estradiol benzoate (OVAHORMON INJECTION; Aska Animal Health, Tokyo, Japan) was inoculated intramuscularly, and peripheral blood samples were collected from these cattle before inoculation (Day 0) and at 1 and 6 hours and 1, 2, 3, 4, and 7 days after inoculation. PGE₂ and estradiol concentrations in serum were measured by ELISA, as described above. Whole-blood culture and PBMC culture were conducted to evaluate BLV-nonspecific and BLV-specific Th1 responses, respectively.

Statistical analysis

Differences were assessed using the Wilcoxon signed-rank test, Welch's *t* test, and the Steel-Dwass test using the EZR software [22]. A *p* value of less than 0.05 was considered statistically significant.

Results

The relationship between parturition and the onset of bovine leukosis

Previous studies have shown that a stressful situation, such as parturition, is considered one of the risk factors for the onset of bovine leukosis [10, 11]. In this study, to examine the relationship between parturition and the onset, we analyzed the onset time of 68 cases which were diagnosed with bovine leukosis. As shown in Fig 1, 28 of the 68 specimens (41.18%) developed bovine leukosis during two months before and after delivery (Fig 1A and 1B). These results suggest the association of parturition with the progression to the clinical stage in BLV-infected cattle, although the detailed mechanisms are still unclear.




Fig 1. The relationship between the onset of bovine leukosis and parturition. The numbers (a) and the percentages (b) of cattle diagnosed as bovine leukosis ($n = 67$). The day of delivery was considered as day 0.

<https://doi.org/10.1371/journal.pone.0263660.g001>

Inhibition of Th1 responses in BLV-infected pregnant cattle

To reveal the association of parturition with the progression of BLV infection, in this study, we focused on the immunosuppressive function of PGE₂, which is also an important factor for the induction of parturition in pregnant animals. To investigate whether PGE₂ is involved in immune dysfunction in pregnant cattle, we examined the PGE₂ kinetics and Th1 responses of BLV-infected pregnant cattle (Fig 2A–2C and S1 Fig). Compared to the samples collected about 2 weeks before the parturition (Pre, day –17–day –12), PGE₂ concentrations in sera were significantly increased on the day of parturition (day 0, Fig 2A). In contrast, Th1 responses against Con A were significantly suppressed on day 0 (Fig 2B). Th1 responses against BLV antigen tended to be suppressed on day 0, although differences were not statistically significant ($p = 0.086$, Fig 2C). Taken together, these results suggest that PGE₂ induction before delivery might associated with the inhibition of Th1 responses in pregnant cattle.




Fig 2. Analyses of BLV-infected pregnant cattle ($n = 5$, animals #1–#5). (a) The concentrations of PGE₂ in the sera were determined by ELISA. (b and c) Whole-blood cultures or PBMC cultures were performed to evaluate IFN- γ production in response to Con A or gp51 peptide mix, respectively. (d) The concentrations of estradiol in the sera were determined by ELISA. (a–d) Statistical significance was determined by Welch's t test. Pre: day –17 for animal #1, day –16 for animals #2 and #5, day –14 for animal #3, and day –12 for animal #4. Day 0: the day of delivery, $n.s.$: not significance.

<https://doi.org/10.1371/journal.pone.0263660.g002>




Fig 3. Immunosuppressive function of estradiol via PGE₂ induction. (a and b) PBMCs from BLV-uninfected or BLV-infected cattle were cultured with estradiol in the presence of anti-CD3 mAb and anti-CD28 mAb or FLK-BLV, respectively (a: BLV-uninfected cattle, $n = 6$, b: BLV-infected cattle, $n = 6$). DMSO was used as a vehicle control of estradiol. IFN- γ concentrations in culture supernatants were quantitated by ELISA. (c and d) PBMCs from BLV-uninfected cattle were cultivated with estradiol, and COX2 expression (c, $n = 8$) and PGE₂ concentration (d, $n = 7$) were measured by real-time PCR and ELISA, respectively. (e) PBMCs from BLV-uninfected cattle ($n = 10$) were pretreated with each EP antagonist for one hour, and then cultured with estradiol in the presence of anti-CD3 mAb and anti-CD28 mAb. IFN- γ production in culture supernatant was determined by ELISA. (a–d) Statistical significances were determined by the Wilcoxon signed-rank test. (e) Data are presented as mean \pm SD, and statistical significance was determined by the Steel-Dwass test.

<https://doi.org/10.1371/journal.pone.0263660.g003>

Immunosuppressive function of estradiol via PGE₂ induction *in vitro*

To investigate the mechanism of PGE₂ induction in blood before calving, estradiol concentrations in the sera of BLV-infected pregnant cattle were measured using ELISA. Interestingly, in the same manner as PGE₂ kinetics in the pregnant cattle, estradiol concentrations tended to be increased on day 0 (Fig 2D and S1 Fig). Previous studies also have shown that the production of estrogen in blood is induced soon before parturition [23–25]. On the basis of these findings, we hypothesized that estradiol plays important roles in the induction of PGE₂ production, which causes the suppression of Th1 responses, in pregnant cattle. In order to examine this hypothesis, PBMCs from BLV-uninfected cattle were cultivated with estradiol in the presence of T-cell stimulation. IFN- γ concentrations in culture supernatants were reduced by the treatment of estradiol (Fig 3A). Additionally, estradiol impaired IFN- γ production from PBMCs of BLV-infected cattle in the presence of BLV antigen (Fig 3B). To assess whether estradiol induces PGE₂ production, PBMCs from BLV-uninfected cattle were cultivated with estradiol. After incubation, both COX2 expression and PGE₂ production were significantly increased by

treatment with estradiol (Fig 3C and 3D). Furthermore, PBMCs from BLV-uninfected cattle were pretreated with each EP antagonist, and then cultured with estradiol in the presence of T-cell stimulation. The suppressive effect of estradiol was not observed when estradiol incubation was performed in combination with EP4 antagonist (Fig 3E), but not with antagonists of other EP receptors, thus suggesting that estradiol inhibits Th1 responses via PGE₂/EP4 signaling.

Estradiol-induced impairment of Th1 responses *in vivo*

To assess the immunosuppressive function of estradiol *in vivo*, the administration of estradiol was performed using 5 BLV-infected cattle (animals #6–#10, Fig 4 and S2 Fig). We confirmed that the concentrations of estradiol in sera were increased by estradiol administration (Fig 4A). PGE₂ concentrations in sera from these infected cattle were significantly increased for 3 days after estradiol administration (Fig 4B). By contrast, IFN- γ production from whole blood cells stimulated with Con A was suppressed by estradiol administration (Fig 4C). Additionally, as shown in Fig 4D, BLV-specific Th1 responses were also impaired by estradiol administration. These results suggest that estradiol inhibits BLV-specific Th1 responses via PGE₂ production *in vivo*.

Discussion

PGE₂ is known as an important factor that is related to the induction of parturition in cattle [18]. Additionally, our previous studies have shown that PGE₂ has suppressive effects on immune responses, particularly those involving Th1 [16, 17, 26]. However, the association of PGE₂ induction during parturition with the suppression of immune responses is still unknown. Therefore, in the current study, we examined whether PGE₂ was associated with the suppression of immune responses in periparturient period. Our data showed that PGE₂ was induced, and that Th1 responses were inhibited before parturition in pregnant cattle. In addition, both BLV-specific and BLV-nonspecific Th1 responses were suppressed before parturition in pregnant cattle infected with BLV. During BLV infection, the suppression of Th1 responses against BLV antigen is associated with disease progression [27, 28]. In the advanced stages of BLV infection, BLV-specific Th1 responses are downregulated, thus leading to further disease progression and possible EBL [29]. Thus, PGE₂ mediated inhibition of BLV-specific Th1 responses around parturition might be a mechanism underlying progression to the clinical stage of BLV infection. In this study, no significant change of BLV proviral load was observed in periparturient period in BLV-infected pregnant cattle (data not shown). However,




Fig 4. Functional analysis of estradiol using BLV-infected cattle ($n = 5$, animals #6–#10). (a) Serum estradiol concentrations were measured by ELISA. (b) Serum PGE₂ concentrations were measured by ELISA. (c and d) Whole-blood cultures or PBMC cultures were performed to evaluate IFN- γ production in response to Con A or gp51 peptide mix, respectively. (a–d) Statistical significance was determined by Welch's t test (a–d).

<https://doi.org/10.1371/journal.pone.0263660.g004>

previous reports have shown that PGE₂/EP4 signaling facilitates BLV viral gene transcription via the activation of cyclic-AMP/protein kinase A/cAMP-response element signaling [17, 30]. Thus, additional experiments with larger numbers of BLV-infected pregnant cattle are needed to examine the effect of PGE₂ on BLV proviral load during parturition.

In this study, we identified a novel mechanism of PGE₂ upregulation via estradiol. Previous studies have shown that treatment with estradiol induces NF-κB activation in human and rat models [31, 32]; moreover, NF-κB activation is essential for COX-2 induction [14, 33]. Therefore, estradiol presumably induces PGE₂ production in the blood via NF-κB activation. In addition, our data showed that PGE₂/EP4 signaling involved at least in the suppressive effect of estradiol. By contrast, we did not observe a significant difference between the group treated with estradiol alone and the group treated with estradiol after pretreatment with the EP2 antagonist. Generally, EP2 and EP4 are receptors, which are associated with the immunosuppressive activity of PGE₂. However, EP4 represents a high-affinity receptor, whereas EP2 requires significantly higher concentrations of PGE₂ for effective signaling [15]. Therefore, the differential effects of EP2 and EP4 blockades are presumably due to the affinity of each receptor.

Tregs are critical for maternal tolerance [4]. Several reports have demonstrated that estrogen drives the induction of Tregs in murine models [34, 35]. Additionally, our previous reports have shown that PGE₂ upregulates the expression levels of *Foxp3* and *TGFβ1* in bovine PBMCs [17], and suppressive function of Tregs is associated with the disease progression during BLV infection [36]. Collectively, the available data suggest that during parturition, Tregs induced by the estradiol/PGE₂ pathway might suppress Th1 responses and contribute to the progression of BLV infection. Further studies are warranted to investigate the involvement of Tregs in the suppression of Th1 responses during parturition.

Our previous studies have shown that PGE₂ upregulation is also involved in the progression of other bovine chronic diseases, such as *Mycoplasma bovis* infection and Johne's disease, which is caused by *Mycobacterium avium* subsp. *paratuberculosis* [16, 25]. Additionally, in terms of Johne's disease, it has been reported that the stress related to parturition is a risk factor for eliciting the onset of the clinical stage of the disease [37]. However, no report has been examined whether PGE₂ induction during parturition inhibits *M. bovis*-specific and *M. avium* subsp. *paratuberculosis*-specific Th1 responses. The involvement of the estradiol/PGE₂ pathway in Th1 suppression during parturition should be determined in further experiments by using pregnant cattle with these diseases.

In conclusion, during parturition, BLV-specific Th1 responses are suppressed via estradiol-induced PGE₂, which contributes to the disease progression of BLV-infection. Additionally, estradiol-induced PGE₂ also inhibited BLV-nonspecific Th1 responses, thus suggesting that PGE₂ might be involved in susceptibility to opportunistic infections in the periparturient period. To the best of our knowledge, this is the first study to show the suppressive function of the estradiol/PGE₂/EP4 pathway. Further studies will open up new avenues for the control of infections during parturition both in pregnant cattle and in pregnant women.

Supporting information

S1 Checklist. The ARRIVE guidelines 2.0: Author checklist.

(PDF)

S1 Fig. Analyses of BLV-infected pregnant cattle. (a) The concentrations of PGE₂ in the sera were determined by ELISA. (b and c) Whole-blood culture or PBMC culture was performed to evaluate IFN-γ production in response to Con A or gp51 peptide mix, respectively. (d) The concentrations of estradiol in the sera were determined by ELISA.

(PPTX)

S2 Fig. Functional analysis of estradiol using BLV-infected cattle. (a–d) Estradiol administration was performed using BLV-infected cattle (animals #6–#10). (a) Serum estradiol concentrations were measured by ELISA. (b) Serum PGE₂ concentrations were measured by ELISA. (c and d) Whole-blood culture or PBMC culture was performed to evaluate IFN- γ production in response to Con A or gp51 peptide mix, respectively.

(PPTX)

S1 Table. The numbers of cattle diagnosed as bovine leukosis, related to Fig 1.

(PPTX)

S2 Table. The original data of analyses of BLV-infected pregnant cattle (animals #1–#5), related to Fig 2. (a) The concentrations of PGE₂ in the sera. (b and c) IFN- γ production in response to Con A (b) or gp51 peptide mix (c) in the whole-blood cultures (b) or PBMC cultures (c). (d) The concentrations of estradiol in the sera.

(PPTX)

S3 Table. The original data of immunosuppressive assay using estradiol, related to Fig 3. (a and b) IFN- γ production in the cultures of PBMCs from BLV-uninfected (a) and infected cattle (b) cultivated with anti-CD3 and anti-CD28 mAbs (a) and FLK-BLV (b), respectively. (c and d) COX2 expression (c) and PGE₂ concentration (d) in PBMCs from BLV-uninfected cattle cultivated with estradiol. (e) IFN- γ production in PBMCs from BLV-uninfected cattle cultivated with each EP antagonist, estradiol, and anti-CD3 and anti-CD28 mAbs.

(PPTX)

S4 Table. The original data of functional analysis of estradiol using BLV-infected cattle (animals #6–#10), related to Fig 4. (a) The concentrations of estradiol in the sera. (b) The concentrations of PGE₂ in the sera. (c and d) IFN- γ production in response to Con A (c) or gp51 peptide mix (d) in the whole-blood cultures (c) or PBMC cultures (d).

(PPTX)

Acknowledgments

We are grateful to Dr. Hideyuki Takahashi, Dr. Tomio Ibayashi, and Dr. Yasuyuki Mori for valuable advice and discussions. We thank Enago (www.enago.jp) for the English language review.

Author Contributions

Conceptualization: Yamato Sajiki, Satoru Konnai, Tomohiro Okagawa, Naoya Maekawa, Shiro Murata, Kazuhiko Ohashi.

Data curation: Yamato Sajiki, Satoru Konnai, Tomohiro Okagawa, Naoya Maekawa, Shinya Goto, Junko Kohara, Atsushi Nitani, Hirofumi Takahashi, Kentaro Kubota, Hiroshi Takeda, Shiro Murata, Kazuhiko Ohashi.

Formal analysis: Yamato Sajiki, Satoru Konnai, Tomohiro Okagawa, Naoya Maekawa, Shinya Goto, Junko Kohara, Hirofumi Takahashi, Kentaro Kubota, Hiroshi Takeda, Shiro Murata, Kazuhiko Ohashi.

Funding acquisition: Satoru Konnai.

Investigation: Yamato Sajiki, Satoru Konnai, Tomohiro Okagawa, Shinya Goto, Junko Kohara, Atsushi Nitani, Hirofumi Takahashi, Kentaro Kubota, Hiroshi Takeda.

Methodology: Yamato Sajiki, Satoru Konnai, Tomohiro Okagawa, Junko Kohara, Atsushi Nitani, Hirofumi Takahashi, Kentaro Kubota, Hiroshi Takeda.

Project administration: Satoru Konnai.

Resources: Satoru Konnai.

Supervision: Satoru Konnai.

Validation: Satoru Konnai.

Visualization: Satoru Konnai.

Writing – original draft: Yamato Sajiki, Satoru Konnai, Naoya Maekawa, Shiro Murata, Kazuhiko Ohashi.

Writing – review & editing: Shiro Murata, Kazuhiko Ohashi.

References

1. Goff JP, Horst RL. Physiological changes at parturition and their relationship to metabolic disorders. *J Dairy Sci.* 1997; 80: 1260–1268. [https://doi.org/10.3168/jds.S0022-0302\(97\)76055-7](https://doi.org/10.3168/jds.S0022-0302(97)76055-7) PMID: 9241588
2. Oliveira LJ, Barreto RS, Perecin F, Mansouri-Attia N, Pereira FT, Meirelles FV. Modulation of maternal immune system during pregnancy in the cow. *Reprod Domest Anim.* 2012; 47 Suppl 4: 384–393. <https://doi.org/10.1111/j.1439-0531.2012.02102.x> PMID: 22827396
3. Saito S, Nakashima A, Shima T, Ito M. Th1/Th2/Th17 and regulatory T-cell paradigm in pregnancy. *Am J Reprod Immunol.* 2010; 63: 601–610. <https://doi.org/10.1111/j.1600-0897.2010.00852.x> PMID: 20455873
4. Figueiredo AS, Schumacher A. The T helper type 17/regulatory T cell paradigm in pregnancy. *Immunology* 2016; 148: 13–21. <https://doi.org/10.1111/imm.12595> PMID: 26855005
5. Feinberg BB, Tan NS, Walsh SW, Brath PC, Gonik B. Progesterone and estradiol suppress human mononuclear cell cytotoxicity. *J Reprod Immunol.* 1997; 21: 139–148. [https://doi.org/10.1016/0165-0378\(92\)90020-5](https://doi.org/10.1016/0165-0378(92)90020-5)
6. Robinson DP, Klein SL. Pregnancy and pregnancy-associated hormones alter immune responses and disease pathogenesis. *Horm Behav.* 2012; 62: 263–271. <https://doi.org/10.1016/j.yhbeh.2012.02.023> PMID: 22406114
7. Maeda Y, Ohtsuka H, Tomioka M, Oikawa M. Effect of progesterone on Th1/Th2/Th17 and regulatory T cell-related genes in peripheral blood mononuclear cells during pregnancy in cows. *Vet Res Commun.* 2013; 37: 43–49. <https://doi.org/10.1007/s11259-012-9545-7> PMID: 23203561
8. Frie MC, Coussens PM. Bovine leukemia virus: a major silent threat to proper immune responses in cattle. *Vet Immunol Immunopathol.* 2015; 163: 103–114. <https://doi.org/10.1016/j.vetimm.2014.11.014> PMID: 25554478
9. Schwartz I, Levy D. Pathobiology of bovine leukemia virus. *Vet Res.* 1994; 25: 521–536. PMID: 7889034
10. Hopkins SG, DiGiacomo RF. Natural transmission of bovine leukemia virus in dairy and beef cattle. *Vet Clin North Am Food Anim Pract.* 1997; 13: 107–128. [https://doi.org/10.1016/s0749-0720\(15\)30367-4](https://doi.org/10.1016/s0749-0720(15)30367-4) PMID: 9071749
11. Monti GE, Frankena K, DeJong MC. Evaluation of transmission of bovine leukemia virus with in dairy herds of Argentina. *Epidemiol Infect.* 2007; 135: 228–237. <https://doi.org/10.1017/S0950268806006637> PMID: 16780609
12. Phipps RP, Stein SH, Roper RL. A new view of prostaglandin E regulation of the immune response. *Immunol Today* 1991; 12: 349–352. [https://doi.org/10.1016/0167-5699\(91\)90064-Z](https://doi.org/10.1016/0167-5699(91)90064-Z) PMID: 1958288
13. Morita I. Distinct functions of COX-1 and COX-2. *Prostaglandins Other Lipid Mediat.* 2002;68–69: 165–175. [https://doi.org/10.1016/s0090-6980\(02\)00029-1](https://doi.org/10.1016/s0090-6980(02)00029-1) PMID: 12432916
14. Subbaramaiah K, Telang N, Ramonetti TJ, Araki R, DeVito B, Weksler BB, et al. Transcription of cyclooxygenase-2 is enhanced in transformed mammary epithelial cells. *Cancer Res.* 1996; 56: 4424–4429. PMID: 8813136
15. Kalinski P. Regulation of Immune Responses by Prostaglandin E₂. *J Immunol.* 2012; 188: 21–28. <https://doi.org/10.4049/jimmunol.1101029> PMID: 22187483

16. Sajiki Y, Konnai S, Okagawa T, Nishimori A, Maekawa N, Goto S, et al. Prostaglandin E₂ Induction Suppresses the Th1 Immune Responses in Cattle with Johne's Disease. *Infect Immun*. 2018; 86: e00910–17. <https://doi.org/10.1128/IAI.00910-17> PMID: 29483289
17. Sajiki Y, Konnai S, Okagawa T, Nishimori A, Maekawa N, Goto S, et al. Prostaglandin E₂-Induced Immune Exhaustion and Enhancement of Antiviral Effects by Anti-PD-L1 Antibody Combined with COX-2 Inhibitor in Bovine Leukemia Virus Infection. *J Immunol*. 2019; 203: 1313–1324. <https://doi.org/10.4049/jimmunol.1900342> PMID: 31366713
18. Hirsbrunner G, Zanolari P, Althaus H, Hüsler J, Steiner A. Influence of prostaglandin E₂ on parturition in cattle. *Vet Rec*. 2007; 161: 414–417. <https://doi.org/10.1136/vr.161.12.414> PMID: 17890770
19. Sajiki Y, Konnai S, Nishimori A, Okagawa T, Maekawa N, Goto S, et al. Intrauterine infection with bovine leukemia virus in pregnant dam with high viral load. *J Vet Med Sci*. 2017; 79: 2036–2039. <https://doi.org/10.1292/jvms.17-0391> PMID: 29109356
20. Mager A, Masengo R, Mammerickx M, Letesson JJ. T cell proliferative response to bovine leukaemia virus (BLV): identification of T cell epitopes on the major core protein (p24) in BLV-infected cattle with normal haematological values. *J Gen Virol*. 1994; 75: 2223–2231. <https://doi.org/10.1099/0022-1317-75-9-2223> PMID: 7521391
21. Ikebuchi R, Konnai S, Okagawa T, Yokoyama K, Nakajima C, Suzuki Y, et al. Blockade of bovine PD-1 increases T cell function and inhibits bovine leukemia virus expression in B cells *in vitro*. *Vet Res*. 2013; 44: 59. <https://doi.org/10.1186/1297-9716-44-59> PMID: 23876077
22. Kanda Y. Investigation of the freely available easy-to-use software 'EZR' for medical statistics. *Bone Marrow Transplant*. 2013; 48: 452–458. <https://doi.org/10.1038/bmt.2012.244> PMID: 23208313
23. Smith VG, Edgerton LA, Hafs HD, Convey EM. Bovine Serum Estrogens, Progestins and Glucocorticoids during Late Pregnancy, Parturition and Early Lactation. *J Anim Sci*. 1973; 36: 391–396. <https://doi.org/10.2527/jas1973.362391x> PMID: 4687097
24. Robertson HA. Changes in the Concentration of Unconjugated Oestrone, oestradiol-17alpha and oestradiol-17beta in the Maternal Plasma of the Pregnant Cow in Relation to the Initiation of Parturition and Lactation. *J Reprod Fertil*. 1974; 36: 1–7. <https://doi.org/10.1530/jrf.0.0360001> PMID: 4809699
25. Stellflug JN, Han DK, Randel RD, Moody EL. Plasma estrogens in the periparturient cow. *Theriogenology* 1978; 10: 269–273.
26. Goto S, Konnai S, Hirano Y, Kohara J, Okagawa T, Maekawa N, et al. Upregulation of PD-L1 Expression by Prostaglandin E₂ and the Enhancement of IFN-g by Anti-PD-L1 Antibody Combined With a COX-2 Inhibitor in *Mycoplasma bovis* Infection. *Front Vet Sci*. 2020; 7: 12. <https://doi.org/10.3389/fvets.2020.00012> PMID: 32154274
27. Orlik O, Splitter GA. Progression to persistent lymphocytosis and tumor development in bovine leukemia virus (BLV)-infected cattle correlates with impaired proliferation of CD4+ T cells in response to gag- and env-encoded BLV proteins. *J Virol*. 1996; 70: 7584–7593. <https://doi.org/10.1128/JVI.70.11.7584-7593.1996> PMID: 8892878
28. Kabeya H, Ohashi K, Onuma M. Host immune responses in the course of bovine leukemia virus infection. *J Vet Med Sci*. 2001; 63: 703–708. <https://doi.org/10.1292/jvms.63.703> PMID: 11503896
29. Konnai S, Murata S, Ohashi K. Immune exhaustion during chronic infections in cattle. *J Vet Med Sci*. 2017; 79: 1–5. <https://doi.org/10.1292/jvms.16-0354> PMID: 27725355
30. Pyeon D, Diaz FJ, Splitter GA. Prostaglandin E₂ increases bovine leukemia virus tax and pol mRNA levels via cyclooxygenase 2: regulation by interleukin-2, interleukin-10, and bovine leukemia virus. *J Virol*. 2000; 74: 5740–5745. <https://doi.org/10.1128/jvi.74.12.5740-5745.2000> PMID: 10823885
31. Hamilton KL, Gupta S, Knowlton AA. Estrogen and regulation of heat shock protein expression in female cardiomyocytes: cross-talk with NF kappa B signaling. *J Mol Cell Cardiol*. 2004; 36: 577–584. <https://doi.org/10.1016/j.yjmcc.2004.02.005> PMID: 15081317
32. Hirano S, Furutama D, Hanafusa T. Physiologically high concentrations of 17beta-estradiol enhance NF-kappaB activity in human T cells. *Am J Physiol Regul Integr Comp Physiol*. 2007; 292: R1465–1471. <https://doi.org/10.1152/ajpregu.00778.2006> PMID: 17194723
33. Jobin C, Morteau O, Han DS, Balfour Sartor R. Specific NF-kappaB blockade selectively inhibits tumour necrosis factor-alpha-induced COX-2 but not constitutive COX-1 gene expression in HT-29 cells. *Immunology* 1998; 95: 537–543. <https://doi.org/10.1046/j.1365-2567.1998.00646.x> PMID: 9893042
34. Polanczyk MJ, Carson BD, Subramanian S, Afentoulis M, Vandembark AA, Ziegler SF, et al. Cutting edge: estrogen drives expansion of the CD4+CD25+ regulatory T cell compartment. *J Immunol*. 2004; 173: 2227–2230. <https://doi.org/10.4049/jimmunol.173.4.2227> PMID: 15294932
35. Tai P, Wang J, Jin H, Song X, Yan J, Kang Y, et al. Induction of regulatory T cells by physiological level estrogen. *J Cell Physiol*. 2008; 214: 456–464. <https://doi.org/10.1002/jcp.21221> PMID: 17654501

36. Ohira K, Nakahara A, Konnai S, Okagawa T, Nishimori A, Maekawa N, et al. Bovine leukemia virus reduces anti-viral cytokine activities and NK cytotoxicity by inducing TGF- β secretion from regulatory T cells. *Immun Inflamm Dis*. 2016; 4: 52–63. <https://doi.org/10.1002/iid3.93> PMID: 27042304
37. Clarke CJ. The pathology and pathogenesis of paratuberculosis in ruminants and other species. *J Comp Pathol*. 1997; 116: 217–261. [https://doi.org/10.1016/s0021-9975\(97\)80001-1](https://doi.org/10.1016/s0021-9975(97)80001-1) PMID: 9147244