



Taxonomic Paper

Revision of *Metaplagia* Coquillett (Diptera: Tachinidae) with description of five new species from Área de Conservación Guanacaste in northwestern Costa Rica

AJ Fleming[‡], D. Monty Wood[‡], M. Alex Smith[§], Winnie Hallwachs[¶], Daniel Janzen[|]

[‡] Agriculture Agri-Food Canada, Ottawa, Canada

[§] University of Guelph, Guelph, Canada

[|] Department of Biology, University of Pennsylvania, Philadelphia, United States of America

Corresponding author: AJ Fleming (aifleming604@gmail.com)

Academic editor: Torsten Dikow

Received: 12 May 2021 | Accepted: 16 Jul 2021 | Published: 29 Jul 2021

Citation: Fleming A, Wood DM, Smith MA, Hallwachs W, Janzen D (2021) Revision of *Metaplagia* Coquillett (Diptera: Tachinidae) with description of five new species from Área de Conservación Guanacaste in northwestern Costa Rica. Biodiversity Data Journal 9: e68598. <https://doi.org/10.3897/BDJ.9.e68598>

ZooBank: [urn:lsid:zoobank.org:pub:2085FED0-9C53-46EB-B8AC-BFD99139FC69](https://urn.nbn.se/resolve?urn=urn:lsid:zoobank.org:pub:2085FED0-9C53-46EB-B8AC-BFD99139FC69)

Abstract

Background

We revise the genus *Metaplagia* Coquillett, 1895 and describe five new species from Área de Conservación Guanacaste (ACG) in northwestern Costa Rica. All new species were reared from an ongoing inventory of wild-caught caterpillars spanning a variety of species within the family Sphingidae (Lepidoptera: Sphingidae). Our study provides a concise description of each new species using morphology, life history, molecular data and photographic documentation. In addition to the new species, the authors provide a re-description of the genus and a revised key to the species of *Metaplagia*.

New information

The following five new species of *Metaplagia* are described: *Metaplagia leahdenniseae* Fleming & Wood **sp. n.**, *Metaplagia lindarobinsonae* Fleming & Wood **sp. n.**, *Metaplagia paulinesaribasae* Fleming & Wood **sp. n.**, *Metaplagia robinsherwoodae* Fleming & Wood **sp. n.** and *Metaplagia svetlanakozikae* Fleming & Wood **sp. n.**.

The following is proposed by Fleming & Wood as new combination of *Plagiomima* Brauer & Bergenstamm, 1891: *Plagiomima latifrons* (Reinhard, 1956) **comb. n.**

Keywords

tropical rain forest, tropical dry forest, cloud forest, parasitoid flies, host-specificity, caterpillars, ACG, Dexiinae, Voriini

Introduction

The tribe Voriini is distributed globally; the vast tachinid fauna of the Neotropical Region and the huge number of genera have proven to be one of the most significant hurdles to understanding the tribal boundaries surrounding the Voriini. Most attempts to understand the synapomorphies or of the tribe have, rather than answered questions, led researchers to even more questions as to what the voriines truly are. Presently, the tribe possesses several diagnostic features which the authors consider as typical to the tribe: conical head profile (longer at level of pedicel than at vibrissa); proclinate, divergent and well-developed ocellar setae; frons wide; proclinate and reclinate orbital setae present in both sexes; facial ridge bare; prosternum bare; anepimeral seta absent or poorly developed appearing hair-like; infrasquamal setae present; apical scutellar setae strong and decussate; dm-m crossvein oblique, making posterior section of M_4 subequal to anterior section; R_{4+5} setulose at least to crossvein r-m and sometimes beyond; middorsal depression of ST1+2 reaching posterior margin; and aedeagus elongate and frequently ribbon-like (Tschorasnig 1985, Cortés and González 1989). Voriini lay flattened membranous incubated eggs directly on the cuticle of the host (Herting 1956, Guimarães 1977). Unfortunately, some of these diagnostic features can be observed across several other taxa within the Dexiinae and, as such, our recognition of the tribe can be seen as a combination of these features used to exclude other genera and define the Voriini; much variation exists within the catch-all tribe and the "gestalt" of the tribe is still somewhat nebulous. The Voriini are a problematic tribe and still the subject of much debate. Recent work has suggested that the Voriini are a vast unifying polyphyletic clade encompassing much of the Dexiinae (Stireman et al. 2019), only further emphasizing the catch-all nature of the tribe; this evidence, however, is still the subject of discussion.

The monotypic genus *Metaplagia* Coquillett, 1895 (Dexiinae: Voriini) was initially erected for the type species *Metaplagia occidentalis* Coquillett, 1895 (Coquillett 1895). Coquillett based the new species on a single male specimen he collected in San Diego Co.

California. The genus was previously known exclusively from the Nearctic; however, the data we present herein suggests the genus reaches into the Neotropics. A significant obstacle to revising a genus like *Metaplagia* is that the interspecific morphological differences are often subtle and, in many cases, variable. However, despite their morphological similarities, when examined at a genetic level, they display a great deal of interspecific variation, allowing the individual species to be further discriminated.

Our goal is to systematically revise and analyze the known members of the New World tachinid genus *Metaplagia* Coquillett and add to the existing taxa five new species from Área de Conservación Guanacaste (ACG). Our species concepts are based on differences in morphology, life history and by comparison of COI (coxI or cytochrome c oxidase I) gene sequences. We also provide a key to the species of *Metaplagia* inclusive of the Nearctic and Mesoamerican Regions. This paper is part of a broader effort to name and catalog all of the tachinid species collected from the ACG inventory, an effort which has already presented new information within the tribe Voriini (Fleming et al. 2017, Fleming et al. 2019). This series of taxonomic papers will represent a foundation for later, detailed ecological and behavioral accounts and studies, extending across ACG ecological groups, whole ecosystems and taxonomic assemblages much larger than a genus. All new information described here derives from the ongoing inventory of the tri-trophic relationships between caterpillars, their food plants and their parasitoids within the dry, rain and cloud forests of the terrestrial portion of ACG (Janzen et al. 2009, Janzen et al. 2011, Janzen and Hallwachs 2011, Janzen et al. 2020). This inventory started in 1978 and has yielded an unprecedented amount of invaluable ecological information on the tri-trophic relationships between parasitoids, hosts and associated food plants.

Materials and methods

Project aims and rearing intensity

All reared specimens were obtained from host caterpillars collected in ACG (Janzen and Hallwachs 2016). ACG's 125,000+ terrestrial hectares span the Provinces of Alajuela and Guanacaste, along the dry forested northwestern coast of Costa Rica and inland to the Caribbean lowland rain forest. ACG comprises several different biomes and intergrades, ranging from sea level up to 2,000 m. The tachinid rearing methods are described at http://janzen.bio.upenn.edu/caterpillars/methodology/how/parasitoid_husbandry.htm. Since its inception, this inventory has reared over 750,000 wild-caught ACG caterpillars. Any frequencies of parasitization reported here need to be considered against this background inventory. Comparative details of the parasitization ecology of these flies will be treated separately in later papers, in the context of the study of all parasitization rates of tachinids on ACG caterpillars, once the overall alpha taxonomy of ACG caterpillar-attacking tachinids is more complete than at present.

Voucher specimen management

The management of voucher specimens has been detailed in previous papers in this series (Fleming et al. 2014). In brief, all caterpillars reared from the ACG efforts receive a unique voucher code in the format yy-SRNP-xxxx. Any parasitoid emerging from a caterpillar receives the same voucher code as a record of the rearing event. If and when the parasitoid is later dealt with individually, it receives a second voucher code unique to it, in the format DHJPARxxxxxx. These voucher codes, assigned to both host and parasitoids, may be used to obtain the individual rearing record at <http://janzen.bio.upenn.edu/caterpillars/database.lasso>.

To date, all DHJPARxxxxx-coded tachinids have had one leg removed for DNA barcoding at the Biodiversity Institute of Ontario (BIO) in Guelph, ON, Canada. All successful barcodes and collateral data are first deposited in the Barcode of Life Data System (BOLD, www.boldsystems.org) (Ratnasingham and Hebert 2007) and later migrated to GenBank. Each barcoded specimen is also assigned unique accession codes from both the Barcode of Life Data System (BOLD) and GenBank, respectively.

Inventoried Tachinidae were collected under Costa Rican government research permits issued to DHJ and exported from Costa Rica to Philadelphia, en route to their final depository in the Canadian National Insect collection in Ottawa, Canada (CNC). Tachinid identifications for the inventory were done by DHJ in coordination with: a) visual inspection by AJF and DMW, b) DNA barcode sequence examination by MAS and DHJ and c) correlation with host caterpillar identifications by DHJ and WH through the inventory itself. Dates of collection, cited for each ACG specimen, are the dates of eclosion of the fly, not the date of capture of the caterpillar since the fly eclosion date is much more representative of the time when that fly species is on the wing than is the time of capture of the host caterpillar. The collector listed on the label is the parataxonomist who found the caterpillar, rather than the person who retrieved the newly-eclosed fly from its rearing container. The holotypes and paratypes of the species newly-described herein are all deposited at CNC.

Descriptions and imaging

Species accounts and descriptions are deliberately brief and concise, complemented by a series of color photos of every species, used to illustrate the morphological differences amongst them. The morphological terminology used follows Cumming and Wood (2017). All dissections and photography were carried out following the methods detailed in Fleming et al. (2014). If only one male were available, it was designated as the holotype and not subjected to dissection.

Interim names of undescribed host species

Names of undescribed host species follow a standardized, interim naming system used for taxonomic units considered as distinct species and identified by DNA barcodes. The interim names are given in the format "*Manduca sexta*DHJ02" or "*Manduca sexta*DHJ03",

where the "species epithet" is either composed of the name of the taxonomist who identified the species and a number or the name of a species-group, followed by a code. This prevents confusion with already-described species, while maintaining traceability of each undescribed species within the ACG project.

DNA barcoding

We generated DNA extracts from single legs using a glass fibre protocol (Ivanova et al. 2006), using the standard DNA barcode region (5' cytochrome c oxidase I (COI) gene) for all specimens of ACG *Metaplagia*. The DNA barcodes (658 bp near the 5' terminus of the COI gene) were amplified using general insect primers, using standard protocols for both production and quality control (Smith et al. 2006, Smith et al. 2007, Smith et al. 2008, Smith et al. 2009, Smith 2012). DNA sequences, trace files and accessions were deposited in the Barcode of Life Data System (BOLD) (Ratnasingham and Hebert 2007). Metadata (including GenBank accession codes) associated with each sequence can be consulted on BOLD by using the persistent DOI (<https://doi.org/10.5883/DS-ASMETAPL>).

The phylogeny (Fig. 1) was created using the Maximum Likelihood method, based on the Tamura 3-parameter model (Tamura 1992) created using holo- and paratype DNA barcode sequences. Aligned with each tip of the tree (species name|DHJPARxxxxxx accession) is a pictorial representation of the DNA barcode of that specimen and the lateral image of the holotype for that species. The best nucleotide substitution model and the phylogeny itself were created using MEGA X (Kumar et al. 2018) and the figure created using the R (R Core Team 2019) package ggtree (Yu et al. 2017).




Figure 1. doi

A phylogeny of the type series of *Metaplagia* species described from Area de Conservacion Guanacaste. Tip photos are the holotypes of the species.

Acronyms for depositories

CNC, Canadian National Collection of Insects, Arachnids and Nematodes, Ottawa, Canada

UCDC, University of California, Davis, Bohart Museum of Entomology, Davis, California, USA

USNM, United States National Museum of Natural History, Washington, D.C., USA

Taxon treatments

Metaplagia Coquillett, 1895

Nomenclature

Metaplagia Coquillett, 1895: 102 [also 1895: 97]. Type species: *Metaplagia occidentalis* Coquillett, 1895, by original designation.

Metavoria Townsend, 1915: 101. Type species: *Metavoria orientalis* Townsend, 1915, by original designation. Synonymy proposed by Wood 1987: 1250, and further clarified in O'Hara and Wood 1998: 755.

Agathomyia Reinhard, 1959: 228 (preocc. by Verrall, 1901). Type species: *Agathomyia cordata* Reinhard, 1959, by original designation. Synonymy proposed by Wood 1987: 1250, and further clarified in O'Hara and Wood 1998: 763.

Anzamyia Reinhard, 1960: 103. (nomen novum for *Agathomyia*, 1959 Reinhard).

Other species included in *Metaplagia* Coquillett

Metaplagia brevicornis Brooks, 1945: 81. Holotype male (CNC), by original designation. Type locality: Canada, Manitoba, Teulon.

Agathomyia cordata Reinhard, 1959: 229. Holotype female (UCDC) (1 female paratype in CNC), by original designation. Type locality: USA, California, Riverside County, Anza.

Metavoria facialis Reinhard, 1956: 123. Holotype female (CNC). Type locality: USA, Utah, Fruitland.

Metaplagia occidentalis Coquillett, 1895: 103. Holotype male (USNM, Type No. [USNM ENT01519725](#)), by original designation. Type locality: USA, California, San Diego County.

Metavoria orientalis Townsend, 1915: 101. Holotype female (USNM, Type No. [USNME NT01519726](#)), by original designation. Type locality: USA, Virginia, Arlington.

Type species

Metaplagia occidentalis Coquillett, 1895

Description

Male, **head**: frontal vitta wide, 1/3-1/6th width of front-orbital plate; with 2-3 proclinate orbital setae and 1-2 reclinate orbital setae; ocellar setae proclinate slightly divergent; eye bare, not descending beyond the level of the vibrissa; fronto-orbital plate coloration ranging from shining silver to gold; fronto-orbital plate with short black setulae interspersed amongst frontal setae; fronto-orbital setae not extending below lower margin of pedicel, with fronto-orbital setulae sometimes extending below lower margin of pedicel; pedicel orange, with a black-dark brown post pedicel; arista bare and subequal to slightly shorter than postpedicel, distinctly-thickened on basal 1/2-2/3, ranging from dark orange to dark brown-black; lower margin of face almost level with vibrissa, not visible in profile; facial ridge bare, but with setulae along parafacial sometimes so close to facial ridge as to be confused with facial ridge setulae; palps either straight or with a slight club at apex, sparsely setulose. **Thorax**: tomentosity ranging from pale beige-gray to dark grey or silver over a black ground color; thorax black ground color tomentum of thorax ranging from pale brassy to silver grey; prosternum bare; 3-4 postpronotal setae arranged in a straight line; supra-alar setae 1-2:3; intra-alar setae 2-3:3; dorsocentral setae 3:3-4; acrostichal setae 3:3; katepisternum with 3 setae. Scutellum black ground color, with tomentum ranging from gray to pale brassy; with five pairs marginal setae; apical scutellar setae 1/2-1/3 as long as subapical scutellars, sub-erect, slightly above marginal plane; basal scutellar subequal in length to subapical setae; subapical setae straight, ranging from divergent to convergent. Legs: dark reddish-brown to black ground color; tarsal claws and pulvilli ranging from shorter than to longer than last tarsomere. Wings: slightly longer than abdomen; translucent slightly hyaline; R₁ and R₄₊₅ can be setulose, setulae of R₄₊₅ ranging from node to crossvein r-m or beyond. **Abdomen**: ground color black, pale silver tomentum in varying degrees on T3-T5, ST1+2 typically glabrous; middorsal depression on ST1+2 reaching to hind margin of tergite; median marginal setae present on T3 and complete rows on T4 and T5; median discal setae absent on all tergites; sex patch absent.

Terminalia: posterior margin of sternite 5 with a deeply excavated and wide U-shaped (sometimes sculptured) median cleft; lateral lobes of sternite apically rounded, often with setae along caudal margin; basal section 1/5 the length of apical lobes. Epandrum often with 3-4 long, strong setae along anal edge. Cerci, in posterior view, medially separated, but parallel and often touching so as to appear fused, with a few short setae on basal half. In lateral view, bowed and sharply tapered apically. Surstyli well-developed, stout basally in lateral view, like a stout broadly rounded triangle terminating in a small knob, appearing hooked or slightly beaked apically; in posterior view, basally enlarged and apically straight.

Female as in male, except in the following aspects: **head**: tomentum of fronto-orbital plate and parafacial can sometimes differ from that of the conspecific male; fronto-orbital plate and parafacial up to 0.5x wider than in males.

Diagnosis

Metaplagia can be distinguished by the following combination of traits: head distinctly conical; males with upper frontal setae reclinate; proclinate orbital setae in both sexes; frontal setae descending below level of pedicel; both sexes with well-developed lateral vertical setae; eye bare; parafacial setulose, but not with strong stout setae, only hair-like setulae; genal dilation very slightly developed; prementum shorter than height of head with an enlarged labellum; prosternum bare; three postsutural supra-alar setae, the anteriormost reduced and much weaker than first postsutural dorsocentral seta; scutellum with four pairs of marginal setae and one pair of erect to semi-erect apical setae; one or two pairs of sub-erect discal setae on scutellum, in line with subapical setae; vein M_1 ending separately in wing margin; anepimeral seta short, not extending beyond edge of lower calypter; wing vein R_{4+5} setulose.

Distribution

From Manitoba, Canada east to Newfoundland and south to Costa Rica.

Ecology

Within the ACG inventory, *Metaplagia* has only been reared from the Lepidoptera family Sphingidae: Lepidoptera throughout the diverse ecosystems of the research area.

Taxon discussion

Our present revision of *Metaplagia* increases the range of the genus, taking it from being a Nearctic endemic genus to a complete New World distribution, inclusive of both the Nearctic and Neotropical Regions.

Both *Plagiomima* Brauer & Bergenstamm, 1891 and *Metaplagia* are extremely similar genera; however, these can be very easily distinguished by the conformation of their mouthparts (Wood and Zumbado 2010). After careful examination of the holotype of *Metaplagia latifrons* Reinhard and based on the generic concepts and diagnostic features of both *Metaplagia* and *Plagiomima*, the authors hereby propose that species *Metaplagia latifrons* Reinhard, 1956 be transferred into the genus *Plagiomima*, as *Plagiomima latifrons* (Reinhard) **comb. n.** We base this taxonomic action on the similarities presented between the slender elongated mouthparts and the slender apically pointed surstyli present in *Metaplagia latifrons*, both diagnostic features observed in the genus *Plagiomima*, yet absent from the genus *Metaplagia*.

***Metaplagia leahdenniseae* Fleming & Wood, sp. n.**

- ZooBank [A4C3E7CC-DCCF-4EA3-B963-FA2BF3A99017](#)

Materials

Holotype:

- a. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 10/Jul/98; individualCount: 1; sex: Male; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0018627; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASTAI1274-07| 98-SRNP-8248| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Paratypes:

- a. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 16/Jul/98; individualCount: 1; sex: Male; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0018631; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASTAI1278-07| 98-SRNP-8346| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- b. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 6/May/99; individualCount: 1; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0018724; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASTAI1371-07| 98-SRNP-8240| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- c. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country:

- Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 5/May/99; individualCount: 1; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0018727; recordedBy: D.H. Janzen, W. Hallwachs & Jessica DiMauro; occurrenceStatus: Present; otherCatalogNumbers: ASTAI1374-07| 98-SRNP-8247| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- d. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Pueblo Cuajiniquil; verbatimElevation: 5; verbatimLatitude: 10.9404; verbatimLongitude: -85.6804; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 6/Jul/13; individualCount: 1; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0052501; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASHYM1855-13| 13-SRNP-15519| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- e. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Pueblo Cuajiniquil; verbatimElevation: 5; verbatimLatitude: 10.9404; verbatimLongitude: -85.6804; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 4/Jul/13; individualCount: 1; sex: Female; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0052509; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASHYM1863-13| 13-SRNP-15522| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- f. scientificName: *Metaplagia leahdenniseae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *leahdenniseae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Pueblo Cuajiniquil; verbatimElevation: 5; verbatimLatitude: 10.9404; verbatimLongitude: -85.6804; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 3/Jul/13; individualCount: 1; lifeStage: adult; preparations: Pinned; catalogNumber: DHJPAR0052512; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: Present; otherCatalogNumbers: ASHYM1866-13| 13-SRNP-15521| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 2), **head** (Fig. 2c, d): fronto-orbital plate wide, coloration silver with gold tomentum over uppermost 50%, appearing as pale brassy gold along upper 50% of occipital margin of eye; vitta wide, 1/3 width of fronto-orbital plate; ocellar setae lateraloclinate with a slight proclinate skew; with 3 proclinate orbital setae (middle proclinate orbital seta most often shorter and thinner than the other two) and 1 reclinate orbital seta fronto-orbital plate with short black setulae interspersed amongst frontal setae; fronto-orbital setulae extending below lower margin of the pedicel, well into the angle of parafacial; parafacial wholly silver tomentose; palps either slightly spathulate at tips, apically acutely inwardly curved, sparsely setulose along outer margins, inner margin appearing bare; arista dark brown-black. **Thorax** (Fig. 2a, b): black ground color with pale grey to brassy tomentum; 4 postpronotal setae; supra-alar setae 2:3; intra-alar setae 3:3; dorsocentral setae 3:3; acrostichal setae 3:3; katepisternum with 3 setae. Infrasquamal setae present. Scutellum black ground color, with tomentum concolorous with thorax; with five pairs marginal setae; apical scutellar setae 1/2 as long as subapical scutellars, sub-erect, arising above plane of marginal setae; basal scutellar subequal in length to subapical setae; subapical setae convergent, lateral scutellar setae can be convergent or strongly divergent. Legs: black ground color; mid tibia with 2–3 strong anteroventral setae; tarsal claws and pulvilli shorter than last tarsomere. Wings: R_1 bare, R_{4+5} setulose from node to crossvein r-m. **Abdomen** (Fig. 2a, b): ground color black; ST1+2 glabrous black, pale silver tomentum, occupying anterior 60% of T3–T5; tomentum not extending to ventral surface of abdomen; median marginal setae present on T3 and complete rows on T4 and T5.

Terminalia (Fig. 3): posterior margin of sternite 5 with a deeply excavated U-shaped median cleft (Fig. 3d); lateral lobes of sternite apically rounded, with a vestiture of short setae along disc, 3x as long along caudal margin; basal section 1/5 the length of apical lobes. Cerci, in posterior view, medially separated and parallel, not often touching medially, with a few short setae on basal half (Fig. 3a). In lateral view, bowed and sharply tapered apically (Fig. 3b). Surstylus well-developed, stout basally in lateral view, like a broadly rounded triangle, appearing hooked or slightly beaked apically (Fig. 3b, c); in posterior view, basally enlarged and apically straight.

Female (Fig. 4) as in male, except in the following aspects: **head** (Fig. 4c, d): frontal vitta 1/2 width of fronto-orbital plate overall brown with a light gold sheen of tomentum; overall head of females 1/5 wider than that of males; setulae interspersed amongst frontal setae more dense than in males; 3–4 rows of parafacial setulae. Legs (Fig. 4a, b): mid-tibia with 3–4 irregularly-sized strong anteroventral setae.

Diagnosis

Metaplagia leahdenniseae sp. n. can be distinguished from all other *Metaplagia* by the following combination of traits: fronto-orbital plate at least 50% gold, with a silver parafacial and setulae on vein R_{4+5} not extending beyond crossvein r-m. *Metaplagia leahdenniseae* sp. n. can be separated from *Metaplagia occidentalis* by the presence of

gold on the fronto-orbital plate. *Metaplagia leahdenniseae* sp. n. is clearly distinguished by its COI sequence clustered within the Barcode Identification Number (BIN) [BOLD:AAE2876](#).




Figure 2.

Metaplagia leahdenniseae sp. n. habitus images a-d: male, holotype n. DHJPAR0018627

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Etymology

Metaplagia leahdenniseae sp. n. is named in honor of Leah Dennis for her many years of coordinating and administrating a plethora of problems and events in the Academic Office of the Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania.



Figure 3.

Metaplagia leahdennise sp. n. terminalia images a-d: male, paratype n. DHJPAR0018631

a: caudal view [doi](#)

b: lateral view [doi](#)

c: oblique view [doi](#)

d: sternite 5, ventral view [doi](#)

Distribution

Costa Rica, ACG, Guanacaste Province, 5–275 m elevation.

Ecology

Metaplagia leahdennise sp. n. has been reared eight times from one species of Lepidoptera in the family Sphingidae: *Agrius cingulata* (Fabricius, 1775) in dry forest.




Figure 4.

Metaplagia leahdenniseae sp. n. habitus images a-d: female, paratype n. DHJPAR0052509

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Metaplagia lindarobinsonae Fleming & Wood, sp. n.

- ZooBank [0F060758-EE04-4BED-8A0A-37E904D0B7C8](#)

Materials

Holotype:

- scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca*

sexta; verbatimEventDate: 15-Jun-2016; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059091; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5508-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Paratypes:

- a. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ03; verbatimEventDate: 21-Jun-1994; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018642; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1289-07| 94-SRNP-3161| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- b. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ03; verbatimEventDate: 30-Jun-1994; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018643; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1290-07| 94-SRNP-3160| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- c. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Pocosol; locality: Area de Conservacion Guanacaste, Quebrada Aserradero; verbatimElevation: 160; verbatimLatitude: 10.8986; verbatimLongitude: -85.5642; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ03; verbatimEventDate: 22-May-1990; individualCount: 1; sex: Female; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018644; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1291-07| 89-SRNP-173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- d. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Pocosol;

- locality: Area de Conservacion Guanacaste, Quebrada Aserradero; verbatimElevation: 160; verbatimLatitude: 10.8986; verbatimLongitude: -85.5642; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ03; verbatimEventDate: 22-May-1990; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018645; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1292-07| 89-SRNP-173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- e. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Horizontes; locality: Area de Conservacion Guanacaste, Vado Esperanza; verbatimElevation: 85; verbatimLatitude: 10.7894; verbatimLongitude: -85.551; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ02; verbatimEventDate: 04-Jun-2007; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0019809; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAB357-07| 06-SRNP-17663| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- f. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059089; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5506-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- g. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059090; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5507-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- h. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa;

- locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*DHJ03; verbatimEventDate: 30-Jun-1994; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018641; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1288-07| 94-SRNP-3160| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- i. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059092; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5509-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- j. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059093; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5510-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- k. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059094; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5511-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- l. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295;

- verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059095; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5512-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- m. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059096; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5513-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- n. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059097; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5514-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- o. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059098; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA5515-16| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- p. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca*

- sexta**; verbatimEventDate: 22-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0059440; recordedBy: D.H. Janzen, W. Hallwachs & Winnie Hallwachs; occurrenceStatus: present; otherCatalogNumbers: ACGBA5857-16| 16-SRNP-10201| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- q. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: May-2017; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0060750; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA7171-17| 16-SRNP-10172| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- r. scientificName: *Metaplagia lindarobinsonae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *lindarobinsonae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca sexta*; verbatimEventDate: 15-Jun-2016; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0060751; recordedBy: D.H. Janzen, W. Hallwachs & Toby Hammer; occurrenceStatus: present; otherCatalogNumbers: ACGBA7172-17| 16-SRNP-10173| [BOLD:AAD5563](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 5), **head** (Fig. 5c, d): fronto-orbital plate wide, coloration silver to pale grey; vitta wide, 1/3 width of fronto-orbital plate ocellar setae lateraloclinate with a slight proclinate skew; with 3–4 proclinate orbital setae (middle proclinate orbital seta most often shorter and thinner than the other two) and 1 reclinate orbital seta fronto-orbital plate with short black setulae interspersed amongst frontal setae; fronto-orbital setulae extending below lower margin of pedicel, well into angle of parafacial; parafacial wholly silver tomentose with a distinct row of parafacial setulae along inner margin of parafacial approaching facial ridge, seemingly continuous with fronto-orbital setulae; palps either slightly spatulate at tips, apically acutely inwardly curved, extremely sparsely setulose along inner margin, remainder appearing bare; arista dark brown-black. **Thorax** (Fig. 5a, b): black ground color with pale grey tomentum; 3–4 postpronotal setae; supra-alar setae 2:3; intra-alar setae 3:3; dorsocentral setae 3:3; acrostichal setae 3:3; katepisternum with 3 setae. Infrasquamal setae present. Scutellum black ground color, with tomentum concolorous with thorax; with five pairs marginal setae; apical scutellar setae 1/2 as long as subapical scutellars, sub-erect,

arising above plane of marginal setae; basal scutellar subequal in length to subapical setae; subapical setae convergent, lateral scutellar setae can be convergent or strongly divergent. Legs: black ground color; mid tibia with 2–3 strong anteroventral setae; tarsal claws and pulvilli shorter than last tarsomere. Wings: R_1 bare, R_{4+5} setulose from node up to, but not beyond, crossvein $r\text{-}m$. **Abdomen** (Fig. 5a, b): ground color black; ST $1+2$ glabrous black, pale silver tomentum occupying anterior 50% of T $3\text{--}T5$; tomentum not extending to ventral surface of abdomen; median marginal setae present on T 3 , and complete rows on T 4 and T 5 .




Figure 5.

Metaplagia lindarobinsonae sp. n. habitus images a–d: male, holotype n. DHJPAR0059091

- a: dorsal view [doi](#)
- b: lateral view [doi](#)
- c: frontal view [doi](#)
- d: three quarters view [doi](#)

Terminalia (Fig. 6): posterior margin of sternite 5 with a deep and narrow excavated U-shaped median cleft (Fig. 6d); lateral lobes of sternite apically rounded, with a vestiture of short setae along disc, 3x as long along caudal margin; basal section 1/3 the length of apical lobes. Cerci, in posterior view, medially separated and parallel not touching

medially, with a few short setae on basal half (Fig. 6a); in lateral view, bowed and sharply tapered to a point apically (Fig. 6b, c). Surstylus well-developed, stout basally in lateral view, rounded basally and appearing hooked or slightly beaked apically; in posterior view, basally enlarged and apically straight.



Figure 6.

Metaplagia lindarobinsonae sp. n. terminalia images a–d: male, paratype n. DHJPAR0019809

a: caudal view [doi](#)

b: lateral view [doi](#)

c: oblique view [doi](#)

d: sternite 5, ventral view [doi](#)

Female (Fig. 7) as in male, except in the following aspects: **head** (Fig. 7c, d): fronto-orbital just under 1/2 of head width, vitta 1/3 width of fronto-orbital plate; overall head of females 1/5 wider than that of males; legs (Fig. 7b): mid-tibia with 3–4 irregularly-sized strong anteroventral setae.




Figure 7.

Metaplagia lindarobinsonae sp. n. habitus images a–d: female, paratype n. DHJPAR0018644

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Diagnosis

Metaplagia lindarobinsonae sp. n. can be distinguished from all other *Metaplagia* by the following combination of traits: frontal vitta indistinct around ocellar triangle, infrasquamal setae present, fronto-orbital plate at pale gray or silver, with a silver parafacial and setulae on vein R_{4+5} not extending beyond crossvein r-m, surstyli sharp at tip. *Metaplagia lindarobinsonae* sp. n. can be separated from *Metaplagia occidentalis* by the presence of infrasquamal setae and from *Metaplagia svetlanakozikae* sp. n. by the presence of a distinct row of setulae along the facial ridge. *Metaplagia lindarobinsonae* sp. n. is clearly distinguished by its COI sequence clustered within the Barcode Identification Number (BIN) [BOLD:AAD5563](#).

Etymology

Metaplagia lindarobinsonae sp. n. is named in honor of Linda Robinson for her many years of coordinating and administrating a plethora of problems and events for the undergraduate biology teaching laboratories of the Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania.

Distribution

Costa Rica, ACG, Guanacaste Province, 85–300 m elevation.

Ecology

Metaplagia lindarobinsonae sp. n. has been reared 17 times from two species of Lepidoptera in the family Sphingidae: *Manduca sexta* (Linnaeus, 1763), *Manduca sexta* DHJ02 and *Manduca sexta* DHJ03, in dry forest.

Metaplagia paulinesaribasae Fleming & Wood, sp. n.

- ZooBank [E421A1D5-215A-4353-BCA6-20B42098E5C6](#)

Material

Holotype:

- a. scientificName: *Metaplagia paulinesaribasae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *paulinesaribasae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca lefeburi*; verbatimEventDate: 24-Aug-1982; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018646; recordedBy: D.H. Janzen, W. Hallwachs; occurrenceStatus: present; otherCatalogNumbers: ASTAI1293-07| 82-SRNP-382| [BOLD:AAX4230](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 8), **head** (Fig. 8c, d): fronto-orbital plate wide, gold tomentum over its entirety; vitta wide, 1/5 width of fronto-orbital plate; ocellar setae lateraloclinate with a slight proclinate skew; with 2 proclinate orbital setae and 1 reclinate orbital seta fronto-orbital plate with short black setulae interspersed amongst frontal setae; fronto-orbital setulae extending below lower margin of the pedicel, well into angle of parafacial; parafacial wholly gold tomentose; palps either slightly spatulate at tips, apically acutely inwardly curved, 3–4 setulae along lower margin, inner margin appearing bare; arista light brown-dark orange, distinctly lighter color than postpedicel. **Thorax** (Fig. 8a, b): black ground color with pale brassy tomentum; 4 postpronotal setae; supra-alar

setae 2:3; intra-alar setae 3:3; dorsocentral setae 3:3; acrostichal setae 3:3; katepisternum with 3 setae. Infrasquamal setae present. Scutellum black ground color, with tomentum concolorous with thorax; with five pairs marginal setae; apical scutellar setae 1/2 as long as subapical scutellars, sub-erect, arising above plane of marginal setae; basal scutellar subequal in length to subapical setae; subapical setae convergent, lateral scutellar setae can be convergent or strongly divergent; 2 pairs of scutellar discal setae set wider apart than subapical setae. Legs: black ground color; mid tibia with 2–3 strong anteroventral setae; tarsal claws and pulvilli shorter than last tarsomere. Wings: R_1 bare, R_{4+5} setulose from node to crossvein r-m. **Abdomen** (Fig. 8a, b): ground color black; ST1+2 glabrous black, silver-pale brassy tomentum occupying anterior 90% of T3–T5; tomentum thin to absent along ventral surface of abdomen; median marginal setae present on T3 and complete rows on T4 and T5.



Figure 8.

Metaplagia paulinesaribasae sp. n. habitus images a–d: male, holotype n. DHJPAR0018646

- a: dorsal view [doi](#)
- b: lateral view [doi](#)
- c: frontal view [doi](#)
- d: three quarters view [doi](#)

Terminalia: holotype male not dissected.

Female: unknown at this time.

Diagnosis

Metaplagia paulinesaribasae sp. n. can be easily distinguished from all other *Metaplagia* by the following combination of traits: both fronto-orbital and parafacial gold and setulae on vein R₄₊₅ not extending beyond crossvein r-m. It is distinguishable from all other congeners by the presence of an entirely gold fronto-orbital plate and parafacial. *Metaplagia paulinesaribasae* sp. n. is clearly distinguished by its COI sequence clustered within the Barcode Identification Number (BIN) [BOLD:AAX4230](#).

Etymology

Metaplagia paulinesaribasae sp. n. is named in honor of Pauline Saribas for her many years of coordinating and administrating a plethora of problems and events in the Academic Office of the Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania.

Distribution

Costa Rica, ACG, Guanacaste Province, 280 m elevation.

Ecology

Metaplagia paulinesaribasae sp. n. has been reared once from one species of Lepidoptera in the family Sphingidae: *Manduca lefeburii* (Guérin-Méneville, 1844), in dry forest.

Metaplagia robinsherwoodae Fleming & Wood, sp. n.

- ZooBank [F6B26F03-C22B-4168-A675-39D20DDFA87C](#)

Materials

Holotype:

- scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 03-Jul-1992; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018640; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1287-07| 92-SRNP-2162| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; dateIdentified:

2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Paratypes:

- a. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Cebollines; verbatimElevation: 270; verbatimLatitude: 10.8222; verbatimLongitude: -85.6434; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 24-Jul-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018629; recordedBy: D.H. Janzen, W. Hallwachs & Ruth Franco; occurrenceStatus: present; otherCatalogNumbers: ASTAI1276-07| 98-SRNP-8635| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- b. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 22-Jul-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018632; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1279-07| 98-SRNP-8715| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- c. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 03-Jul-1993; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018633; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1280-07| 93-SRNP-1643| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- d. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 26-Jun-1993; individualCount: 1; lifeStage: adult;

- preparations: pinned; catalogNumber: DHJPAR0018634; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1281-07| 93-SRNP-1659| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- e. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 03-Jul-1992; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018635; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1282-07| 92-SRNP-2162| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- f. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 03-Jul-1993; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018636; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1283-07| 93-SRNP-1643| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- g. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque Humedo; verbatimElevation: 290; verbatimLatitude: 10.8514; verbatimLongitude: -85.608; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 28-Jul-1991; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018637; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1284-07| 91-SRNP-1170| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- h. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem:

- Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 04-Nov-1990; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018638; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1285-07| 90-SRNP-647| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- i. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 03-Jul-1992; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018639; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1286-07| 92-SRNP-2162| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- j. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 31-Oct-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018628; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1275-07| 98-SRNP-9161| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- k. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 07-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018713; recordedBy: D.H. Janzen, W. Hallwachs & Daniel H. Janzen; occurrenceStatus: present; otherCatalogNumbers: ASTAI1360-07| 98-SRNP-8704| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- l. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa;

- locality: Area de Conservacion Guanacaste, Quebrada Cebollines; verbatimElevation: 270; verbatimLatitude: 10.8222; verbatimLongitude: -85.6434; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018714; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1361-07| 98-SRNP-8687| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- m. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018715; recordedBy: D.H. Janzen, W. Hallwachs & Elieth Cantillano; occurrenceStatus: present; otherCatalogNumbers: ASTAI1362-07| 98-SRNP-8707| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- n. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018716; recordedBy: D.H. Janzen, W. Hallwachs & Ruth Franco; occurrenceStatus: present; otherCatalogNumbers: ASTAI1363-07| 98-SRNP-8815| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- o. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Chirigón; verbatimElevation: 250; verbatimLatitude: 10.8388; verbatimLongitude: -85.601; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018717; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1364-07| 98-SRNP-8789| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- p. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*;

- scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 07-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018718; recordedBy: D.H. Janzen, W. Hallwachs & Adrian Guadamuz; occurrenceStatus: present; otherCatalogNumbers: ASTAI1365-07| 98-SRNP-9068| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- q. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Laguna Escondida; verbatimElevation: 285; verbatimLatitude: 10.8487; verbatimLongitude: -85.6305; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018719; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1366-07| 98-SRNP-8893| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- r. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Cebollines; verbatimElevation: 270; verbatimLatitude: 10.8222; verbatimLongitude: -85.6434; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 24-Jul-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018720; recordedBy: D.H. Janzen, W. Hallwachs & Ruth Franco; occurrenceStatus: present; otherCatalogNumbers: ASTAI1367-07| 98-SRNP-8635| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- s. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 31-Oct-1998; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018725; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1372-07| 98-SRNP-9161| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

- t. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 28-Mar-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018726; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1373-07| 98-SRNP-9080| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- u. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 22-Jul-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018728; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1375-07| 98-SRNP-8715| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- v. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Chirigon; verbatimElevation: 250; verbatimLatitude: 10.8388; verbatimLongitude: -85.601; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018729; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1376-07| 98-SRNP-8789| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- w. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Bosque San Emilio; verbatimElevation: 300; verbatimLatitude: 10.8439; verbatimLongitude: -85.6138; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 08-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018730; recordedBy: D.H. Janzen, W. Hallwachs & Elieth Cantillano; occurrenceStatus: present; otherCatalogNumbers: ASTAI1377-07| 98-SRNP-8707| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified:

- 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- x. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 15-Apr-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018731; recordedBy: D.H. Janzen, W. Hallwachs & Ruth Franco; occurrenceStatus: present; otherCatalogNumbers: ASTAI1378-07| 98-SRNP-8815| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
 - y. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0060291; recordedBy: D.H. Janzen, W. Hallwachs & Tanner Frank; occurrenceStatus: present; otherCatalogNumbers: ACGBA6712-17| 16-SRNP-10177| [BO LD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
 - z. scientificName: *Metaplagia robinsherwoodae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *robinsherwoodae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Area Administrativa; verbatimElevation: 295; verbatimLatitude: 10.8376; verbatimLongitude: -85.6187; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Manduca rustica*; verbatimEventDate: 09-Jun-2018; individualCount: 1; sex: Female; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0062861; recordedBy: D.H. Janzen, W. Hallwachs & Erasmo Coronado; occurrenceStatus: present; otherCatalogNumbers: ACGBA9195-18| 17-SRNP-10052| [BOLD:AAB3286](#); identifiedBy: AJ Fleming; datelidentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 9), **head** (Fig. 9c, d): fronto-orbital plate wide, coloration silver with gold tomentum over uppermost 50%, appearing as pale brassy gold along upper 25% of occipital margin of eye; frontal vitta wide, 1/5 width of fronto-orbital plate almost entirely covered in gold tomentum; ocellar setae strongly proclinate slightly divergent; with 3 proclinate orbital setae (middle proclinate orbital seta most often almost hair-like reduced) and 1 reclinate orbital seta fronto-orbital plate with only a sparse few short

black setulae interspersed amongst frontal setae; sparse fronto-orbital setulae extending below lower margin of the pedicel (very few), these not intermingled with those in angle of parafacial; parafacial wholly silver tomentose with a single row of parafacial setulae mid-way along parafacial near facial ridge; palps slightly spathulate at tips almost oar-shaped, apically acutely inwardly curved, setulose along outer margins, inner margin appearing bare; arista dark brown-black. **Thorax** (Fig. 9a, b): black ground color with pale grey with a slightly brassy tinged tomentum, laterally this color transitions to a darkened grey tomentum; 4–5 postpronotal setae; supra-alar setae 2:3; intra-alar setae 3:3; dorsocentral setae 3:3–4; acrostichal setae 3:3; katepisternum with 3 setae. Infrasquamal setae absent. Scutellum black ground color, with tomentum concolorous with disc of thorax; with four pairs marginal setae; apical scutellar setae absent; basal scutellar 2/3 length of subapical setae; subapical setae convergent, lateral scutellar setae strongly divergent. Legs: black ground color; mid tibia with 2–3 strong anteroventral setae; tarsal claws and pulvilli shorter than last tarsomere. Wings: R₁ bare, R₄₊₅ setulose from node slightly beyond crossvein r-m. **Abdomen** (Fig. 9a, b): ground color black; ST1+2 glabrous black, pale silver tomentum occupying anterior 60% of T3–T5; tomentum extending to ventral surface of abdomen on T4 and T5; median marginal setae present on T3 and complete rows on T4 and T5.

Terminalia (Fig. 10): posterior margin of sternite 5 with a deeply excavated and sculptured U-shaped median cleft (Fig. 10d); lateral lobes of sternite apically rounded, widened at mid-point and tapering again towards base, with a vestiture of short setae along disc, 3x as long along caudal margin; basal section slightly less than 1/5 the length of apical lobes. Cerci, in posterior view, medially separated, with a few short setae along entire length, setae lengthening posteriorly towards anal operculum (Fig. 10a). In lateral view, bowed and sharply tapered apically (Fig. 10b, c). Surstyli well-developed, stout basally in lateral view, like a broadly rounded triangle, appearing like a rounded ellipsoid leaf shape; in posterior view, basally enlarged and apically straight.

Female (Fig. 11) as in male, except in the following aspects: **head** (Fig. 11c, d): fronto-orbital just under 1/2 of head width, vitta 1/4 width of fronto-orbital plate; ocellar setae lateralocliniate; post-pedicel slightly orange along basal 1/5. Legs (Fig. 11b): mid-tibia with 3–4 irregularly-sized strong anteroventral setae.

Diagnosis

Metaplagia robinsherwoodae sp. n. can be distinguished from all other *Metaplagia* by the following combination of traits: fronto-orbital plate at least 75% gold, with a silver parafacial and setulae on vein R₄₊₅ extending well beyond crossvein r-m. *Metaplagia robinsherwoodae* sp. n. can be separated from *M. leahdenniseae* and *M. paulinesaribasae* by the presence of setulae on R₄₊₅ extending well beyond crossvein r-m and it is separated from *M. facialis* and *M. cordata* by the presence of gold on its fronto-orbital plate. *Metaplagia robinsherwoodae* sp. n. is clearly distinguished by its COI sequence clustered within the Barcode Identification Number (BIN) [BOLD:AAB3286](#).




Figure 9.

Metaplagia robinsherwoodae sp. n. habitus images a–d: male, holotype n. DHJPAR0018640

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Etymology

Metaplagia robinsherwoodae sp. n. is named in honor of Robin Sherwood for her many years of coordinating and administrating a plethora of problems and events in the Academic Office of the Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania.

Distribution

Costa Rica, ACG, Guanacaste Province, 250–300 m elevation.



Figure 10.

Metaplagia robinsherwoodae sp. n. terminalia images a-d: male, holotype n. DHJPAR0018725

a: caudal view [doi](#)

b: lateral view [doi](#)

c: oblique view [doi](#)

d: sternite 5, ventral view [doi](#)

Ecology

Metaplagia robinsherwoodae sp. n. has been reared 28 times from one species of Lepidoptera in the family Sphingidae: *Manduca rustica* (Fabricius, 1775), in dry forest.




Figure 11.

Metaplagia robinsherwoodae sp. n. habitus images a-d: female, paratype n. DHJPAR0062861

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Metaplagia svetlanakozikae Fleming & Wood, sp. n.

- ZooBank [42C752D8-B95A-4450-B7D2-0AEB10008472](https://zoobank.org/42C752D8-B95A-4450-B7D2-0AEB10008472)

Materials

Holotype:

- scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Vado Rio Calera; verbatimElevation: 10; verbatimLatitude: 10.8027; verbatimLongitude: -85.6742; verbatimCoordinateSystem:

Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 12-Nov-2001; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018732; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1379-07| 01-SRNP-13896| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Paratypes:

- a. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 06-May-1999; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018721; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1368-07| 98-SRNP-8103| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- b. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 08-Jun-1998; individualCount: 1; sex: Female; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018722; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1369-07| 98-SRNP-8104| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- c. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 10-Jul-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018723; recordedBy: D.H. Janzen, W. Hallwachs & Manuel Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1370-07| 98-SRNP-8248| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- d. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country:

- Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Quebrada Costa Rica; verbatimElevation: 275; verbatimLatitude: 10.8274; verbatimLongitude: -85.6365; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 08-Jun-1998; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018630; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1277-07| 98-SRNP-8104| [BOLD:AAE2876](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- e. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Cafetal; verbatimElevation: 280; verbatimLatitude: 10.8583; verbatimLongitude: -85.6109; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 29-May-2002; individualCount: 1; sex: Male; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018733; recordedBy: D.H. Janzen, W. Hallwachs & gusaneros; occurrenceStatus: present; otherCatalogNumbers: ASTAI1380-07| 01-SRNP-14594| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- f. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Vado Rio Calera; verbatimElevation: 10; verbatimLatitude: 10.8027; verbatimLongitude: -85.6742; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 12-Nov-2001; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018734; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1381-07| 01-SRNP-13896| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen
- g. scientificName: *Metaplagia svetlanakozikae*; phylum: Arthropoda; class: Insecta; order: Diptera; family: Tachinidae; genus: *Metaplagia*; specificEpithet: *svetlanakozikae*; scientificNameAuthorship: Fleming & Wood, 2021; continent: Central America; country: Costa Rica; countryCode: CR; stateProvince: Guanacaste; county: Sector Santa Rosa; locality: Area de Conservacion Guanacaste, Vado Rio Calera; verbatimElevation: 10; verbatimLatitude: 10.8027; verbatimLongitude: -85.6742; verbatimCoordinateSystem: Decimal degrees; samplingProtocol: Reared from the larva of the Sphingidae, *Agrius cingulata*; verbatimEventDate: 12-Nov-2001; individualCount: 1; lifeStage: adult; preparations: pinned; catalogNumber: DHJPAR0018735; recordedBy: D.H. Janzen, W. Hallwachs & Guillermo Pereira; occurrenceStatus: present; otherCatalogNumbers: ASTAI1382-07| 01-SRNP-13896| [BOLD:AAD5456](#); identifiedBy: AJ Fleming; dateIdentified: 2021; language: en; institutionCode: CNC; collectionCode: Insects; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 12), **head** (Fig. 12c, d): fronto-orbital plate wide, coloration silver with pale brassy tomentum over uppermost 10%, appearing as pale brassy gold along upper 25% of occipital margin of eye; vitta wide, 1/3 width of fronto-orbital plate; ocellar setae lateraloclinate with a slight proclinate skew; with 3 proclinate orbital setae (middle proclinate orbital seta shorter and thinner than the other two) and 1 reclinate (often outwardly lateraloclinate) orbital seta; fronto-orbital plate with a sparse vestiture of short black setulae interspersed amongst frontal setae; fronto-orbital setulae not extending below pedicel; parafacial wholly silver tomentose, upper 1/3 of parafacial with small grouping of short black setulae in line with frontal setae, but separated from fronto-orbital plate; palps very slightly spathulate at tips, apically acutely inwardly curved, sparsely setulose along ventral surface, inner and upper surfaces appearing bare; arista dark brown-black. **Thorax** (Fig. 12a, b): black ground color with pale grey tomentum, brassy tones can occur under certain light; 4 postpronotal setae, anteriomost seta thin and weak almost hair-like; supra-alar setae 2:3; intra-alar setae 3:3; dorsocentral setae 3:3; acrostichal setae 3:3; katepisternum with 3 setae. Infrasquamal setae present. Scutellum black ground color, with tomentum concolorous with thorax; with five pairs marginal setae; apical scutellar setae 1/2 as long as subapical scutellars, sub-erect, arising above plane of marginal setae; basal scutellar 2/3 length of subapical setae; subapical setae convergent, lateral scutellar setae can be convergent or strongly divergent. Legs: black ground color; mid tibia with 2–3 strong anteroventral setae; tarsal claws and pulvilli as long as last tarsomere. Wings: R_1 bare, R_{4+5} setulose from node to crossvein r-m. **Abdomen** (Fig. 12a, b): ground color black; ST1+2 glabrous black, pale silver tomentum occupying anterior 75% of T3 and T4, T5 pale brassy-grey tomentose over entirety; tomentum extending to ventral surface of abdomen, but not completely; median marginal setae present on T3 and complete rows on T4 and T5.

Terminalia (Fig. 13): posterior margin of sternite 5 with a deeply excavated U-shaped median cleft (Fig. 13d); lateral lobes of sternite apically rounded, with a vestiture of short setae along disc, 2x as long along caudal margin; basal section 1/2 the length of apical lobes. Cerci, in posterior view, sharply pointed and slightly medially separated and parallel, almost touching medially, with a few short setae on basal half (Fig. 13a). In lateral view, curved and sharply tapered apically (Fig. 13b). Surstyli well-developed, stout basally in lateral view, broadly rounded; in posterior view, basally enlarged and apically straight.

Female (Fig. 14) as in male, except in the following aspects: **head** (Fig. 14c, d): fronto-orbital wider than males, just over 1/2 of head width entirely bright silver tomentose, frontal vitta 2/5 width of fronto-orbital plate; overall head of females 1.5x wider than that of males; legs (Fig. 14b): mid-tibia with 3–4 irregularly-sized strong anteroventral setae.




Figure 12.

Metaplagia svetlanakozikae sp. n. habitus images a-d: male, holotype n. DHJPAR0018732

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Diagnosis

Metaplagia svetlanakozikae sp. n. can be distinguished from all other *Metaplagia* by the following combination of traits: frontal vitta indistinct around ocellar triangle, infrasquamal setae present, fronto-orbital plate at pale gray or silver, with a silver parafacial and setulae on vein R_{4+5} not extending beyond crossvein r-m, surstyli rounded at tip. *Metaplagia svetlanakozikae* sp. n. can be separated from *Metaplagia occidentalis* by the presence of infrasquamal setae and from *Metaplagia lindarobinsonae* sp. n. by the presence of a small circular formation of setulae on the parafacial and not distinct row of setulae along the facial ridge. *Metaplagia svetlanakozikae* sp. n. is clearly distinguished by its COI sequence clustered within the Barcode Identification Number (BIN) [BOLD:AAD5456](#).




Figure 13.

Metaplagia svetlanakozikae sp. n. terminalia images a-d: male, paratype n. DHJPAR0018733

a: caudal view [doi](#)

b: lateral view [doi](#)

c: something [doi](#)

d: sternite 5, ventral view [doi](#)

Etymology

Metaplagia svetlanakozikae sp. n. is named in honor of Svetlana Kozik for her years of coordinating and administrating a plethora of problems and events for the undergraduate biology teaching laboratories of the Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania.

Distribution

Costa Rica, ACG, Guanacaste Province, 10–280 m elevation.

Ecology

Metaplagia svetlanakozikae sp. n. has been reared seven times from one species of Lepidoptera in the family Sphingidae: *Agrius cingulata*, in dry forest.

Identification keys

Key to the *Metaplagia* of North and Mesoamerica

1	Setulae present on wing vein R ₄₊₅ not extending beyond crossvein r-m	2
-	Setulae present on wing vein R ₄₊₅ extending beyond crossvein r-m	8
2	Fronto-orbital plate and parafacial entirely and uniformly gold	<i>paulinesaribasae</i> sp. n.
-	Fronto-orbital plate at most with some gold present and parafacial not gold	3
3	Fronto-orbital plate bearing some gold	<i>leahdenniseae</i> sp. n.
-	Fronto-orbital plate pale grey or silver (can include pale brassy) distinctly not gold	4
4	Infrasquamal setae absent	<i>occidentalis</i> Coquillett, 1895
-	Infrasquamal setae present	5
5	Frontal vitta indistinct around ocellar triangle, not reaching occiput	6
-	Frontal vitta wide and prominent extending around ocellar triangle to occiput	7
6	Parafacial setulae descending along parafacial in a row near facial ridge; surstyli sharp at tip	<i>lindarobinsonae</i> sp. n.
-	Parafacial setulae present as a small circular grouping along upper 1/3 of parafacial, not in a row near facial ridge; surstyli rounded at tip	<i>svetlanakozikae</i> sp. n.
7	Postpedicel just over 3 times as long as pedicel; claws and pulvilli shorter than last tarsomere	<i>brevicornis</i> Brooks, 1945
-	Postpedicel 2 times as long as pedicel; claws and pulvilli longer than last tarsomere	<i>orientalis</i> Townsend, 1915

8	Fronto-orbital plate gold on at least upper two thirds	<i>robinsherwoodae</i> sp. n.
-	Fronto-orbital plate entirely silver	9
9	Wing vein R ₁ bare throughout	<i>facialis</i> Reinhard, 1956
-	Wing vein R ₁ setulose throughout	<i>cordata</i> Reinhard, 1960




Figure 14.

Metaplagia svetlanakozikae sp. n. habitus images a-d: female, paratype n. DHJPAR0018722

a: dorsal view [doi](#)

b: lateral view [doi](#)

c: frontal view [doi](#)

d: three quarters view [doi](#)

Acknowledgements

We gratefully acknowledge the unflagging support of the team of ACG parataxonomists (Janzen et al. 2009, Janzen & Hallwachs 2011) who found and reared the specimens used in this study and the team of biodiversity managers who protect and manage the ACG forests that host these tachinids and their caterpillar hosts. The study has been supported by U.S. National Science Foundation grants BSR 9024770 and DEB 9306296, 9400829, 9705072, 0072730, 0515699 and grants from the Wege Foundation, International Conservation Fund of Canada, Jessie B. Cox Charitable Trust, Blue Moon Fund, Guanacaste Dry Forest Conservation Fund, Area de Conservación Guanacaste, Permian Global and University of Pennsylvania (DHJ & WH) and in-kind support from INBio and the Museo Nacional de Costa Rica. This study has also been supported by the Government of Canada through its ongoing support of the Canadian National Collection, Genome Canada, the Biodiversity Institute of Ontario and the Ontario Genomics Institute (2008–0GI–ICI–03) (MAS) and by a Discovery Grant from Natural Sciences and Engineering Research Council of Canada (MAS). AJF and DMW would also like to extend a very special thank you Dr. Torsten Dikow (USNM), Dr. Nigel Wyatt (BMNH), Dr. David Grimaldi (AMNH) and his technician Courtney Richenbacher who provided invaluable information relative to types in their museums. Acknowledgements to Dr. James E. O'Hara and Shannon Mahony Henderson, for allowing us to make use of their indispensable literature library and World Tachinidae Database (MW & AJF).

References

- Coquillett DW (1895) New genera and species of Tachinidae. Journal of the New York Entomological Society 3 (3): 97-10.
- Cortés RP, González CRA (1989) Generos Voríinos de Taquinidos chilenos (Diptera: Tachinidae, Voríini). Memórias do Instituto Oswaldo Cruz 84: 115-123. <https://doi.org/10.1590/s0074-02761989000800025>
- Cumming J, Wood DM (2017) Adult Morphology and Terminology. In: Kirk-Spriggs AH, Sinclair BJ (Eds) Manual of Afrotropical Diptera. Volume 1. Introductory chapters and keys to Diptera families. 1. South African National Biodiversity Institute, Pretoria, 89-133 pp.
- Fleming A, Wood DM, Smith MA, Hallwachs W, Janzen D (2014) Revision of the New World species of *Houghia* Coquillett (Diptera, Tachinidae) reared from caterpillars in Area de Conservación Guanacaste, Costa Rica. Zootaxa 3858 (1). <https://doi.org/10.11646/zootaxa.3858.1.1>
- Fleming A, Wood DM, Smith A, Dapkey T, Hallwachs W, Janzen D (2017) A new species of *Voria* Robineau-Desvoidy (Diptera: Tachinidae) from Area de Conservación Guanacaste in northwestern Costa Rica. Biodiversity Data Journal 5 URL: <http://zoobank.org/61701A00-E0C6-4415-B46C-D6D2EFF9FAB6>
- Fleming A, Wood DM, Smith A, Dapkey T, Hallwachs W, Janzen D (2019) A new species of *Trismegistomyia* Reinhard (Diptera: Tachinidae) from Area de Conservación Guanacaste in northwestern Costa Rica. Biodiversity Data Journal 7 URL: <http://zoobank.org/668D5B8B-4A97-4D27-8563-7DBF2BEB82D2>

- Guimarães JH (1977) Host-parasite and parasite-host catalogue of South American Tachinidae (Diptera). Arquivos de Zoologia 28 (3). <https://doi.org/10.11606/issn.2176-7793.v28i3p1-131>
- Herting B (1956) Das weibliche postabdomen der calypraten fliegen (Diptera) und sein merkmalswert für die systematik der gruppe. Zeitschrift für Morphologie und Ökologie der Tiere 45 (5): 429-461. <https://doi.org/10.1007/bf00389872>
- Ivanova NV, DeWaard JR, Hebert PN (2006) An inexpensive, automation-friendly protocol for recovering high-quality DNA. Molecular Ecology Notes 6 (4): 998-1002. <https://doi.org/10.1111/j.1471-8286.2006.01428.x>
- Janzen D, Hallwachs W, Blandin P, Burns J, Cadiou J, Chacon I, Dapkey T, Deans A, Epstein M, Espinoza B, Franclemont J, Haber W, Hajibabaei M, Hall JW, Hebert PN, Gauld I, Harvey D, Hausmann A, Kitching I, Lafontaine D, Landry J, LeMaire C, Miller J, Miller J, Miller L, Miller SE, Montero J, Munroe E, Green SR, Ratnasingham S, Rawlins J, Robbins R, Rodriguez J, Rougerie R, Sharkey M, Smith MA, Solis MA, Sullivan JB, Thiacourt P, Wahl D, Weller S, Whitfield J, Willmott K, Wood DM, Woodley N, Wilson J (2009) Integration of DNA barcoding into an ongoing inventory of complex tropical biodiversity. Molecular Ecology Resources 9: 1-26. <https://doi.org/10.1111/j.1755-0998.2009.02628.x>
- Janzen D, Hallwachs W (2011) Joining inventory by parataxonomists with DNA barcoding of a large complex tropical conserved wildland in Northwestern Costa Rica. PLoS One 6 (8). <https://doi.org/10.1371/journal.pone.0018123>
- Janzen D, Hallwachs W, Burns J, Hajibabaei M, Bertrand C, Hebert PN (2011) Reading the complex skipper butterfly fauna of one tropical place. PLoS One 6 (8). <https://doi.org/10.1371/journal.pone.0019874>
- Janzen D, Hallwachs W (2016) DNA barcoding the Lepidoptera inventory of a large complex tropical conserved wildland, Area de Conservacion Guanacaste, northwestern Costa Rica. Genome 59 (9): 641-660. <https://doi.org/10.1139/gen-2016-0005>
- Janzen D, Hallwachs W, Pereira G, Blanco R, Masis A, Chavarria MM, Chavarria F, Guadamuz A, Araya M, Smith MA, Valerio J, Guido H, Sanchez E, Bermudez S, Perez K, Manjunath R, Ratnasingham S, St. Jacques B, Milton M, DeWaard J, Zakharov E, Naik S, Hajibabaei M, Hebert PN, Hasegawa M (2020) Using DNA-barcoded Malaise trap samples to measure impact of a geothermal energy project on the biodiversity of a Costa Rican old-growth rain forest. Genome 63 (9): 407-436. <https://doi.org/10.1139/gen-2020-0002>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. Molecular Biology and Evolution 35 (6): 1547-1549. <https://doi.org/10.1093/molbev/msy096>
- O'Hara JE, Wood DM (1998) Tachinidae (Diptera): Nomenclatural review and changes, primarily for America North of Mexico. Canadian Entomologist (130) 751-774. <https://doi.org/10.4039/Ent130751-6>
- Ratnasingham S, Hebert PN (2007) BARCODING: bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). Molecular Ecology Notes 7 (3): 355-364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>
- Smith MA, Woodley NE, Janzen DH, Hallwachs W, Hebert PDN (2006) DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of

- parasitoid flies (Diptera: Tachinidae). Proceedings of the National Academy of Sciences 103 (10): 3657-3662. <https://doi.org/10.1073/pnas.0511318103>
- Smith MA, Wood DM, Janzen DH, Hallwachs W, Hebert PDN (2007) DNA barcodes affirm that 16 species of apparently generalist tropical parasitoid flies (Diptera, Tachinidae) are not all generalists. Proceedings of the National Academy of Sciences 104 (12): 4967-4972. <https://doi.org/10.1073/pnas.0700050104>
 - Smith MA, Rodriguez JJ, Whitfield JB, Deans AR, Janzen DH, Hallwachs W, Hebert PDN (2008) Extreme diversity of tropical parasitoid wasps exposed by iterative integration of natural history, DNA barcoding, morphology, and collections. Proceedings of the National Academy of Sciences of the United States of America 105 (34): 12359-64. <https://doi.org/10.1073/pnas.0805319105>
 - Smith MA, Fernandez-Triana J, Roughley R, Hebert PDN (2009) DNA barcode accumulation curves for understudied taxa and areas. Molecular Ecology Resources 9 Suppl s1: 208-16. <https://doi.org/10.1111/j.1755-0998.2009.02646.x>
 - Smith MA (2012) Hyperparasitoid wasps (Hymenoptera, Trigonaliidae) reared from dry forest and rain forest caterpillars of Area de Conservacion Guanacaste, Costa Rica. Plazi.org taxonomic treatments database <https://doi.org/10.15468/v4scdj>
 - Stireman J, Cerretti P, O'Hara J, Blaschke J, Moulton J (2019) Molecular phylogeny and evolution of world Tachinidae (Diptera). Molecular Phylogenetics and Evolution 139 <https://doi.org/10.1016/j.ympev.2018.12.002>
 - Tamura K (1992) Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G + C-content biases. Molecular Biology and Evolution 9: 678-687. <https://doi.org/10.1093/oxfordjournals.molbev.a040752>
 - Tschorchnig VHP (1985) Taxonomie forstlich wichtiger Parasiten: Untersuchungen zur Struktur des männlichen Postabdomens der Raupenfliegen (Diptera, Tachinidae)*. Stuttgarter Beiträge zur Naturkunde Serie A (Biologie) A (383): 1-137.
 - Wood DM (1987) Tachinidae. In: McAlpine JF, Peterson BV, Shewell GE (Eds) Manual of Nearctic Diptera. 2. Agriculture Canada, Ottawa, Ontario, 1193–1269. pp.
 - Wood DM, Zumbado M (2010) Tachinidae (Tachinid Flies, Parasitic Flies). In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE, Zumbado MA (Eds) Manual of Central American Diptera Vol. 2. 2. National Research Council of Canada, Ottawa, 728 pp. [In English]. [ISBN 978-0-660-19958-0].
 - Yu G, Smith D, Zhu H, Guan Y, Lam TT (2017) ggtree: an rpackage for visualization and annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology and Evolution 8 (1): 28-36. <https://doi.org/10.1111/2041-210x.12628>