



REVIEW

Recurrence Rates and Pharmacological Treatment for Hemorrhoidal Disease: A Systematic Review

Varut Lohsiriwat · Parvez Sheikh · Robert Bandolon ·

Dong-Lin Ren · April Camilla Roslani · Kassandra Schaible ·

Andreas Freitag · Monique Martin · Pelin Yaltirik · Philippe Godeberge

Received: July 8, 2022 / Accepted: October 5, 2022 / Published online: November 4, 2022
© The Author(s) 2022

ABSTRACT

Introduction: Hemorrhoidal disease (HD) is characterized by prolapse of the inflamed and bleeding vascular tissues of the anal canal. Although HD is associated with a high recurrence rate, there is a lack of understanding around interventions that can reduce

recurrence and improve outcomes for patients. As such, a systematic literature review (SLR) was conducted to summarize evidence on epidemiology, recurrence, and efficacy of interventions in HD.

Methods: Real-world evidence (RWE) studies evaluating the incidence, prevalence, or recurrence of HD, as well as SLRs including a meta-analytic component reporting on the efficacy of systemic or topical pharmacological treatments for adults with HD, were included. Systematic

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12325-022-02351-7>.

V. Lohsiriwat (✉)
Department of Surgery, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand
e-mail: bolloon@hotmail.com

P. Sheikh
Department of Colorectal Surgery, Saifee Hospital, Mumbai, India

R. Bandolon
Southern Philippines Medical Center, Davao, Philippines

D.-L. Ren
Department of Colorectal Surgery, The Sixth Affiliated Hospital (Gastrointestinal and Anal Hospital), Sun Yat-Sen University, Guangzhou, China

A. C. Roslani
Department of Surgery, Faculty of Medicine, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

K. Schaible
Evidera, Waltham, MA, USA

A. Freitag · M. Martin
Evidera, London, UK

P. Yaltirik
Internal Medicine and Neuropsychiatry, Global Medical and Patient Affairs (GMPA), Servier, Suresnes, France

P. Godeberge
Unité d'endoscopie et de proctologie, Clinique du Trocadéro, Paris, France

searches were conducted in MEDLINE, Embase, Cumulative Index to Nursing and Allied Health Literature, and the Cochrane Database of Systematic Reviews.

Results: The SLR identified 44 eligible publications. Consistent data were limited on the epidemiology of HD or HD recurrence. Specifically, incidence and prevalence reported across geographies were impacted by differences in data collection. Reported risk factors for HD were sedentary behavior, constipation, male gender, and age. Twenty-three RWE studies and one meta-analysis reported HD recurrence rates ranging from 0 to 56.5% following surgery or phlebotonics, with most ($n = 19$) reporting rates

of 20% or less. In addition to time since treatment, risk factors for recurring disease were similar to those for HD in general. With respect to treatment, micronized purified flavonoid fractions significantly improved the main symptoms of HD compared to other pharmacological treatments.

Conclusion: The SLRs did not identify any RWE studies reporting recurrence in patients receiving systemic or topical treatments, highlighting the need for future research in this area. Further, more studies are needed to understand the optimum duration of medical treatment to prevent recurrence.

Graphical abstract:

Advances in Therapy

PEER-REVIEWED FEATURE

RECURRENCE RATES AND PHARMACOLOGICAL TREATMENT FOR HEMORRHOIDAL DISEASE: A SYSTEMATIC REVIEW

Varut Lohsiriwat, Parvez Sheikh, Robert Bandolon, Dong-Lin Ren, April Camilla Roslani, Kassandra Schaible, Andreas Freitag, Monique Martin, Pelin Yaltrik, Philippe Godeberge

Introduction

HD is a common anorectal condition, though data on the rates of recurrence and interventions that can reduce recurrence are limited

Objective

To conduct a systematic literature review in order to summarize evidence on the epidemiology and recurrence of HD in adults, and the effectiveness of available interventions

Methods

A systematic literature search of databases including MEDLINE, Embase, and the Cochrane Database was conducted on 13 August 2019 to identify RWE studies on the epidemiology of HD and SLRs with meta-analyses on the efficacy/effectiveness of systemic or topical pharmacological treatment for HD. PRISMA and Cochrane reporting guidelines were followed

Results

2,037
records identified

1,579
screened

443
full text assessed

+

69
assessed (other sources)

44
records included

39

RWE studies

Up to 56.5%

recurrence rate*

5

meta-analyses

≤ 20%

recurrence rate in 19/23 studies

Treatments

- > HD recurrence rates following systemic treatment were infrequently reported
- > Lower recurrence rates and superior efficacy (bleeding, discharge/leakage, pain) with pharmacological treatment versus placebo or no treatment
- > In one meta-analysis, MPFFs were the only phlebotonic agents to statistically significantly reduce recurrence risk vs no treatment/placebo (low-quality evidence)

Risk factors for recurrence

- > Time since treatment
- > Similar to HD risk factors:
 - ✓ Sedentary behavior
 - ✓ Constipation
 - ✓ Chronic venous disease
 - ✓ CEAP class
 - ✓ Male gender
 - ✓ Age

Conclusion

Among venoactive drugs, MPFF has shown some evidence supporting a benefit in the prevention of HD recurrence. Further research on the risk of recurrence and the impact of treatments in the long term is needed.

CEAP: Classification System for Chronic Venous Disorders; HD, hemorrhoidal disease; MPFFs, micronized purified flavonoid fractions; RWE, real world evidence; SLR, systematic literature review. *After surgery/phlebotonic treatment; data from 23 RWE studies and 1 meta-analysis

This graphical abstract represents the opinions of the authors. For a full list of declarations, including funding and author disclosure statements, and copyright information, please see the full text online.

PLAIN LANGUAGE SUMMARY

Patients with hemorrhoidal disease (HD) can experience recurring disease following a period of improvement or remission. It is not well established how often this might happen, who is at greatest risk, or which treatments can reduce this risk. In this study, a systematic literature review (SLR) was conducted to summarize evidence on the occurrence and recurrence of HD, as well as treatment effectiveness. Several literature databases were searched for articles that described real-world evidence (RWE) studies reporting the epidemiology or recurrence of HD as well as published SLRs that combined the results of multiple studies (meta-analyses) on treatment for adults with HD. Forty of 2037 articles identified by the search were considered relevant, and four others identified by clinicians were also included (total = 44; 39 RWE, 5 meta-analyses). Review of the RWE articles revealed that HD epidemiology was determined differently between studies. Only 23 reported recurrence rates (up to 56.5%) after surgery or treatment with phlebotonic drugs (drugs that improve blood flow in veins). Most (19/23) reported recurrence rates of 20% or less. Risk factors for recurrence were similar to usual HD risk factors (e.g., constipation, male gender, age) in addition to time since treatment. Phlebotonic agents, including those made from plant extracts (micronized purified flavonoid fractions, MPFFs) improved hemorrhoidal symptoms compared with placebo or no treatment. In one meta-analysis, MPFF was the only phlebotonic to significantly reduce recurrence risk versus no treatment or placebo. Overall, more research is needed to compare treatments and determine optimal treatment duration to prevent recurrence.

Keywords: Hemorrhoids; MPFF; Phlebotonics; Recurrence; Risk factors

Key Summary Points

A systematic literature review was conducted to summarize evidence on the epidemiology and recurrence of hemorrhoidal disease.

Consistent data were limited on the topics of interest.

Recurrence rates varied across the included real-world evidence studies, ranging from 0% to 56.5% following surgery or other procedural treatment.

Additional research is needed to evaluate comparative efficacy for reduction in recurrence between two active treatments.

DIGITAL FEATURES

This article is published with digital features, including video abstract and infographic, to facilitate understanding of the article. To view digital features for this article go to <https://doi.org/10.6084/m9.figshare.21277593>.

INTRODUCTION

Hemorrhoidal disease (HD) is characterized by prolapse of the inflamed and bleeding vascular tissues of the anal canal and is associated with a high recurrence rate as well as a considerable burden to individuals.

International consensus on individualized, effective disease management has not been reached. Treatment options for HD range from conservative methods (e.g., dietary and lifestyle modifications) and medical management (e.g., venoactive drugs [also known as phlebotonics]) to non-invasive procedures such as sclerotherapy, rubber band ligation, infrared coagulation, radiofrequency ablation, or invasive surgery procedures. Although recurrence commonly occurs in people with HD, there is a lack of understanding around the rates of recurrence,

and interventions that can reduce this and improve outcomes for patients. A further understanding of recurrence rates in HD could support clinical decision-making around the best treatments.

The objective of this systematic literature review (SLR) was to summarize studies evaluating the epidemiology of HD, including incidence, prevalence, recurrence rates, and risk factors for HD and recurrence of HD, and to assess the efficacy/effectiveness of various systemic or topical pharmacological treatments for adults with HD on the basis of previously published SLRs that included a meta-analytic component.

METHODS

This SLR was conducted following standards from the Cochrane Collaboration [1] and reporting standards of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [2]. The topics of interest reported herein were part of a broader SLR focused on the overall burden of HD, which included literature on diagnosis, epidemiology, humanistic burden, economic burden, and disease management, as well as clinical efficacy and safety of treatment for HD.

This article is based on previously conducted studies; none of the authors performed any new studies involving human participants or animals. No analyses were conducted.

Eligibility Criteria

Studies meeting the following criteria were included: (1) Real-world studies reporting epidemiology (including incidence, prevalence, recurrence rates, and risk factors) of adults with HD as reported for key regions of interest in Europe, North America, Asia–Pacific, as well as Brazil, Egypt, and Türkiye (see Appendix in the supplementary material); (2) SLRs that included a meta-analytic component reporting on the efficacy/effectiveness of systemic or topical pharmacological treatments (alone or in combination with other therapies) for adults with HD.

SLRs focusing on invasive or surgical procedures were excluded. There were no restrictions with respect to disease stage. Epidemiology studies were limited to those published in 2009 or later; there was no time limit for published SLRs.

Further details regarding the population, intervention, comparator, outcomes, and study designs of interest are presented in the Appendix.

Study Identification

Systematic database searches were conducted via Ovid SP on 13 August 2019, including MEDLINE, Embase, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Cochrane Database of Systematic Reviews. The search strategies (Tables S1, S2, and S3 in the supplementary material) were designed using a combination of Medical Subject Heading, Emtree, and free-text terms for HD paired with validated filters to identify studies reporting on all topics of interest in the broader review (epidemiology, diagnostics, quality of life, and clinical and economic burden). The search strategies are provided in the online Appendix.

Study Selection

Each title and abstract were reviewed by one investigator to determine the article's suitability for inclusion in the review, according to the defined inclusion and exclusion criteria (Table S4 in the supplementary material). A second, independent investigator reviewed 10% of the abstracts as a quality check. Discrepancies between the first and second reviewers were resolved by a third, senior investigator. For abstracts that were deemed relevant, the corresponding full-text articles were retrieved for further evaluation. Each full-text article was reviewed by one investigator. All rejected articles were assigned a reason for exclusion and were validated by a second, independent investigator. Discrepancies were resolved by a third, senior investigator.

Data Extraction and Reporting

Data extraction was conducted by one investigator with all extractions validated by a second investigator. Study, population, and treatment characteristics, as well as outcomes of interest, from selected full-text articles were extracted into a pre-defined data extraction table. Results were summarized qualitatively; no analyses were performed.

The quality of the included real-world evidence (RWE) studies was assessed using the modified Downs and Black checklist, and each was scored out of 28 [3].

Assessment of Risk of Bias

The methodological quality of the included SLRs was assessed using the A MeaSurement Tool to Assess systematic Reviews (AMSTAR) 2 tool [4]. The AMSTAR 2 checklist includes 16 items to assess specific elements of the conduct of SLRs, seven of which are considered critical. The quality of included RWE studies reporting

on recurrence rates was assessed using the modified Downs and Black checklist [3].

RESULTS

PRISMA

Systematic searches for the review identified 2037 records from electronic literature databases. After the removal of 458 duplicates, 1579 remained for title and abstract screening. Of these, 443 publications were deemed eligible for screening at the full-text level, of which 40 met the inclusion criteria of the review. An additional four records were identified by clinical experts, resulting in 44 total records ultimately included in the review (Fig. 1).

Characteristics of Included Studies

Thirty-nine RWE studies were identified which reported on the epidemiology of HD, of which two reported incidence, five reported




Fig. 1 PRISMA diagram. *CDSR* Cochrane Database of Systematic Reviews, *RWE* real-world evidence, *SLR* systematic literature review

prevalence, and 14 reported risk factors of HD. A snapshot of included studies is reported in Table 1. Five SLRs provided data on the clinical efficacy or safety of treatment for HD (Table 2). Twenty-three of the included RWE studies also reported on recurrence of disease, all of which reported recurrence following procedural treatments. A single SLR reported recurrence rates following systemic treatment.

Twenty-one RWE studies took place in Europe, five in China, five in the USA, two in Australia, two in Türkiye, and one each in Taiwan and Brazil. One RWE study was international, and one was conducted in the USA and Puerto Rico (Fig. 2). The included SLRs were published between 2006 and 2018. Databases searched within the SLRs included PubMed/MEDLINE ($n = 4$), Embase ($n = 4$), Cochrane Library ($n = 4$), and CINAHL ($n = 3$). All SLRs included only clinical trial evidence.

The quality of most RWE studies was in the 10- to-15-point range out of 28. Five SLRs were assessed per AMSTAR 2 criteria—one was judged to be viewed with low confidence, two with critically low confidence, and two with moderate confidence. Three of the five SLRs failed to assess the impact of publication bias on their results (AMSTAR 2 criterion 15) and did not have a protocol available (AMSTAR 2 criterion 3).

Epidemiology

Thirty-nine RWE studies reporting on the incidence, prevalence, risk factors, and recurrence of HD were included in the SLR (Table S5 in the supplementary material). The characteristics of 10 of the most representative epidemiology studies, by geography, are reported in Table 1.

Prevalence and Incidence of HD

Prevalence based on 2011 data from a convenience sample of 39 general practices across metropolitan France and some overseas territories was 5.84% [5]. The annual prevalence rate was 1.2% among 18.9 million adults with at least one HD-related claim enrolled in the US Truven claims database in 2014 [14].

Prevalence rates were greater in at-risk individuals. Increased sitting (≥ 9 h/day) was associated with a nonsignificant increase in HD prevalence in a multivariate analysis (odds ratio [OR] 1.18; 95% CI 0.92, 1.50; $p = 0.08$) [12]. In Brazil, HD was approximately 4.5 times more prevalent in women with vaginal deliveries compared to those who underwent caesarean deliveries [10]. Caesarean deliveries were found to be protective against hemorrhoids with a prevalence ratio of 0.16 (95% CI 0.04, 0.57) when adjusted for age, type of provider, number of prenatal consultations, parity, and hypertension and diabetes during pregnancy [10].

Abramowitz et al. presented incident cases diagnosed and reported by 39 primary care physicians in France during 2.5 days of consultation in 2010. Among 1079 patients, 74 presented with a new diagnosis of HD (6.9% or 69 per 1000 persons) [5]. Lin et al. reported on the new HD diagnosis (International Classification of Diseases, Ninth Revision, Clinical Modification: 455) among patients with and without chronic obstructive pulmonary disease (COPD) as reported in a Taiwanese national database between 2000 and 2003 and followed up until 2011 [9]. Overall incidence among patients with COPD was approximately 1.6 times greater than in patients without COPD (average annual incidence 11.5 vs. 7.1 per 1000 person-years) [9]. These findings were consistent between men and women, and across different age groups [9]. Neither study was designed specifically to measure HD incidence in the general population, and thus estimates are likely to be imperfect. The dissimilar estimates may be due to methodology of data collection, or may reflect true differences; additional information is needed to understand the real incidence (69/1000 vs. 7.1/1000) of HD.

Risk Factors for HD

A cross-sectional study was conducted by Peery et al. in patients recruited from 11 centers in the USA and Puerto Rico from 2004 to 2008 with satisfactory preparation for colonoscopy; results found that constipation, straining during bowel movements, incomplete bowel emptying, and

Table 1 Snapshot of representative epidemiology studies

Author, year	Country	Sample size	Period	Study type	Patients	Incidence	Prevalence	Recurrence rates	Risk factors	Quality score
Abramowitz et al. 2014 [5]	France	1079	2010 (February–September)	Cross-sectional	All visiting GP during period	X	X			10
Conaghan and Farouk 2009 [6]	England	203	September 2004–June 2007	Prospective cohort study	With symptomatic grade 2 or 3 hemorrhoids who had failed rubber band ligation and were treated with DGHAL			X		13
Godeberge et al. 2020 [7]	International ^a	5617	January 2015–December 2016	Cross-sectional	Adults attending a consultation for hemorrhoidal complaints (questionnaire completed by physician)			X	X	14
Ince et al. 2012 [8]	Türkiye	200	2011 (January–December)	Retrospective case control study	Women aged < 25 years, with stage II–IV HD who underwent a surgical operation in 2011				X	9
Lin et al. 2017 [9]	Taiwan	154,518	From 2000 and 2003 to 2011	Case control study	COPD cohort: all new diagnoses; COPD (ICD-9-CM: 490–492, 494, 496). Non-COPD matching by age group, gender, and index date	X			X	15
Mascarello et al. 2018 [10]	Brazil	4244	2004–2010	Prospective cohort study	All births occurring in Pelotas city		X		X	17
Peery et al. 2015 [11]	USA and Puerto Rico	1074	2004–2008	Cross-sectional	Underwent colonoscopy as part of a colorectal adenoma prevention trial				X	15
Peeters et al. 2013 [12]	Australia	10,286 (2004) 10,128 (2007) 9529 (2010)	1996–2010	Population-based prospective cohort study	Participants in the Australian Longitudinal Study on Women's Health completed mail surveys		X		X	13
Ratto et al. 2010 [13]	Italy	170	July 2005–October 2008	Prospective cohort study	With HD undergoing THD			X		15

Table 1 continued

Author, year	Country	Sample size	Period	Study type	Patients	Incidence	Prevalence	Recurrence rates	Risk factors	Quality score
Yang et al. 2019 [14]	USA	227,638	2014 (January–December)	Cross-sectional study	With one hemorrhoid-related claim in claims database		X			10

COPD chronic obstructive pulmonary disease, *DGHAL* Doppler-guided hemorrhoid artery ligation, *GP* general practitioner, *HD* hemorrhoidal disease, *ICD-9-CM* International Classification of Diseases, Ninth Revision, Clinical Modification, *THD* transanal hemorrhoidal dearterialization
^aRussia, India, Pakistan, Mexico, Belgium, Slovenia, and Thailand

hard stools were associated with increased risk of hemorrhoids [11]. Additionally, while not statistically significant, high total fiber intake was associated with a reduced risk of hemorrhoids (OR 0.80; 95% CI 0.64, 1.01) [11]. While neither physical activity nor weight was associated with decreased risk, the highest quartile of sedentary behavior (mean 656 min/day) was associated with a lower age- and sex-adjusted risk than the lowest quartile (mean 176 min/day) [11]. An international multivariate analysis by Godeberge et al. examining risk factors in 5617 adults seeking consultation for hemorrhoidal complaints demonstrated that hemorrhoids were significantly associated with age, chronic venous disease (CVD) Classification System for Chronic Venous Disorders (CEAP) class, constipation, and male gender ($p < 0.0001$ for all) [7].

Clinical Efficacy and Treatments

Five SLRs and meta-analyses of clinical trials reported on the clinical efficacy and safety of treatments for HD including: one SLR for micronized purified flavonoid fractions (MPFF); one SLR for metronidazole; two SLRs for phlebotonics in general in patients with HD; and one SLR for conservative treatments including dietary modifications, bowel transit modifications, local treatments, and drugs in patients with symptomatic hemorrhoids during pregnancy and the puerperium [15–19].

Controls typically included placebo, no treatment, or usual care. One SLR included phlebotonics as a comparator [16] and one SLR did not specify the control group [17]. SLRs differed in terms of the number of trials and patients included, ranging from two to 24 randomized controlled trials (RCT; or 150–2344 patients). Overall, the available evidence across the SLRs was limited and heterogenous, with no single meta-analysis reporting all outcomes of interest.

The SLR by Wanis et al. examined the effect of metronidazole for the treatment of HD pain and found inconsistent efficacy [19]. While metronidazole resulted in statistically significantly lower pain scores than placebo or usual

Table 2 Characteristics of included SLRs

Author, year	Treatments compared	Outcomes	# of studies included	# of patients included	PubMed/MEDLINE	Embase	Cochrane Library	CINAHL	Search cut-off point	Quality
Alonso-Coello et al. 2006 [15]	Phlebotonics vs. placebo/no treatment	Global improvement, bleeding, pain, symptoms, recurrence, other	14 RCTs	1514	✓	✓	✓	✓	April 2005	Low
Aziz et al. 2018 [16]	MPEF vs. any control	Bleeding, pain, symptoms, safety	10 RCTs	1164	✓	✓	✓	✓	April 2018	Critically low
Perera et al. 2012 [17]	Phlebotonics vs. placebo/control interventions	Safety, symptoms, pain, compliance	24 RCTs	2344	✓	✓	✓	✓	September 2011	Moderate
Quijano and Abalos 2012 [18]	Conservative treatments vs. placebo/no treatment	Treatment response, safety, maternal outcomes	2 RCTs	150			✓ ^a		March 2010	Moderate
Wanis et al. 2017 [19]	Metronidazole vs. placebo/usual care	Postoperative pain, time to return to work	5 RCTs	337	✓	✓	✓		July 2016	Critically low

CINAHL Cumulative Index of Nursing and Allied Health Literature, *MPEF* micronized purified flavonoid fraction, *RCT* randomized controlled trial, *SLR* systematic literature review

^aCochrane Pregnancy and Childbirth Group's Trials Register



Fig. 2 Geographical distribution of included RWE studies

care on the first and fourth days following surgery (standard mean difference [SMD] -0.87 ; 95% CI $-1.73, -0.015$; $p = 0.046$ and -1.43 ; 95% CI $-2.83, -0.037$; $p = 0.044$, respectively) as well as a shorter time to return to normal activities (SMD -0.76 ; 95% CI $-1.43, -0.088$; $p = 0.027$), these results became not significant when the largest trial with the highest risk of bias was removed from the pooled analysis [19].

The SLR by Quijano and Abalos assessed the effectiveness of conservative treatments including dietary modifications, bowel transit modifications, local treatments, and drugs in patients with symptomatic hemorrhoids during pregnancy and the puerperium. Only two studies with rutosides were eligible for the analysis. Rutosides were more likely to produce a treatment response, defined as improvement in symptoms, at 4 weeks post pregnancy when compared with placebo or no treatment (risk ratio [RR] 0.07; 95% CI 0.03, 0.20; $p < 0.0001$) [18]. However, as there was no difference in maternal outcomes, and the safety data were insufficient to rule out clinically important harms, the authors did not recommend the use of rutosides until further evidence was available [18].

The meta-analysis by Alonso-Coello et al. compared phlebotonics including MPFFs, troxerutin, diosmin, hydroxyethylrutosides with no treatment, or placebo in the treatment of

symptomatic HD. Patients treated with phlebotonics were more likely to experience a reduction in bleeding (RR 0.33; 95% CI 0.19, 0.57; $p < 0.001$) [15] and a significant reduction in pain (RR 0.35; 95% CI 0.18, 0.69; $p < 0.001$) and itching (RR 0.65; 95% CI 0.44, 0.97; $p = 0.01$). There was no difference in effect for safety outcomes [15].

Another SLR on phlebotonics by Perera et al., which included mainly studies with MPFFs (but also studies with rutosides, diosmin, troxerutin, pycnogenol, and calcium dobesilate), found a potential benefit in the treatment of acute HD crises as well as a benefit in alleviating post-hemorrhoidectomy symptoms [17]. In this analysis of 24 trials, phlebotonics, particularly MPFFs, statistically significantly improved bleeding (OR 0.12; 95% CI 0.04, 0.37; $p = 0.0002$), bleeding post-hemorrhoidectomy (OR 0.18; 95% CI 0.06, 0.58; $p = 0.004$), discharge and leakage (OR 0.12; 95% CI 0.04, 0.42; $p = 0.0008$), pruritus (OR 0.23; 95% CI 0.07, 0.79; $p = 0.02$), and overall symptoms (OR 15.99; 95% CI 5.97, 42.84; $p < 0.00001$) [17]. The use of phlebotonics was associated with a beneficial but not statistically significant effect for pain (OR 0.11; 95% CI 0.01, 1.11; $p = 0.06$) and for pain scores post hemorrhoidectomy (SMD -1.04 ; 95% CI $-3.21, 1.12$; $p = 0.35$). They were also associated with less analgesic consumption (OR 0.54; 95% CI 0.30, 0.99;

$p = 0.05$) as a reflection of their benefit in alleviating pain caused postoperatively.

Finally, the SLR by Aziz et al. specifically examined MPFFs and found a statistically significant improvement in bleeding compared with placebo (RR 1.46; 95% CI 1.10, 1.93; $p = 0.008$), without a difference between MPFFs and placebo for pain improvement [16]. Itching was improved significantly with MPFFs in two of three studies. As a result of heterogeneity in study designs, endpoints were limited for pooled meta-analyses. Aziz et al. did not perform a pooled analysis of efficacy outcomes comparing MPFFs to other phlebotonics [16].

Recurrence Rates

Twenty-three RWE studies and one meta-analysis [15] reported HD recurrence rates following surgery or phlebotonics. Recurrence rates in the included studies typically ranged from 0 [20, 21] to 56.5% [7] with most ($n = 19$) reporting rates of 20% or less. Time since procedure was the main driver of this range. Of the studies reporting recurrence, only two prospectively evaluated recurrence rates over time. Ratto et al. found that recurrence of HD requiring surgery after transanal hemorrhoidal dearterialization occurred in 4.1% of patients over a median follow-up of 11.5 months [13]. Conaghan and Farouk found that, among patients with symptomatic grade 2 or 3 HD who failed rubber band ligation and were treated with Doppler-guided hemorrhoid artery ligation, 20% had recurrent symptoms at a median follow-up of 18 months [6].

In addition to time since treatment, risk factors for recurring disease were similar to risk factors for HD in general. In their study of 5617 patients presenting with HD complaints in clinical practice across multiple countries, Godeberge et al. found 56.5% of adults attending a consultation for HD complaints had a previous history of HD [7]. The factors associated with HD recurrence after previous consultation for HD in multivariate analysis included constipation (OR 2.20; 95% CI 1.96, 2.46), age group (OR 2.11; 95% CI 1.68, 2.65 for the comparison of 18–34 years vs. > 65 years), CVD

CEAP class (OR 3.75; 95% CI 1.30, 10.90 for the comparison of CEAP C0a vs. C6), body mass index (BMI) category (OR 2.34; 95% CI 1.51, 3.64 for the comparison of 12–18 years vs. ≥ 31 kg/m²), and male gender (OR 1.25; 95% CI 1.11, 1.41) [7]. Among women, having given birth and number of births were the most important risk factors ($p < 0.0001$), followed by constipation ($p < 0.001$), age group ($p < 0.0001$), presence of CVD ($p = 0.0089$), and BMI category ($p = 0.0123$) [7]. Among men, constipation was the most important risk factor ($p < 0.0001$) followed by age group ($p < 0.0001$), BMI category ($p = 0.0011$), and presence of CVD ($p < 0.0001$) [7]. Gender had no influence on likelihood of hemorrhoid recurrence in univariate analysis ($p = 0.1539$) [7].

In the meta-analysis conducted by Alonso-Coello et al., phlebotonics such as MPFFs, troxerutin, diosmin, hydroxyethylrutinosides were compared with no therapy or placebo in the treatment of symptomatic HD. The risk of recurrence was statistically significantly reduced across four studies only for those treated with MPFFs. These recurrences were evaluated for short-to-medium follow-up (2–6 months). The pooled estimate was 47% RR reduction in favor of MPFFs (RR 0.53; 95% CI 0.41, 0.69) [15].

DISCUSSION

This SLR provided evidence regarding the epidemiology of HD and the efficacy of various pharmacological treatments for the condition. Consistent data were limited on the epidemiology of HD or HD recurrence. Specifically, incidence and prevalence reported across geographies were impacted by differences in how data were collected and recorded and sampling methodology. There was also a lack of epidemiology data for HD, and the studies included were heterogeneous in terms of design and target populations. These differences across epidemiological studies hindered the ability to accurately assess the current burden of disease, which is generally acknowledged to be substantial.

A number of studies reported on independent and statistically significant risk factors for HD in general, many of which examined well-established risk factors (e.g., comorbidities and lifestyle aspects). Similar risk factors were identified for HD recurrence, such as constipation and age.

Five included SLRs reported limited information on the efficacy and safety of treatments as a result of heterogeneity of study designs. Generally speaking, metronidazole and rutosides failed to demonstrate superior efficacy when compared with placebo or no treatment, while among phlebotonics, MPFFs significantly improved the main symptoms of HD such as bleeding, discharge/leakage, or pruritus.

These outcomes were supported by a recent meta-analysis which was published after the search period. It assessed 11 RCTs in which MPFF treatment was compared to placebo or no treatment for acute HD or for relief of symptoms after patients had undergone medical management or a surgical procedure to remove hemorrhoids. On the basis of findings from qualitative analysis, MPFFs were reported in most studies to be beneficial in treating bleeding, pain, pruritus, anal discharge/leakage, and tenesmus, and in overall improvement. Quantitative meta-analysis indicated that MPFF treatment provided significant benefits for bleeding (OR 0.082; 95% CI 0.027, 0.250; $p < 0.001$), discharge/leakage (OR 0.12; 95% CI 0.04, 0.42; $p < 0.001$), and overall improvement according to patients (OR 5.25; 95% CI 2.58, 10.68; $p < 0.001$) and investigators (OR 5.51; 95% CI 2.76, 11.00; $p < 0.001$). MPFF also tended to decrease pain (OR 0.11; 95% CI 0.01, 1.11; $p = 0.06$). These results suggested MPFF treatment can improve the most important symptoms of HD [22].

While safety events were not a main outcome of interest, three SLRs that focused on phlebotonics ($n = 2$) or conservative treatment in general ($n = 1$) found no significant difference between active treatment and placebo with respect to adverse events. Overall, systemic treatment was generally well tolerated with no safety concerns.

Recurrence rates across the included RWE studies ranged from 0 to 56.5% following

surgery or other procedural treatment. Factors such as history of HD, time from procedure, and type of procedure played a significant role in HD recurrence. However, differences among populations observed across the RWE studies precluded the ability to draw any strong, meaningful conclusions from the literature. A better understanding of the risk factors and the pathophysiology that trigger the recurrence of HD will help to alleviate further burden on patients.

An included meta-analysis of RCT evidence which involved MPFFs, troxerutin, diosmin, hydroxyethylrutosides found that the risk of recurrence was statistically significantly reduced for those treated with MPFFs compared to placebo or no treatment with 47% risk reduction in favor of MPFFs. This was based on data from four studies out of 14 included RCTs. MPFFs were the only phlebotonic with proven efficacy on the reduction of recurrence of hemorrhoidal symptoms. This finding was further supported by a recently published consensus document which states that the use of MPFFs can help prevent HD recurrence, especially recurrent bleeding [23].

This SLR identified a few gaps in the literature that require additional research. Though 39 RWE studies were included, the SLRs did not identify any reporting recurrence in patients receiving systemic or topical treatments, highlighting the need for more research. Among phlebotonics, MPFF was the most investigated product with the highest quality of evidence. Additional research is also needed to evaluate comparative efficacy for reduction in recurrence between two active treatments. Further, more studies are needed to understand the optimum duration of medical treatment to prevent recurrence.

This research has several limitations; first, there was a general lack of high-quality evidence for RWE studies and limited data available from published SLR/meta-analyses of HD. In particular, only one SLR examined the rate of recurrence of HD after treatment. Second, considerable differences were found across the included studies in how outcome data were defined, collected, measured, and reported, making a direct comparison of treatments or

geographic areas difficult. For this reason, consistent measurements, or a set of core outcomes, for efficacy or effectiveness should be used in future studies. Third, the evidence summarized in this report was selected on the basis of a number of additional selection criteria (e.g., sample size, geographic area of interest) to support a meaningful synthesis of results. Finally, the evidence was limited to studies from 2009 onward, with the exception of clinical trials, limiting the ability to identify long-term trends in the data.

This SLR brings added value with several strengths. First, the SLR was conducted in alignment with guidance by the Cochrane Collaboration along with reporting guidelines established by PRISMA. The search strategy used for this SLR was paired with gray literature searches to ensure all relevant evidence was identified. The SLR approach allows for synthesis of evidence across multiple geographies and time periods, creating a broad overview of recurrence rates in HD.

CONCLUSIONS

This SLR identified risk factors for HD and HD recurrence that were already well established. Recurrence rates were not commonly reported, though time from treatment and type of treatment were considered important factors. Patients receiving pharmacological treatment generally had lower recurrence rates, along with superior efficacy outcomes such as improvements in bleeding, discharge/leakage, or pain when compared to patients with non-pharmacological or no treatment. MPFF was the most investigated product among phlebotonics and had the highest quality of evidence, demonstrating that MPFF treatment improves the most important symptoms of HD. Among venoactive drugs, MPFF has shown some evidence supporting a benefit in the prevention of HD recurrence. However, further research on the risk of recurrence and the impact of treatments in the long term is needed.

ACKNOWLEDGEMENTS

Funding. This study and the journal's Rapid Service Fee were sponsored by Servier.

Authorship. All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published.

Author Contributions. Andreas Freitag, Cassandra Schaible, Monique Martin, and Pelin Yaltirik contributed to the concept and design of the study as well as drafting of the manuscript. April Camilla Roslani, Dong-Lin Ren, Philippe Godeberge, Parvez Sheikh, Robert Bandolon, and Varut Lohsiriwat contributed to the drafting of the manuscript. All authors read and approved the final version of the manuscript.

Disclosures. Pelin Yaltirik is an employee of Servier. Andreas Freitag was an employee of Evidera at the time the work was conducted and is now an employee of Cytel. Monique Martin and Cassandra Schaible are employees of Evidera. April Camilla Roslani, Dong-Lin Ren, Philippe Godeberge, Parvez Sheikh, Robert Bandolon, and Varut Lohsiriwat have nothing to disclose.

Compliance with Ethics Guidelines. This article is based on previously conducted studies; none of the authors performed any new studies involving human participants or animals.

Data Availability. Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Open Access. This article is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and

the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Higgins JPT, Thomas J, Chandler J, et al., editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021). <https://training.cochrane.org/handbook>. Accessed Aug 2021.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372: n71.
- Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998;52(6):377–84.
- Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358: j4008.
- Abramowitz L, Benabderrahmane M, Pospait D, Philip J, Laouenan C. The prevalence of proctological symptoms amongst patients who see general practitioners in France. *Eur J Gen Pract*. 2014;20(4): 301–6.
- Conaghan P, Farouk R. Doppler-guided hemorrhoid artery ligation reduces the need for conventional hemorrhoid surgery in patients who fail rubber band ligation treatment. *Dis Colon Rectum*. 2009;52(1):127–30.
- Godeberge P, Sheikh P, Zagriadskii E, et al. Hemorrhoidal disease and chronic venous insufficiency: concomitance or coincidence; results of the CHORUS study (Chronic venous and HemORrhoidal diseases evaluation and Scientific research). *J Gastroenterol Hepatol*. 2020;35(4):577–85.
- Ince M, Ozdemir Y, Kucukerdonmez O, Levhi Akin M. Etiological risk factors of hemorrhoidal disease in young Anatolians. *J Clin Investig*. 2012;6(2): 109–12.
- Lin LH, Siu JJ, Liao PC, et al. Association of chronic obstructive pulmonary disease and hemorrhoids: a nationwide cohort study. *Medicine (Baltimore)*. 2017;96(10):e6281.
- Mascarello KC, Matijasevich A, da Silva dos Santos I, Silveira MF. Complicações puerperais precoces e tardias associadas à via de parto em uma coorte no Brasil. *Rev Bras Epidemiol*. 2018. <https://doi.org/10.1590/1980-549720180010>.
- Peery AF, Sandler RS, Galanko JA, et al. Risk factors for hemorrhoids on screening colonoscopy. *PLoS ONE*. 2015;10(9):e0139100.
- Peeters GM, Burton NW, Brown WJ. Associations between sitting time and a range of symptoms in mid-age women. *Prev Med*. 2013;56(2):135–41.
- Ratto C, Donisi L, Parello A, Litta F, Doglietto GB. Evaluation of transanal hemorrhoidal dearterialization as a minimally invasive therapeutic approach to hemorrhoids. *Dis Colon Rectum*. 2010;53(5):803–11.
- Yang JY, Peery AF, Lund JL, Pate V, Sandler RS. Burden and cost of outpatient hemorrhoids in the United States employer-insured population, 2014. *Am J Gastroenterol*. 2019;114(5):798–803.
- Alonso-Coello P, Zhou Q, Martinez-Zapata MJ, et al. Meta-analysis of flavonoids for the treatment of haemorrhoids. *Br J Surg*. 2006;93(8):909–20.
- Aziz Z, Huin WK, Badrul Hisham MD, Tang WL, Yaacob S. Efficacy and tolerability of micronized purified flavonoid fractions (MPFF) for haemorrhoids: a systematic review and meta-analysis. *Complement Ther Med*. 2018;39:49–55.
- Perera N, Liolitsa D, Iype S, et al. Phlebotonics for haemorrhoids. *Cochrane Database Syst Rev*. 2012. <https://doi.org/10.1002/14651858.CD004322.pub3>.
- Quijano CE, Abalos E. Conservative management of symptomatic and/or complicated haemorrhoids in pregnancy and the puerperium. *Cochrane Database Syst Rev*. 2005. <https://doi.org/10.1002/14651858.CD004077.pub2>.
- Wanis KN, Emmerton-Coughlin HM, Coughlin S, Foley N, Vinden C. Systemic metronidazole may not reduce posthemorrhoidectomy pain: a meta-analysis of randomized controlled trials. *Dis Colon Rectum*. 2017;60(4):446–55.

-
20. Naldini G, Martellucci J, Rea R, et al. Tailored prolapse surgery for the treatment of haemorrhoids and obstructed defecation syndrome with a new dedicated device: TST STARR Plus. *Int J Colorectal Dis.* 2014;29(5):623–9.
 21. Kendirci M, Şahiner İT, Şahiner Y, Güney G. Comparison of effects of vessel-sealing devices and conventional hemorrhoidectomy on postoperative pain and quality of life. *Med Sci Monit.* 2018;24:2173–9.
 22. Sheikh P, Lohsiriwat V, Shelygin Y. Micronized purified flavonoid fraction in hemorrhoid disease: a systematic review and meta-analysis. *Adv Ther.* 2020;37(6):2792–812.
 23. Godeberge P, Sheikh P, Lohsiriwat V, Jalife A, Shelygin Y. Micronized purified flavonoid fraction in the treatment of hemorrhoidal disease. *J Comp Eff Res.* 2021;10(10):801–13.