



Received: 2018/03/01, Revised: 2018/04/28,  
 Accepted: 2018/05/27, Published: 2018/06/30

©2018 Yoshifumi Tsuchiya et al; License Journal of Exercise Nutrition and Biochemistry. This is an open access article distributed under the terms of the creative commons attribution license (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*Corresponding author: Kazushige Goto

Faculty of Sports and Health Science, Ritsumeikan University, Shiga, Japan

Phone: +81-77-599-4127 / Fax: +81-77-599-4127

E-mail: [kagoto@fc.ritsumei.ac.jp](mailto:kagoto@fc.ritsumei.ac.jp)

©2018 The Korean Society for Exercise Nutrition

## Irisin response to downhill running exercise in humans

Yoshifumi Tsuchiya<sup>1,2</sup> / Sahiro Mizuno<sup>1,2</sup> / Kazushige Goto<sup>\*1,3</sup>

1. Graduate School of Sport and Health Science, Ritsumeikan University, Shiga, Japan.

2. Research Fellowship for Young Scientists of the Japan Society for the Promotion of Science, Tokyo, Japan.

3. Faculty of Sports and Health Science, Ritsumeikan University, Shiga, Japan.

**[Purpose]** To determine the effects of exercise-induced muscle damage, we examined irisin responses during level running (LR), with less muscle damage, and downhill running (DHR), with greater muscle damage under equivalent exercise duration and oxygen consumption ( $\text{VO}_2$ ) conditions.

**[Methods]** Fifteen healthy men (age:  $21.6 \pm 2.0$  y, height:  $170 \pm 1.3$  cm, weight:  $64.8 \pm 2.7$  kg) were randomly assigned to either the LR group ( $n = 8$ ) or the DHR group ( $n = 7$ ). Subjects in the LR group performed treadmill running at 70% of maximum oxygen uptake ( $\text{VO}_{2\text{max}}$ ) for 30 min on a 0% gradient. In contrast, subjects in the DHR group performed the same exercise on a -10% gradient. Blood samples were collected before exercise, immediately after exercise, and 1, 3, and 24 h after exercise.

**[Results]** No significant interaction (group  $\times$  time) or main effect of group or time was observed for changes in plasma irisin concentrations over time ( $P > 0.05$ ). However, the area under the curve of plasma irisin concentrations during a 3-h post-exercise period was significantly greater in the DHR ( $239,197 \pm 8,166$  ng/mL) group than in the LR ( $92,293 \pm 8,755$  ng/mL) group ( $P < 0.05$ ). The blood lactate, serum cortisol, myoglobin, and plasma interleukin-6 concentrations were significantly higher in the DHR group than in the LR group after exercise ( $P < 0.05$  for all variables).

**[Conclusion]** DHR associated with marked muscle damage promoted a greater increase in exercise-induced irisin did LR after the same duration under identical  $\text{VO}_2$  conditions.

**[Key words]** Irisin, downhill running, exercise-induced muscle damage, FNDC5, myoglobin, IL-6

## INTRODUCTION

Myokines are physiologically active substances released from skeletal muscle<sup>1</sup>. Irisin is a relatively recently discovered myokine that enhances energy expenditure (EE) by increasing UCP1 expression in white and/or beige adipocytes<sup>2,3</sup>. Several studies have described the irisin response to a single bout of exercise<sup>4-10</sup>. However, the effect of endurance exercise on the irisin response remains unclear, although the mode of exercise may affect the response. The plasma irisin concentration increased significantly at 54 min during 90 min of running<sup>11</sup>, but not after 60 min of pedaling<sup>12</sup>. In contrast, a single bout of resistance exercise (6–8 exercises) significantly increased the irisin response<sup>13,14</sup>. Notably, resistance exercise caused a greater increase in irisin concentration than did the same duration of endurance exercise (pedaling), although the EE was much lower during resistance exercise<sup>12</sup>. Differences in the exercise-induced irisin response among various studies may be explained by the different exercise modalities, with varying degrees of eccentric contraction. During eccentric contraction, fast-twitch fibers are more prominently recruited than during concentric contraction<sup>15-18</sup>. An increase in exercise intensity promotes recruitment of fast-twitch fibers, leading to an augmented irisin response<sup>19</sup>.

Downhill running (DHR) especially highlights eccentric contraction in the quadriceps femoris compared with level running (LR), causing severe exercise-induced muscle damage<sup>20-22</sup> and inflammation<sup>23</sup> during the post-exercise period. These factors cause damage to neutrophils and monocytes due to exercise-induced oxidative stress<sup>24</sup>. Huh et al. showed that exercise-induced elevation of irisin was observed with a concomitant increase in creatine kinase (CK) (an indirect muscle damage marker)<sup>25, 26</sup>. However, it remains unclear whether the magnitude of exercise-induced muscle damage influences the irisin response. Therefore, we examined the irisin responses in 2 different types of running exercise: DHR and LR under similar EE conditions. We hypothesized that the irisin response would be augmented more after DHR than after LR.

## METHODS


### Subjects

Fifteen healthy, physically active men [mean age  $\pm$  standard error (SE):  $21.6 \pm 2.0$  y, height:  $170 \pm 1.3$  cm, weight:  $64.8 \pm 2.7$  kg, body mass index:  $22.4 \pm 0.8$  kg/m<sup>2</sup>] participated in the present study. All participants had undergone several years of exercise training and exercised at least weekly. They were informed of the study purpose, experimental procedures, and possible risks of the study, and they provided written informed consent. All experimental procedures were approved by the Ethics Committee for Human Experiments at Ritsumeikan University (IRB-2014-004) and all complied with the Helsinki Declaration.

### Experimental setting

The subjects visited the laboratory twice during the experimental period. On the first day, maximum oxygen consumption ( $\dot{V}O_{2\max}$ ) during treadmill running was determined. On the second day (main experiment day), all subjects completed either a 30-min LR or a DHR protocol on a treadmill (Valiant Lode B. V., Groningen, the Netherlands). Exercise-induced irisin and metabolic responses were monitored during a 24-h post-exercise period.

Fifteen subjects were randomly divided into either LR ( $n = 8$ ) or DHR ( $n = 7$ ) groups to match the  $\dot{V}O_2$  level between the groups. In the LR group, subjects ran for 30 min on a treadmill with a 0% gradient, whereas subjects in the DHR group ran with a -10% gradient. The exercise intensity was set to achieve 70% of  $\dot{V}O_{2\max}$  during incremental running on a treadmill (0% gradient). The running velocity in the DHR group was adjusted individually during the initial 5 min of exercise to achieve 70% of the  $\dot{V}O_{2\max}$ . The DHR protocol was based on those of previous studies [27, 28] and our pilot study. All subjects were instructed to refrain from caffeine and alcohol and strenuous physical activity for 3 days before the trial day. They were also asked to minimize physical activity for 24 h after the exercise. Prescribed meals were provided at 3 h and 9 h after exercise (Figure 1). All subjects were allowed to consume only water ad libitum on the day of the trial. The sleep duration was set from 22:00 to 07:00.



**Fig.1.** Time frame of main experiment.

### $\dot{V}O_{2\max}$

Approximately 1 week prior to the main experiment, the  $\dot{V}O_{2\max}$  was evaluated to determine running velocity.

Respiratory gas samples during exercise were collected with breath-by-breath method using an automatic gas analyzer (AE300S Minato Medical Science, Osaka, Japan). All data were averaged every 30 s. The treadmill gradient during the  $\dot{V}O_{2\max}$  test was set at 0%. The  $\dot{V}O_{2\max}$  test commenced with running at 6.0, 8.0, and 10.0 km/h for 4 min each. The running speed was then increased by 2.0 km/h every 3 min to 12.0 km/h and 14.0 km/h. When treadmill running at 14.0 km/h was completed, the speed was progressively increased every minute by 0.6 km/h to exhaustion. Exhaustion was defined as 1)  $\dot{V}O_{2\max}$  plateau; 2) a heart rate equal to the maximum age-predicted value (220 minus age); or, 3) a respiratory exchange ratio (RER) of 1.1. When a subject met at least 2 of these 3 criteria, the test was concluded.

### Respiratory gas sampling during exercise

On main experiment days, respiratory gas samples during 30 min (8:40-9:10) of treadmill running were evaluated with the breath-by-breath method using an automatic gas analyzer (AE300S Minato Medical Science, Osaka, Japan). Respiratory gas samples were used for continuous determination of  $\dot{V}O_2$ , carbon dioxide output ( $\dot{V}CO_2$ ), minute ventilation ( $\dot{V}E$ ), and the RER. All data were averaged every 30 s.

### Blood samples

The subjects presented at 8:00. First, a venous blood sample (i.e., pre-exercise sample) was collected after 30 min of rest. The subjects began prescribed exercise at 8:40. Additional blood samples were collected immediately after exercise (9:10), and at 1 h (10:10), 3 h (13:10), and 24 h (09:10) after exercise (Figure 1). The collected blood samples were centrifuged for 10 min (3,000 rpm, 4°C), and the serum and plasma samples were stored at -60°C prior to analysis. Blood samples were assessed immediately after collection with automated glucose (Freestyle; Nipro Co., Osaka, Japan) and lactate (Lactate Pro2 LT-1730; Arkray Inc., Kyoto, Japan) analyzers. Serum cortisol, myoglobin, and CK concentrations were measured at a clinical laboratory (SRL Inc., Tokyo, Japan). Plasma interleukin (IL)-6 and irisin concentrations were determined using enzyme-linked immunosorbent assay (ELISA) kits [HS600B; R&D Systems, Minneapolis, MN, USA (IL-6); EK-067-52 (Lot 605767); Phoenix Pharmaceuticals, Inc., Darmstadt, Germany (irisin)]. All samples for ELISA were assayed in duplicate and the results were averaged. The intra-assay coefficients of variation were: 2.5% for serum cortisol, 2.4% for serum myoglobin, 2.3% for serum CK, 2.5% for plasma IL-6, and 4.3% for plasma irisin.

### Statistical analysis

All experimental data are shown as means  $\pm$  SE. An unpaired t-test was used to explore the significance of differences between groups for respiratory gas data and the area under the curve (AUC) of plasma irisin concentrations. Moreover, two-way (group  $\times$  time) repeated analysis of variance (ANOVA) was applied to identify significant in-

teractions and main effects. When a significant interaction and main effect were observed, the data were subjected to post-hoc analysis (Tukey-Kramer test).  $P < 0.05$  was considered to reflect significance. Cohen's  $d$  (for the unpaired t-test) and the  $\eta^2$  value (for 2-way ANOVA) were calculated, as appropriate, to evaluate effect sizes.

## RESULTS

### Respiratory gas variables during exercise

Table 1 presents the respiratory gas variables during the 30 min of exercise in each group. The average  $\dot{V}O_2$ ,  $\dot{V}CO_2$ ,  $\dot{V}E$ , and energy expenditure (EE) during exercise did not differ significantly between the groups ( $P = 0.35$ ,  $d = 0.50$  for  $\dot{V}O_2$ ;  $P = 0.60$ ,  $d = 0.28$  for  $\dot{V}CO_2$ ;  $P = 0.62$ ,  $d = 0.26$  for  $\dot{V}E$ ; and  $P = 0.39$ ,  $d = 0.46$  for EE, respectively). However, the average RER during exercise was significantly higher in the DHR group ( $P = 0.03$ ,  $d = 1.23$ ).

### Blood variables

Figure 2 a presents the time-course changes in plasma irisin concentrations. No significant interaction (group  $\times$  time,  $P = 0.159$ ,  $\eta^2 = 0.07$ ) or main effect for group ( $P = 0.25$ ,  $\eta^2 = 0.04$ ) and time ( $P = 0.30$ ,  $\eta^2 = 0.04$ ) was evident. In the DHR group, and the plasma irisin concentration increased approximately 3-fold by 3 h after exercise (before:  $466.1 \pm 78.8$  ng/ml; 3-h:  $1,358.0 \pm 666.7$  ng/mL), but the difference did not reach significance ( $P = 0.20$ ,  $d = 0.74$ ). When the exercise-induced irisin responses over the 3-h post-exercise period were compared, the AUC was significantly greater in the DHR group than in the LR


**Table 1.** Respiratory gas variables during exercise in each group.

Variables	LR(n=8)	DHR(n=7)	P Value
$\dot{V}O_2$ (mL/min)	$2,443 \pm 192$	$2,205 \pm 139$	0.35
$\dot{V}CO_2$ (mL/min)	$2,268 \pm 188$	$2,143 \pm 129$	0.60
$\dot{V}E$ (mL/min)	$66.3 \pm 7.5$	$71.4 \pm 6.5$	0.62
RER	$0.93 \pm 0.01$	$0.97 \pm 0.02^\dagger$	0.03
EE(kcal)	$361 \pm 29$	$329 \pm 20$	0.39

Values are means  $\pm$  SE.  $^\dagger$ :  $P < 0.05$  vs. LR group.

**Table 2.** Time-course changes in blood variables in each group. Values are means  $\pm$  SE. \*:  $P < 0.05$  vs. Pre.  $^\dagger$ :  $P < 0.05$  vs. LR group.

Variables		Pre	0 h	1 h	3 h	24 h
Glucose (mg/dL)	LR	$91 \pm 1$	$108 \pm 10$	$86 \pm 2$	$86 \pm 2$	$90 \pm 1$
	DHR	$87 \pm 2$	$133 \pm 8^*$	$81 \pm 3$	$87 \pm 2$	$91 \pm 2$
Lactate (mmol/L)	LR	$1.5 \pm 0.2$	$3.5 \pm 0.8^*$	$1.5 \pm 0.1$	$1.5 \pm 0.2$	$1.4 \pm 0.1$
	DHR	$1.8 \pm 0.2$	$6.1 \pm 1.4^{*\dagger}$	$2.1 \pm 0.1$	$1.6 \pm 0.1$	$1.6 \pm 0.1$
Cortisol ( $\mu$ g/dL)	LR	$17.2 \pm 2.2$	$18.1 \pm 2.4$	$16.4 \pm 2.5$	$11.8 \pm 1.9^*$	$17.5 \pm 1.5$
	DHR	$21.1 \pm 3.0$	$28.0 \pm 2.3^\dagger$	$26.7 \pm 2.8^\dagger$	$13.9 \pm 1.5^*$	$19.6 \pm 2.5$
Myoglobin (ng/mL)	LR	$39.3 \pm 5.6$	$63.3 \pm 9.0$	$87.3 \pm 9.9$	$79.1 \pm 10.1$	$40.9 \pm 3.3$
	DHR	$60.0 \pm 29.7$	$207.7 \pm 62.1^{*\dagger}$	$564.0 \pm 115.9^{*\dagger}$	$483.7 \pm 109.5^{*\dagger}$	$68.2 \pm 23.4$
CK (U/L)	LR	$220 \pm 55$	$241 \pm 62$	$275 \pm 54$	$261 \pm 53$	$246 \pm 30$
	DHR	$606 \pm 467$	$712 \pm 514$	$751 \pm 525$	$877 \pm 505$	$1070 \pm 307$




**Fig. 2.** (a) Plasma irisin concentrations before and after each exercise. Shaded box: exercise period. (b) Area under the curve (AUC) of plasma irisin concentration during the 3-h post-exercise period. Values are means  $\pm$  SE.  $^\dagger$ :  $P < 0.05$ .

group ( $P = 0.04$ ,  $d = 0.99$ ; Figure 2b).

As shown in Figure 3a, there was no significant interaction (group  $\times$  time;  $P = 0.122$ ,  $\eta^2 = 0.06$ ) or main effect for group ( $P = 0.105$ ,  $\eta^2 = 0.10$ ) for the plasma IL-6 concentration. Plasma IL-6 concentrations increased significantly immediately (0 h) and 1 h after exercise in both groups (main effect for time;  $P = 0.001$ ,  $\eta^2 = 0.37$ ). Moreover, the DHR group showed a significant increase at 3 h after exercise ( $P = 0.002$ ,  $d = 1.51$ ). The AUC over the 3-h post-exercise period for plasma IL-6 tended to be greater in the DHR group than in the LR group ( $P = 0.061$ ,  $d = 1.06$ , Figure 3b).

Table 2 shows the time-course changes in blood variables. No significant interaction (group  $\times$  time) was evident for blood glucose ( $P = 0.07$ ,  $\eta^2 = 0.09$ ), lactate ( $P = 0.16$ ,  $\eta^2$



**Fig. 3.** (a) Plasma IL-6 concentrations before and after each exercise. Shaded box: exercise period. (b) AUC of plasma IL-6 concentration during the 3-h post-exercise period. Values are means  $\pm$  SE. \*:  $P < 0.05$  vs. Pre.

= 0.06), or serum CK concentrations ( $P = 0.22$ ,  $\eta^2 = 0.1$ ). There was no significant main effect for group for blood glucose ( $P = 0.20$ ,  $\eta^2 = 0.01$ ), serum cortisol ( $P = 0.06$ ,  $\eta^2 = 0.14$ ), or CK concentrations ( $P = 0.22$ ,  $\eta^2 = 0.15$ ). The blood glucose concentration exhibited a significant main effect for time ( $P < 0.001$ ,  $\eta^2 = 0.54$ ), and there was a significant increase in the DHR group ( $P < 0.001$ ,  $d = 1.19$ ), but not in the LR group. The blood lactate concentration exhibited a significant main effect for time ( $P < 0.05$ ,  $\eta^2 = 0.44$ ) and a tendency toward significant main effect for group ( $P = 0.06$ ,  $\eta^2 = 0.04$ ). Blood lactate concentration increased significantly immediately after exercise in both groups (LR:  $P = 0.002$ ,  $d = 1.19$ ; DHR:  $P < 0.001$ ,  $d = 1.62$ ). Furthermore, the blood lactate concentration immediately after exercise was significantly higher in the DHR group than in the LR group ( $P < 0.001$ ,  $d = 0.85$ ). The serum cortisol concentration exhibited a significant interaction (group  $\times$  time,  $P = 0.005$ ,  $\eta^2 = 0.06$ ) and a main effect for time ( $P < 0.001$ ,  $\eta^2 = 0.21$ ). The DHR group revealed significantly higher values both immediately and 1 h after exercise (0 h:  $P < 0.001$ ,  $d = 1.53$  and 1 h:  $P < 0.001$ ,  $d = 1.43$ ). The serum myoglobin concentration exhibited a significant interaction (group  $\times$  time,  $P < 0.003$ ,  $\eta^2 = 0.22$ ) and main effects for group ( $P < 0.001$ ,  $\eta^2 = 0.18$ ) and time ( $P = 0.003$ ,  $\eta^2 = 0.26$ ). In the DHR group, the myoglobin concentration was significantly higher than in the LR group immediately ( $P = 0.01$ ,  $d = 1.28$ ), and 1 ( $P < 0.001$ ,  $d = 2.28$ ) and 3 h after exercise ( $P < 0.001$ ,  $d = 1.82$ ). The serum CK concentration did not exhibit a main effect for time ( $P = 0.12$ ,  $\eta^2 = 0.17$ ).

## DISCUSSION

This is the first study to explore the effects of exercise-induced muscle damage on the irisin response. As expected, exercise-induced elevation of serum myoglobin (an indirect muscle damage marker) and plasma IL-6 (an inflammatory cytokine) were significantly greater in the DHR group than in the LR group, although EE during 30 min of exercise was not significantly different. Consequently, the plasma irisin concentration increased approximately 3-fold 3 h after completion of DHR (Pre:  $466.1 \pm 78.8$  ng/mL, 3 h after exercise:  $1,358.0 \pm 666.7$  ng/mL). Additionally, the exercise-induced irisin response over 3 h after DHR was significantly greater than that after LR, suggesting that the augmented irisin response may be associated with magnitude of muscle damage.

Exercise intensity is a primary factor in irisin production<sup>19,26</sup>, and increased recruitment of fast twitch fibers may play an important role in the irisin response. During DHR, eccentric muscle contraction (which preferentially recruits fast twitch fibers) is highlighted during the landing phase<sup>15-18</sup>. Furthermore, running velocity was significantly higher in the DHR group ( $14.0 \pm 0.2$  km/h) than in the LR group ( $10.8 \pm 0.3$  km/h), to match EE during exercise. We previously showed that higher-velocity running was associated with a greater irisin response than lower-velocity running under matched EE conditions<sup>19</sup>. Therefore, enhanced irisin response (revealed by the AUC over 3 h) after DHR might be, at least in part, explained by higher running velocity with increased recruitment of fast twitch fibers.

We sought to identify associations between the levels of exercise-induced muscle damage markers (myoglobin and CK) in blood and the irisin response. Several studies suggested a relationship between these 2 factors in both humans<sup>26</sup> and rats<sup>29</sup>. However, the above studies involved limitations, including low frequencies of blood collection. In the present study, we monitored the time course of changes in the levels of plasma irisin and muscle damage markers over a 24-h post-exercise period. The serum myoglobin level was markedly higher in the DHR group than in the LR group, indicating that exercise-induced muscle damage was profound in the DHR group. Moreover, exercise-induced plasma IL-6 elevation was profound in the DHR group (until 3 h post-exercise). The AUC value over a 3-h post-exercise period tended to be greater in the DHR group than in the LR group ( $P = 0.06$ ). Similarly, Vyver et al. showed that both serum myoglobin and IL-6 concentrations increased concomitantly after completion of DHR<sup>30</sup>. Therefore, significantly higher plasma IL-6 concentration during the post-exercise period in the DHR group may be related to proinflammatory action following ultrastructural muscle damage. The exercise-induced elevation of lactate was also greater in the DHR group than in the LR group, suggesting that DHR promoted muscle glycogen utilization. Previous studies suggested that a high level of IL-6 was seen with low glycogen levels, and that the lower glycogen level induces PGC-1 $\alpha$  (an upstream factor in irisin production) gene expression<sup>31-33</sup>. Additionally, Huh et al.



found that the mRNA levels of PGC-1 $\alpha$  (an upstream factor in irisin production) and FNDC5 (the precursor of irisin) were increased by IL-6 addition in vitro<sup>13</sup>. Thus, further reduction of muscle glycogen levels caused by DHR may promote the PGC-1 $\alpha$ -FNDC5-irisin axis via IL-6 production. However, it is difficult to clarify completely the association between IL-6 and irisin, since IL-6 has multiple sources (e.g., leucocytes, myocytes) and actions (pro- and anti-inflammatory). Taken together, the present data suggest that the exercise-induced irisin response may be associated with muscle damage (reflected by the elevation in myoglobin concentrations) and/or inflammation (reflected by the elevation in IL-6 concentrations).

In the present study, post-exercise increases in irisin levels were modest compared to those of previous studies<sup>11</sup>. Although the intensity (70% of VO<sub>2max</sub>) and duration (30 min) during DHR exercise appeared to be sufficient to stimulate irisin production, strenuous muscle contraction can trigger myokine proteolysis<sup>34</sup>. Therefore, it is possible that excessive muscle damage caused by DHR attenuated the transient irisin response.

Several limitations should be noted when interpreting the results. First, we did not use a crossover design, because we attempted to avoid “repeated bout effects” for exercise-induced muscle damage<sup>35</sup>. Thus, we divided the 15 subjects into DHR and LR groups, and inter-individual variation in the irisin response should be considered. Second, our results may be specific to healthy subjects. Indeed, obese subjects and patients with type 2 diabetes exhibited irisin responses different from those of healthy individuals<sup>34</sup>. Third, the existence of irisin and its role in humans is controversial<sup>37-39</sup>. However, Jedrychowski et al. used quantitative mass spectrometry to precisely measure plasma irisin levels in healthy humans in both the resting and post-exercise state<sup>40</sup>. Furthermore, the ELISA that we used in the present study was the most commonly employed in previous studies<sup>25,36,41-44</sup>.

From a practical viewpoint, our findings will aid in the design of exercises that efficiently reduce the risk of metabolic disorder and associated disease. We speculate that exercise-induced irisin secretion would be influenced by muscle damage and inflammation. A future study should address the physiological implications of transient increases in irisin levels on the resting metabolic rate, insulin sensitivity, and body composition in various populations.

In conclusion, we found that the exercise-induced irisin response over the 3 h period after DHR was significantly greater than that after LR, although the EE was similar.

## ACKNOWLEDGEMENTS

We gratefully thank the subjects for commitment in the present study. We also thank laboratory members for help with the experiment and analyses of blood samples.

## REFERENCES

1. Steensberg A, van Hall G, Osada T, Sacchetti M, Saltin B, Pedersen BK. Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *J Physiol*. 2000;529:237–42.
2. Boström P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Boström EA, Choi JH, Long JZ, Kajimura S, Zingaretti MC, Vind BF, Tu H, Cinti S, Højlund K, Gygi SP, Spiegelman BM. A PGC1- $\alpha$ -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature*. 2012;481:463–8.
3. Wu J, Boström P, Sparks LM, Ye L, Choi JH, Giang AH, Khandekar M, Virtanen KA, Nuutila P, Schaart G, Huang K, Tu H, van Marken Lichtenbelt WD, Hoeks J, Enerbäck S, Schrauwen P, Spiegelman BM. Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell*. 2012;150:366–76.
4. Huh JY, Panagiotou G, Mougios V, Brinkoetter M, Vamvini MT, Schneider BE, Mantzoros CS. FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. *Metabolism*. 2012;61:1725–38.
5. Pekkala S, Wiklund PK, Hulmi JJ, Ahtiainen JP, Horttanainen M, Pöllänen E, Mäkelä KA, Kainulainen H, Häkkinen K, Nyman K, Alén M, Herzig KH, Cheng S. Are skeletal muscle FNDC5 gene expression and irisin release regulated by exercise and related to health? *J Physiol*. 2013;591:5393–400.
6. Norheim F, Langleite TM, Hjorth M, Holen T, Kielland A, Stadheim HK, Gulseth HL, Birkeland KI, Jensen J, Drevon CA. The effects of acute and chronic exercise on PGC-1 $\alpha$ , irisin and browning of subcutaneous adipose tissue in humans. *FEBS J*. 2014;281:739–49.
7. Daskalopoulou SS, Cooke AB, Gomez YH, Mutter AF, Filipaivos A, Mesfum ET, Mantzoros CS. Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *Eur J Endocrinol*. 2014;171:343–52.
8. Anastasilakis AD, Polyzos SA, Saridakis ZG, Kynigopoulos G, Skouvaklidou EC, Molyvas D, Vasiloglou MF, Apostolou A, Karagiozoglou-Lampoudi T, Siopi A, Mougios V, Chatzistavridis P, Panagiotou G, Filippaivos A, Delaroudis S, Mantzoros CS. Circulating irisin in healthy, young individuals: day-night rhythm, effects of food intake and exercise, and associations with gender, physical activity, diet, and body composition. *J Clin Endocrinol Metab*. 2014;9:324755.
9. Brenmoehl J, Albrecht E, Komolka K, Schering L, Langhammer M, Hoeflich A, Maak S. Irisin is elevated in skeletal muscle and serum of mice immediately after acute exercise. *Int J Biol Sci*. 2014;10:338–49.
10. Bell MA, Levine CB, Downey RL, Griffiths C, Mann S, Fryeb CW, Wakshlag JJ. Influence of endurance and sprinting exercise on plasma adiponectin, leptin and irisin concentrations in racing Greyhounds and sled dogs. *Aus Vet J*. 2016;94:154–9.
11. Kraemer RR, Shockett P, Webb ND, Shah U, Castracane VD. A transient elevated irisin blood concentration in response to prolonged, moderate aerobic exercise in young men and women. *Horm Metab Res*. 2014;46:150–4.

12. Tsuchiya Y, Ando D, Takamatsu K, Goto K. Resistance exercise induces a greater irisin response than endurance exercise. *Metabolism*. 2015;64:1042–50.
13. Huh JY, Mougios V, Kabasakalis A, Fatouros I, Siopi A, Douroudos II, Filippaios A, Panagiotou G, Park KH, Mantzoros CS. Exercise-induced irisin secretion is independent of age or fitness level and increased irisin may directly modulate muscle metabolism through AMPK activation. *J Clin Endocrinol Metab*. 2014;99:E2154–61.
14. Nygaard H, Slettakken G, Vegge G, Hollan I, Whist JE, Strand T, Rønnestad BR, Ellefsen S. Irisin in blood increases transiently after single sessions of intense endurance exercise and heavy strength training. *PLoS One*. 2015;10:e0121367.
15. Fridén J, Sjöström M, Ekblom B. Myofibrillar damage following intense eccentric exercise in man. *Int J Sports Med*. 1983; 4:170–6.
16. Warren G, Hayes D, Lowe D, Williams J, Armstrong R. Eccentric contraction-induced injury in normal and hindlimb-suspended mouse soleus and edl muscles. *J Appl Physiol*. 1994;77:1421–30.
17. Macaluso F, Isaacs AW, Myburgh KH. Preferential type II muscle fiber damage from plyometric exercise. *J Athl Train*. 2012;47:414–20.
18. Choi S.J. Differential susceptibility on myosin heavy chain isoform following eccentric-induced muscle damage. *J Exerc Rehabil*. 2014;10:344–8.
19. Tsuchiya Y, Ando D, Goto K, Kiuchi M, Yamakita M, Koyama K. High-intensity exercise causes greater irisin response compared with low-intensity exercise under similar energy consumption. *Tohoku J Exp Med*. 2014;233:135–40.
20. Chen TC, Nosaka K, Wu CC. Effects of a 30-min running performed daily after downhill running on recovery of muscle function and running economy. *J Sci Med Sport*. 2007;11:271–9.
21. Jamurtas AZ, Garyfallopoulou A, Theodorou AA, Zalavras A, Paschalis V, Deli CK, Nikolaidis MG, Fatouros IG, Koutedakis Y. A single bout of downhill running transiently increases HO-MA-IR without altering adipokine response in healthy adult women. *Eur J Appl Physiol*. 2013;113:2925–32.
22. Yu T, Chang Y, Gao XL, Li H, Zhao P. Dynamic Expression and the Role of BDNF in Exercise-induced Skeletal Muscle Regeneration. *Int J Sports Med*. 2017;38:1–8.
23. Peake J, Nosaka K, Suzuki K. Characterization of inflammatory responses to eccentric exercise in humans. *Exerc Immunol*. 2005;11:64–85.
24. Liao P, Zhou J, Ji LL, Zhang Y. Eccentric contraction induces inflammatory responses in rat skeletal muscle : role of tumor necrosis factor- $\alpha$ . *Am J Physiol Regul Integr Comp Physiol*. 2010;298:599–607.
25. Huh JY, Mantzoros CS. Irisin physiology, oxidative stress, and thyroid dysfunction: What next? *Metabolism*. 2015;64:765–7.
26. Huh JY, Siopi A, Mougios V, Park KH, Mantzoros CS. Irisin in response to exercise in humans with and without metabolic syndrome. *J Clin Endocrinol Metab*. 2015;100:1–7.
27. Chen TC, Nosaka K, Lin M-J, Chen H-L, Wu C-J. Changes in running economy at different intensities following downhill running. *J Sports Sci*. 2009;27:1137–44.
28. Mizuno S, Morii I, Tsuchiya Y, Goto K. Wearing Compression Garment after Endurance Exercise Promotes Recovery of Exercise Performance. *Int J Sports Med*. 2016;37:870–7.
29. Samy DM, Ismail CA, Nassra RA. Circulating irisin concentrations in rat models of thyroid dysfunction - Effect of exercise. *Metabolism*. 2015;64:804–13.
30. van de Vyver M, Myburgh KH. Variable inflammation and intramuscular STAT3 phosphorylation and myeloperoxidase levels after downhill running. *Scand J Med Sci Sport*. 2014;24:360–1.
31. Gavin JP, Myers SD, Willems ME. Effect of eccentric exercise with reduced muscle glycogen on plasma interleukin-6 and neuromuscular responses of musculus quadriceps femoris. *J Appl Physiol*. 2016;121:173–84.
32. Psilander N, Frank P, Flockhart M, Sahlin K. Exercise with low glycogen increases PGC-1 $\alpha$  gene expression in human skeletal muscle. *Eur J Appl Physiol*. 2013;113:951–63.
33. Pedersen BK, Akerstrom TC, Nielsen AR, Fischer CP. Role of myokines in exercise and metabolism. *J Appl Physiol*. 2007;103:1093–8.
34. Manabe Y, Miyatake S, Takagi M, Nakamura M, Okeda A, Nakano T, Hirshman MF, Goodyear LJ, Fujii NL. Characterization of an Acute Muscle Contraction Model Using Cultured C2C12 Myotubes. *PLoS One*. 2012;7:e52592.
35. Nosaka K, Sakamoto K, Newton M SP. How long does the protective effect on eccentric exercise-induced muscle damage last? *Med Sci Sport Exerc*. 2001;33:1490–5.
36. Löffler D, Müller U, Scheuermann K, Friebe D, Gesing J, Bielitz J, Erbs S, Landgraf K, Wagner IV, Kiess W, Kömer A. Serum irisin levels are regulated by acute strenuous exercise. *J Clin Endocrinol Metab*. 2015;100:1289–99.
37. Erickson HP. Irisin and FNDC5 in retrospect. *Adipocyte*. 2013;2:289–93.
38. Sanchis-Gomar F, Alis R, Lippi G. Circulating irisin detection: does it really work? *Trends Endocrinol Metab*. 2015;26:335–6.
39. Albrecht E, Norheim F, Thiede B, Holen T, Ohashi T, Schering L, Lee S, Brenmoehl J, Thomas S, Drevon CA, Erickson HP, Maak S. Irisin – a myth rather than an exercise-inducible myokine. *Sci Rep*. 2015;5:8889.
40. Jedrychowski MP, Wrann CD, Paulo JA, Gerber KK, Szpyt J, Robinson MM, Nair KS, Gygi SP, Spiegelman BM. Detection and quantitation of circulating human irisin by tandem mass spectrometry. *Cell Metab*. 2015;22:734–40.
41. Hew-Butler T, Landis-Piowar K, Byrd G, Seimer M, Seigneurie N, Byrd B, Muzik O. Plasma irisin in runners and nonrunners: no favorable metabolic associations in humans. *Physiol Rep*. 2015;3:e12262.
42. Reza MM, Subramaniam N, Sim CM, Ge X, Sathiakumar D, McFarlane C, Sharma M, Kambadur R. Irisin is a pro-myogenic factor that induces skeletal muscle hypertrophy and rescues denervation-induced atrophy. *Nat Commun*. 2017;8:1104.
43. Buscemi S, Corleo D, Buscemi C, Giordano C. Does iris(in) bring bad news or good news? *Eat Weight Disord*. 2017;1–12.
44. Colaianni G, Cuscito C, Mongelli T, Pignataro P, Buccoliero C, Liu P Lu P, Sartini L, Di Comite M, Mori G, Di Benedetto A, Brunetti G, Yuen T, Sun L, Reseland JE, Colucci S, New MI, Zaidi M, Cinti S, Grano M. The myokine irisin increases cortical bone mass. *Proc Natl Acad Sci*. 2015;112:12157–62.