

Received: 2013.09.24  
Accepted: 2013.11.22  
Published: 2014.01.27

# Co-occurrence of carbapenem and aminoglycoside resistance genes among multidrug-resistant clinical isolates of *Acinetobacter baumannii* from Cracow, Poland

Authors' Contribution:  
Study Design A  
Data Collection B  
Statistical Analysis C  
Data Interpretation D  
Manuscript Preparation E  
Literature Search F  
Funds Collection G

ABCDEF **Paweł Nowak**  
ABCDEF **Paulina Paluchowska**  
ADEFG **Alicja Budak**

Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Jagiellonian University Medical College, Cracow, Poland

**Corresponding Author:** Alicja Budak, e-mail: [budak@cm-uj.krakow.pl](mailto:budak@cm-uj.krakow.pl)

**Source of support:** Scientific work was partially supported by the Polish government as a research grant (no. N N405 672140) in the years 2011–12 and by the Jagiellonian University Medical College with funds from maintenance of the research potential of the Department of Pharmaceutical Microbiology JU MC

**Background:** *Acinetobacter baumannii* is a significant hospital pathogen, possessing a considerable degree of antimicrobial resistance. *A. baumannii* resistance to carbapenems and aminoglycosides is mostly conferred by class D OXA carbapenemases and aminoglycoside-modifying enzymes, respectively. The aim of this study was to determine the prevalence of selected genes encoding OXA carbapenemases and aminoglycoside-modifying enzymes in multidrug-resistant strains of *A. baumannii*.

**Material/Methods:** The study included 61 carbapenem-resistant and aminoglycoside-nonsusceptible *A. baumannii* isolates, collected between 2009 and 2011 in Cracow, Poland. Selected resistance genes, including: *bla*<sub>OXA-51-like</sub>, *bla*<sub>OXA-23-like</sub>, *bla*<sub>OXA-40-like</sub>, *bla*<sub>OXA-58-like</sub>, *aac(6)-Ih*, *aac(3)-Ia*, *aac(3)-IIa*, *aac(6)-Ib*, *aph(3')-Ia* and *aph(3')-VI*, were detected by PCR method.

**Results:** The *bla*<sub>OXA-51-like</sub> genes were detected in all isolates, while acquired carbapenemase encoding genes were found in 96.7% of tested strains. Presence of *bla*<sub>OXA-40-like</sub> and *bla*<sub>OXA-23-like</sub> genes was observed among 65.6% and 27.9% of isolates, respectively. Assayed aminoglycoside resistance genes were found to harbor 98.4% of isolates. Among tested strains, we observed the following percentages of resistance determinants: *aac(3)-Ia* – 78.7%, *aph(3')-VI* – 78.7% and *aph(3')-Ia* – 27.9%. Analysis of co-occurrence of carbapenem and aminoglycoside resistance genes revealed the highest percentage of strains possessing *bla*<sub>OXA-40-like</sub>, *aac(3)-Ia*, and *aph(3')-VI* genes (44.3%).

**Conclusions:** The *bla*<sub>OXA-40-like</sub> and *aac(3)-Ia/aph(3')-VI* were the most prevalent genes encoding acquired OXA carbapenemases and aminoglycoside-modifying enzymes, respectively, among *A. baumannii* strains in Cracow, Poland. Genes conferring resistance to carbapenems and aminoglycosides coexisted in the clinical strains of *A. baumannii*. The phenomenon of *A. baumannii* resistance indicates the necessity of monitoring for the presence of the resistance genes.


**Keywords:** *Acinetobacter baumannii* • carbapenem resistance detection • aminoglycoside resistance detection

**Full-text PDF:** <http://www.basic.medscimonit.com/download/index/idArt/889811>

 2411

 2

 2

 29

## Background

Over the last 30 years the incidence of hospital infections caused by *Acinetobacter* spp. has increased greatly. It is connected with the growing number of highly specialized medical procedures that affect patient immunity. The incidence and seriousness of *Acinetobacter* spp. nosocomial infections is furthermore associated with the simplicity of its nutritional requirements and low susceptibility to adverse environmental conditions, as well as the remarkable ability of these bacteria to acquire resistance determinants to a wide range of antimicrobial agents [1–3]. The most clinically important species belonging to the *Acinetobacter* genus and responsible for over 90% of infections is *Acinetobacter baumannii*. Infections caused by this bacterium are associated with significant mortality [3]. The main problems caused by *A. baumannii* isolates in the hospital environment concern patients hospitalized in intensive care and burn units [1,2,4].

*A. baumannii* strains exhibit a wide spectrum of antimicrobial resistance mechanisms, both intrinsic and acquired. These mechanisms include: outer membrane permeability modifications, efflux pumps, production of drug inactivating/modifying enzymes, and target-site modifications. *A. baumannii* can become resistant due to modification of existing genes, their regulation mechanisms, or acquisition of exogenous DNA by horizontal gene transfer. Diversity and coexistence of mechanisms of resistance in this bacterium result in multiple drug resistance and can cause difficulties in treatment [5–8].

Among antimicrobials frequently used in treatment of *A. baumannii* infections, carbapenems and aminoglycosides are notable. Carbapenems produce synergistic bactericidal activity in combination with aminoglycosides; therefore, carbapenems are often used in combination therapy with aminoglycosides.

Carbapenem resistance in *A. baumannii* principally involves the molecular class D OXA-type serine oxacillinases, but may also be mediated by other mechanisms such as porin modification or loss, and efflux of antimicrobial and modification of penicillin-binding proteins [9]. The most widespread acquired carbapenemases in *A. baumannii* are the class D oxacillinases, represented by 3 main phylogenetic subgroups: OXA-23, OXA-40, and OXA-58, which are often encoded by genes localized in chromosomes or on plasmids. Furthermore, carbapenem resistance can be mediated by intrinsic *A. baumannii*, chromosome-harbored OXA-51-like enzymes, but only when the insertion sequence IS*Aba1* element is inserted upstream of the gene [9]. The ubiquitous nature of OXA-51-like encoding genes in *A. baumannii* resulted in use of *bla*<sub>OXA-51-like</sub> gene detection in identification of this bacterium to the species level [10]. However, recent studies by Lee et al. [11] revealed the presence of *bla*<sub>OXA-51-like</sub> genes in carbapenem-resistant non-*baumannii* species of *Acinetobacter*: *Acinetobacter nosocomialis*

and *Acinetobacter* genomic species “Close to 13TU”. With regard to these findings, the identification of *A. baumannii* strains cannot be based only on the detection of *bla*<sub>OXA-51-like</sub> gene, but must be supported by other identification methods [11].

Mechanisms responsible for *A. baumannii* resistance to aminoglycoside antibiotics include decreased antibiotic uptake and accumulation, modification of the ribosomal target, efflux of antibiotic, and enzymatic modification of aminoglycosides. One of the most important mechanisms attributed to aminoglycoside resistance in *A. baumannii* is the production of aminoglycoside-modifying enzymes, including aminoglycoside acetyltransferases (AAC), phosphoryltransferases (APH), and adenylyltransferases (ANT) [12].

The genes encoding resistance to carbapenems and aminoglycosides may be located in mobile genetic elements such as transposons, integrons, and/or plasmids [8].

The aim of this study was to determine the prevalence of selected genes encoding OXA carbapenemases and aminoglycoside-modifying enzymes in multidrug-resistant strains of *A. baumannii*.

## Material and Methods

### Bacterial strains

A collection of 61 non-repetitive (1 per patient) carbapenem-resistant (imipenem and meropenem) and aminoglycoside-nonsusceptible (amikacin, gentamicin, and tobramycin) *A. baumannii* strains were isolated in 2009 (n=27), 2010 (n=20), and 2011 (n=14), respectively, from patients hospitalized in the Specialized Hospital in Cracow, Poland. The isolates were identified as *A. baumannii* by the Vitek 2 Compact automatic system (bioMérieux, Poland) and PCR amplification of the *bla*<sub>OXA-51-like</sub> gene. Most tested isolates originated from Intensive Care Unit patients (36; 59.0%) and Burn Therapy Unit patients (18; 29.5%). The isolates were recovered from various clinical specimens including: (in descending frequency) respiratory tract samples (30; 49.2%), wound swabs (13; 21.3%), urine (9; 14.8%), blood (6; 9.8%), and other specimens (3; 4.9%). While phenotypic identification and antibiotic susceptibility testing, as well as molecular detection of *bla*<sub>OXA-51-like</sub>, *bla*<sub>OXA-23-like</sub>, *bla*<sub>OXA-40-like</sub>, and *bla*<sub>OXA-58-like</sub> genes, among 27 and 20 strains recovered in 2009 and 2010, respectively were described previously [13], 14 additional isolates derived from the Specialized Hospital in Cracow, Poland in 2011 were examined in the current study.

### Susceptibility testing

*A. baumannii* susceptibility testing concerning minimal inhibitory concentration (MIC) were performed using the VITEK-2

Compact system and relevant antimicrobial susceptibility testing cards (bioMérieux, Poland), interpreted according to standard CLSI criteria [14]. *A. baumannii* ATCC 19606, *P. aeruginosa* ATCC 27853, and *E. coli* ATCC 25922 were used as reference strains for quality control. The isolates were stored in a freezer at  $-80^{\circ}\text{C}$  for further analysis.

### DNA extraction

Genomic DNA was extracted using the Sherlock AX (A&A Biotechnology, Poland) isolation kit according to the manufacturer's recommendations. DNA quantification was performed by spectrophotometry at 260 nm. The purity of DNA was evaluated by the ratio of the absorbance at 260 nm and 280 nm (A260/A280) (Biometra, Germany).

### Molecular determination of carbapenem resistance

Detection of *bla*<sub>OXA-51-like</sub>, *bla*<sub>OXA-23-like</sub>, *bla*<sub>OXA-40-like</sub>, and *bla*<sub>OXA-58-like</sub> genes was performed by multiplex PCR, as described previously [15]. The PCR was carried out in a thermocycler T personal (Biometra, Germany). A single reaction mixture contained: 30 ng of genomic DNA, 20 pM of each primer (Sigma Aldrich, Poland), 10  $\mu\text{l}$  GoTaq<sup>®</sup> buffer, 3  $\mu\text{l}$  25 mM MgCl<sub>2</sub>, 1  $\mu\text{l}$  dNTPs and 0.25  $\mu\text{l}$  GoTaq<sup>®</sup> DNA Polymerase (Promega, USA) in a final volume of 50  $\mu\text{l}$ . Initial denaturation at  $94^{\circ}\text{C}$  for 3 min was followed by 30 cycles of amplification consisting of  $94^{\circ}\text{C}$  for 25 s,  $52^{\circ}\text{C}$  for 40 s,  $72^{\circ}\text{C}$  for 50 s, and a final extension step at  $72^{\circ}\text{C}$  for 5 min.

### Molecular determination of aminoglycoside resistance

Detection of genes encoding aminoglycoside-modifying enzymes was carried out using: 1) triplex PCR for *aac(6')-Ih*, *aac(3)-Ia*, *aac(3)-IIa*; 2) duplex PCR for *aac(6')-Ib*, *aph(3')-Ia*; 3) monoplex PCR for *aph(3')-VI*, described previously by Noppe-Leclercq et al. [16]. PCR reactions were performed in a final volume of 50  $\mu\text{l}$  containing: 10  $\mu\text{l}$  of GoTaq<sup>®</sup> buffer, 4  $\mu\text{l}$  of MgCl<sub>2</sub> solution, 1  $\mu\text{l}$  of PCR nucleotide mix, 2  $\mu\text{l}$  of each primers, 0.25  $\mu\text{l}$  GoTaq<sup>®</sup> DNA Polymerase, and 5  $\mu\text{l}$  of DNA (Promega, USA). Amplification reactions were performed in a T Personal thermal cycler (Biometra, Germany) with the following parameters: 1) triplex PCR at  $-95^{\circ}\text{C}$  for 2 min, following 35 cycles: of 30 s at  $95^{\circ}\text{C}$ , 40 s at  $58^{\circ}\text{C}$ , 90 s at  $72^{\circ}\text{C}$ , and 5 min at  $72^{\circ}\text{C}$ ; 2) duplex PCR at  $-95^{\circ}\text{C}$  for 2 min, following 35 cycles: of 30 s at  $95^{\circ}\text{C}$ , 40 s at  $55^{\circ}\text{C}$ , 90 s at  $72^{\circ}\text{C}$ , and 5 min at  $72^{\circ}\text{C}$ , and 3) single PCR at  $-95^{\circ}\text{C}$  for 2 min, following 35 cycles: of 30 s at  $95^{\circ}\text{C}$ , 40 s at  $49^{\circ}\text{C}$ , 90 s at  $72^{\circ}\text{C}$ , and 5 min at  $72^{\circ}\text{C}$ .

### PCR products detection

All obtained amplicons were subjected to electrophoresis with 2% agarose gel stained with ethidium bromide, visualized, and




Figure 1. Occurrence of *A. baumannii* strains (n=61) including OXA encoding genes.

digitalized. The size of PCR products was compared with the molecular weight standard O'Gene Ruler 100 bp DNA Ladder Plus (Thermo Scientific, USA).

## Results

### OXA carbapenemases genes

Phenotypic identification of 14 *A. baumannii* strains (originated from 2011) was confirmed in all isolates by presence of *bla*<sub>OXA-51-like</sub> genes. Furthermore, analysis of occurrence of acquired OXA carbapenemases genes among 14 *A. baumannii* isolates (derived from 2011) revealed the presence of *bla*<sub>OXA-40-like</sub> gene in 10 (71.4%) strains, and *bla*<sub>OXA-23-like</sub> gene was present in 3 (21.4%) strains. Moreover, only 1 strain did not carry any assayed acquired OXA carbapenemase genes. Figure 1 shows results of a previously performed analysis [13] and the present analysis of occurrence of OXA encoding genes over the selected period of time (2009, 2010, 2011). The most prevalent genes encoding OXA carbapenemases among strains isolated in 2009, 2010, and 2011 were *bla*<sub>OXA-40-like</sub> genes.

### Aminoglycoside resistance genes

All 61 *A. baumannii* strains were screened for occurrence of selected genes encoding aminoglycoside-modifying enzymes (*aac(6')-Ih*, *aac(3)-Ia*, *aac(3)-IIa*, *aac(6')-Ib*, *aph(3')-Ia*, *aph(3')-VI*). Our studies revealed the presence of at least 1 gene in 60 strains (98.4%). We observed the following percentages of genes *aac(3)-Ia* – 78.7%, *aph(3')-VI* – 78.7% and *aph(3')-Ia* – 27.9% among analyzed strains. Furthermore, 47 (77%) isolates

**Table 1.** Analysis of occurrence of aminoglycoside resistance genotypes and susceptibility patterns to amikacin, gentamicin and tobramycin among 61 *A. baumannii* clinical strains.

| Resistance genotype |                   |                   | Isolates |      | No. of isolates resistant (R)/intermediate (I) |            |            |
|---------------------|-------------------|-------------------|----------|------|--|------------|------------|
|                     |                   |                   | No.      | %    | Amikacin                                       | Gentamicin | Tobramycin |
| <i>aac(3)-Ia</i>    | –                 | <i>aph(3')-VI</i> | 30       | 49.2 | 26/4   | 30/0       | 16/14      |
| –                   | <i>aph(3')-Ia</i> | <i>aph(3')-VI</i> | 11       | 18.1 | 5/6  | 11/0       | 3/8        |
| <i>aac(3)-Ia</i>    | –                 | –                 | 12       | 19.7 | 11/1   | 12/0       | 10/2       |
| <i>aac(3)-Ia</i>    | <i>aph(3')-Ia</i> | <i>aph(3')-VI</i> | 6        | 9.8  | 4/2  | 6/0        | 3/3        |
| –                   | –                 | <i>aph(3')-VI</i> | 1        | 1.6  | 1/0  | 1/0        | 1/0        |
| –                   | –                 | –                 | 1        | 1.6  | 1/0  | 1/0        | 1/0        |


simultaneously possessed more than 1 gene encoding aminoglycoside-modifying enzymes. We noted genotypes with different combinations of genes encoding aminoglycoside-modifying enzymes, such as: 1) *aac(3)-Ia*, *aph(3')-Ia*, *aph(3')-VI*; 2) *aac(3)-Ia*, *aph(3')-VI*; 3) *aph(3')-Ia*, *aph(3')-VI*, respectively, in 6 (9.8%), 30 (49.2%), and 11 (18.1%) strains (Table 1). The frequency of occurrence of analyzed genes encoding aminoglycoside-modifying enzymes over the selected period of time (2009, 2010, and 2011) is presented in Figure 2. Taking into account the year of strain isolation, we observed in 2009 and 2010 the prevalence of *aac(3)-Ia*, *aph(3')-VI* genotype in 63% (17) and 50%, respectively (10). Among strains isolated in 2011, the dominant aminoglycoside resistance genotype consisted of *aac(3)-Ia* gene, which was present in 50% (7). Analysis of aminoglycoside resistance genotypes and susceptibility patterns to amikacin, gentamicin, and tobramycin is presented in Table 1. The most prevalent aminoglycoside resistance genotype (*aac(3)-Ia*, *aph(3')-VI*) was present in 30 (49.2%) strains, which were characterized by resistance to amikacin – 26 isolates, gentamicin – 30 isolates, and tobramycin – 16 isolates.

**Co-occurrence of OXA carbapenemases and aminoglycoside resistance genes**

Analysis of coexistence of genes encoding OXA carbapenemases and aminoglycoside-modifying enzymes is presented in Table 2. Our studies revealed the highest percentage of *A. baumannii* strains carrying simultaneously *bla<sub>OXA-40-like</sub>*, *aac(3)-Ia*, and *aph(3')-VI* genes – 44.3% (27). We also noticed the prevalence of isolates possessing 1) *bla<sub>OXA-23-like</sub>*, *aph(3')-Ia*, *aph(3')-VI* and 2) *bla<sub>OXA-40-like</sub>*, *aac(3)-Ia*, which accounted for 18.1% (11) of both genotypes.

**Discussion**

*A. baumannii* is an opportunistic pathogen often associated with nosocomial infections worldwide. Strains of this species have high ability to develop resistance to different groups of



**Figure 2.** Occurrence of *A. baumannii* strains (n=61) including genes encoding AAC(3)-Ia, APH(3')-Ia, APH(3')-VI enzymes.

antimicrobial drugs both by upregulation of intrinsic resistance mechanisms and acquisition of determinants of resistance. Multidrug-resistant *A. baumannii* strains are responsible for severe infections with high rates of morbidity and mortality, mainly affecting immunocompromised patients [6,10,17,18]. Carbapenems and aminoglycosides are antimicrobial drugs frequently used in treatment *A. baumannii* infections, often in combination therapy. Although the prevalence of resistance to these agents has become an increasing therapeutic problem, there is limited data concerning Poland [4,19].

The *bla<sub>OXA-51-like</sub>* genes naturally occur in the genome of *A. baumannii* but have been recently detected in carbapenem-resistant non-*baumannii* species of *Acinetobacter*. The detection of *bla<sub>OXA-51-like</sub>* gene can be used as a supplementary tool to identify the organism to the species level, confirmed by additional methods [10,11].

**Table 2.** Co-occurrence of genes encoding OXA carbapenemases and aminoglycoside-modifying enzymes among *A. baumannii* strains (n=61).

| Co-occurrence of genes mediating  |                                   |                               |                   |                   | Isolates |      |
|-----------------------------------|-----------------------------------|-------------------------------|-------------------|-------------------|----------|------|
| Resistance to carbapenems         |                                   | Resistance to aminoglycosides |                   |                   | No.      | %    |
| <i>bla</i> <sub>OXA-23-like</sub> | <i>bla</i> <sub>OXA-40-like</sub> | <i>aac(3)-Ia</i>              | –                 | <i>aph(3')-VI</i> | 1        | 1.6  |
| <i>bla</i> <sub>OXA-23-like</sub> | –                                 | –                             | <i>aph(3')-Ia</i> | <i>aph(3')-VI</i> | 11       | 18.1 |
| <i>bla</i> <sub>OXA-23-like</sub> | –                                 | <i>aac(3)-Ia</i>              | <i>aph(3')-Ia</i> | <i>aph(3')-VI</i> | 5        | 8.2  |
| <i>bla</i> <sub>OXA-23-like</sub> | –                                 | <i>aac(3)-Ia</i>              | –                 | <i>aph(3')-VI</i> | 2        | 3.3  |
| –                                 | <i>bla</i> <sub>OXA-40-like</sub> | <i>aac(3)-Ia</i>              | –                 | <i>aph(3')-VI</i> | 27       | 44.3 |
| –                                 | <i>bla</i> <sub>OXA-40-like</sub> | <i>aac(3)-Ia</i>              | –                 | –                 | 11       | 18.1 |
| –                                 | <i>bla</i> <sub>OXA-40-like</sub> | –                             | –                 | –                 | 1        | 1.6  |
| –                                 | <i>bla</i> <sub>OXA-40-like</sub> | –                             | –                 | <i>aph(3')-VI</i> | 1        | 1.6  |
| –                                 | –                                 | <i>aac(3)-Ia</i>              | <i>aph(3')-Ia</i> | <i>aph(3')-VI</i> | 1        | 1.6  |
| –                                 | –                                 | <i>aac(3)-Ia</i>              | –                 | –                 | 1        | 1.6  |

In our studies concerning the group of 61 *A. baumannii* strains (47 described previously and 14 currently), all carried *bla*<sub>OXA-51-like</sub> genes [13]. A study from Poland, conducted by Wroblewska et al., revealed the presence of intrinsic OXA carbapenemases among all tested strains [19]. The most frequent enzymatic mechanism of carbapenem resistance in *A. baumannii* is the production of oxacillinases encoded by genes of the *bla*<sub>OXA-23-like</sub>, *bla*<sub>OXA-40-like</sub>, and *bla*<sub>OXA-58-like</sub> lineages [10]. We have reported that genes encoding acquired carbapenemases belonging to OXA-40 group were the most frequent (40; 65.5%) among carbapenem-resistant, aminoglycoside-nonsusceptible *A. baumannii* isolates (n=61) obtained from the Specialized Hospital in Cracow. While *bla*<sub>OXA-40-like</sub> oxacillinases were dominant in Spain and Taiwan [17,20], *bla*<sub>OXA-23-like</sub> were the most prevalent in Greece and Germany [21,22].

Aminoglycoside resistance in *A. baumannii* is predominantly associated with modifications of these antimicrobials by enzymes, such as acetyltransferases (AAC), phosphotransferases (APH), and adenylyltransferases (ANT), often coexisting in a single isolate. Our analysis of the distribution of genes encoding aminoglycoside-modifying enzymes among assayed carbapenem-resistant, aminoglycoside-nonsusceptible *A. baumannii* isolates (n=61) revealed the dominance of *aac(3)-Ia* (48; 78.7%) and *aph(3')-VI* (48; 78.7%). Studies by Moniri et al. also highlighted high prevalence of *aac(3)-Ia* and *aph(3')-VI*, in 63.3% and 65%, respectively, of isolates [23]. Nemeč et al. investigated the occurrence of aminoglycoside resistance genes, and reported high percentages of strains carrying *aac(3)-Ia* (64%) and *aph(3')-VI* (52%) [24]. In our study, 77% (47) of isolates possessed 3 combinations of more than 1 gene encoding aminoglycoside-modifying enzymes, with

the predominance of *aac(3)-Ia*, *aph(3')-VI* genotype. Among 11 combinations of aminoglycoside-modifying enzymes encoding genes, Bakour et al. found 11.3% of strains possessing both *aac(3)-Ia* and *aph(3')-VI* genes [25]. Analysis of correlation between the content of resistance genes and resistance phenotypes revealed that strains with *aac(3)-Ia*, *aph(3')-VI* genotype (30) were resistant to gentamicin, and had resistant/intermediate susceptibility to amikacin (26/4) and tobramycin (16/14). Strains carrying both *aph(3')-Ia* and *aph(3')-VI* genes (11) were resistant to gentamicin and resistant/intermediate susceptible to amikacin (5/6) and tobramycin (3/8). We also noted that 1 isolate did not carry any of the mechanisms assayed, but presented an aminoglycoside-resistant phenotype. To conclude, in some strains there was no correlation between the presence of assayed resistance genes and susceptibility phenotype, which can be explained by the occurrence of other aminoglycoside resistance genes, overexpression of an efflux pump such as AdeABC, or additional resistance mechanisms. These findings are also supported by Salazar De Vegas et al. [18] and Nemeč et al. [24].

This study analyzed the occurrence of various genes that confer resistance to carbapenems and aminoglycosides in the group of *A. baumannii* strains resistant to carbapenems and nonsusceptible to aminoglycosides in Cracow, Poland. Our research revealed the co-occurrence of selected carbapenem and aminoglycoside resistance genes in 8 combinations, with the dominance of isolates carrying *bla*<sub>OXA-40-like</sub>, *aac(3)-Ia* and *aph(3')-VI* genes (44.3%). The coexistence of genes accountable for resistance to carbapenems and aminoglycosides in isolates of *A. baumannii* was also found by Salazar de Vegas et al., Endimiani et al., Nigro et al., and Sung et al. [18,26–28].

*A. baumannii* multidrug resistance appears to be a consequence of accumulation of mutations and the acquisition of resistance determinants by the transfer of plasmids, transposons, and integrons, which may lead to the formation of clusters of resistance genes, termed “resistance islands” [6]. The high genetic plasticity of the *A. baumannii* genome contributes to high adaptation, virulence capacities, and multidrug resistance results in its ability to persist in the hospital environment and survive antibiotic treatment [29].

## Conclusions

1. *bla*<sub>OXA-40-like</sub> were the most prevalent genes encoding acquired OXA carbapenemases among *A. baumannii* strains isolated in Cracow, Poland.

## References:

1. Garnacho-Montero J, Amaya-Villar R: Multiresistant *Acinetobacter baumannii* infections: epidemiology and management. *Curr Opin Infect Dis*, 2010; 23: 332–39
2. Towner KJ: *Acinetobacter*: an old friend, but a new enemy. *J Hosp Infect*, 2009; 73: 355–63
3. Dijkshoorn L, Nemec A, Seifert H: An increasing threat in hospitals: multidrug-resistant *Acinetobacter baumannii*. *Nat Rev Microbiol*, 2007; 5: 939–51
4. Szejbach A, Mikucka A, Bogiel T, Gospodarek E: Usefulness of phenotypic and genotypic methods for metallo-beta-lactamases detection in carbapenem-resistant *Acinetobacter baumannii* strains. *Med Sci Monit Basic Res*, 2013; 19: 32–36
5. Neonakis IK, Spandidos DA, Petinaki E: Confronting multidrug-resistant *Acinetobacter baumannii*: a review. *Int J Antimicrob Agents*, 2011; 37: 102–9
6. Poirel L, Bonnin RA, Nordmann P: Genetic basis of antibiotic resistance in pathogenic *Acinetobacter* species. *IUBMB Life*, 2011; 63: 1061–67
7. Gordon NC, Wareham DW: Multidrug-resistant *Acinetobacter baumannii*: mechanisms of virulence and resistance. *Int J Antimicrob Agents*, 2010; 35: 219–26
8. Zavascki AP, Carvalhaes CG, Picao RC, Gales AC: Multidrug-resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: resistance mechanisms and implications for therapy. *Expert Rev Anti Infect Ther*, 2010; 8: 71–93
9. Kempf M, Rolain JM: Emergence of resistance to carbapenems in *Acinetobacter baumannii* in Europe: clinical impact and therapeutic options. *Int J Antimicrob Agents*, 2012; 39: 105–14
10. Howard A, O'Donoghue M, Feeney A, Sleator RD: *Acinetobacter baumannii*: an emerging opportunistic pathogen. *Virulence*, 2012; 3: 243–50
11. Lee YT, Kuo SC, Chiang MC et al: Emergence of carbapenem-resistant non-*baumannii* species of *Acinetobacter* harboring a *bla*OXA-51-like gene that is intrinsic to *A. baumannii*. *Antimicrob Agents Chemother*, 2012; 56: 1124–27
12. Vakulenko SB, Mobashery S: Versatility of aminoglycosides and prospects for their future. *Clin Microbiol Rev*, 2003; 16: 430–50
13. Nowak P, Paluchowska P, Budak A: Distribution of *bla*OXA genes among carbapenem-resistant *Acinetobacter baumannii* nosocomial strains in Poland. *New Microbiol*, 2012; 35: 317–25
14. Clinical and Laboratory Standards Institute. 2008. Performance standards for antimicrobial susceptibility testing; 18<sup>th</sup> informational supplement. CLSI document M100-S18. Clinical and Laboratory Standards Institute, Wayne, PA
15. Woodford N, Ellington MJ, Coelho JM et al: Multiplex PCR for genes encoding prevalent OXA carbapenemases in *Acinetobacter* spp. *Int J Antimicrob Agents*, 2006; 27: 351–53
16. Noppe-Leclercq I, Wallet F, Haentjens S et al: PCR detection of aminoglycoside resistance genes: a rapid molecular typing method for *Acinetobacter baumannii*. *Res Microbiol*, 1999; 150: 317–22
17. Ben RJ, Yang MC, Hsueh JC et al: Molecular characterisation of multiple drug-resistant *Acinetobacter baumannii* isolates in southern Taiwan. *Int J Antimicrob Agents*, 2011; 38: 403–8
18. Salazar De Vegas EZ, Nievesm B, Ruiz M et al: Molecular epidemiology and characterization of resistance mechanisms to various antimicrobial agents in *Acinetobacter baumannii* isolated in Mérida, Venezuela. *Med Sci Monit*, 2007; 13(4): BR89–94
19. Wroblewska MM, Towner KJ, Marchel H, Luczak M: Emergence and spread of carbapenem-resistant strains of *Acinetobacter baumannii* in a tertiary-care hospital in Poland. *Clin Microbiol Infect*, 2007; 13: 490–96
20. Villalón P, Valdezate S, Medina-Pascual MJ et al: Epidemiology of the *Acinetobacter*-derived cephalosporinase, carbapenem-hydrolysing oxacillinase and metallo-β-lactamase genes, and of common insertion sequences, in epidemic clones of *Acinetobacter baumannii* from Spain. *J Antimicrob Chemother*, 2013; 68: 550–53
21. Schleicher X, Higgins PG, Wisplinghoff H et al: Molecular epidemiology of *Acinetobacter baumannii* and *Acinetobacter nosocomialis* in Germany over a 5-year period (2005–2009). *Clin Microbiol Infect*, 2013; 19: 737–42
22. Liakopoulos A, Miriagou V, Katsifas EA et al: Identification of OXA-23-producing *Acinetobacter baumannii* in Greece, 2010 to 2011. *Euro Surveill*, 2012; 17(11): pii: 20117
23. Moniri R, Farahani RK, Shajari G et al: Molecular epidemiology of aminoglycosides resistance in *Acinetobacter* spp. with emergence of multidrug-resistant strains. *Iran J Public Health*, 2010; 39: 63–68
24. Nemec A, Dolzani L, Brisse S et al: Diversity of aminoglycoside-resistance genes and their association with class 1 integrons among strains of pan-European *Acinetobacter baumannii* clones. *J Med Microbiol*, 2004; 53: 1233–40
25. Bakour S, Touati A, Sahli F et al: Antibiotic resistance determinants of multidrug-resistant *Acinetobacter baumannii* clinical isolates in Algeria. *Diagn Microbiol Infect Dis*, 2013; 76: 529–31
26. Endimiani A, Hujer KM, Hujer AM et al: *Acinetobacter baumannii* isolates from pets and horses in Switzerland: molecular characterization and clinical data. *J Antimicrob Chemother*, 2011; 66: 2248–54
27. Nigro SJ, Post V, Hall RM: Aminoglycoside resistance in multiply antibiotic-resistant *Acinetobacter baumannii* belonging to global clone 2 from Australian hospitals. *J Antimicrob Chemother*, 2011; 66: 1504–9
28. Sung JY, Kwon KC, Cho HH, Koo SH: Antimicrobial resistance determinants in imipenem-nonsusceptible *Acinetobacter calcoaceticus-baumannii* complex isolated in Daejeon, Korea. *Korean J Lab Med*, 2011; 31: 265–70
29. Imperi F, Antunes LC, Blom J et al: The genomics of *Acinetobacter baumannii*: insights into genome plasticity, antimicrobial resistance and pathogenicity. *IUBMB Life*, 2011; 63: 1068–74

## Acknowledgements

The Microbiological Laboratory of the Department of Laboratory Diagnostics of the Specialized Rydygiel's Hospital in Cracow, Poland is gratefully acknowledged for providing the isolates used in this study.