

Acute digit ischemia due to a ruptured digital collateral artery aneurysm in a patient of hemophilia: A case report

Ruben Dukan¹, Maxime Lacroix², Emilie Bertin¹, Guillaume Reverdito², Franck Fitoussi³, Adeline Cambon Binder⁴

Learning Point of the Article:

In the case of a hemophilic patient, surgical management is necessary and medical treatment alone is not sufficient.

Abstract

Introduction: Digital artery aneurysms are a rare event but cases have been reported in the literature. The hemostasis disorders make these aneurysms particularly dangerous with potentially irreversible hand complications: Compression of adjacent vascular and nervous structures, embolization of associated thrombi, finger ischemia, and necrosis.

Case Report: We reported a case of digital ischemia due to a ruptured aneurysm of a digital collateral artery, leading to the diagnosis of congenital hemophilia A. Hematoma evacuation allowed finger revascularization. Complete symptom resolution required ligature excision associated with Factor VIII supplementation.

Conclusion: Aneurysms of the digital collateral arteries are a rare event. In the case of a hemophilic patient, surgical management is necessary, and medical treatment alone is not sufficient. The consequences of a ruptured aneurysm in this type of patient can be serious. A close monitoring is required to ensure the absence of early recurrence.

Keywords: Hemophilia, digital aneurysm, ischemia.

Introduction

Digital artery aneurysms are a rare event and cases have been reported in the literature [1]. Most of these aneurysms were pseudoaneurysms, corresponding to perforation of the arterial wall, related to direct blunt trauma (wound or vascular procedure) or repeated microtrauma. The hemostasis disorders make these aneurysms particularly dangerous with potentially irreversible hand complications: Compression of adjacent vascular and nervous structures, embolization of associated thrombi, finger ischemia, and necrosis. Several authors in the literature reported the management of these lesions by aneurysm

excision and ligation in the majority of cases versus excision and grafting/direct suture [2-8]. In case of associated hemophilia, surgical management is usually associated with coagulation factor deficiency supplementation [9]. None of the authors reported any initial or post-operative complication in the hand. The diagnosis of hemophilia was known preoperatively. We reported the case of a patient with a pseudoaneurysm of a digital collateral artery complicated by acute ischemia due to aneurysm rupture that lead to diagnosis of hemophilia.

The patient gave his consent for the publication of this clinical case.

Author's Photo Gallery



Dr. Ruben Dukan



Dr. Maxime Lacroix



Dr. Emilie Bertin



Dr. Guillaume Reverdito



Dr. Franck Fitoussi



Dr. Adeline Cambon Binder

Access this article online

Website:
www.jocr.co.in

DOI:
<https://doi.org/10.13107/jocr.2023.v13.i11.4010>

¹Department of Orthopedic Surgery, Hand, Upper Limb and Peripheral Nerve Surgery, Georges-Pompidou European Hospital (HEGP), Paris, France,

²Department of Radiology, Georges-Pompidou European Hospital (HEGP), Paris, France,

³Department of Pediatric Orthopedics, Sorbonne University, Paris, France,

⁴Department of Orthopedic and Hand Surgery, Sorbonne University, Saint Antoine Hospital, Paris, France.

Address of Correspondence:

Dr. Dukan Ruben,
Hand, Upper Limb and Peripheral Nerve Surgery, Georges-Pompidou European Hospital (HEGP), Paris, France.
E-mail: ruben.dukan@aphp.fr

Submitted: 22/07/2023; Review: 29/08/2023; Accepted: Oct 2023; Published: November 2023

DOI: <https://doi.org/10.13107/jocr.2023.v13.i11.4010>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License <https://creativecommons.org/licenses/by-nc-sa/4.0/>, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms



Figure 1: Radial digital collateral artery false aneurysm.

Case Report

A 36-year-old man, right-handed, non-smoker, sportive, manual worker, and without any medical background, presented with a stab wound of the left index in palmar zone 2 International Federation of Societies for Surgery of the Hand Classification [10]. Surgical procedure reported an opening of the flexor sheath without tendon injury and a continuous digital radial collateral artery with no active bleeding.

After 13-day interval, the patient presented with scar swelling for 2 days. On clinical examination, vital signs were normal, without fever. There was a hypoesthesia of the radial fingertip. Palpation showed a non-compressible, pulseless but inflammatory mass. We hypothesized a surgical site infection with abscessed collection. An ultrasound assessment was performed and reported a cystic mass without any sign of blood flow inside.

A surgical revision was performed. An aneurysm of the digital radial collateral artery with compression of the radial collateral nerve was identified (Fig. 1). A clamping test was performed using a clamp on the digital radial collateral artery. The finger remained vascularized on the digital ulnar collateral artery. A ligation-excision of the aneurysm was performed. Anatomopathological analysis concluded to a false aneurysm. Bacteriological examination was negative. Sensitivity of the radial fingertip was recovered immediately after surgery.

After a 15-day interval, the patient presented with intense pain (visual analog scale [VAS] 8/10) in the finger with anesthesia of

the radial fingertip and reappearance of a mass at the level of the scar in less than a day. An upper limb computed tomography (CT) angio was performed in emergency and showed the presence of an aneurysm below the previous ligation excision (Fig. 2). VAS was suddenly at 10/10. Clinical examination revealed acute ischemia of the finger. An emergency revision was performed.

We noticed a ruptured aneurysm extending to the origin of the digital radial collateral artery from the common digital artery 1–2. Evacuation of the hematoma allowed immediate revascularization of the finger. The aneurysm was excised, and ligation was made. Sensory recovery was immediate postoperatively.

Due to the recurrence, the patient was referred to the department of hematology and vascular medicine.

The imaging assessment (CT angio) did not show any other aneurysmal lesions. The patient had no previous bleeding history. There was no family history of bleeding or thrombosis. The bleeding assessment tool (BAT comprises 14 categories for assessing bleeding symptoms retrospectively and it has been shown that a high bleeding score is associated with the presence of an inherited bleeding disorder) score was 4 (A score ≥ 4 is considered “abnormal”) and the partial thromboplastin time (PTT) ratio was 1.46 (corresponding to an intrinsic coagulation pathway defect, and thus elevated PTT). Endogenous factors were investigated and a minor factor VIII (FVIII) deficiency of 20% was found. The diagnosis of minor hemophilia A was made.

A protocol of supplementation was performed: Immediately 3000 IU of FVIII and then 2000 IU every 8 h until FVIII value correction. Supplementation of FVIII deficiency and the specialized management allowed hemostasis and avoidance of early aneurysm recurrence.

The patient recovered full hand function at 2 weeks postoperatively. At 1 year, we did not observe any aneurysm recurrence.

Discussion

We reported a rare case of acute finger ischemia by compression of the intact digital collateral artery following a rupture of a




Figure 2: Hand computed tomography (CT) scan showing a recurrence of a ruptured radial digital collateral artery false aneurysm with a compressive hematoma.

digital collateral artery aneurysm resulting in the diagnosis of minor hemophilia A.

Hemophilia is an X-linked disorder due to clotting factor VIII deficiency. Congenital hemophilia A is not known to be associated with arterial aneurysms. However, these lesions can be particularly dangerous in the hand with the risk of ischemia and necrosis of the finger in case of rupture as we have reported.

We believed that the heavy bleeding of hemophilia causes extended intimal lesions with a perivascular hematoma in continuity with the vessel lumen, and pseudocavity surrounded by the adventitia, perivascular tissue, and fibrotic tissue originating from the hematoma as explained by Mann [11]. This is why we observed in our patient the aneurysm several days after the initial intimal injury. Various authors in the literature agree that hemophilia constitutes an excess risk in case of aneurysm [3, 7, 12]. The chronology of our case report is consistent with the various cases in the literature on post-traumatic hand pseudoaneurysm [3, 13]. There is an intimal lesion that will progressively dissect due to the continuous bleeding. This kind of complication must be differentiated from the risk of embolization of arterial aneurysms that can also lead to digital ischemia [4]. We can consider this risk to be lower in the hemophilia patient.

To date, the prevalence of pseudoaneurysms in patients with hemophilia is not known and the optimal strategy has yet to be defined. Although Drew et al. [8] suggested that factor replacement was an adequate option in healthy hemophilia

Author	Localization	Patient age	Type of aneurysm	Mecanism	Known pathology	Discovery mode	Post-traumatic delay	Management	Recurrance
Lokey et al. [13]	Hand – first web space	37 y.o.	FALSE	Stab wound	Hemophilia	Asymptomatic pulsatile mass	10 days	1. FVIII infusion (failure) 2. Excision ligation+FVIII infusion	None
Ballas et al. [3]	Hand - Fingertip	40 y.o.	FALSE	Struck with a hammer	Hemophilia A (partial deficiency VIII 25%)	Asymptomatic pulseless mass	4 months	Excision ligation	None
Adant et al. [7]	Hand - Thumb	55 y.o.	TRUE	-	Hemophilia B (Factor IX deficit)	Painful pulseless mass	N.A.	FIX infusion+excision ligation	None
Drew et al. [8]	Elbow - antecubital fossa	7 y.o.	FALSE	Veinpuncture	Mild Hemophilia A (FVIII=6%)	Asymptomatic pulseless mass	1 month	18 days daily FVIII infusion	None
Wisniewski and Szopinski [6]	Elbow - antecubital fossa	26 y.o.	TRUE	-	Mild Hemophilia A (FVIII=2%)	Pulseless mass - pricking sensation, numbness, dysaesthesia in the fingers	N.A.	FVIII infusion+Excision +vein graft anastomose	None
Harkin et al. [5]	Wrist - radial artery	57 y.o.	FALSE	Arterial puncture	Hemophilia A	Pulsatile swelling – skin necrosis – septic cellulitis	12 days	FVIII infusion+Excision ligation	None
Fields et al. [2]	Wrist - radial artery	2 weeks	FALSE	Arterial puncture	-	Pulsatile mass	Immediate	Compression - thrombosis	None
Filis et al. [4]	Wrist - radial artery	45 y.o.	TRUE	-	Mild Hemophilia A	Painful pulsatile mass	N.A.	FVIII infusion+excision +anastomose	None

FVIII: Factor VIII

Table 1: Upper limb aneurysm – literature review.



patients without inhibitors when pseudoaneurysms are small, we believe that these patients should be systematically managed with either excision/ligation or direct suture/grafting, considering the risk of acute ischemia by rupture and compression. Given the extensive intimal lesions that we have reported, we believe that the ligation solution is the safest strategy to avoid the development of an aneurysm proximal to the graft. It is obvious that in case of insufficient vascularization by a single artery, an anastomose with graft is required [3]. Lokey et al. [13] reported a case of failure of medical treatment, alone with continuous aneurysm growth until rupture and the need for emergency hemostasis and ligation.

A blood hemostasis test is required in case of digital aneurysm (true or pseudo aneurysm). The resolution of the symptoms in our case was only observed after FVIII supplementation. Elective surgical repair under proper FVIII concentrate infusion is safe, whereas surgery in an emergency setting may be hazardous for the patient.

In the upper limb, cases of aneurysm are sporadic, and it is not possible to know their frequency (Table 1). The main

difference with our case is the absence of a diagnosis of hemophilia at the time of management, except for the case reported by Fields concerning a 2-month-old child [2].

Conclusion

Aneurysms of the digital collateral arteries are a rare event. In the case of a hemophilic patient, surgical management is necessary and medical treatment alone is not sufficient. The consequences of a ruptured aneurysm in this type of patient can be fatal. A close monitoring is required to ensure the absence of early recurrence.

Clinical Message

Digital aneurysm is a rare event. Patient should be screened for hemophilia. Medical treatment is a mandatory complement to surgery to avoid recurrence.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given the consent for his/ her images and other clinical information to be reported in the journal. The patient understands that his/ her names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Conflict of interest: Nil **Source of support:** None

References

1. Sheikh Z, Selvakumar S, Goon P. True aneurysm of the digital artery: A case report and systematic literature review. *J Surg Case Rep* 2020;2020:rjz400.
2. Fields JM, Saluja S, Schwartz DS, Touloukian RJ, Keller MS. Hemophilia presenting in an infant as a radial artery pseudoaneurysm following arterial puncture. *Pediatr Radiol* 1997;27:763-4.
3. Ballas KD, Rafailidis SE, Kouskouras KA, Fardellas J, Alatsakis MB, Sakadamis AK. False aneurysm of the proper palmar digital artery--report of a case. *Acta Chir Belg* 2006;106:249-51.
4. Filis K, Arhontovassilis F, Theodorou D, Theodossiades G, Manouras A. True radial artery aneurysm in a mild haemophilia A patient. *Haemophilia* 2007;13:440-2.
5. Harkin DW, Connolly D, Chandrasekar R, Anderson M, Blair PH, Hood JM, et al. Radial artery mycotic pseudoaneurysm in a haemophiliac: A potentially fatal complication of arterial catheterization. *Haemophilia* 2002;8:721-4.
6. Wiszniewski A, Szopiński P. Surgical treatment of arteriovenous fistula and brachial artery aneurysm in a patient with mild haemophilia A. *Haemophilia* 2011;17:548-50.
7. Adant JP, Grattagliano B, Fissette J. True aneurysm of digital artery in hemophilia. A case report. *Ann Chir Main Memb Super* 1994;13:278-81.
8. Drew M, Yu JC, Michaels LA. Arterial pseudoaneurysms in haemophilia: Conservative treatment with observation and factor replacement. *Haemophilia* 2011;17:550-2.
9. Diplaris KT, Karfis EA, Ampatzidou F, Ananiadou OG, Vakalopoulou S, Madesis A, et al. Acute type-a dissection in a patient with severe hemophilia A. *J Cardiothorac Vasc Anesth* 2012;26:660-3.
10. Castel LC, Hurst SA, Masmajejan E, Gregory TM. Hand wounds: An analysis of topography and related damage to

underlying structures. *Injury* 2022;53:4048-53.

11. Mann HA, Goddard NJ, Lee CA, Brown SA. Periarticular aneurysm following total knee replacement in hemophilic arthropathy. A case report. *J Bone Joint Surg Am* 2003;85:2437-40.

12. Sleightholm R, Wahlmeier S, Carson JS, Drincic A,

Lazenby A, Foster JM. Massive adrenal vein aneurysm mimicking an adrenal tumor in a patient with hemophilia A: A case report and review of the literature. *J Med Case Rep* 2016;10:343.

13. Lokey H, Phelps DB, Boswick JA. Traumatic false aneurysm of the hand in hemophilia. *J Trauma* 1978;18:283-4.

Conflict of Interest: Nil

Source of Support: Nil

Consent: The authors confirm that informed consent was obtained from the patient for publication of this case report

How to Cite this Article

Dukan R, Lacroix M, Bertin E, Reverdito G, Fitoussi F, Binder AC. Acute digit ischemia due to a ruptured digital collateral artery aneurysm in a patient of hemophilia: A case report. *Journal of Orthopaedic Case Reports* 2023 November;13(11): 70-74.

