CORRECTION

Correction: The effect of national antenatal care guidelines and provider training on obstetric danger sign counselling: a propensity score matching analysis of the 2014 Ethiopia service provision assessment plus survey

Tebikew Yeneabat[1,2]*, Andrew Hayen[2], Theodros Getachew[3] and Angela Dawson[2]

Correction to: Reproductive Health (2022) 19:132
https://doi.org/10.1186/s12978-022-01442-6

After publication of this article [1], the authors reported that in the section 'Sample size and selection process' in the first sentence, the number '1237' should have read '1327'.

Moreover, reference [50] should have been replaced by [53]. References [54–87] were renumbered.

The original article [1] has been corrected.

Author details
[1]Department of Midwifery, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia. [2]School of Public Health, Faculty of Health, University of Technology Sydney, Sydney, Australia. [3]Health System and Reproductive Health Research Directorate, Ethiopian Public Health Institute, Addis Ababa, Ethiopia.

Published online: 17 August 2022

The original article can be found online at https://doi.org/10.1186/s12978-022-01442-6.

*Correspondence: wait4myfather@gmail.com
[1] Department of Midwifery, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia
Full list of author information is available at the end of the article

Reference

1. Yeneabat T, Hayen A, Getachew T, Dawson A. The effect of national antenatal care guidelines and provider training on obstetric danger sign counselling: a propensity score matching analysis of the 2014 Ethiopia service provision assessment plus survey. Reprod Health. 2022;19:132. https://doi.org/10.1186/s12978-022-01442-6.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.