

A new species of *Neoperla* from China, with a redescription of the female of *N. mnong* Stark, 1987 (Plecoptera, Perlidae)

Zhi-Teng Chen¹, Yu-Zhou Du¹

¹ School of Horticulture and Plant Protection & Institute of Applied Entomology, Yangzhou University, Yangzhou 225009, China

Corresponding author: Yu-Zhou Du (yzdu@yzu.edu.cn)

Academic editor: P. Stoev | Received 24 April 2016 | Accepted 24 August 2016 | Published 12 September 2016

<http://zoobank.org/7C4379BB-BD14-4486-86F5-C10C90197042>

Citation: Chen Z-T, Du Y-Z (2016) A new species of *Neoperla* from China, with a redescription of the female of *N. mnong* Stark, 1987 (Plecoptera, Perlidae). ZooKeys 616: 103–113. doi: 10.3897/zookeys.616.8996

Abstract

A new species of the *Neoperla clymene* group (Plecoptera, Perlidae), *N. chebalinga* sp. n. from Guangdong Province of southern China is described, illustrated, and compared with related taxa. The new species is characterized by the slender aedeagal tube, strongly sclerotized dorsally, and weakly sclerotized ventrally with an upcurved, medial, finger-like membranous lobe. Additionally the aedeagal sac gradually tapers to a blunt apex with a dorsoapical patch of spines. A supplementary description of the female of *N. mnong* Stark, 1987 from Guangdong Province, China is also given.

Keywords

China, *Neoperla*, new species, Perlidae, Plecoptera

Introduction

Neoperla Needham (1905) is the most species-rich perlid genus in China with at least 57 species (DeWalt et al. 2016). Contributions to Chinese species of *Neoperla* were made by Chu (1929), Wu and Claassen (1934), Wu (1935, 1938, 1948, 1962, 1973), Yang and Yang (1992, 1995, 1998), Du (1998, 1999, 2000a, 2000b), Du et al. (1999,

2001), Du and Sivec (2004, 2005), Du and Wang (2005, 2007), Sivec and Zwick (1987), Li and Wang (2011), Li et al. (2011a, 2011b, 2012a, 2012b, 2013a, 2013b, 2014a, 2014b), Qin et al. (2013), Wang et al. (2013a, 2013b), Li and Zhang (2014), Kong et al. (2014), Wang et al. (2014), Chen and Du (2015, 2016). Herein, a new species of this genus is described from Guangdong Province, a coastal region of southern China. Additionally, a supplementary description and new illustrations for the female of *N. mnong* Stark, 1987 from Guangdong Province are provided to indicate the variation of the female subgenital plate and head pattern as compared to the original description of Vietnamese material.

Material and methods

Specimens used in this study were collected from riparian areas by hand and preserved in 75% ethanol. Abdomens were cleared in 10% NaOH. Details of the morphology were studied with a Leica MZAPO microscope, and color illustrations were taken by Leica SZ45 and S4800 FESEM, both in Yangzhou, Jiangsu Province, China. All studied material, including the holotype and paratypes of the new species, are deposited in the Insect Collection of Yangzhou University, China.

Taxonomy

Neoperla chebalinga sp. n.

<http://zoobank.org/960227BF-2234-483D-9363-3C7A40D0F32D>

Figs 1–10

Type material. Holotype male: China: Guangdong Province, Chebaling Nature Reserve, 114.1320E, 24.7128N, 17 June, 2009, leg. Hai-Yang Xue and Bo Yu. Paratypes: 4 males and 2 females, same data as holotype.

Diagnosis. Distinguishing characteristics of this species include a slender aedeagal tube that has an upcurved, medial, finger-like, membranous lobe on the ventral surface, and a short, blunt, ventrally curved aedeagal sac with a dorsoapical patch of spines.

Adult habitus. General body color yellow patterned with brown. Head slightly wider than pronotum with a brown stigma on anterior frons and a subtriangular dark brown area covering ocelli; compound eyes dark and antennae brown (Figs 1, 9–10). Pronotum disc yellow with pale brown rugosities (Fig. 1). Wing membrane subhyaline, veins brown; legs yellow, femora grading to brown distally. Cerci pale (Figs 9–10).

Male (Figs 1–5, 9). Forewing length 13.5–14.0 mm, hindwing length 11.5–12.0 mm ($n = 5$). The posterior margin of tergum 7 with a raised process densely covered with sensilla basiconica. Tergum 8 bears a recurved oval process with small spines at the distal margin (Figs 2–3). Tergum 9 without sensilla basiconica. Hemitergal processes of tergum 10 slightly curved anteriorly (Fig. 2). Aedeagal tube slender, strongly



Figure 1. *Neoperla chebalinga* sp. n. Male head and pronotum, dorsal view.



Figures 2–3. *Neoperla chebalinga* sp. n. Male terminalia: **2** dorsal view **3** lateral view.

sclerotized dorsally, weakly sclerotized ventrally except for the basal bulb and an up-curved, ventromedial, finger-like membranous lobe. Aedeagal sac short and gradually tapered to a blunt apex, curved ventrally with a dorsoapical patch of spines (Figs 4–5).

Female (Figs 6–7, 10). Forewing length 15.5–16.0 mm, hindwing length 13.0–13.5 mm ($n = 2$). General color and pattern similar to males. Subgenital plate of sternum 8 slightly produced and lightly sclerotized posteromedially (Figs 6, 10). Va-



Figure 4–5. *Neoperla chebalinga* sp. n. Aedeagus: **4** *In situ*, lateral view **5** Everted, lateral view.




Figure 6. *Neoperla chebalinga* sp. n. Female terminalia, ventral view.




Figure 7. *Neoperla chebalinga*, sp. n. Female genitalia, vagina and spermathecum, lateral view.




Figure 8. *Neoperla chebalinga* sp. n. Egg, lateral view.

gina elongate-oval, lined sparsely along lateral margins. Spermathecal stalk short, spermatheca oval and curled at tip (Fig. 7).

Eggs (Fig. 8). Outline oval. Collar short and surrounded by 1–2 irregular rows of follicle cell impressions. Primary striae join rim on sides of collar, narrow and widely spaced; each pair of primary striae enclose many secondary striae; sulci with dense pits. Lid small consisting of follicle cell impressions (Fig. 8).

Etymology. The species is named after the type locality, Chebaling Nature Reserve.

Remarks. The new species is a member of the *N. clymene* group possessing a slender and mostly sclerotized aedeagal tube. The processes of terga 7–8 and aedeagus of the new species are similar to those of *N. furcostyla* Li & Qin, 2013 (in: Li et al. 2013b) and *N. forcipata* Yang & Yang, 1992 (Yang and Yang 1992). However, in *N. furcostyla* and *N. forcipata*, the aedeagal tube has a subapical fork instead of the medial finger-like lobe as in *N. chebalinga*. In addition, the aedeagal sacs of *N. furcostyla* and *N. forcipata* are not curved ventrally (see Figs 1–2 in: Li et al. 2013b and fig. 1 in Yang and Yang 1992).




Figure 9–10. *Neoperla chebalinga* sp. n. Habitus: **9** Male habitus, dorsal view **10** Female habitus, dorsal view.

***Neoperla mnong* Stark, 1987**

Figs 11–13

Neoperla mnong Stark, 1987: 48; Stark & Sivec 2008: 33; Du 1998: 393; Li et al. 2012: 22; Wang et al. 2013b: 87.

Material examined. 1 male, 4 females, China: Guangdong Province, Chebalong Nature Reserve, 114.1320E, 24.7128N, 17 June, 2009, leg. Hai-Yang Xue and Bo Yu.

Description of the female Chinese specimens. Female (Figs 11–13). Forewing length 12.0–12.5 mm, hindwing length 10.0–10.5 mm ($n = 4$). General color yel-



Figure 11. *Neoperla mnong* Stark. Female habitus, dorsal view.




Figure 12. *Neoperla mnong* Stark. Female terminalia, ventral view.




Figure 13. *Neoperla mnong* Stark. Female genitalia, vagina and spermathecum, lateral view.

low to brown. Head nearly as wide as pronotum, with a subtriangular dark brown area covering ocelli; compound eyes dark and antennae pale brown. Pronotum disc yellow with pale brown rugosities. Wing membrane subhyaline, veins pale brown; legs yellow without darker markings. Cerci pale brown (Fig. 11). Subgenital plate of sternum 8 forming a small rounded sclerite with a posteromedial notch (Fig. 12). Vagina oval, lined densely around base of spermathecal stalk. Spermathecal stalk and spermatheca short, slender and curled at tip; a single accessory gland occurs near spermatheca tip (Fig. 13).

Distribution. China (Guangdong Province: Conghua County, Liuxi River; Shixing County, Chebaling Nature Reserve. Guangxi Province: Jinxiu suburb; Tianlin County, Pinglang). Vietnam (Di Linh, Dak Song, Gia Lai, Nghe An, Pleiku). Thailand (Chanthaburia).

Remarks. This species is expected to be widespread in southern Asia. The male of *N. mnong* has been described from Vietnam by Stark (1987) and well-illustrated by Stark and Sivec (2008) from Vietnamese material. This species has been previously recorded from China by Du (1998) and Li et al. (2012b). Females of *N. mnong* from Vietnam were described as *Javanita costalis* by Navás (1932) and later described by Stark and Sivec (2008). In this study, females collected from Guangdong Province of China in 2009 are described and illustrated for the first time. The subgenital plate of these specimens is smaller and pigment patch over the ocelli is larger (for comparison see Figs 40, 43 in Stark and Sivec 2008).

Acknowledgments

We are very grateful to Dr. Boris C. Kondratieff for assistance with the English syntax. This research was supported by the National Natural Science Foundation of China (No. 31071958).

References

- Chu YT (1929) Descriptions of four new species and one new genus of stone-flies in the family Perlidae from Hangchow. *The China Journal* 10: 88–92.
- Chen ZT, Du YZ (2015) The first record of a species of Perlidae from Jiangsu Province, China: a new species of *Neoperla* (Plecoptera: Perlidae). *Zootaxa* 3974(3): 424–430. doi: 10.11646/zootaxa.3974.3.9
- Chen ZT, Du YZ (2016) A new species of *Neoperla* (Plecoptera: Perlidae) from Zhejiang Province of China. *Zootaxa* 4093(4): 589–594. doi: 10.11646/zootaxa.4093.4.12
- DeWalt RE, Maehr MD, Neu-Becker U, Stueber G (2016) Plecoptera Species File Online. <http://Plecoptera.SpeciesFile.org> [accessed 23 Mar 2016]
- Du YZ (1998) Two new record species of genus *Neoperla* Needham (Plecoptera: Perlidae) from China. *Journal of Zhejiang Agricultural University* 24(4): 393.
- Du YZ (1999) Plecoptera. In: Huang BK (Ed.) *Fauna of Insects in Fujian Province of China*. Vol. 3. Fujian Science and Technology Publishing house, Fujian, 301–335.
- Du YZ (2000a) *Neoperla magisterchoui*, a new species of the genus *Neoperla* Needham (Plecoptera: Perlidae) from China. In: Zhang YL (Ed.) *Systematic and Faunistic Research on Chinese Insects. Proceedings of the 5th National Congress of Insect Taxonomy*. China Agriculture Press, Beijing, 1–3.
- Du YZ (2000b) Two new species of the genus *Neoperla* Needham (Plecoptera: Perlidae: Perlinae) from Guizhou, China. *Entomotaxonomia* 22(1): 1–5.
- Du YZ, Sivec I (2004) Plecoptera: Perlidae, Nemouridae, Leuctridae. In: Yang XK (Ed.) *Insects from Mt. Shiwanashan Area of Guangxi*. China Forestry Publishing House, Beijing, 39–45.
- Du YZ, Sivec I (2005) Plecoptera. In: Yang XK (Ed.) *Insect Fauna of Middle-west Qinling Range and South Mountains of Gansu Province*. Science Press, Beijing, 38–54.
- Du YZ, Sivec I, He JH (1999) A checklist of the Chinese species of the family Perlidae (Plecoptera: Perloidea). *Acta Entomologica Slovenica* 7: 59–67.
- Du YZ, Sivec I, Zhao MS (2001) Plecoptera. In: Wu H, Pan CW (Eds) *Insects of Tianmushan National Nature Reserve*. Science Press, Beijing, 69–80.
- Du YZ, Wang ZJ (2005) Plecoptera: Leuctridae, Nemouridae, Perlidae and Peltoperlidae. In: Yang MF, Jin DC (Eds) *Insects from Dashahe Nature Reserve of Guizhou*. Guizhou People Press, Guizhou, 51–57.
- Du YZ, Wang ZJ (2007) Nemouridae and Perlidae. In: Li ZZ, Yang MF, Jin DC (Eds) *Insects from Mountain Leigongshan Landscape of Guizhou*. Guizhou Science and Technology Publishing House, Guizhou, 84–90.
- Kong FB, Lv JF, Li WH (2014) Three new species of *Neoperla* in the *montivaga* group (Plecoptera: Perlidae) from China. *Zootaxa* 3841(3): 429–438. doi: 10.11646/zootaxa.3841.3.7
- Li WH, Liang HY, Li WL (2013a) Review of *Neoperla* (Plecoptera: Perlidae) from Zhejiang Province, China. *Zootaxa* 3652: 353–369. doi: 10.11646/zootaxa.3652.3.4
- Li WH, Li S, Feng GW, Wang YB (2014a) Species of *Neoperla* (Plecoptera: Perlidae) from Henan Province, China. *Zootaxa* 3838(2): 174–182. doi: 10.11646/zootaxa.3838.2.2
- Li WH, Murányi D, Wang RF (2014b) Species of *Neoperla* (Plecoptera: Perlidae) from Yunnan Province, China. *Zootaxa* 3753(1): 1–9. doi: 10.11646/zootaxa.3753.1.1

- Li WH, Wang GQ, Lu WY (2012a) Species of *Neoperla* (Plecoptera: Perlidae) from Hubei, China. Zootaxa 3478: 32–37.
- Li WH, Wang GQ, Li WL, Murányi D (2012b) Review of *Neoperla* (Plecoptera: Perlidae) from Guangdong Province of China. Zootaxa 3597(4): 15–24.
- Li WH, Wang GQ, Qin XF (2013b) Two new species of *Neoperla* (Plecoptera, Perlidae) from China. ZooKeys 290: 21–30. doi: 10.3897/zookeys.290.4568
- Li WH, Wang HL, Lu WY (2011a) Species of the genus *Neoperla* (Plecoptera: Perlidae) from Henan, China. Zootaxa 2735: 57–63.
- Li WH, Wu LM, Zhang HR (2011b) A new species of the genus *Neoperla* (Plecoptera: Perlidae) from Henan, China. Acta Zootaxonomica Sinica 36(1): 33–35.
- Li WH, Wang RF (2011) A new species of *Neoperla* (Plecoptera: Perlidae) from China. Entomological News 122(3): 261–264. doi: 10.3157/021.122.0308
- Li WH, Zhang SQ (2014) Two new species of *Neoperla* (Plecoptera, Perlidae) from Dabie Mountains of China. ZooKeys 438: 45–55. doi: 10.3897/zookeys.438.8230
- Needham JG (1905) New genera and species of Perlidae. Proceedings of the Biological Society of Washington 17: 107–110.
- Navas L (1932) Insecta Orientalis. Memorie della Pontificia Accademia Romana dei Nuovi Lincei 16: 917–918, 923–926, 952–954.
- Qin XF, Murányi D, Wang GQ, Li WH (2013) Stoneflies of the genus *Neoperla* (Plecoptera, Perlidae) from Wuyi Mountain National Nature Reserve, Fujian of China. ZooKeys 326: 1–16. doi: 10.3897/zookeys.326.5911
- Stark BP (1987) Records and descriptions of oriental Neoperlini (Plecoptera: Perlidae). Aquatic insects 9(1): 45–50. doi: 10.1080/01650428709361270
- Stark BP, Sivec I (2008) New species and records of *Neoperla* (Plecoptera: Perlidae) from Vietnam. Illiesia 4: 19–54.
- Sivec I, Zwick P (1987) Some *Neoperla* (Plecoptera) from Taiwan. Beiträge zur Entomologie 37: 391–405.
- Wu CF (1935) New species of stoneflies from East and South China. (Order Plecoptera). Bulletin of the Peking Society of Natural History 9(3): 227–243.
- Wu CF (1938) Plecopterorum sinensium: A monograph of stoneflies of China (Order Plecoptera). Yenching University, Beijing, 225 pp.
- Wu CF (1948) Fourth supplement to the stoneflies of China (Order Plecoptera). Peking Natural History Bulletin 17(1): 75–82.
- Wu CF (1962) Results of the Zoologico-Botanical expedition to Southwest China, 1955–1957 (Plecoptera). Acta Entomologica Sinica 11: 139–153.
- Wu CF (1973) New species of Chinese stoneflies (Order Plecoptera). Acta Entomologica Sinica 16(2): 97–118.
- Wu CF, Claassen PW (1934) New species of Chinese stoneflies. Bulletin of the Peking Society of Natural History 9: 111–129.
- Wang GQ, Li WH, Yang J (2013a) New species and records of the stonefly genus *Neoperla* (Plecoptera, Perlidae) from Jinhuacha Nature Reserve, Guangxi of China. ZooKeys 351: 37–48. doi: 10.3897/zookeys.351.6259

- Wang HL, Wang GQ, Li WH (2013b) Two new species in the subfamily Perlinae (Plecoptera, Perlidae) from China. *ZooKeys* 313: 81–90. doi: 10.3897/zookeys.313.5460
- Wang GQ, Murányi D, Li WH (2014) Two new species of dark *Neoperla* (Plecoptera: Perlidae) from Oriental Realm of China. *Zootaxa* 3872(1): 22–30. doi: 10.11646/zootaxa.3872.1.2
- Yang D, Yang CK (1992) Plecoptera: Perlidae. In: Huang FS (Ed.) *Insects of Wuling Mountains area, Southwestern China*. Science Press, Beijing, 62–64.
- Yang D, Yang CK (1995) Plecoptera: Perlidae. In: Wu H (Ed.) *Insects of Baishanzu Mountain, Eastern China*. China Forestry Publishing House, Beijing, 59–60.
- Yang D, Yang CK (1998) Plecoptera: Styloperlidae, Perlidae and Leuctridae. In: Wu H (Ed.) *Insects of Longwangshan*. China Forestry Publishing House, Beijing, 40–46.