

Supplemental Table 1: Demographics and Baseline Characteristics of the Intent-to-Treat Participants

Cohort	Avacincaptad Pegol 2 mg n = 67	Sham* n = 110	Avacincaptad Pegol 4 mg n = 83	Sham† n = 84
Mean Age, Years	78.8	78.2	79.2	78.2
Female Gender, Number (%)	45 (67.2)	79 (71.8)	58 (69.9)	61 (72.6)
Active Smoker, Number (%)	25 (37.3)	36 (32.7)	26 (31.3)	29 (34.5)
Noncenterpoint involving GA‡, Number (%)	62 (92.5)	104 (94.5)	81 (97.6)	82 (97.6)
Mean GA Area, mm²	7.33	7.42	7.90	7.45
Mean SQ Root GA Area, mm	2.62	2.63	2.72	2.64
Bilateral GA, Number (%)	67 (100)	108 (98.2)	83 (100)	83 (98.8)
Hyper FAF (%)	66 (98.5)	109 (99.1)	82 (98.8)	83 (98.8)
Trickling Pattern, Number (%)	12 (18.2)	20 (18.3)	13 (15.9)	17 (20.5)
Mean BCVA (ETDRS Letters)	70.2	69.0	69.5	68.3
Mean LL-BCVA (ETDRS Letters)	36.7	34.5	36.8	33.9
LL-VD (ETDRS Letters)	33.5	34.5	32.7	34.4

BCVA = best-corrected visual acuity; GA = geographic atrophy; LL-BCVA = low-luminance best-corrected visual acuity; LL-VD = low-luminance visual acuity deficit (BCVA minus LL-BCVA); SD = standard deviation.

*Sham for 2 mg arm.

†Sham for 4 mg arm.

‡ includes disease in the fovea and outside the fovea but does not involve the foveal center point; refers to observations of baseline images at the Reading Center after participants had been enrolled