

Original Article
Obstetrics & Gynecology



Diagnostic Test Accuracy Review of Cytology for Squamous Intraepithelial Lesion and Squamous Cell Carcinoma of Uterine Cervix

Jung-Soo Pyo ,¹ Guhyun Kang ,² Hye Kyoung Yoon ,³ and Hyun Jung Kim ²

¹Department of Pathology, Eulji University Hospital, Eulji University School of Medicine, Daejeon, Korea

²Department of Pathology, Sanggye Paik Hospital, Inje University College of Medicine, Seoul, Korea

³Department of Pathology, Busan Paik Hospital, Inje University College of Medicine, Busan, Korea



Received: May 23, 2018

Accepted: Oct 25, 2018

Address for Correspondence:

Hyun Jung Kim, MD, PhD

Department of Pathology, Inje University
Sanggye Paik Hospital, 1342 Dongil-ro, Nowon-
gu, Seoul 01757, Korea.

E-mail: hjkim@paik.ac.kr

© 2019 The Korean Academy of Medical
Sciences.

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License ([https://
creativecommons.org/licenses/by-nc/4.0/](https://creativecommons.org/licenses/by-nc/4.0/))
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Jung-Soo Pyo

<https://orcid.org/0000-0003-0320-8088>

Guhyun Kang

<https://orcid.org/0000-0001-7730-1686>

Hye Kyoung Yoon

<https://orcid.org/0000-0003-0714-8537>

Hyun Jung Kim

<https://orcid.org/0000-0002-6617-4578>

Funding

This study was supported by a research
grant received from the Korean Society of
Cytopathology in 2018.

Disclosure

The authors have no potential conflicts of
interest to disclose.

<https://jkms.org>

ABSTRACT

Background: Even though cervico-vaginal smears have been used as a primary screening test for cervical carcinoma, the diagnostic accuracy has been controversial. The present study aimed to evaluate the diagnostic accuracy of cytology for squamous intraepithelial lesion (SIL) and squamous cell carcinoma (SqCC) of the uterine cervix through a diagnostic test accuracy (DTA) review.

Methods: A DTA review was performed using 38 eligible studies that showed concordance between cytology and histology. In the DTA review, sensitivity, specificity, diagnostic odds ratio (OR), and the area under the curve (AUC) on the summary receiver operating characteristic (SROC) curve were calculated.

Results: In the comparison between abnormal cytology and histology, the pooled sensitivity and specificity were 93.9% (95% confidence interval [CI], 93.7%–94.1%) and 77.6% (95% CI, 77.4–77.8%), respectively. The diagnostic OR and AUC on the SROC curve were 8.90 (95% CI, 5.57–14.23) and 0.8148, respectively. High-grade squamous intraepithelial lesion (HSIL) cytology had a higher sensitivity (97.6%; 95% CI, 94.7%–97.8%) for predicting HSIL or worse histology. In the comparison between SqCC identified on cytology and on histological analysis, the pooled sensitivity and specificity, diagnostic OR, and AUC were 92.7% (95% CI, 87.3%–96.3%), 87.5% (95% CI, 87.2%–87.8%), 865.81 (95% CI, 68.61–10,925.12), and 0.9855, respectively. Geographic locations with well-organized screening programs had higher sensitivity than areas with insufficient screening programs.

Conclusion: These results indicate that cytology had a higher sensitivity and specificity for detecting SIL and SqCC of the uterine cervix during primary screening.

Keywords: Cytology; Diagnostic Test Accuracy Review; Squamous Intraepithelial Lesion; Squamous Cell Carcinoma; Uterine Cervix

INTRODUCTION

A cervico-vaginal smear, including the conventional smear and liquid-based cytology, is a simple and inexpensive test for the prediction of squamous intraepithelial lesion (SIL)

Author Contributions

Conceptualization: Pyo JS, Kang G. Data curation: Pyo JS, Yoon HK. Formal analysis: Pyo JS, Kang G. Investigation: Pyo JS, Kim HJ. Methodology: Pyo JS, Kang G. Writing - original draft: Pyo JS. Writing - review & editing: Pyo JS, Yoon HK, Kim HJ.

or squamous cell carcinoma (SqCC) of the uterine cervix.¹ These tests have contributed to a decrease in the incidence of cervical cancer, especially in geographic areas supported by well-organized screening programs.¹ Although several studies have reported on the diagnostic accuracy of the cervico-vaginal smear, results showed a wide range of estimated sensitivity compared to the specificity.¹⁻³⁸ Because the diagnostic accuracy can be affected by variable factors, including study time, geographic area, and population,¹⁻³⁸ it should be fully elucidated based on these standardized parameters, including the diagnostic grades of cytology. We tried to establish the universally acceptable value beyond the limitations of individual studies. A diagnostic test accuracy (DTA) review should be performed to confirm the cytology test outcomes of the uterine cervix.

To evaluate the diagnostic accuracy of cytology, the concordance rates between cytology and histology of the uterine cervix were investigated. In addition, the present study aimed to evaluate the diagnostic accuracy of cytology for SIL and SqCC of the uterine cervix through DTA review. A subgroup analysis based on the number of patients and study location was also conducted.

METHODS**Published study search and selection criteria**

Relevant articles were obtained by searching the PubMed databases through January 31, 2018. There was no time limit for the start. These databases were searched using the following key words: '(Uterine Cervical Neoplasms OR Uterine Cervical Dysplasia OR Cervical Intraepithelial Neoplasia OR ((cervix OR cervical OR cervico*) AND (cancer* OR carcinoma OR adenocarcinoma OR neoplas* OR dysplas* OR dyskaryos*)) OR (CIN OR CINII* OR CIN2* OR CINIII* OR CIN3*) AND (SIL OR HSIL OR H-SIL OR LSIL OR L-SIL OR ASCUS OR ASC-US).' The titles and the abstracts of all searched articles were screened for exclusion. Review articles, including the previous meta-analysis, were also screened to obtain additional eligible studies. Search results were then reviewed and articles were included if the study investigated the uterine cervix and there was information regarding the concordance between cytology and histology. The articles were excluded when they were case reports or non-original articles or non-English language publications.

Data extraction

Data from all eligible studies¹⁻³⁸ were extracted by two independent authors. Extracted data included the following: first author's name, year of publication, study location, dates of the research, methodology of cytologic examination, and number of patients analyzed. For the meta-analysis, we extracted all data associated with the concordance between cytology and histology in various categories of comparison.

Statistical analyses

The review of DTA was performed using the Meta-Disc program (version 1.4; Unit of Clinical Biostatistics, the Ramon y Cajal Hospital, Madrid, Spain). In order to calculate the pooled sensitivity and specificity, individual data were collected from each eligible study in various categories of comparison. The summary receiver operating characteristic (SROC) curve was initially constructed by plotting 'sensitivity' and '1-specificity' of each study, and the curve fitting was performed through linear regression using the Littenberg and Moses linear model. Because the data were heterogeneous owing to differences in various methodology and populations, the accuracy data were pooled by fitting a SROC curve and measuring the

value of the area under the curve (AUC). An AUC close to 1 indicates a strong test and an AUC close to 0.5 is considered as a poor test. In addition, the diagnostic odds ratio (OR) was calculated by the Meta-Disc program. The estimated values were those that predict abnormal histology of abnormal cytology. In addition, the estimated values of cytologic low-grade squamous intraepithelial lesion (LSIL), high-grade squamous intraepithelial lesion (HSIL), and SqCC were predicted to histologic LSIL, HSIL or worse, and SqCC. To obtain the detailed information, a subgroup analysis based on number of patients, was conducted.

To obtain the results of concordance between abnormal cytology and histology, the Comprehensive Meta-Analysis software package was used (Biostat, Englewood, NJ, USA). The concordance was measured by agreement rates between HSIL identified with cytology and histology and between SqCC identified with cytology and histology. Because the eligible studies used various cytologic methods, including conventional and liquid-based preparations, in various populations, a random-effects model was more suitable than a fixed-effects model. Heterogeneity between the studies was checked using the Q and I^2 statistics and presented using P values. To assess publication bias, Begg's funnel plot and Egger's test were used. The results were considered statistically significant at $P < 0.05$.

RESULTS

Selection and characteristics

A total of 3,314 reports were searched and screened in the database. Due to insufficient information on concordance, 3,155 reports were excluded. An additional 48 reports were excluded owing to results reported on other diseases, 45 were excluded because they were non-original, and 28 articles were excluded because they were written in non-English language. Finally, 38 studies were included in the present analysis (**Fig. 1** and **Table 1**), which provided data on 302,148 patients. Information on the concordance between abnormal cytology and histology test results is shown in **Table 1**.




Fig. 1. Flow chart of study search and selection methods.

Table 1. Main characteristics of the eligible studies

Study	Location	Duration	Method	No.	No. of patients ^a			
					TP	FP	FN	TN
Agorastos et al. ²	Greece	2000–2001	CC	1,296	8	14	8	22
Agorastos et al. ³	Greece	2011–2013	LBC	3,993	62	18	63	45
Alanbay et al. ⁴	Turkey	2013–2015	CC	52	23	17	9	0
Beerman et al. ⁵	Netherland	1997–2002	CC	86,469	347	498	30	49,826
Belinson et al. ⁶	China	ND	LBC	8,497				
Benedet et al. ⁷	Canada	1986–2000	CC	84,244	44,847	15,561	628	1,163
Bigras and de Marval ⁸	Switzerland	ND	LBC	13,842	209	150	285	884
Blumenthal et al. ⁹	Zimbabwe	1995–1997	CC	2,199				
Canda et al. ¹⁰	Turkey	2005	CC	5,835	6	4	1	2
		2006–2009	LBC		13	4	2	6
Cárdenas-Turanzas et al. ¹¹	USA/Canada	ND	CC	963	30	47	104	782
Castle et al. ¹²	USA	2008–2009	LBC	7,823	482	1,704	539	5,098
Chung et al. ¹³	Korea	2004	CC	1,221	27	2	9	17
			LBC		32	2	3	17
Chute et al. ¹⁴	USA	2003	CC	530	155	133	11	231
Cuzick et al. ¹⁵	UK	1992–1994	CC	1,985	64	54	43	43
Cuzick et al. ¹⁶	UK	ND	CC	10,358	117	280	39	551
Depuydt et al. ¹⁷	Belgium	2005–2007	LBC	2,905	45	27	42	153
Ferreccio et al. ¹⁸	Chile	ND	CC	8,265				
Guo et al. ¹⁹	USA	2000–2001	LBC	788	551	63	65	103
Hovland et al. ²⁰	Congo	ND	CC	301				
			LBC					
Hutchinson et al. ²¹	Costa Rica	ND	CC	8,636	219	357	101	7,956
			LBC		284	811	39	7,502
Iftner et al. ²²	Germany	ND	LBC	9,451				
Kim et al. ¹	Korea	2005–2012	LBC	3,141	623	152	47	2,319
Li et al. ²³	China	2004–2005	LBC	2,562				
Mahmud et al. ²⁴	Congo	2003–2004	CC	1,366	16	33	24	441
McAdam et al. ²⁵	Vanuatu	2006	LBC	519	38	13	13	6
Monsonogo et al. ²⁶	France	2008–2009	LBC	4,429	268	117	344	378
Negri et al. ²⁷	Italy	2000–2002	CC	214	27	2	9	17
			LBC		36	5	1	3
Pan et al. ²⁸	China	ND	LBC	1,780	174	339	39	1,441
Parakevaidis et al. ²⁹	Greece	1997–1999	CC	977	64	179	11	34
Petry et al. ³⁰	Germany	1998–2000	CC	8,466				
Rahimi et al. ³¹	Italy	ND	CC	461	16	2	2	0
			LBC		14	3	4	1
Salmerón et al. ³²	Mexico	1999	CC	7,732	77	59	72	213
Sankaranarayanan et al. ³³	India	1999–2003	CC	24,915	718	1,285	638	20,018
Schneider et al. ³⁴	Germany	1996–1998	CC	5,455	24	2	140	193
Sigurdsson ³⁵	Iceland	2007–2011	CC	61,574	1,603	206	24	18
			LBC		1,081	111	7	57
Sykes et al. ³⁶	New Zealand	2004–2006	CC	913	250	60	16	35
			LBC		253	59	23	41
Wu et al. ³⁷	China	ND	LBC	2,098				
Zhu et al. ³⁸	Sweden	ND	CC	137	84	25	23	5
			LBC		89	23	18	7

TP = true positive, FP = false positive, FN = false negative, TN = true negative, CC = conventional cytology, LBC = liquid-based cytology, ND = no description.

^aConcordance between abnormal cytology and abnormal histology.

DTA review of cytology

A DTA review was conducted to elucidate the diagnostic accuracy using cytology in uterine cervix. In the comparison between abnormal cytology and histology, the pooled sensitivity and specificity values were 93.9% (95% confidence interval [CI], 93.7%–94.1%) and 77.6% (95% CI, 77.4%–77.8%), respectively (**Fig. 2**). The diagnostic OR and AUC on the SROC curve were 8.90 (95% CI, 5.57–14.23) and 0.8112, respectively (**Fig. 3**). A subgroup analysis based on the number of included patients of each eligible study ($\geq 1,000$ and $< 1,000$) and study

locations (areas with well-organized versus insufficient screening programs) was conducted. In the subgroup that included the larger number of patients, the pooled sensitivity and specificity, diagnostic OR and AUC on the SROC curve were 94.9% (95% CI, 94.8%–95.1%), 77.8% (95% CI, 77.5%–78.0%), 22.91 (95% CI, 10.70–49.04), and 0.8963, respectively. However, the pooled sensitivity and specificity of the subgroup with a smaller number of patients was 71.1% (95% CI, 69.3%–72.9%) and 73.6% (95% CI, 72.2%–75.0%), respectively. Next, in the subgroup analysis based on study location, areas with well-organized screening programs had a higher sensitivity than areas with insufficient screening programs (94.9% vs. 71.1%).

In the comparison between LSIL identified with cytology and LSIL identified with histology, the pooled sensitivity and specificity, diagnostic OR, and AUC were 80.5% (95% CI, 78.7%–81.2%), 80.6% (95% CI, 80.2%–81.0%), 11.80 (95% CI, 5.30–26.29), and 0.8339, respectively (Table 2). For predicting HSIL or worse histology, the sensitivity and specificity of LSIL cytology were




Fig. 2. The forest plots for the sensitivity and specificity of abnormal cytology in predicting SIL or SqCC in uterine cervix. (A) Sensitivity. (B) Specificity. SIL = squamous intraepithelial lesion, SqCC = squamous cell carcinoma, CI = confidence interval.

(continued to the next page)




Fig. 2. (Continued) The forest plots for the sensitivity and specificity of abnormal cytology in predicting SIL or SqCC in uterine cervix. (A) Sensitivity. (B) Specificity. SIL = squamous intraepithelial lesion, SqCC = squamous cell carcinoma, CI = confidence interval.

97.6% (95% CI, 97.4%–97.8%) and 71.7% (95% CI, 71.3%–72.0%), respectively. The diagnostic OR and AUC were 64.49 (95% CI, 29.04–143.20) and 0.9444, respectively. The pooled sensitivity and specificity, diagnostic OR, and AUC of cytologic SqCC were 92.7% (95% CI, 87.3%–96.3%), 87.5% (95% CI, 87.2%–87.8%), 865.81 (95% CI, 68.61–10,925.12), and 0.9855 for predicting SqCC in histology. In the subgroup analysis, those that used conventional cytology and well-organized screening programs had a higher sensitivity and lower specificity than subgroups that used liquid-based cytology and lacked screening programs.

DISCUSSION

In daily practice, screening tests use cytology and/or the human papillomavirus (HPV) test to predict SIL and SqCC of the uterine cervix. However, it is difficult to obtain information on diagnostic accuracy of cytology and the HPV test from individual studies. Previous




Fig. 3. SROC curve of abnormal cytology in predicting SIL or SqCC in uterine cervix. SROC = summary receiver operating characteristic, SIL = squamous intraepithelial lesion, SqCC = squamous cell carcinoma, AUC = area under the curve, SE = standard error, Q* = the point where sensitivity and specificity are equal.

Table 2. Sensitivity, specificity, diagnostic OR and AUC of SROC curve in cases with histologic confirmation

Comparison	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Diagnostic OR (95% CI)	AUC on SROC
LSIL in cytology vs. LSIL in histology	80.5 (78.7–81.2)	80.6 (80.2–81.0)	11.80 (5.30–26.29)	0.8339
HSIL in cytology vs. HSIL+ in histology	97.6 (97.4–97.8)	71.7 (71.3–72.0)	64.49 (29.04–143.20)	0.9444
SqCC in cytology vs. SqCC in histology	92.7 (87.3–96.3)	87.5 (87.2–87.8)	865.81 (68.61–10,925.12)	0.9855

OR = odds ratio, AUC = area under curve, SROC = summary receiver operating characteristic, CI = confidence interval, LSIL = low-grade squamous intraepithelial lesion, HSIL = high-grade squamous intraepithelial lesion, HSIL+ = HSIL or worse, SqCC = squamous cell carcinoma.

studies show that the ranges of sensitivities and specificities of cytology and HPV test varied widely.³⁹ In the eligible studies, sensitivities and specificities of cytology ranged from 22.4% to 99.4% and 0.0% to 99.0%, respectively.¹⁻³⁸ Therefore, it is useful to assess the diagnostic accuracy of a screening test to predict the presence of SIL and SqCC in the uterine cervix by performing a meta-analysis, including a DTA review. To the best of our knowledge, the present study is the first to assess the diagnostic accuracy of cytology for predicting SIL and SqCC in the uterine cervix.

In the present DTA review, regardless of the diagnostic grade of cytology, its diagnostic accuracy was initially evaluated for the prediction of abnormal histology. The sensitivity and specificity of cytology were 93.9% and 77.6%, respectively. In a subgroup analysis based on the number of patients, the larger subgroup showed a higher sensitivity than the smaller subgroup (94.9% vs. 71.1%). Eligible studies with a small number of patients might affect the sensitivity and specificity, since patient cohort sizes ranged from 13 to 50,701. In addition, experiences of cytopathologists and cytotechnologists may be important for the diagnostic accuracy of cytologic examination. Recent automated cytoscreening systems can also be helpful for effective screening. Results of this DTA review show that cytology is a useful screening test in the prediction of SIL or SqCC histology.

In the DTA review for the diagnostic accuracy of cytology, index should be cytology and comparator test should be histology. However, in previous studies, colposcopy was included in the comparator test.³⁹ Cases with negative colposcopic findings were considered as true negative in these studies.³⁹ However, because colposcopy is not a confirmative examination, specificity might be overestimated due to the increase in true negative cases. Therefore,

cytology and histology should be compared to properly evaluate the diagnostic accuracy. The present study included only patients with histologic confirmation, but not those who underwent colposcopic examination.

In a previous DTA review, the sensitivity of cytology and HPV test were 65.87%–75.51% and 92.60%–95.13%, respectively.³⁹ However, in this study, cytology was compared between atypical squamous cells of undetermined significance (ASC-US) or worse cytology and HSIL histology. The true positive rate and sensitivity were decreased because patients who underwent LSIL histology were considered false positives in abnormal cytology. The sensitivity of cytology was higher in our study compared to the previous DTA review. Therefore, overestimation of specificity could be possibly considered. In addition, the previous DTA review only included studies that assessed both cytology and HPV tests. The estimated value for diagnostic OR and AUC on SROC, which are useful in comparing various tests, were not shown. In summary, the superiority of the HPV test for accurately diagnosing SIL or SqCC in the uterine cervix cannot be proven in the previous DTA review. In addition, in other DTA review,⁴⁰ the pooled sensitivity of cytology with HSIL or worse was 79.4% for predicting cancer. However, this review did not show results for other parameters, such as specificity, diagnostic OR, AUC on SROC. The estimated values of overall abnormal cytology and LSIL were not found in the previous review.⁴⁰

In practice, ASC-US cytology usually requires a repeat smear and/or an HPV test. An ancillary test, such as the HPV test, may be useful because the confirmative information in the repeat smear cannot be obtained. However, the gradient correlation between HPV test and histology is unclear. The advantage of cytology is its ability to predict histologic abnormalities which can help with patient management, compared to that of an HPV test. After a cytologic preparation, HPV tests using the remaining cytologic specimen can be performed. The presence of ASC-US cytology groups, which can increase the false-positive rate and decrease sensitivity. In the previous study, the rate of ASC-US cytology was less than 5.0%.¹² However, in the Republic of Korea which has a well-organized screening system, the rate of ASC-US were 0.045% in 432,691 women who had screening tests.¹ Therefore, an ancillary HPV test can be more useful in patients with ASC-US cytology. In areas with insufficient screening systems, the effectiveness of a cytologic examination is not fully elucidated. In addition, in areas with a well-organized screening system, the usefulness of an HPV test as the primary screening test is unclear. Primary screening tests should not be selected by simply considering the sensitivity. Availability of screening systems may be important for choosing the screening method to help diagnose SIL or SqCC of the uterine cervix.

In a subanalysis of the ATHENA study, co-testing using cytology and the HPV test has no advantage compared with the HPV test alone.¹² However, this study did not enroll patients without an HPV test. This criterion could decrease the sensitivity and true positive cases of cytology. In addition, this report compared ASC-US and worse cytology with HSIL or worse confirmed with histology. Therefore, because sensitivity can differ by patient populations, the diagnostic accuracy of the screening test in the general population can differ between individual studies. The results showed that sensitivity of cytology in our results (96.9%) was higher than that of the HPV test sensitivity for HSIL or worse with histology as shown in Castle's report (88.2%). In addition, in our study, the estimated concordance rates were 93.1% (95% CI, 84.7%–97.1%) and 98.8% (95% CI, 69.0%–100.0%) for HSIL and SqCC cytology, respectively.

There are some limitations in the current DTA review. First, the comparisons between various cytologic abnormalities and histologic abnormalities were conducted in the present DTA review. ASC-US/atypical squamous cells, cannot exclude HSIL (ASC-H) cytology belongs to the heterogeneous diagnostic category. However, the diagnostic accuracy of ASC-H could not be performed due to insufficient information included in the eligible studies. Second, the aim of the present DTA review was to elucidate the diagnostic accuracy of cytology. Thus, the effectiveness between cytology and HPV test was compared with the results of previous reports.^{12,39} Third, the number of patients in the individual studies did not apply to exclusion criteria in the present DTA review. The eligible studies with a smaller number of patients showed far from average estimation. However, the effects of studies with a smaller number of patients on overall estimated values were insignificant. Therefore, the diagnostic accuracy of cytology using individual studies with a smaller number of patients should be accurately interpreted. Fourth, histologic examinations include a punch biopsy, loop electrocautery excision procedure, conization, or hysterectomy in the uterine cervix. Sampling error can occur with histologic examinations, such as a punch biopsy. However, in the present DTA review, a detailed evaluation based on histologic methodology could not be conducted due to insufficient information on eligible studies.

In conclusion, our results show that cytology has higher sensitivity and specificity for the prediction of SIL or SqCC, regardless of the diagnostic grade of cytology. The diagnostic accuracy of cytology as a primary screening test was re-confirmed in the present DTA review. Therefore, cytology is one of the most sensitive and confirmative primary screening tests for SIL and SqCC.

REFERENCES

1. Kim SH, Lee JM, Yun HG, Park US, Hwang SU, Pyo JS, et al. Overall accuracy of cervical cytology and clinicopathological significance of LSIL cells in ASC-H cytology. *Cytopathology* 2017;28(1):16-23.
[PUBMED](#) | [CROSSREF](#)
2. Agorastos T, Dinas K, Lloveras B, de Sanjose S, Kornegay JR, Bonti H, et al. Human papillomavirus testing for primary screening in women at low risk of developing cervical cancer. The Greek experience. *Gynecol Oncol* 2005;96(3):714-20.
[PUBMED](#) | [CROSSREF](#)
3. Agorastos T, Chatzistamatiou K, Katsamagkas T, Koliopoulos G, Daponte A, Constantinidis T, et al. Primary screening for cervical cancer based on high-risk human papillomavirus (HPV) detection and HPV 16 and HPV 18 genotyping, in comparison to cytology. *PLoS One* 2015;10(3):e0119755.
[PUBMED](#) | [CROSSREF](#)
4. Alanbay İ, Öztürk M, Fıratlıgil FB, Karaşahin KE, Yenen MC, Bodur S. Cytohistological discrepancies of cervico-vaginal smears and HPV status. *Ginekol Pol* 2017;88(5):235-8.
[PUBMED](#) | [CROSSREF](#)
5. Beerman H, van Dorst EB, Kuenen-Boumeester V, Hogendoorn PC. Superior performance of liquid-based versus conventional cytology in a population-based cervical cancer screening program. *Gynecol Oncol* 2009;112(3):572-6.
[PUBMED](#) | [CROSSREF](#)
6. Belinson JL, Qiao YL, Pretorius RG, Zhang WH, Rong SD, Huang MN, et al. Shanxi Province cervical cancer screening study II: self-sampling for high-risk human papillomavirus compared to direct sampling for human papillomavirus and liquid based cervical cytology. *Int J Gynecol Cancer* 2003;13(6):819-26.
[PUBMED](#) | [CROSSREF](#)
7. Benedet JL, Maticic JP, Bertrand MA. An analysis of 84244 patients from the British Columbia cytology-colposcopy program. *Gynecol Oncol* 2004;92(1):127-34.
[PUBMED](#) | [CROSSREF](#)

8. Bigras G, de Marval F. The probability for a Pap test to be abnormal is directly proportional to HPV viral load: results from a Swiss study comparing HPV testing and liquid-based cytology to detect cervical cancer precursors in 13,842 women. *Br J Cancer* 2005;93(5):575-81.
[PUBMED](#) | [CROSSREF](#)
9. Blumenthal PD, Gaffikin L, Chirenje ZM, McGrath J, Womack S, Shah K. Adjunctive testing for cervical cancer in low resource settings with visual inspection, HPV, and the Pap smear. *Int J Gynaecol Obstet* 2001;72(1):47-53.
[PUBMED](#) | [CROSSREF](#)
10. Canda MT, Demir N, Sezer O, Doganay L, Ortac R. Clinical results of the liquid-based cervical cytology tool, Liqui-PREP, in comparison with conventional smears for detection of squamous cell abnormalities. *Asian Pac J Cancer Prev* 2009;10(3):399-402.
[PUBMED](#)
11. Cárdenas-Turanzas M, Nogueras-Gonzalez GM, Scheurer ME, Adler-Storthz K, Benedet JL, Beck JR, et al. The performance of human papillomavirus high-risk DNA testing in the screening and diagnostic settings. *Cancer Epidemiol Biomarkers Prev* 2008;17(10):2865-71.
[PUBMED](#) | [CROSSREF](#)
12. Castle PE, Stoler MH, Wright TC Jr, Sharma A, Wright TL, Behrens CM. Performance of carcinogenic human papillomavirus (HPV) testing and HPV16 or HPV18 genotyping for cervical cancer screening of women aged 25 years and older: a subanalysis of the ATHENA study. *Lancet Oncol* 2011;12(9):880-90.
[PUBMED](#) | [CROSSREF](#)
13. Chung JH, Park EJ, Choi YD, Kim HS, Lee YJ, Ko HM, et al. Efficacy assessment of CellSlide in liquid-based gynecologic cytology. *Gynecol Oncol* 2005;99(3):597-602.
[PUBMED](#) | [CROSSREF](#)
14. Chute DJ, Covell J, Pambuccian SE, Stelow EB. Cytologic-histologic correlation of screening and diagnostic Papanicolaou tests. *Diagn Cytopathol* 2006;34(7):503-6.
[PUBMED](#) | [CROSSREF](#)
15. Cuzick J, Szarewski A, Terry G, Ho L, Hanby A, Maddox P, et al. Human papillomavirus testing in primary cervical screening. *Lancet* 1995;345(8964):1533-6.
[PUBMED](#) | [CROSSREF](#)
16. Cuzick J, Szarewski A, Cubie H, Hulman G, Kitchener H, Luesley D, et al. Management of women who test positive for high-risk types of human papillomavirus: the HART study. *Lancet* 2003;362(9399):1871-6.
[PUBMED](#) | [CROSSREF](#)
17. Depuydt CE, Makar AP, Ruymbeke MJ, Benoy IH, Vereecken AJ, Bogers JJ. BD-ProExC as adjunct molecular marker for improved detection of CIN2+ after HPV primary screening. *Cancer Epidemiol Biomarkers Prev* 2011;20(4):628-37.
[PUBMED](#) | [CROSSREF](#)
18. Ferreccio C, Barriga MI, Lagos M, Ibáñez C, Poggi H, González F, et al. Screening trial of human papillomavirus for early detection of cervical cancer in Santiago, Chile. *Int J Cancer* 2013;132(4):916-23.
[PUBMED](#) | [CROSSREF](#)
19. Guo M, Hu L, Baliga M, He Z, Hughson MD. The predictive value of p16(INK4a) and hybrid capture 2 human papillomavirus testing for high-grade cervical intraepithelial neoplasia. *Am J Clin Pathol* 2004;122(6):894-901.
[PUBMED](#) | [CROSSREF](#)
20. Hovland S, Arbyn M, Lie AK, Ryd W, Borge B, Berle EJ, et al. A comprehensive evaluation of the accuracy of cervical pre-cancer detection methods in a high-risk area in East Congo. *Br J Cancer* 2010;102(6):957-65.
[PUBMED](#) | [CROSSREF](#)
21. Hutchinson ML, Zahniser DJ, Sherman ME, Herrero R, Alfaro M, Bratti MC, et al. Utility of liquid-based cytology for cervical carcinoma screening: results of a population-based study conducted in a region of Costa Rica with a high incidence of cervical carcinoma. *Cancer* 1999;87(2):48-55.
[PUBMED](#) | [CROSSREF](#)
22. Iftner T, Becker S, Neis KJ, Castanon A, Iftner A, Holz B, et al. Head-to-head comparison of the RNA-based aptima human papillomavirus (HPV) assay and the DNA-based hybrid capture 2 HPV test in a routine screening population of women Aged 30 to 60 years in Germany. *J Clin Microbiol* 2015;53(8):2509-16.
[PUBMED](#) | [CROSSREF](#)
23. Li N, Shi JF, Franceschi S, Zhang WH, Dai M, Liu B, et al. Different cervical cancer screening approaches in a Chinese multicentre study. *Br J Cancer* 2009;100(3):532-7.
[PUBMED](#) | [CROSSREF](#)
24. Mahmud SM, Sangwa-Lugoma G, Nasr SH, Kayembe PK, Tozin RR, Drouin P, et al. Comparison of human papillomavirus testing and cytology for cervical cancer screening in a primary health care setting in the Democratic Republic of the Congo. *Gynecol Oncol* 2012;124(2):286-91.
[PUBMED](#) | [CROSSREF](#)

25. McAdam M, Sakita J, Tarivonda L, Pang J, Frazer IH. Evaluation of a cervical cancer screening program based on HPV testing and LLETZ excision in a low resource setting. *PLoS One* 2010;5(10):e13266.
[PUBMED](#) | [CROSSREF](#)
26. Monsonogo J, Hudgens MG, Zerat L, Zerat JC, Syrjänen K, Halfon P, et al. Evaluation of oncogenic human papillomavirus RNA and DNA tests with liquid-based cytology in primary cervical cancer screening: the FASE study. *Int J Cancer* 2011;129(3):691-701.
[PUBMED](#) | [CROSSREF](#)
27. Negri G, Menia E, Egarter-Vigl E, Vittadello F, Mian C. ThinPrep versus conventional Papanicolaou smear in the cytologic follow-up of women with equivocal cervical smears. *Cancer* 2003;99(6):342-5.
[PUBMED](#) | [CROSSREF](#)
28. Pan Q, Belinson JL, Li L, Pretorius RG, Qiao YL, Zhang WH, et al. A thin-layer, liquid-based pap test for mass screening in an area of China with a high incidence of cervical carcinoma. A cross-sectional, comparative study. *Acta Cytol* 2003;47(1):45-50.
[PUBMED](#) | [CROSSREF](#)
29. Paraskevaides E, Malamou-Mitsi V, Koliopoulos G, Pappa L, Lolis E, Georgiou I, et al. Expanded cytological referral criteria for colposcopy in cervical screening: comparison with human papillomavirus testing. *Gynecol Oncol* 2001;82(2):355-9.
[PUBMED](#) | [CROSSREF](#)
30. Petry KU, Menton S, Menton M, van Loenen-Frosch F, de Carvalho Gomes H, Holz B, et al. Inclusion of HPV testing in routine cervical cancer screening for women above 29 years in Germany: results for 8466 patients. *Br J Cancer* 2003;88(10):1570-7.
[PUBMED](#) | [CROSSREF](#)
31. Rahimi S, Carnovale-Scalzo C, Marani C, Renzi C, Malvasi A, Votano S. Comparison of conventional Papanicolaou smears and fluid-based, thin-layer cytology with colposcopic biopsy control in central Italy: a consecutive sampling study of 461 cases. *Diagn Cytopathol* 2009;37(1):1-3.
[PUBMED](#) | [CROSSREF](#)
32. Salmerón J, Lazcano-Ponce E, Lorincz A, Hernández M, Hernández P, Leyva A, et al. Comparison of HPV-based assays with Papanicolaou smears for cervical cancer screening in Morelos State, Mexico. *Cancer Causes Control* 2003;14(6):505-12.
[PUBMED](#) | [CROSSREF](#)
33. Sankaranarayanan R, Chatterji R, Shastri SS, Wesley RS, Basu P, Mahe C, et al. Accuracy of human papillomavirus testing in primary screening of cervical neoplasia: results from a multicenter study in India. *Int J Cancer* 2004;112(2):341-7.
[PUBMED](#) | [CROSSREF](#)
34. Schneider A, Hoyer H, Lotz B, Leistritz S, Kühne-Heid R, Nindl I, et al. Screening for high-grade cervical intra-epithelial neoplasia and cancer by testing for high-risk HPV, routine cytology or colposcopy. *Int J Cancer* 2000;89(6):529-34.
[PUBMED](#) | [CROSSREF](#)
35. Sigurdsson K. Is a liquid-based cytology more sensitive than a conventional Pap smear? *Cytopathology* 2013;24(4):254-63.
[PUBMED](#) | [CROSSREF](#)
36. Sykes PH, Harker DY, Miller A, Whitehead M, Neal H, Wells JE, et al. A randomised comparison of SurePath liquid-based cytology and conventional smear cytology in a colposcopy clinic setting. *BJOG* 2008;115(11):1375-81.
[PUBMED](#) | [CROSSREF](#)
37. Wu R, Belinson SE, Du H, Na W, Qu X, Wu R, et al. Human papillomavirus messenger RNA assay for cervical cancer screening: the Shenzhen Cervical Cancer Screening Trial I. *Int J Gynecol Cancer* 2010;20(8):1411-4.
[PUBMED](#)
38. Zhu J, Norman I, Elfgrén K, Gaberi V, Hagmar B, Hjerpe A, et al. A comparison of liquid-based cytology and Pap smear as a screening method for cervical cancer. *Oncol Rep* 2007;18(1):157-60.
[PUBMED](#)
39. Koliopoulos G, Nyaga VN, Santesso N, Bryant A, Martin-Hirsch PP, Mustafa RA, et al. Cytology versus HPV testing for cervical cancer screening in the general population. *Cochrane Database Syst Rev* 2017;8:CD008587.
[PUBMED](#)
40. Castanon A, Landy R, Michalopoulos D, Bhudia R, Leaver H, Qiao YL, et al. Systematic review and meta-analysis of individual patient data to assess the sensitivity of cervical cytology for diagnosis of cervical cancer in low- and middle-income countries. *J Glob Oncol* 2017;3(5):524-38.
[PUBMED](#) | [CROSSREF](#)