## Heliyon



Received: 17 July 2017 Accepted: 17 July 2017

Cite as: Toshinari Takamura, Yuki Kita, Masatoshi Nakagen, Masaru Sakurai, Yuki Isobe, Yumie Takeshita, Kohzo Kawai, Takeshi Urabe, Shuichi Kaneko. Corrigendum to 'Weight-adjusted lean body mass and calf circumference are protective against obesity-associated insulin resistance and metabolic abnormalities'. Heliyon 3 (2017) e00363. doi: 10.1016/j.heliyon.2017. e00363



## Corrigendum

## Corrigendum to 'Weight-adjusted lean body mass and calf circumference are protective against obesity-associated insulin resistance and metabolic abnormalities'

Toshinari Takamura <sup>a,\*</sup>, Yuki Kita <sup>a</sup>, Masatoshi Nakagen <sup>d</sup>, Masaru Sakurai <sup>c</sup>, Yuki Isobe <sup>a</sup>, Yumie Takeshita <sup>a</sup>, Kohzo Kawai <sup>d</sup>, Takeshi Urabe <sup>d</sup>, Shuichi Kaneko <sup>b</sup>

DOI of original article: http://dx.doi.org/10.1016/j.heliyon.2017.e00347

E-mail address: ttakamura@m-kanazawa.jp (S. Kaneko).

In the original published version of this article, a typographical error was present in Table 5 displaying '2.451' in 'Column: Female, Coefficient ( $\beta$ ); Row: Hip Circumference' instead of the correct value of '0.322'. The authors apologize for the mistake. The corrected version of Table 5 is displayed below.

**Table 5.** Multiple linear regression analysis showing variables independently associated with lean body mass.

	All			Male			Female		
	Coefficient (β)	t-statistic	P	Coefficient (β)	t-statistic	P	Coefficient (β)	t-statistic	P
Arm circumference (cm)	0.222	2.564	0.011	0.200	2.446	0.017	-0.230	-2.214	0.029
Waist circumference (cm)	0.183	1.605	0.110	-0.227	-1.656	0.102	0.100	0.876	0.383
Hip circumference (cm)	-0.154	-1.298	0.196	0.627	4.405	0.000	0.322	2.451	0.016
Thigh circumference (cm)	-0.149	-1.352	0.178	0.056	0.465	0.643	-0.029	-0.248	0.805
Calf circumference (cm)	0.718	6.736	0.000	0.306	2.874	0.005	0.714	5.778	0.000

Multiple linear regression was used for the analysis, adjusted with each other.

<sup>&</sup>lt;sup>a</sup>Department of Endocrinology and Metabolism, Kanazawa University Graduate School of Medical Sciences, Ishikawa, Japan

<sup>&</sup>lt;sup>b</sup> Department of System Biology, Kanazawa University Graduate School of Medical Sciences, Ishikawa, Japan

<sup>&</sup>lt;sup>c</sup> Social and Environmental Medicine, Kanazawa Medical University, Ishikawa, Japan

<sup>&</sup>lt;sup>d</sup> Department of Internal Medicine, Public Central Hospital of Matto Ishikawa, Hakusan, Ishikawa, Japan

<sup>\*</sup> Corresponding author.