

Genome Sequences of Five *Oenococcus oeni* Strains Isolated from Nero Di Troia Wine from the Same Terroir in Apulia, Southern Italy

Vittorio Capozzi,^a Pasquale Russo,^a Antonella Lamontanara,^b Luigi Orrù,^b Luigi Cattivelli,^b Giuseppe Spano^a

Department of Agriculture, Food and Environment Sciences, University of Foggia, Foggia, Italy^a; Consiglio, Per la Ricerca e Sperimentazione in Agricoltura (CRA), Genomics Research Centre, Fiorenzuola d'Arda, Italy^b

***Oenococcus oeni* is the principal lactic acid bacterium responsible for malolactic fermentation in wine. Here, we announce the genome sequences of five *O. oeni* strains isolated from Nero di Troia wine undergoing spontaneous malolactic fermentation, and we report, for the first time, several genome sequences of strains isolated from the same terroir.**

Received 11 September 2014 Accepted 16 September 2014 Published 23 October 2014

Citation Capozzi V, Russo P, Lamontanara A, Orrù L, Cattivelli L, Spano G. 2014. Genome sequences of five *Oenococcus oeni* strains isolated from Nero di Troia wine from the same terroir in Apulia, southern Italy. *Genome Announc.* 2(5):e01077-14. doi:10.1128/genomeA.01077-14.

Copyright © 2014 Capozzi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Giuseppe Spano, giuseppe.spano@unifg.it.

Oenococcus oeni is the main species of lactic acid bacteria (LAB) responsible of driving malolactic fermentation (MLF) in wine. The key biochemical stage of MLF is the microbial decarboxylation of L-malic acid that leads to the production of L-lactic acid and CO₂. MLF leads to a decrease in wine acidity and an improved microbial stability and sensorial quality (1–3). Increasing attention to the selection of autochthonous microbes from spontaneous fermentation is warranted to aid the design of specific starter cultures used in fermented foods and beverages with a geographical indication status (4–6). This is particularly true for the grape/wine environment, in which a relationship exists among cultivars, vintages, climates, and wine grape microbial biogeography, suggesting a possible dimension of the so-called microbial terroir (7, 8).

While the only fully complete genome sequence is available for the *O. oeni* PSU-1 strain (9), an increasing number of *O. oeni* assembled genome sequences also have been deposited in the GenBank database (10–12). Moreover, for *O. oeni* strain ATCC BAA-1163, a proteome reference map is also available, which is useful for the validation of annotated genes (13).

Here, we present the genome sequences of five *O. oeni* strains (OM22, OT25, OT4, OT5, and OT3) isolated from Nero di Troia wine (a typical Apulian red wine obtained from uva di Troia, an autochthonous Apulian black grape variety) undergoing spontaneous malolactic fermentation (Table 1) (14).

These new assembled complete genomes represent an important opportunity for assessing the molecular basis of (i) some

safety aspects (15–17), (ii) tolerance to the harsh wine conditions (18, 19), and (iii) the contribution to wine sensorial quality (2, 3, 20, 21). To the best of our knowledge, it is the first time that five strains isolated from the same terroir are sequenced.

Two micrograms of genomic DNA was subjected to library preparation using the TruSeq DNA sample prep kit FC-121-1001, according to the manufacturer's instructions. Whole-genome sequencing was performed using the Illumina GAIIx platform. Prior to assembly, raw reads were filtered using the PrinSeq version 0.20.3 software (22) to remove low-quality 3' ends ($Q < 25$), reads containing a percentage of uncalled bases (Ns) of $\geq 10\%$, and duplicated sequences. The genome sequences of *O. oeni* OM22 were *de novo* assembled using the Ray version 2.2.0 assembly program (23), with a *k*-mer size of 71. The genome sequences of *O. oeni* OT25, *O. oeni* OT4, *O. oeni* OT5, and *O. oeni* OT3 were *de novo* assembled using CLC Genomics Workbench 7.0, with a *k*-mer size of 64. The sequence was annotated by the National Center for Biotechnology Information (NCBI) Prokaryotic Genomes Annotation Pipeline. The genome information for each strain is summarized in Table 1.

Nucleotide sequence accession numbers. The draft genome sequences of the *O. oeni* strains sequenced in this study have been deposited as whole-genome shotgun projects at DDBJ/EMBL/GenBank under the accession numbers [JPEK000000000](https://www.ncbi.nlm.nih.gov/nuccore/JPEK000000000) (*O. oeni* OM22), [JPEM000000000](https://www.ncbi.nlm.nih.gov/nuccore/JPEM000000000) (*O. oeni* OT25), [JPEL000000000](https://www.ncbi.nlm.nih.gov/nuccore/JPEL000000000) (*O. oeni* OT4), [JPEJ000000000](https://www.ncbi.nlm.nih.gov/nuccore/JPEJ000000000) (*O. oeni* OT5), and [JOOH000000000](https://www.ncbi.nlm.nih.gov/nuccore/JOOH000000000) (*O. oeni* OT3).

TABLE 1 Summary of information for the whole genomes of the five *Oenococcus oeni* strains OM22, OT25, OT4, OT5, and OT3

<i>O. oeni</i> strain	G+C content (%)	Genome size (bp)	No. of genes	No. of proteins	Accession no.	No. of contigs
OM22	38	1,862,817	1,895	1,772	JPEK000000000	23
OT25	37.9	1,834,661	1,845	1,731	JPEM000000000	61
OT4	37.9	1,779,962	1,776	1,654	JPEL000000000	55
OT5	37.8	1,767,097	1,775	1,655	JPEJ000000000	60
OT3	37.8	1,769,724	1,780	1,658	JOOH000000000	61

ACKNOWLEDGMENTS

This research was partially supported by a grant from the project PON02_00186_3417512, “Si.Mi.SA,” and by the Apulian Region in the framework of the “OenoMicroManagement” project (PIF—Progetti Integrati di Filiera no. 94750304571).

REFERENCES

- Lonvaud-Funel A. 1999. Lactic acid bacteria in the quality improvement and depreciation of wine. *Antonie Van Leeuwenhoek* 76:317–331. <http://dx.doi.org/10.1023/A:1002088931106>.
- Bartowsky EJ, Borneman AR. 2011. Genomic variations of *Oenococcus oeni* strains and the potential to impact on malolactic fermentation and aroma compounds in wine. *Appl. Microbiol. Biotechnol.* 92:441–447. <http://dx.doi.org/10.1007/s00253-011-3546-2>.
- Bartowsky E. 2005. *Oenococcus oeni* and malolactic fermentation—moving into the molecular arena. *Aust. J. Grape Wine Res.* 11:174–187. <http://dx.doi.org/10.1111/j.1755-0238.2005.tb00286.x>.
- Capozzi V, Spano G. 2011. Food microbial biodiversity and “microbes of protected origin.” *Front. Microbiol.* 2:237. <http://dx.doi.org/10.3389/fmicb.2011.00237>.
- Capozzi V, Spano G, Fiocco D. 2012. Transdisciplinarity and microbiology education. *J. Microbiol. Biol. Educ.* 13:70–73. <http://dx.doi.org/10.1128/jmbe.v13i1.365>.
- Capozzi V, Russo P, Spano G. 2012. Microbial information regimen in EU Geographical Indications. *World Pat. Inf.* 34:229–231. <http://dx.doi.org/10.1016/j.wpi.2012.04.001>.
- Bokulich NA, Thorngate JH, Richardson PM, Mills DA. 2014. Microbial biogeography of wine grapes is conditioned by cultivar, vintage, and climate. *Proc. Natl. Acad. Sci. U. S. A.* 111:E139–E148. <http://dx.doi.org/10.1073/pnas.1317377110>.
- Gilbert JA, van der Lelie D, Zarraonaindia I. 2014. Microbial terroir for wine grapes. *Proc. Natl. Acad. Sci. U. S. A.* 111:5–6. <http://dx.doi.org/10.1073/pnas.1320471110>.
- Mills DA, Rawsthorne H, Parker C, Tamir D, Makarova K. 2005. Genomic analysis of *Oenococcus oeni* PSU-1 and its relevance to winemaking. *FEMS Microbiol. Rev.* 29:465–475. <http://dx.doi.org/10.1016/j.fm-rre.2005.04.011>.
- Borneman AR, McCarthy JM, Chambers PJ, Bartowsky EJ. 2012. Comparative analysis of the *Oenococcus oeni* pan genome reveals genetic diversity in industrially-relevant pathways. *BMC Genomics* 13:373. <http://dx.doi.org/10.1186/1471-2164-13-373>.
- Dimopoulou M, Vuillemin M, Campbell-Sills H, Lucas PM, Ballestra P, Miot-Sertier C, Favier M, Coulon J, Moine V, Doco T, Roques M, Williams P, Petrel M, Gontier E, Moulis C, Remaud-Simeon M, Dols-Lafargue M. 2014. Exopolysaccharide (EPS) synthesis by *Oenococcus oeni*: from genes to phenotypes. *PLoS One* 9:e98898. <http://dx.doi.org/10.1371/journal.pone.0098898>.
- Lamontanara A, Orrù L, Cattivelli L, Russo P, Spano G, Capozzi V. 2014. Genome sequence of *Oenococcus oeni* OM27, the first fully assembled genome of a strain isolated from an Italian wine. *Genome Announc.* 2(4):e00658-14. <http://dx.doi.org/10.1128/genomeA.00658-14>.
- Mohedano MDLL, Russo P, de los Rios V, Capozzi V, Spano G, López P. 2014. A proteome reference map of the wine lactic acid bacteria *Oenococcus oeni* ATCC BAA-1163. *Open Biol.* 4 1–10. <http://dx.doi.org/10.1098/rsob.130154>.
- Capozzi V, Russo P, Beneduce L, Weidmann S, Grieco F, Guzzo J, Spano G. 2010. Technological properties of *Oenococcus oeni* strains isolated from typical southern Italian wines. *Lett. Appl. Microbiol.* 50:327–334. <http://dx.doi.org/10.1111/j.1472-765X.2010.02795.x>.
- Spano G, Russo P, Lonvaud-Funel A, Lucas P, Alexandre H, Grandvalet C, Coton E, Coton M, Barnavon L, Bach B, Rattray F, Bunte A, Magni C, Ladero V, Alvarez M, Fernández M, Lopez P, de Palencia PF, Corbi A, Trip H, Lolkema JS. 2010. Biogenic amines in fermented foods. *Eur. J. Clin. Nutr.* 64(Suppl 3):95–100. <http://dx.doi.org/10.1038/ejcn.2010.218>.
- Capozzi V, Ladero V, Beneduce L, Fernández M, Alvarez MA, Benoit B, Laurent B, Grieco F, Spano G. 2011. Isolation and characterization of tyramine-producing *Enterococcus faecium* strains from red wine. *Food Microbiol.* 28:434–439. <http://dx.doi.org/10.1016/j.fm.2010.10.005>.
- García-Moruno E, Muñoz R. 2012. Does *Oenococcus oeni* produce histamine? *Int. J. Food Microbiol.* 157:121–129. <http://dx.doi.org/10.1016/j.ijfoodmicro.2012.05.013>.
- Spano G, Massa S. 2006. Environmental stress response in wine lactic acid bacteria: beyond *Bacillus subtilis*. *Crit. Rev. Microbiol.* 32:77–86. <http://dx.doi.org/10.1080/10408410600709800>.
- Capozzi V, Russo P, Ladero V, Fernández M, Fiocco D, Alvarez MA, Grieco F, Spano G. 2012. Biogenic amines degradation by *Lactobacillus plantarum*: toward a potential application in wine. *Front Microbiol.* 3:122. <http://dx.doi.org/10.3389/fmicb.2012.00122>.
- Malherbe S, Menichelli E, du Toit M, Tredoux A, Muller N, Naes T, Nieuwoudt H. 2013. The relationships between consumer liking, sensory and chemical attributes of *Vitis vinifera* L. cv. Pinotage wines elaborated with different *Oenococcus oeni* starter cultures. *J. Sci. Food Agric.* 93:2829–2840. <http://dx.doi.org/10.1002/jsfa.6115>.
- Lerm E, Engelbrecht L, du Toit M. 2010. Malolactic fermentation: the ABC’s of MLF. *S. Afr. J. Enol. Vitic.* 31:186–212.
- Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864. <http://dx.doi.org/10.1093/bioinformatics/btr026>.
- Boisvert S, Laviolette F, Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of high-throughput sequencing technologies. *J. Comput. Biol.* 17:1519–1533. <http://dx.doi.org/10.1089/cmb.2009.0238>.