





Elsevier has created a [Monkeypox Information Center](#) in response to the declared public health emergency of international concern, with free information in English on the monkeypox virus. The Monkeypox Information Center is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its monkeypox related research that is available on the Monkeypox Information Center - including this research content - immediately available in publicly funded repositories, with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the Monkeypox Information Center remains active.



LETTER TO THE EDITOR

[Translated article] Anti-*Vaccinia* Immunoglobulin and Post-exposure Prophylaxis with *Vaccinia*-based Vaccine for Management of the Monkeypox Outbreak[☆]

Inmunoglobulina anti-*Vaccinia* y profilaxis postexposición mediante vacuna basada en *Vaccinia* para el control del brote de viruela símica (*Monkeypox*)

To the Editor,

Some time ago, we published a brief review of treatments available or in development that could be of use in the control of the then incipient outbreak of monkeypox.¹ In that review, we highlighted the role of tecovirimat as the only antiviral drug approved by the European Medicines Agency (EMA) for the virus, given its good safety profile and proven efficacy in reducing mortality in animal models.²

In response to that publication, Dr Sookaromdee and Dr Wiwanitkit published a letter entitled *Treatments for Monkeypox*,³ in which they shared their opinion on the use of tecovirimat and mentioned intravenous immunoglobulin as another potentially effective treatment for monkeypox, now declared a *Public Health Emergency of International Concern* by the World Health Organization (WHO).⁴

One of the affirmations by Dr Sookaromdee and Dr Wiwanitkit is that, although tecovirimat has demonstrated efficacy, it is not a widely used drug in countries in which the *Monkeypox virus* has traditionally been considered endemic.³ A possible reason for the limited use of tecovirimat in those countries is that their healthcare resources and structure are not comparable to those in European countries.⁵ In addition,

we should add that approval by the EMA² is not applicable in Africa, and to date, it is the only health agency to have approved the drug for the indication of monkeypox.

The second aspect that we wanted to comment on is the possibility of using intravenously-administered anti-*Vaccinia* immunoglobulin, proposed by Sookaromdee and Wiwanitkit.³ This immunoglobulin is indicated only for the treatment of certain complications such as *eczema vaccinatum*, progressive *Vaccinia*, and severe generalized *Vaccinia* caused by administration of the vaccine derived from the *Vaccinia virus*.⁶ The regimen administered is 6000 U/kg as soon as possible after onset of the first symptoms of the disease, with possible dose repetitions according to the severity of the condition and response to the initial dose (doses can be increased to 9000 U/kg if the patient has not responded to the first dose).⁷ Although there are factors to support its use against monkeypox, such as the similarity of *Orthopoxvirus* genomes,⁸ to date, there have been no human trials on its use in this indication.

Finally, it is necessary to highlight the use of the *Vaccinia*-based vaccine as post-exposure prophylaxis. The WHO currently recommends administration of a second- or third-generation vaccine for case contacts in the first 4 days after exposure.⁹ It is estimated that the vaccine could provide cross-immunity against the *Monkeypox virus* with an efficacy of approximately 80%–85%,¹⁰ given the aforementioned genomic similarity among *Orthopoxvirus*.⁸

Funding

None declared.

Conflict of interests

None declared.

References

1. Rodríguez-Cuadrado FJ, Pinto-Pulido EL, Fernández-Parrado M. Potential treatments for monkeypox. *Actas Dermosifiliogr.* 2022;S0001-7310:00601-609, <http://dx.doi.org/10.1016/j.ad.2022.06.013>.
2. European Medicines Agency. Tecovirimat SIGA (tecovirimat): an overview of tecovirimat SIGA and why it is authorised in the EU. European Medicines Agency; 2022. Available from:

DOI of original article:

<https://doi.org/10.1016/j.ad.2022.08.017>

[☆] Please cite this article as: Rodríguez-Cuadrado FJ, Pinto-Pulido EL, Fernández-Parrado M. Inmunoglobulina anti-*Vaccinia* y profilaxis postexposición mediante vacuna basada en *Vaccinia* para el control del brote de viruela símica (*Monkeypox*). *Actas Dermosifiliogr.* 2022. <https://doi.org/10.1016/j.ad.2022.08.017>.

<https://doi.org/10.1016/j.ad.2022.08.026>

0001-7310/© 2022 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: F.J. Rodríguez-Cuadrado, E.L. Pinto-Pulido and M. Fernández-Parrado, [Translated article] Anti-*Vaccinia* Immunoglobulin and Post-exposure Prophylaxis with *Vaccinia*-based Vaccine for Management of the Monkeypox Outbreak, *ACTAS Dermo-Sifiliográficas*, <https://doi.org/10.1016/j.ad.2022.08.026>

- <https://www.ema.europa.eu/en/medicines/human/EPAR/tecovirimat-siga#:~:text=the%20European%20Union.-,Overview,happen%20following%20vaccination%20against%20smallpox> [accessed 20.07.22].
3. Sookaromdee P, Wiwanitkit V. Treatments for monkeypox. *Actas Dermosifiliogr.* 2022, <http://dx.doi.org/10.1016/j.ad.2022.07.022>. S0001-7310(22)00769-4.
 4. World Health Organization. Second meeting of the International Health Regulations (2005) (IHR) Emergency Committee regarding the multi-country outbreak of monkeypox. 2022. Available from: [https://www.who.int/news/item/23-07-2022-second-meeting-of-the-international-health-regulations-\(2005\)-\(ihr\)-emergency-committee-regarding-the-multi-country-outbreak-of-monkeypox](https://www.who.int/news/item/23-07-2022-second-meeting-of-the-international-health-regulations-(2005)-(ihr)-emergency-committee-regarding-the-multi-country-outbreak-of-monkeypox) [accessed 24.07.22].
 5. Tambo E, Al-Nazawi AM. Combating the global spread of poverty-related Monkeypox outbreaks and beyond. *Infect Dis Poverty.* 2022;11:80, <http://dx.doi.org/10.1186/s40249-022-01004-9>.
 6. Wittek R. Vaccinia immune globulin: current policies, preparedness, and product safety and efficacy. *Int J Infect Dis.* 2006;10:193–201, <http://dx.doi.org/10.1016/j.ijid.2005.12.001>.
 7. Rizk JG, Lippi G, Henry BM, Forthal DN, Rizk Y. Prevention and treatment of monkeypox. *Drugs.* 2022;82:957–63, <http://dx.doi.org/10.1007/s40265-022-01742-y>.
 8. Shchelkunov SN, Totmenin AV, Babkin IV, Safronov PF, Ryazankina OI, Petrov NA, et al. Human monkeypox and smallpox viruses: genomic comparison. *FEBS Lett.* 2001;509:66–70, [http://dx.doi.org/10.1016/s0014-5793\(01\)03144-1](http://dx.doi.org/10.1016/s0014-5793(01)03144-1).
 9. World Health Organization. Vaccines and immunization for monkeypox: Interim guidance. 2022. Available from: <https://www.who.int/publications/i/item/who-mpx-immunization-2022.1> [accessed 24.07.22].
 10. Rimoin AW, Mulembakani PM, Johnston SC, Lloyd Smith JO, Kivalu NK, Kinkela TL, et al. Major increase in human monkeypox incidence 30 years after smallpox vaccination campaigns cease in the Democratic Republic of Congo. *Proc Natl Acad Sci U S A.* 2010;107:16262–7, <http://dx.doi.org/10.1073/pnas.1005769107>.

F.J. Rodríguez-Cuadrado^{a,*}, E.L. Pinto-Pulido^b,
M. Fernández-Parrado^c

^a *Servicio de Dermatología, Hospital Universitario Puerta de Hierro Majadahonda, Madrid, Spain*
^b *Servicio de Dermatología, Hospital Universitario Príncipe de Asturias, Madrid, Spain*
^c *Servicio de Dermatología, Hospital Universitario de Navarra, Pamplona, Spain*

* Corresponding author.
E-mail address:
franciscojose.rodriguezcuadrado@gmail.com
(F.J. Rodríguez-Cuadrado).