

RESEARCH ARTICLE

# Commensal *Streptococcus salivarius* Modulates PPAR $\gamma$ Transcriptional Activity in Human Intestinal Epithelial Cells

Benoît Couvigny<sup>1,2</sup>, Tomas de Wouters<sup>1,2</sup>, Ghaliya Kaci<sup>1,2</sup>, Elsa Jacouton<sup>1,2</sup>, Christine Delorme<sup>1,2</sup>, Joël Doré<sup>1,2,3</sup>, Pierre Renault<sup>1,2</sup>, Hervé M. Blottière<sup>1,2,3</sup>, Eric Guédon<sup>1,2</sup><sup>\*</sup>, Nicolas Lapaque<sup>1,2</sup><sup>\*</sup>

**1** INRA, UMR 1319 MICALIS, Domaine de Vilvert, Jouy-en-Josas, France, **2** AgroParisTech, UMR Micalis, Jouy-en-Josas, France, **3** INRA, US 1367 MetaGenoPolis, Jouy-en-Josas, France

<sup>\*</sup> These authors contributed equally to this work.

<sup>\*</sup> [eric.guedon@jouy.inra.fr](mailto:eric.guedon@jouy.inra.fr) (EG); [nicolas.lapaque@jouy.inra.fr](mailto:nicolas.lapaque@jouy.inra.fr) (NL)



**OPEN ACCESS**

**Citation:** Couvigny B, de Wouters T, Kaci G, Jacouton E, Delorme C, Doré J, et al. (2015) Commensal *Streptococcus salivarius* Modulates PPAR $\gamma$  Transcriptional Activity in Human Intestinal Epithelial Cells. PLoS ONE 10(5): e0125371. doi:10.1371/journal.pone.0125371

**Academic Editor:** Raymond Schuch, ContraFect Corporation, UNITED STATES

**Received:** November 14, 2014

**Accepted:** March 23, 2015

**Published:** May 6, 2015

**Copyright:** © 2015 Couvigny et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by Institut National de la Recherche Agronomique and financed by grants from the European Community's Seventh Framework Program (FP7/2007-2013), MetaHIT, grant agreement HEALTH-F4-2007-201052, the ANR project MetaGenoPolis (ANR-11-DPBS-0001) and the ANR Blanc FunMetaGen. BC was a recipient of a grant from the ministère de la Recherche et de l'Éducation Nationale (ED425).

## Abstract

The impact of commensal bacteria in eukaryotic transcriptional regulation has increasingly been demonstrated over the last decades. A multitude of studies have shown direct effects of commensal bacteria from local transcriptional activity to systemic impact. The commensal bacterium *Streptococcus salivarius* is one of the early bacteria colonizing the oral and gut mucosal surfaces. It has been shown to down-regulate nuclear transcription factor (NF- $\kappa$ B) in human intestinal cells, a central regulator of the host mucosal immune system response to the microbiota. In order to evaluate its impact on a further important transcription factor shown to link metabolism and inflammation in the intestine, namely PPAR $\gamma$  (peroxisome proliferator-activated receptor), we used human intestinal epithelial cell-lines engineered to monitor PPAR $\gamma$  transcriptional activity in response to a wide range of *S. salivarius* strains. We demonstrated that different strains from this bacterial group share the property to inhibit PPAR $\gamma$  activation independently of the ligand used. First attempts to identify the nature of the active compounds showed that it is a low-molecular-weight, DNase-, proteases- and heat-resistant metabolite secreted by *S. salivarius* strains. Among PPAR $\gamma$ -targeted metabolic genes, *I-FABP* and *Angptl4* expression levels were dramatically reduced in intestinal epithelial cells exposed to *S. salivarius* supernatant. Both gene products modulate lipid accumulation in cells and down-regulating their expression might consequently affect host health. Our study shows that species belonging to the salivarius group of streptococci impact both host inflammatory and metabolic regulation suggesting a possible role in the host homeostasis and health.

## Introduction

The Human gastrointestinal tract (GIT) harbors a highly diverse and dense population of commensal microorganisms, commonly named microbiota. Its functions were formerly

**Competing Interests:** The authors have declared that no competing interests exist.

thought to be purely digestive and protective by building a competitive barrier against pathogen colonization. Research performed over the last decades has given rise to an emergent awareness that the function of the GIT along with its microbiota also strongly influences host physiology, locally and at a systemic level contributing largely to the host health and wellbeing (for review see [1]). The interface between commensal bacteria and the host epithelium is crucial for the establishment of this interaction in a homeostatic and mutualistic manner. With a large genetic pool (over 150 times larger than the Human genome), the microbiota is highly adapted for intestinal fermentation of non-digestible foodstuff [2]. On top of this important fermentative role it contributes to the development of the local and systemic immune system, to the regulation of host fat storage and even to behavior [3–9]. Strong correlations between the microbiota, low-grade inflammation and host metabolism have been highlighted recently [4, 10, 11]. However, the understanding of underlying mechanisms by which the gut microbiota could contribute to the host metabolic homeostasis or functions remains fragmentary.

An important role has been attributed to metabolites produced by the microbiota (including short chain fatty acids (SCFA)) in the activation of peroxisome proliferator-activated receptor (PPAR) family of nuclear receptors that initiate transcriptional gene expression linked to metabolic reprogramming and immune functions [12–15]. PPAR $\gamma$  is a well-characterized nuclear receptor for which natural known ligands are endogenous and exogenous lipid moieties along with derivatives of thiazolidinedione [16]. PPAR $\gamma$  forms heterodimers with the retinoid X receptor (RXR) and upon activation stimulates target-gene expression through binding to PPAR-responsive elements (PPREs) [17]. PPAR $\gamma$  is predominantly expressed in adipose tissue and the GIT, and is involved in the metabolic regulation of lipids, glucose homeostasis, cell proliferation and differentiation and local inflammation. Microbiota-induced PPAR $\gamma$  has also a role beyond the gut, as it regulates *Angptl4* (Angiopoietin like protein-4) expression, responsible for lipid storage in the adipose tissues [12, 18]. In addition to its implication in metabolic gene regulation, PPAR $\gamma$  is a well-characterized transcription factor recently reported to act as an E3 ubiquitin ligase implicated in the degradation of the p65 subunit of the pleiotropic nuclear factor NF- $\kappa$ B that consequently reduces inflammatory responses [19, 20].

*Streptococcus salivarius* is one of the primo colonizers of the human oral cavity and upper airways where it remains a predominant commensal inhabitant [21–23]. This bacterium is also a dominant part of the early-life human intestinal microbiota (for review see [24]). In adults, *S. salivarius* colonizes the stomach and intestine mucosal surfaces including the ileum, jejunum and colon [23, 25–28]. Metagenomic and phylogenetic microarray approaches showed that *Streptococcus* species are ubiquitously present in ileum and colon of healthy adults in dominance and sub-dominance respectively [29–32]. Previously, our laboratory demonstrated that *S. salivarius* strains display regulatory effects on the NF- $\kappa$ B pathway in human intestinal epithelial cells via unknown pathways and protect its host during induced-colitis suggesting a potential role in GIT inflammatory homeostasis [33, 34]. Interestingly, pathogen and commensal bacteria have been shown to actively modulate PPAR $\gamma$  and NF- $\kappa$ B activation [13, 20, 35–38]. Altogether, these studies suggest that, NF- $\kappa$ B and PPAR $\gamma$  are transcription factors, influenced by bacteria, with distinct but overlapping roles linking metabolic and immune functions. These observations prompted us to investigate the effect of *S. salivarius* on PPAR $\gamma$  transcriptional activity. In the present study, we showed that *S. salivarius* strains down-regulate PPAR $\gamma$  transcriptional activity and PPAR $\gamma$ -dependent genes expression *via* low molecular weight, heat resistant, non peptidic and non nucleotidic secreted components.

## Materials and Methods

### Epithelial cell culture

The human epithelial cell lines HT-29, Caco-2 and SW-116 were obtained from the American Type Culture Collection (ATCC). HT-29 and SW-116 were cultured in RPMI 1640 and Caco-2 in DMEM, supplemented with 10% heat-inactivated foetal calf serum, 2mM glutamine, 1X non essential amino acids, penicillin (50U/ml) and streptomycin (50U/ml) in a humidified 5% CO<sub>2</sub> atmosphere at 37°C. All culture media and supplements were supplied by Lonza.

### Reagents

All PPAR $\gamma$  ligands and antagonist: Rosiglitazone (10 $\mu$ M), GW9662 (10 $\mu$ M) pioglitazone (10 $\mu$ M), troglitazone (10 $\mu$ M) and ciglitazone (10 $\mu$ M) (Cayman chemical) were prepared in DMSO following the manufacturer's recommendations and diluted in RPMI. Sodium butyrate (But, 2mM) was from Sigma-aldrich.

### Bacterial culture and bacterial supernatant preparation

The *Streptococcus salivarius* strains used in this study are listed in [S1 Table](#). Bacteria were grown overnight at 37°C, *S. salivarius*, *S. vestibularis* and *S. agalactiae* in chemically defined medium (CDM) [39] and *E. coli* in Luriani Broth (LB). Bacterial cultures supernatants were collected after centrifugation and 0.22 $\mu$ m filtration. CDM adjusted to bacterial supernatant pH (pH $\approx$ 5.5) was used as control. When mentioned, bacterial supernatants were size-fractionated using 10kDa and 3kDa cut-off filters (Millipore) or exposed to high temperature (100°C for 10 min) and heat shock assay (100°C/10 min prior to liquid nitrogen-freezing). Bacterial lysates were prepared by mechanical lysis using a FastPrep instrument (MP Biomedicals). 3kDa filtered bacterial supernatants and CDM were treated with proteinase K (100 $\mu$ g/ml, Sigma), DNase I (100 $\mu$ g/ml, Sigma) or trypsin (0,25%) for 2h at 37°C. Then, based on their molecular weight characteristics ranging from 23 to 29 kDa, the enzymes were eliminated by using a 10kDa cut-off filters.

### Luciferase Reporter and cell viability Assays

PPAR $\gamma$  reporter construct pJ3-TK-Luc bearing three repeats of the PPRE (PPAR $\gamma$  responsive element) [40] was used to establishing HT-29-PPAR $\gamma$  and Caco-2-PPAR $\gamma$  reporter cell lines as described [41]. PPAR $\gamma$  reporter system response was characterised using a wide range of known agonists: rosiglitazone, pioglitazone, troglitazone and ciglitazone and its well-characterised antagonist GW9662 ([S3 Fig](#)). ANGPTL4-reporter system containing the pANGPTL4-1.8luc was previously described [12]. Cell reporters were seeded in 96-well plates at 3.10<sup>4</sup> cells per well. After 24h culture, bacterial supernatants or lysates, CDM, PPAR $\gamma$  agonists/antagonists were added at 10% final volume for 12h. For the live bacteria experiments, *S. salivarius* over-night cultures were incubated at a MOI of 40 bacteria per cell for 6h. Luciferase activity was measured in cells lysates using Tecan Infinite M200 device and luciferase assay kit (One Glo, Promega) according to the manufacturer's instructions. PPAR $\gamma$  activation and *Angptl4* expression were normalized to controls, i.e. cells stimulated with activators in addition to CDM control media or CDM control media alone. Experiments were performed in triplicates for at least three independent assays. Cell viability was monitored by MTS measurement using the CellTiter 96 Aqueous One solution (Promega) according to the manufacturer's recommendations.

## Real Time PCR

HT-29 cell line was seeded in 6-well culture plates at densities of  $10^6$  cells per well and cultured for 24h prior stimulation. Total messenger RNA (mRNA) was extracted after 6h incubation with CDM, CDM and rosiglitazone ( $10\mu\text{M}$ ), or supernatant and rosiglitazone using an RNeasy mini-kit (Qiagen), cDNA was synthesized from 2  $\mu\text{g}$  of mRNA using High Capacity cDNA Reverse Transcription Hits (Applied biosystems) according to the manufacturer's instructions. qPCRs were carried out using thermal cycler in a reaction volume of 25 $\mu\text{l}$  and SYBR Green (Applied Biosystems)-based quantitative real-time PCR. Primers were designed and tested according to Applied Biosystems recommendations. Liver-Fatty acid binding protein (L-FABP), 5' -GGAAGCACTTCAAGTTCACCAT-3' / 5' -ACCTTCCAACCTGAACCACTGTC-3' ;  $\beta$ -actin (used for normalization), 5' -AAGACCTGTACGCCAACACAGT-3' / 5' -GGAGCAATGATCTTGATCTTCA-3' ; Angiotensin like protein-4 (*Angptl4*), 5' -AGGCTGGACAGTAATTCAGAGG-3' / 5' -ATGCTATGCACCTTCTCCAGAC-3' ; Peroxisome Proliferator-Activated Receptor- $\gamma$  (PPAR $\gamma$ ), 5' -TCCAGTGGTTGCAGATTACAAG3' / 5' -AGGCTCTTCATGAGGCTTAT TG-3' . The sample setups included biological duplicates and experimental triplicates.

## Western-blot

Protein extracts were run in 10% SDS-PAGE and transferred onto PVDF membranes (Bio-Rad, Transblot). Membranes were blocked overnight in PBS/4% skim milk/0.1% Tween-20 (Sigma-aldrich). Primary (anti-PPAR $\gamma$ , SC-7273, clone E-8; anti-GAPDH, SC-365062, clone G-9; both from Santa Cruz Biotechnology) and secondary (Goat anti-mouse IgG HRP, Dako, P0447) antibodies were successively added in PBS/Tween/milk, each being left for 1h before detection with the Clarity Western ECL Substrate using the Chemidoc MP System (Bio-Rad). Quantifications were performed using the image Lab software (Bio-Rad).

## Statistical analysis

Graphic representation and statistical analyses of PPAR $\gamma$ , *Angptl4* and FABP expressions by RT-qPCR were performed using Graphpad Prism software. Comparisons of distributions were performed using a student's t test with 95% confidence intervals with p values of  $\leq 0.05$  considered to be significant.

## Results

### *S. salivarius* strains inhibit PPAR $\gamma$ activity

We previously demonstrated that *S. salivarius* strains inhibit NF- $\kappa$ B activation in IECs through an unknown mechanism [34]. PPAR $\gamma$  is a well-characterized nuclear receptor recently reported to act as an E3 ubiquitin ligase implicated in the degradation of the p65 sub-unit of NF- $\kappa$ B and consequently reducing inflammatory responses [19]. As NF- $\kappa$ B and PPAR $\gamma$  are transcription factors targeted by bacteria including commensals [20, 34], we aimed to understand if *S. salivarius*-dependent down-modulation of NF- $\kappa$ B was linked to the induction of PPAR $\gamma$  transcriptional activity.

For this purpose, we used a PPAR $\gamma$  reporter system under transcriptional control by three repeats of a PPRE (PPAR $\gamma$ -responsive element) stably expressed in HT-29 intestinal epithelial cells [12]. We screened the effect of bacterial supernatants derived from a wide range of *S. salivarius* and *vestibularis* strains isolated from diverse human sites previously described to modulate NF- $\kappa$ B pathway on the PPAR $\gamma$  reporter system [34]. Using non-activated cells, none of the strains tested showed any effect on PPAR $\gamma$  activation *per se*, suggesting that PPAR $\gamma$  is not involved in *S. salivarius*-dependent NF- $\kappa$ B inhibition (S1 Fig). Interestingly, upon activation of

PPAR $\gamma$  by butyrate supernatants secreted by *S. salivarius* and *S. vestibularis* strains down-regulated activity with an inhibition rate, ranging from 15% to 40%. (Fig 1A). Indirect effects through cell viability were excluded using the MTS assay (S2 Fig). This repression was specific to *S. salivarius* and *S. vestibularis* supernatants as *E. coli* and *S. agalactiae* supernatants did not affect butyrate-induced PPAR $\gamma$  transcriptional activity (Fig 1B).

### The *S. salivarius*–dependent down-regulation of PPAR $\gamma$ transcriptional activity relies on secreted molecules

Growth media are highly enriched in molecules, which are potentially metabolized by bacteria. In order to avoid growth medium-derived effects in PPAR $\gamma$  down-regulation, we used living bacteria from exponentially growing cultures that were washed with PBS prior to application to the PPAR $\gamma$ -HT-29 reporter cell line. In Fig 2, we showed that live *S. salivarius* strains, similarly to their culture supernatants inhibited the activation of butyrate-induced PPAR $\gamma$  on intestinal epithelial cells (Fig 2A and 2B). Moreover, bacterial cell components are known to be released during replication and lysis in exponentially growing cultures. In order to test if the observed PPAR $\gamma$  down-regulation with bacterial supernatants resulted from cytoplasmic or membranous compounds that might have been released during growth, we tested lysates of the washed bacteria on HT-29-PPAR $\gamma$  reporter cell-line but did not find any significant changes in PPAR $\gamma$  activation (Fig 2C).

These results strongly suggest an effect on butyrate-induced PPAR $\gamma$  activation exclusively with secreted components from metabolically active bacteria.


### The *S. salivarius*–dependent down-regulation of PPAR $\gamma$ activity is not dependent of the epithelial cell-line or PPAR $\gamma$ ligands used

*S. salivarius* effects on PPAR $\gamma$  activation was also confirmed in a second epithelial cell line, i.e. Caco-2, carrying the same reporter system (Fig 3A). Supernatants from three *S. salivarius* strains down-regulated butyrate-induced PPAR $\gamma$  activation in Caco-2 cells in a similar range as observed in HT-29 (30 to 40%).

Butyrate is a short chain fatty acid (SCFA) with broad cellular activities through receptor activation such as GPR41 and GPR43 or histone deacetylases (HDAC) inhibition [12]. The observed down-regulation of butyrate mediated PPAR $\gamma$  activation by *S. salivarius* and *S. vestibularis* supernatants could therefore be mediated either through direct PPAR $\gamma$  inhibition or indirectly through inhibition of the activation mechanisms of butyrate. We therefore repeated our experiments with a range of well-characterized specific PPAR $\gamma$  ligands inducing its translocation to the nucleus and consequently enhancing its transcriptional activity (Fig 3B and S3 Fig). Interestingly, bacterial supernatants inhibited PPAR $\gamma$  activation independently of the PPAR $\gamma$  ligand used (rosiglitazone, pioglitazone, sodium butyrate) suggesting a ligand-specific effect and potentially a common downstream regulation of PPAR $\gamma$  activation pathway (Fig 3B).

### The secreted bioactive component is a small, non nucleotidic, non peptidic, heat-resistant molecule


By serial dilution of the bacterial supernatants, we showed that the observed inhibitory effect was dose-dependent and detected at a concentration of 2.5% of supernatant in the culture media suggesting that the bioactive compound was expressed in a significant amount (Fig 4A). Standard size-cutoff columns were used to assess the size of the active(s) compound(s) secreted by *S. salivarius*. Bacterial supernatants were submitted to ultrafiltration through 3 and 10 kDa



**Fig 1. *S. salivarius* and *S. vestibularis*, but not *S. agalactiae* and *E. coli*, culture supernatants down-regulate butyrate-induced PPAR $\gamma$  activity in HT-29/PPAR $\gamma$ .** A- Culture supernatants of a wide range of *S. salivarius* (white bars) and *S. vestibularis* (grey bars) strains were applied on HT-29-PPAR $\gamma$  (10% vol/vol) prior to PPAR $\gamma$ -induction by sodium butyrate (But, 2mM) for 12h. PPAR $\gamma$  transcriptional activity was measured by luciferase activity and expressed as fold increase towards its control: growth-medium + butyrate. Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. The dark grey bars represent the selected *S. salivarius* strains used for the remaining of the study (JIM8772, JIM8777 and K12). B- Culture supernatants of three *S. salivarius* (JIM8772, JIM8777 and K12), *E. coli* and *S. agalactiae* strains were applied on HT-29-PPAR $\gamma$  (10% vol/vol) prior to PPAR $\gamma$ -induction by sodium butyrate (2mM) for 12h. PPAR $\gamma$  expression was measured by luciferase activity and expressed as fold increase towards growth-medium + sodium butyrate (Control). Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. \*\*\*P<0.001 compared with controls (Student's t-test).


doi:10.1371/journal.pone.0125371.g001





**Fig 2. The *S. salivarius*–dependent down-regulation of PPAR $\gamma$  activity relies on secreted molecules.** PPAR $\gamma$  was activated (+ But) or not (- But) in HT-29 reporter cells using sodium butyrate (2mM) in the presence of control medium (Control) or as indicated live bacteria (A), bacterial supernatants (B) or lysates (C). The values represent the luciferase activity normalized towards their respective control. Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. \*\*\*P<0.001, \*\*P<0.005, compared with controls (Student's t-test).


doi:10.1371/journal.pone.0125371.g002



**Fig 3. The *S. salivarius*-dependent down-regulation of PPAR $\gamma$  activity is independent of the epithelial cell-line or the specific PPAR $\gamma$ -ligand used.** A- PPAR $\gamma$  was activated (+ But) or not (- But) in Caco-2 reporter cells (Caco2-PPAR $\gamma$ ) using sodium butyrate (2mM) in the presence of control medium (Control) or supernatants. B- HT-29-PPAR $\gamma$  reporter cell-line was activated with different activators: rosiglitazone (Rosi, 10 $\mu$ M), pioglitazone (Pio, 10 $\mu$ M) and sodium butyrate (But, 2mM) in presence of *S. salivarius* JIM8777 supernatant. The values represent the luciferase activity normalized towards their respective control. Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. \*\*\*P<0.001, \*\*P<0.005, \*P<0,05 compared with controls (Student's t-test).

doi:10.1371/journal.pone.0125371.g003





**Fig 4. Determination of the nature and molecular mass of the secreted bioactive compounds.** A- Serial dilutions of *S. salivarius* JIM8777 supernatant were tested on HT-29-PPAR $\gamma$  cells prior to activation with sodium butyrate (2mM). B- Exposure to high temperature (100°C/10 min) and heat shock were applied to *S. salivarius* JIM8777 supernatant prior to addition to activated HT-29-PPAR $\gamma$  cells. C- Butyrate-activated HT-29-PPAR $\gamma$  cells were incubated with *S. salivarius* supernatants fractions derived from ultrafiltration through 3 and 10 kDa cutoff membranes. >10kDa/>3kDa and <10kDa/<3kDa represent the retained and filtered fractions respectively. Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. \*\*\*P<0.001, \*\*P<0.005, \*P<0,05 compared with controls (Student's t-test).

doi:10.1371/journal.pone.0125371.g004

cutoff membranes. For the three strains tested, the <10 kDa and <3 kDa fractions inhibited PPAR $\gamma$  activity similarly to the unfiltered fractions while fractions > 10kDa and >3 kDa displayed no effect suggesting that the active compound was a small molecule lower than 3kDa (Fig 4B). Moreover, exposure to high temperatures (100°C for 10 min) or to heat shock (100°C for 10 min before freezing in liquid nitrogen) did not affect the inhibitory property of the supernatant (Fig 4C). Furthermore, the treatments of *S. salivarius* supernatants with proteases (proteinase K and trypsin) or DNase I did not impair the inhibition of butyrate-induced PPAR $\gamma$  activity (Fig 5). All together, these results showed that the bioactive compound is a small non-peptidic and non-nucleotidic heat-resistant molecule.

### *S. salivarius* modulates PPAR $\gamma$ -dependent gene expression, I-FABP and *Angptl4*, without affecting PPAR $\gamma$ expression


To assess the impact of *S. salivarius*-dependent inhibition on PPAR $\gamma$  target genes, we quantified by RT-qPCR the expression of *Angptl4* and *I-FABP* (intestinal fatty acid binding protein). The expression of these two well-characterized PPAR $\gamma$ -regulated genes were strongly reduced by *S. salivarius* supernatant in the presence of the PPAR $\gamma$ -specific ligand, rosiglitazone (Fig 6A and 6B). This inhibitory effect was independent on PPAR $\gamma$  expression as its expression level was unchanged when HT-29 cells were incubated with both supernatant and the specific ligand rosiglitazone (Fig 6C). Moreover, *S. salivarius* did not decrease the stability of PPAR $\gamma$  protein as it remained identical after bacterial supernatant treatment (Fig 7). To confirm these observations, we used a cell-line expressing a ANGPTL4 reporter system containing the 1.8 kb of ANGPTL4 promoter devoid of the functional PPRE present in the third intron of the gene [12]. As shown in Fig 8, none of the *S. salivarius* supernatants tested were able to affect the PPAR $\gamma$ -independent *Angptl4* expression induced by sodium butyrate.

Together, these findings show that *S. salivarius* supernatants specifically repressed the PPAR $\gamma$  pathway without affecting PPAR $\gamma$  gene expression and protein stability, and consequently can impact metabolic responses.

## Discussion


Commensal bacteria provide crucial biological functions to their host, including metabolic functions and the control of moderate immunological responses, beneficial for human health. This intimate relationship results from a balanced interaction between the host and its microbiota mediated through direct contact or secreted active compounds. The colonization of the human digestive tract by commensals occurs immediately after birth and has been shown by a growing number of evidence to be the key factor directly impacting host physiology [42–45]. However, while most studies concern adult or predominant gut microbiota, very little is known about the impact of early bacterial colonizers on human physiology. The commensal bacterium *Streptococcus salivarius* is one of the primo-colonizers of oral and gut mucosal surfaces that have been shown to influence inflammation by down-modulating NF- $\kappa$ B activity in human intestinal cells [33, 34]. The present study focused on the direct impact of *S. salivarius* on a key transcription factor in intestinal epithelial cells (IEC) that links metabolism and inflammation, namely PPAR $\gamma$ .

One of PPAR $\gamma$ 's immunomodulatory properties is to inhibit NF- $\kappa$ B activation by directly impairing its translocation into the nucleus or by acting as an E3 ligase targeting the p65 subunit of NF- $\kappa$ B for degradation [19]. We previously showed that *S. salivarius* strains inhibit NF- $\kappa$ B activation in IECs via unknown pathways leading to *in vivo* protection in induced-colitis mice models [33, 34]. Since several studies have linked NF- $\kappa$ B control and PPAR $\gamma$  activity, it was thus conceivable that *S. salivarius*-dependent down-modulation of NF- $\kappa$ B could be linked




**Fig 5. The *S. salivarius* bioactive compounds are not from peptidic or nucleotidic origin.** A- 3kDa-filtered *S. salivarius* JIM8777 supernatants were exposed to proteases (trypsin, left panel; proteinase K, right panel) prior to addition to butyrate-activated HT-29-PPAR $\gamma$  cells. B- 3kDa-filtered *S. salivarius* JIM8777 supernatants were exposed to DNase I prior to addition to butyrate-activated HT-29-PPAR $\gamma$  cells.

doi:10.1371/journal.pone.0125371.g005




**Fig 6. Transcriptional regulation of PPAR $\gamma$  and PPAR $\gamma$ -target genes upon stimulation with rosiglitazone and/or *S. salivarius* supernatant.** The mRNA expression of *I-FABP* (A), *Angptl4* (B) and *PPAR $\gamma$*  (C) were determined by Quantitative real-time PCR on total RNA extracted from HT-29 cells exposed to culture medium (CDM), *S. salivarius* JIM8777 supernatant (Sn) alone or in addition with the PPAR $\gamma$  specific activator rosiglitazone (CDM+Rosi; Sn+Rosi) for 6 h. Expression is represented as fold change compared to the absence of any stimulation (CDM medium only). Data are represented as mean  $\pm$  standard deviation (SD) of 2 to 3 independent repetitions done in triplicates. \*\*P<0.005, \*P<0,05 compared with controls (Student's t-test).

doi:10.1371/journal.pone.0125371.g006



**Fig 7. PPAR $\gamma$  protein level is unchanged upon stimulation with sodium butyrate and/or *S. salivarius* supernatant.** A- The protein level of PPAR $\gamma$  and GAPDH were determined by western-blot on total protein extracted from HT-29 cells exposed to culture medium (CDM), *S. salivarius* JIM8777 supernatant in addition with sodium butyrate (Control + Butyrate; JIM8777 + Butyrate) for 24 h. B- Quantifications of total PPAR $\gamma$  protein normalized to GAPDH protein level. Protein expression is represented as fold change compared to the sodium butyrate stimulation in presence of culture media (Control + Butyrate). Data are represented as mean  $\pm$  standard deviation (SD) of the effect of 5 independent bacterial cultures.

doi:10.1371/journal.pone.0125371.g007



**Fig 8. *S. salivarius* supernatants do not affect PPAR $\gamma$ -independent ANGPTL4 reporter system.** ANGPTL4-reporter system cell-line was activated with sodium butyrate (2mM) with or without *S. salivarius* supernatant (JIM8772, JIM8777 or K12). *Angptl4* expression was measured by luciferase activity and expressed as fold increase towards its control: growth-medium alone (left panel) and growth-medium + sodium butyrate (right panel). Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. \*\*\*P<0.001, \*\*P<0.005, \*P<0,05 compared with controls (Student's t-test).

doi:10.1371/journal.pone.0125371.g008

to the induction of PPAR $\gamma$  transcriptional activity [19, 20]. Using human intestinal epithelial cell-lines engineered to monitor PPAR $\gamma$  transcriptional activity in response to a wide range of *S. salivarius* strains, we demonstrate that different members of the *S. salivarius* group share the capacity to inhibit PPAR $\gamma$  transcriptional activity. This result suggests that PPAR $\gamma$  is not involved in the NF- $\kappa$ B-dependent anti-inflammatory properties of *S. salivarius*. Moreover, these results together with our previous study suggest that species belonging to the streptococcal salivarius group of commensal streptococci are presenting effects both on host inflammatory regulation and on metabolism processes [34].

First attempts to decipher the nature of the active compound showed that it is a low-molecular-weight metabolite (<3kDa) produced and secreted by *S. salivarius*. Moreover, exposure to enzymatic and heat treatments did not affect the inhibitory potential suggesting that it may be a small organic molecule of non-peptidic and non-nucleotidic origin. The phylogenetic proximity of the strains tested suggests a potential related metabolite production pathway inherited from a common ancestor [46]. We and others showed that common bacterial metabolites such as short-chain fatty acids (SCFAs) regulate PPAR $\gamma$  activity [12, 41]. It is unlikely that SCFAs are involved in PPAR $\gamma$  down-modulation as acetate, butyrate and propionate have an opposite effect to *S. salivarius* supernatant by inducing the up-regulation of PPAR $\gamma$  responses (S3 Fig and [41]). Moreover *S. salivarius* is not reported to produce butyrate, propionate and acetate at activating concentrations [34]. We also ruled out that the inhibitory effect was due to lactic acid known to be produced by *S. salivarius*. Indeed, high-pressure liquid chromatography (HPLC) measurement of organics revealed a maximal concentration of lactic acid around 80 mM in *S. salivarius* supernatant [34]. We assess the effect of lactic acid on our reporter systems and observe no effect on PPAR $\gamma$  activity (data not shown). Interestingly, on the contrary to what was observed for NF- $\kappa$ B down-modulation, *S. salivarius* derived compounds inhibiting



PPAR $\gamma$  are resistant to trypsin treatment ruling out a component with a pleiotropic activity on the host.

Recently, pathogen and commensal bacteria have been shown to actively modulate PPAR $\gamma$  transcriptional activity mainly *via* up-regulating its expression (including *Helicobacter pylori*, *Bacteroidetes thetaiotaomicron*, *Brucella abortus*, *Salmonella Typhimurium*, *Mycobacterium tuberculosis*, *Neisseria lactamica* and short chain fatty acids derived from commensals) [12, 13, 20, 35, 37, 47–49]. Alternatively, PPAR $\gamma$  activity is induced by phosphorylation via unknown bacterial molecules derived from *Enterococcus faecalis* or by direct interaction with lipids derived from *M. tuberculosis* [13, 48]. For pathogens, a direct correlation between PPAR $\gamma$  up-regulation and bacterial survival has been observed [48]. On the contrary to what is observed with these pathogens, we showed that PPAR $\gamma$  activity is impaired by *S. salivarius* supernatants. So far, only *Salmonella typhimurium* has been shown to inhibit PPAR $\gamma$  activity in mouse epithelial cells by down-regulating its gene expression, aggravating acute colitis [35]. However, common mechanisms are unlikely to occur as we showed by RT-qPCR and western-blotting that PPAR $\gamma$  gene and protein expression are unchanged in presence of *S. salivarius*-derived supernatant (Figs 6C and 7).

Endogenous ligands of PPAR $\gamma$  such as polyunsaturated fatty acids and their derivatives, along with the synthetic anti-diabetic drugs derived from thiazolidinediones such as rosiglitazone and pioglitazone promote heterodimers formation with the retinoid X receptor (RXR), and subsequent recognition of PPAR-response elements (PPREs) within target gene promoters. Additionally to ligand-dependent activation, PPAR $\gamma$  has been shown to be phosphorylated, SUMOylated and ubiquitinated regulating its transcriptional properties (for review see [50]). These diverse regulatory processes regulating PPAR $\gamma$  activity offer numerous possible mechanisms by which PPAR $\gamma$  transcriptional activity might be inhibited by *S. salivarius* supernatants. Our present study shows that *S. salivarius* supernatant inhibits PPAR $\gamma$  transcriptional activity induced in epithelial cells even by highly specific ligands such as rosiglitazone, pioglitazone or butyrate, a SCFA. Absence of repression on the basal PPAR $\gamma$  activity suggests a putative competition in the activation of the PPAR $\gamma$  pathway (Fig 1A). Butyrate-induced signaling cascade involve an increased expression of PPAR $\gamma$  mRNA whereas synthetic ligands induce its dimerization and phosphorylation suggesting that a common regulation in the early pathway is unlikely. Although the exact mechanisms remain unclear, the fact that *S. salivarius* supernatant inhibits PPAR $\gamma$  independently of the activator used, along with the nature of the inhibitory molecules suggest its possible SUMOylation known to recruit a co-repressor complex impairing PPAR $\gamma$  translocation or the implication of a ligand or regulatory pathways with direct antagonist effects [50].

Considering the multitude of strategies evolved in pathogenic bacteria to modulate the host environment in order to multiply and spread, it is not surprising that commensal microorganisms have co-evolved similarly to survive in highly competitive niches such as mucosal sites. *H. pylori* and *M. tuberculosis* up-regulate host PPAR $\gamma$  expression to suppress exaggerated inflammatory responses, ensuring their survival within the host [48, 49]. In contrast, *S. typhimurium* initiates an acute inflammation by inducing PPAR $\gamma$  down-regulation in the intestinal epithelium, leading to the development of a hostile niche for local competitors that will favor its survival [35]. It is therefore tempting to speculate that PPAR $\gamma$ -inhibition induced by *S. salivarius* is a mechanism to thwart pathogen's strategy or to enhance its own survival. However infection or colonization trials on animal models using *S. salivarius* as a PPAR $\gamma$  inhibitor would be needed to prove this hypothesis. Studies in animals have demonstrated that *S. salivarius* significantly inhibited inflammation in TNBS-induced colitis mouse models suggesting that the PPAR $\gamma$ -independent inhibition of NF- $\kappa$ B counterbalances the inhibition of PPAR $\gamma$  transcriptional activity observed in our study [33].

Despite being a master regulator of inflammation, by antagonizing the activities of the transcription factors AP-1, STAT, and NF- $\kappa$ B and inducing visceral adipose tissue (VAT)-resident regulatory T cells (Treg), PPAR $\gamma$  has a broad range of physiological properties including the regulation of lipid and glucose metabolism [20, 51–53]. Among PPAR $\gamma$ -targeted metabolic genes, *I-FABP* and *Angptl4* levels are dramatically reduced in presence of *S. salivarius*. Both of these gene products modulate lipid accumulation in cells and down-regulating their expression might consequently be detrimental for host health [18, 54–56]. However, the expression of lipid metabolic factors might be important in the weaning or early post-weaning to deal with a highly rich nutritional intake when *S. salivarius* is particularly dominant. It is therefore tempting to speculate that intestinal PPAR $\gamma$  may be one of several transcription factors controlling genes involved in lipid metabolism such as *Angptl4* under the influence of microbiota and recently reported in studies of obese patients [18, 56, 57]. We believe that a combination of commensal bacteria can be involved in transcriptional regulation through factors such as PPAR $\gamma$ , and that the consequence of such regulations are a tightly tuned balance of immune and metabolic genes. Studies of *S. salivarius* in high-fat diet obese or post-weaning mice models would help to decipher this important issue.

Here we demonstrate that strains of *S. salivarius*, an early colonizer of the gastrointestinal tract, can inhibit the activity of the transcription factor PPAR $\gamma$  and the subsequent expression of target genes in intestinal epithelial cell lines. By showing that a commensal bacterium targets such type of nuclear receptor with wide activity, this study reinforces the notion that the microbiota contributes to immune and metabolic regulation that are highly interconnected in intestinal cells.

## Supporting Information

**S1 Fig. *S. salivarius* and *S. vestibularis* culture supernatants do not impair steady-state PPAR $\gamma$  activity in HT-29/PPAR $\gamma$ .** Culture supernatants of a wide range of *S. salivarius* (white bars) and *S. vestibularis* (grey bars) strains were applied on HT-29-PPAR $\gamma$  (10% vol/vol) without any PPAR $\gamma$  activation for 12h. PPAR $\gamma$  transcriptional activity was measured by luciferase activity and expressed as fold increase towards its control: growth-medium alone. Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. The dark grey bars represent the selected *S. salivarius* strains used for the remaining of the study (JIM8772, JIM8777 and K12). (EPS)

**S2 Fig. *S. salivarius* supernatants do not affect cell viability of butyrate-induced HT-29 cells.** Cell viability was monitored on HT-29 cells by MTS measurement after incubation for 12h with butyrate alone (Control), sodium butyrate + growth media (CDM) or sodium butyrate + culture supernatant from *S. salivarius* (JIM8772, JIM8777 or K12). Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. (EPS)

**S3 Fig. HT-29-PPAR $\gamma$  cell-line responses to different PPAR $\gamma$  ligands (rosiglitazone, pioglitazone, tioglitazone, ciglitazone, 10 $\mu$ M).** The values represent the luciferase activity normalized towards the negative control (untreated cells). Data are expressed as means  $\pm$  standard deviations (SD) of triplicate measurements from one representative experiment out of three independent experiments. (EPS)

**S1 Table. Bacterial strains used in this study.**  
(DOCX)

## Acknowledgments

We thank Dr John Tagg for the JIM9125 (DC117A) and JIM9126 (H3) strains, Dr Diana R. Martins for other JIM strains, Malgorzata Nepelska for the Caco-2 PPAR $\gamma$  cell-line; Fabienne Devime and Rozenn Dervyn-Hirel for technical advices.

## Author Contributions

Conceived and designed the experiments: CD JD PR HMB EG NL. Performed the experiments: BC EJ EG NL. Analyzed the data: BC HMB EG NL. Contributed reagents/materials/analysis tools: TdW GK CD. Wrote the paper: TdW HMB EG NL.

## References

1. Sekirov I, Russell SL, Antunes LC, Finlay BB. Gut microbiota in health and disease. *Physiological reviews*. 2010; 90(3):859–904. doi: [10.1152/physrev.00045.2009](https://doi.org/10.1152/physrev.00045.2009) PMID: [20664075](https://pubmed.ncbi.nlm.nih.gov/20664075/).
2. Flint HJ, Scott KP, Louis P, Duncan SH. The role of the gut microbiota in nutrition and health. *Nature reviews Gastroenterology & hepatology*. 2012; 9(10):577–89. doi: [10.1038/nrgastro.2012.156](https://doi.org/10.1038/nrgastro.2012.156) PMID: [22945443](https://pubmed.ncbi.nlm.nih.gov/22945443/).
3. Atarashi K, Honda K. Microbiota in autoimmunity and tolerance. *Curr Opin Immunol*. 2011; 23(6): 761–8. doi: [10.1016/j.coi.2011.11.002](https://doi.org/10.1016/j.coi.2011.11.002) PMID: [22115876](https://pubmed.ncbi.nlm.nih.gov/22115876/).
4. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, et al. Richness of human gut microbiome correlates with metabolic markers. *Nature*. 2013; 500(7464):541–6. doi: [10.1038/nature12506](https://doi.org/10.1038/nature12506) PMID: [23985870](https://pubmed.ncbi.nlm.nih.gov/23985870/).
5. Ley RE, Backhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. Obesity alters gut microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102(31):11070–5. PMID: [16033867](https://pubmed.ncbi.nlm.nih.gov/16033867/).
6. Nishio J, Honda K. Immunoregulation by the gut microbiota. *Cell Mol Life Sci*. 2012; 69(21):3635–50. PMID: [22527722](https://pubmed.ncbi.nlm.nih.gov/22527722/). doi: [10.1007/s00018-012-0993-6](https://doi.org/10.1007/s00018-012-0993-6)
7. Turnbaugh PJ, Gordon JI. The core gut microbiome, energy balance and obesity. *J Physiol*. 2009; 587(Pt 17):4153–8. PMID: [19491241](https://pubmed.ncbi.nlm.nih.gov/19491241/). doi: [10.1113/jphysiol.2009.174136](https://doi.org/10.1113/jphysiol.2009.174136)
8. Diaz Heijtz R, Wang S, Anuar F, Qian Y, Bjorkholm B, Samuelsson A, et al. Normal gut microbiota modulates brain development and behavior. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(7):3047–52. doi: [10.1073/pnas.1010529108](https://doi.org/10.1073/pnas.1010529108) PMID: [21282636](https://pubmed.ncbi.nlm.nih.gov/21282636/); PubMed Central PMCID: PMC3041077.
9. Kelly D, Mulder IE. Microbiome and immunological interactions. *Nutr Rev*. 2012; 70 Suppl 1:S18–30. doi: [10.1111/j.1753-4887.2012.00498.x](https://doi.org/10.1111/j.1753-4887.2012.00498.x) PMID: [22861803](https://pubmed.ncbi.nlm.nih.gov/22861803/).
10. Chassaing B, Gewirtz AT. Gut microbiota, low-grade inflammation, and metabolic syndrome. *Toxicologic pathology*. 2014; 42(1):49–53. doi: [10.1177/0192623313508481](https://doi.org/10.1177/0192623313508481) PMID: [24285672](https://pubmed.ncbi.nlm.nih.gov/24285672/).
11. Nicholson JK, Holmes E, Kinross J, Burcelin R, Gibson G, Jia W, et al. Host-gut microbiota metabolic interactions. *Science*. 2012; 336(6086):1262–7. doi: [10.1126/science.1223813](https://doi.org/10.1126/science.1223813) PMID: [22674330](https://pubmed.ncbi.nlm.nih.gov/22674330/).
12. Korecka A, de Wouters T, Cultrone A, Lapaque N, Pettersson S, Dore J, et al. ANGPTL4 expression induced by butyrate and rosiglitazone in human intestinal epithelial cells utilizes independent pathways. *American journal of physiology Gastrointestinal and liver physiology*. 2013; 304(11):G1025–37. doi: [10.1152/ajpgi.00293.2012](https://doi.org/10.1152/ajpgi.00293.2012) PMID: [23518684](https://pubmed.ncbi.nlm.nih.gov/23518684/).
13. Are A, Aronsson L, Wang S, Greicius G, Lee YK, Gustafsson JA, et al. Enterococcus faecalis from newborn babies regulate endogenous PPAR $\gamma$  activity and IL-10 levels in colonic epithelial cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(6):1943–8. doi: [10.1073/pnas.0711734105](https://doi.org/10.1073/pnas.0711734105) PMID: [18234854](https://pubmed.ncbi.nlm.nih.gov/18234854/); PubMed Central PMCID: PMC2538862.
14. Crawford PA, Crowley JR, Sambandam N, Muegge BD, Costello EK, Hamady M, et al. Regulation of myocardial ketone body metabolism by the gut microbiota during nutrient deprivation. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106(27):11276–81. doi: [10.1073/pnas.0902366106](https://doi.org/10.1073/pnas.0902366106) PMID: [19549860](https://pubmed.ncbi.nlm.nih.gov/19549860/); PubMed Central PMCID: PMC2700149.

15. Wahli W, Devchand PR, A IJ, Desvergne B. Fatty acids, eicosanoids, and hypolipidemic agents regulate gene expression through direct binding to peroxisome proliferator-activated receptors. *Advances in experimental medicine and biology*. 1999; 447:199–209. PMID: [10086196](#).
16. Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, Wahli W. Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell*. 1992; 68(5):879–87. PMID: [1312391](#).
17. A IJ, Jeannin E, Wahli W, Desvergne B. Polarity and specific sequence requirements of peroxisome proliferator-activated receptor (PPAR)/retinoid X receptor heterodimer binding to DNA. A functional analysis of the malic enzyme gene PPAR response element. *The Journal of biological chemistry*. 1997; 272(32):20108–17. PMID: [9242684](#).
18. Backhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, et al. The gut microbiota as an environmental factor that regulates fat storage. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(44):15718–23. doi: [10.1073/pnas.0407076101](#) PMID: [15505215](#); PubMed Central PMCID: PMC524219.
19. Hou Y, Moreau F, Chadee K. PPAR $\gamma$  is an E3 ligase that induces the degradation of NF $\kappa$ B/p65. *Nature communications*. 2012; 3:1300. doi: [10.1038/ncomms2270](#) PMID: [3250430](#).
20. Kelly D, Campbell JI, King TP, Grant G, Jansson EA, Coutts AG, et al. Commensal anaerobic gut bacteria attenuate inflammation by regulating nuclear-cytoplasmic shuttling of PPAR- $\gamma$  and RelA. *Nature immunology*. 2004; 5(1):104–12. doi: [10.1038/ni1018](#) PMID: [14691478](#).
21. Nakajima T, Nakanishi S, Mason C, Montgomery J, Leggett P, Matsuda M, et al. Population structure and characterization of viridans group streptococci (VGS) isolated from the upper respiratory tract of patients in the community. *The Ulster medical journal*. 2013; 82(3):164–8. PMID: [24505152](#); PubMed Central PMCID: PMC3913407.
22. Pearce C, Bowden GH, Evans M, Fitzsimmons SP, Johnson J, Sheridan MJ, et al. Identification of pioneer viridans streptococci in the oral cavity of human neonates. *Journal of medical microbiology*. 1995; 42(1):67–72. PMID: [7739028](#).
23. Park HK, Shim SS, Kim SY, Park JH, Park SE, Kim HJ, et al. Molecular analysis of colonized bacteria in a human newborn infant gut. *Journal of microbiology*. 2005; 43(4):345–53. PMID: [16145549](#).
24. Arrieta MC, Stiemsma LT, Amenyogbe N, Brown EM, Finlay B. The intestinal microbiome in early life: health and disease. *Frontiers in immunology*. 2014; 5:427. doi: [10.3389/fimmu.2014.00427](#) PMID: [25250028](#); PubMed Central PMCID: PMC4155789.
25. Van den Bogert B, Boekhorst J, Herrmann R, Smid EJ, Zoetendal EG, Kleerebezem M. Comparative genomics analysis of *Streptococcus* isolates from the human small intestine reveals their adaptation to a highly dynamic ecosystem. *PloS one*. 2013; 8(12):e83418. doi: [10.1371/journal.pone.0083418](#) PMID: [24386196](#); PubMed Central PMCID: PMC3875467.
26. van den Bogert B, Erkus O, Boekhorst J, de Goffau M, Smid EJ, Zoetendal EG, et al. Diversity of human small intestinal *Streptococcus* and *Veillonella* populations. *FEMS microbiology ecology*. 2013; 85(2):376–88. doi: [10.1111/1574-6941.12127](#) PMID: [23614882](#).
27. Hakalehto E, Vilpponen-Salmela T, Kinnunen K, von Wright A. Lactic Acid bacteria enriched from human gastric biopsies. *ISRN gastroenterology*. 2011; 2011:109183. doi: [10.5402/2011/109183](#) PMID: [21991494](#); PubMed Central PMCID: PMC3168382.
28. Hayashi H, Takahashi R, Nishi T, Sakamoto M, Benno Y. Molecular analysis of jejunal, ileal, caecal and recto-sigmoidal human colonic microbiota using 16S rRNA gene libraries and terminal restriction fragment length polymorphism. *Journal of medical microbiology*. 2005; 54(Pt 11):1093–101. doi: [10.1099/jmm.0.45935-0](#) PMID: [16192442](#).
29. Frank DN, St Amand AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104(34):13780–5. PMID: [17699621](#).
30. Hong PY, Croix JA, Greenberg E, Gaskins HR, Mackie RI. Pyrosequencing-based analysis of the mucosal microbiota in healthy individuals reveals ubiquitous bacterial groups and micro-heterogeneity. *PloS one*. 2011; 6(9):e25042. PMID: [21966408](#). doi: [10.1371/journal.pone.0025042](#)
31. Zoetendal EG, Raes J, van den Bogert B, Arumugam M, Booiijink CC, Troost FJ, et al. The human small intestinal microbiota is driven by rapid uptake and conversion of simple carbohydrates. *The ISME journal*. 2012; 6(7):1415–26. doi: [10.1038/ismej.2011.212](#) PMID: [22258098](#); PubMed Central PMCID: PMC3379644.
32. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*. 2010; 464(7285):59–65. PMID: [20203603](#). doi: [10.1038/nature08821](#)
33. Kaci G, Goudercourt D, Dennin V, Pot B, Dore J, Ehrlich SD, et al. Anti-inflammatory properties of *Streptococcus salivarius*, a commensal bacterium of the oral cavity and digestive tract. *Applied and*

- environmental microbiology. 2014; 80(3):928–34. doi: [10.1128/AEM.03133-13](https://doi.org/10.1128/AEM.03133-13) PMID: [24271166](https://pubmed.ncbi.nlm.nih.gov/24271166/); PubMed Central PMCID: PMC3911234.
34. Kaci G, Lakhdari O, Dore J, Ehrlich SD, Renault P, Blottiere HM, et al. Inhibition of the NF-kappaB pathway in human intestinal epithelial cells by commensal *Streptococcus salivarius*. *Applied and environmental microbiology*. 2011; 77(13):4681–4. doi: [10.1128/AEM.03021-10](https://doi.org/10.1128/AEM.03021-10) PMID: [21602373](https://pubmed.ncbi.nlm.nih.gov/21602373/); PubMed Central PMCID: PMC3127691.
  35. Kundu P, Ling TW, Korecka A, Li Y, D'Arienzo R, Bunte RM, et al. Absence of intestinal PPARgamma aggravates acute infectious colitis in mice through a lipocalin-2-dependent pathway. *PLoS pathogens*. 2014; 10(1):e1003887. doi: [10.1371/journal.ppat.1003887](https://doi.org/10.1371/journal.ppat.1003887) PMID: [24465207](https://pubmed.ncbi.nlm.nih.gov/24465207/); PubMed Central PMCID: PMC3900641.
  36. Raymond B, Young JC, Pallett M, Endres RG, Clements A, Frankel G. Subversion of trafficking, apoptosis, and innate immunity by type III secretion system effectors. *Trends in microbiology*. 2013; 21(8):430–41. doi: [10.1016/j.tim.2013.06.008](https://doi.org/10.1016/j.tim.2013.06.008) PMID: [23870533](https://pubmed.ncbi.nlm.nih.gov/23870533/).
  37. Xavier MN, Winter MG, Spees AM, den Hartigh AB, Nguyen K, Roux CM, et al. PPARgamma-mediated increase in glucose availability sustains chronic *Brucella abortus* infection in alternatively activated macrophages. *Cell host & microbe*. 2013; 14(2):159–70. doi: [10.1016/j.chom.2013.07.009](https://doi.org/10.1016/j.chom.2013.07.009) PMID: [23954155](https://pubmed.ncbi.nlm.nih.gov/23954155/); PubMed Central PMCID: PMC3777723.
  38. Salcedo SP, Marchesini MI, Lelouard H, Fugier E, Jolly G, Balor S, et al. *Brucella* control of dendritic cell maturation is dependent on the TIR-containing protein Btp1. *PLoS pathogens*. 2008; 4(2):e21. doi: [10.1371/journal.ppat.0040021](https://doi.org/10.1371/journal.ppat.0040021) PMID: [18266466](https://pubmed.ncbi.nlm.nih.gov/18266466/); PubMed Central PMCID: PMC2233671.
  39. Sissler M, Delorme C, Bond J, Ehrlich SD, Renault P, Francklyn C. An aminoacyl-tRNA synthetase paralog with a catalytic role in histidine biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America*. 1999; 96(16):8985–90. PMID: [10430882](https://pubmed.ncbi.nlm.nih.gov/10430882/); PubMed Central PMCID: PMC17719.
  40. Vu-Dac N, Schoonjans K, Kosykh V, Dallongeville J, Fruchart JC, Staels B, et al. Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor. *The Journal of clinical investigation*. 1995; 96(2):741–50. doi: [10.1172/JCI118118](https://doi.org/10.1172/JCI118118) PMID: [7635967](https://pubmed.ncbi.nlm.nih.gov/7635967/); PubMed Central PMCID: PMC185258.
  41. de Wouters T, Ledue F, Nepelska M, Dore J, Blottiere HM, Lapaque N. A robust and adaptable high throughput screening method to study host-microbiota interactions in the human intestine. *PloS one*. 2014; 9(8):e105598. doi: [10.1371/journal.pone.0105598](https://doi.org/10.1371/journal.pone.0105598) PMID: [25141006](https://pubmed.ncbi.nlm.nih.gov/25141006/); PubMed Central PMCID: PMC4139392.
  42. El Aidy S, van Baarlen P, Derrien M, Lindenberg-Kortleve DJ, Hooiveld G, Levenez F, et al. Temporal and spatial interplay of microbiota and intestinal mucosa drive establishment of immune homeostasis in conventionalized mice. *Mucosal Immunol*. 2012; 5(5):567–79. PMID: [22617837](https://pubmed.ncbi.nlm.nih.gov/22617837/). doi: [10.1038/mi.2012.32](https://doi.org/10.1038/mi.2012.32)
  43. El Aidy S, Van den Abbeele P, Van de Wiele T, Louis P, Kleerebezem M. Intestinal colonization: how key microbial players become established in this dynamic process: microbial metabolic activities and the interplay between the host and microbes. *BioEssays: news and reviews in molecular, cellular and developmental biology*. 2013; 35(10):913–23. doi: [10.1002/bies.201300073](https://doi.org/10.1002/bies.201300073) PMID: [23946088](https://pubmed.ncbi.nlm.nih.gov/23946088/).
  44. Wang S, Hibberd ML, Pettersson S, Lee YK. *Enterococcus faecalis* from healthy infants modulates inflammation through MAPK signaling pathways. *PloS one*. 2014; 9(5):e97523. doi: [10.1371/journal.pone.0097523](https://doi.org/10.1371/journal.pone.0097523) PMID: [24830946](https://pubmed.ncbi.nlm.nih.gov/24830946/); PubMed Central PMCID: PMC4022717.
  45. Hansen CH, Andersen LS, Krych L, Metzdorff SB, Hasselby JP, Skov S, et al. Mode of delivery shapes gut colonization pattern and modulates regulatory immunity in mice. *Journal of immunology*. 2014; 193(3):1213–22. doi: [10.4049/jimmunol.1400085](https://doi.org/10.4049/jimmunol.1400085) PMID: [24951818](https://pubmed.ncbi.nlm.nih.gov/24951818/).
  46. Delorme C, Bartholini C, Bolotine A, Ehrlich SD, Renault P. Emergence of a cell wall protease in the *Streptococcus thermophilus* population. *Applied and environmental microbiology*. 2010; 76(2):451–60. doi: [10.1128/AEM.01018-09](https://doi.org/10.1128/AEM.01018-09) PMID: [19915034](https://pubmed.ncbi.nlm.nih.gov/19915034/); PubMed Central PMCID: PMC2805209.
  47. Tezera LB, Hampton J, Jackson SK, Davenport V. *Neisseria lactamica* attenuates TLR-1/2-induced cytokine responses in nasopharyngeal epithelial cells using PPAR-gamma. *Cellular microbiology*. 2011; 13(4):554–68. doi: [10.1111/j.1462-5822.2010.01554.x](https://doi.org/10.1111/j.1462-5822.2010.01554.x) PMID: [21105983](https://pubmed.ncbi.nlm.nih.gov/21105983/).
  48. Mahajan S, Dkhar HK, Chandra V, Dave S, Nanduri R, Janmeja AK, et al. *Mycobacterium tuberculosis* modulates macrophage lipid-sensing nuclear receptors PPARgamma and TR4 for survival. *Journal of immunology*. 2012; 188(11):5593–603. doi: [10.4049/jimmunol.1103038](https://doi.org/10.4049/jimmunol.1103038) PMID: [22544925](https://pubmed.ncbi.nlm.nih.gov/22544925/).
  49. Bassaganya-Riera J, Dominguez-Bello MG, Kronsteiner B, Carbo A, Lu P, Viladomiu M, et al. *Helicobacter pylori* colonization ameliorates glucose homeostasis in mice through a PPAR gamma-dependent mechanism. *PloS one*. 2012; 7(11):e50069. doi: [10.1371/journal.pone.0050069](https://doi.org/10.1371/journal.pone.0050069) PMID: [23166823](https://pubmed.ncbi.nlm.nih.gov/23166823/); PubMed Central PMCID: PMC3499487.



50. van Beekum O, Fleskens V, Kalkhoven E. Posttranslational modifications of PPAR-gamma: fine-tuning the metabolic master regulator. *Obesity*. 2009; 17(2):213–9. doi: [10.1038/oby.2008.473](https://doi.org/10.1038/oby.2008.473) PMID: [19169221](https://pubmed.ncbi.nlm.nih.gov/19169221/).
51. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature*. 1998; 391(6662):79–82. doi: [10.1038/34178](https://doi.org/10.1038/34178) PMID: [9422508](https://pubmed.ncbi.nlm.nih.gov/9422508/).
52. Subbaramaiah K, Lin DT, Hart JC, Dannenberg AJ. Peroxisome proliferator-activated receptor gamma ligands suppress the transcriptional activation of cyclooxygenase-2. Evidence for involvement of activator protein-1 and CREB-binding protein/p300. *The Journal of biological chemistry*. 2001; 276(15):12440–8. doi: [10.1074/jbc.M007237200](https://doi.org/10.1074/jbc.M007237200) PMID: [11278336](https://pubmed.ncbi.nlm.nih.gov/11278336/).
53. Cippolletta D, Feuerer M, Li A, Kamei N, Lee J, Shoelson SE, et al. PPAR-gamma is a major driver of the accumulation and phenotype of adipose tissue Treg cells. *Nature*. 2012; 486(7404):549–53. doi: [10.1038/nature11132](https://doi.org/10.1038/nature11132) PMID: [22722857](https://pubmed.ncbi.nlm.nih.gov/22722857/); PubMed Central PMCID: [PMC3387339](https://pubmed.ncbi.nlm.nih.gov/PMC3387339/).
54. Berger JP. Role of PPARgamma, transcriptional cofactors, and adiponectin in the regulation of nutrient metabolism, adipogenesis and insulin action: view from the chair. *International journal of obesity*. 2005; 29 Suppl 1:S3–4. doi: [10.1038/sj.ijo.0802904](https://doi.org/10.1038/sj.ijo.0802904) PMID: [15711580](https://pubmed.ncbi.nlm.nih.gov/15711580/).
55. Atshaves BP, Martin GG, Hostetler HA, McIntosh AL, Kier AB, Schroeder F. Liver fatty acid-binding protein and obesity. *The Journal of nutritional biochemistry*. 2010; 21(11):1015–32. doi: [10.1016/j.jnutbio.2010.01.005](https://doi.org/10.1016/j.jnutbio.2010.01.005) PMID: [20537520](https://pubmed.ncbi.nlm.nih.gov/20537520/); PubMed Central PMCID: [PMC2939181](https://pubmed.ncbi.nlm.nih.gov/PMC2939181/).
56. Robciuc MR, Naukkarinen J, Ortega-Alonso A, Tyynismaa H, Raivio T, Rissanen A, et al. Serum angiopoietin-like 4 protein levels and expression in adipose tissue are inversely correlated with obesity in monozygotic twins. *Journal of lipid research*. 2011; 52(8):1575–82. doi: [10.1194/jlr.P015867](https://doi.org/10.1194/jlr.P015867) PMID: [21596930](https://pubmed.ncbi.nlm.nih.gov/21596930/); PubMed Central PMCID: [PMC3137024](https://pubmed.ncbi.nlm.nih.gov/PMC3137024/).
57. Aronsson L, Huang Y, Parini P, Korach-Andre M, Hakansson J, Gustafsson JA, et al. Decreased fat storage by *Lactobacillus paracasei* is associated with increased levels of angiopoietin-like 4 protein (ANGPTL4). *PLoS one*. 2010; 5(9). doi: [10.1371/journal.pone.0013087](https://doi.org/10.1371/journal.pone.0013087) PMID: [20927337](https://pubmed.ncbi.nlm.nih.gov/20927337/); PubMed Central PMCID: [PMC2948012](https://pubmed.ncbi.nlm.nih.gov/PMC2948012/).