

Can ganglion cell complex assessment on cirrus HD OCT aid in detection of early glaucoma?



Avadhesh Oli*; D. Joshi

Abstract

Context: Ganglion cell complex is damaged early in glaucoma. Does this loss of GCC help in early diagnosis of glaucoma.

Aims: To compare the RNFL thickness and ganglion cell complex (GCC) in diagnosed patients of glaucoma, pre-perimetric glaucoma and normal controls.

Settings and design: Case controlled, observational study.

Methods and material: 33 glaucoma patients, 45 pre-perimetric glaucoma, and 30 controls were enrolled in the study. ONH parameters on cirrus HD OCT like CD ratio, para papillary RNFL thickness and GCC were calculated for each case.

Statistical analysis used: ANOVA test to analyse differences between groups. ROC for ganglion cell layer.

Results: RNFL thickness was 71.6 μ and GCC was 69.19 μ in glaucoma patients. RNFL thickness was 77.31 μ and GCC was 71 μ in pre-perimetric glaucoma and 99.6 μ and 85.16 μ in controls respectively. The difference of mean for RNFL and GCC by ANOVA was statistically significant for controls, glaucoma patients and pre-perimetric glaucoma patients. RNFL ($p < 0.001$) and GCC ($p < 0.001$). Receiver operating characteristic curve for GCC was 0.83 ($p < 0.000$).

Conclusions: The RNFL analysis is increasingly being used as newer tool in diagnosis of glaucoma. In addition, GCC can be used as a supplementary tool in picking up cases of pre-perimetric glaucoma as loss is significant in pre-perimetric glaucoma also.

Keywords: Ganglion cell complex, Optical coherence tomography, Retinal nerve fibre layer, Glaucoma

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of Saudi Ophthalmological Society, King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
<http://dx.doi.org/10.1016/j.sjopt.2015.02.007>

Introduction

Glaucoma is a chronic progressive optic neuropathy with high intra-ocular pressure (IOP) as one of the important risk factors. Apart from IOP, vascular insufficiency and other cellular factors have a definite role to play. The final common pathway is cupping of optic nerve head due to death of ganglion cell axons. The death of axons is associated and may be by ganglion cell death. The characteristic changes of the nerve head (ONH) and visual field defects on perimetry confirm the diagnosis of glaucoma. The loss of retinal ganglion cells is considered as an important step in pathogenesis in

cascade of events leading to retinal nerve fibre (RNFL) loss and changes of optic nerve head. Various modalities for diagnosis of glaucoma have been used but the search for the ideal modality for early diagnosis remains elusive.¹ Newer perimetric methods were launched with great enthusiasm but they seem to have a limited role to play. For visual field defects to be evident on white on white perimetry, it requires at least 30–40% of neuronal cell loss to occur which is irreversible. The problem is confounded by the fact that the patient is also not aware of the problem.¹

Use of Optical Coherence Tomography (OCT) in the diagnosis and follow up of glaucoma has gained universal

Received 26 June 2014; received in revised form 2 February 2015; accepted 25 February 2015; available online 23 March 2015.

Eye Department Command Hospital, Bangalore 560007, India

* Corresponding author. Tel.: +91 9686634002.
e-mail address: olieve@rediffmail.com (A. Oli).



Peer review under responsibility of Saudi Ophthalmological Society, King Saud University



Production and hosting by Elsevier

Access this article online:
www.saudiophthaljournal.com
www.sciencedirect.com

acceptability in recent years. It can give very accurate parameters for optic nerve head assessment along with the retinal nerve fibre layer analysis.² Even so, early diagnosis remains a challenge and requires correlation with other factors. Recent studies on the use of OCT in glaucoma have explored the ability of this technology to detect the loss of ganglion cell complex (GCC). The ganglion cell complex is studied by the OCT which has the maximum density at the macula. Roughly, one lac ganglion cells are found in the human retina and half of this number is centred on the fovea. This anatomic arrangement suggests that a macula scan with GCC analysis could be early indicator of this disease.³

This prospective observational study was designed to study the role of GCC in early diagnosis of glaucoma.

Materials and methods

The Study was conducted at a tertiary care teaching centre between Aug 2012 and Jul 2013 on Asian patients. Institutional ethical clearance was taken and tenants of declaration of Helensiki were adhered to. Participants were given detailed information and consent was taken. IOP was corrected to central corneal thickness wherever required (Average corneal thickness was taken as 530 μm . For pachymetry values above this one mm Hg was reduced for every 14 μm and one mm Hg was added for every 14 μm for values lower than normal). IOP was recorded after wash off period of the drug wherever indicated: 45 pre-perimetric glaucoma patients (IOP > 21 mmHg, with optic nerve head changes or RNFL defects and normal Visual fields) and 33 glaucoma patients (IOP > 21 mmHg, with optic nerve head changes or RNFL defects correlating with visual field changes). 30 normal controls with IOP < 21 mmHg without evidence of glaucomatous optic neuropathy and RNFL defects with negative family history of glaucoma were enrolled in the study. Patients with poor vision and media opacity which hampered OCT examination were excluded. Patients with other retinal or macular pathologies like diabetic retinopathy, retinal vein occlusion, high refractive errors (> +/−6 DS) were excluded from the study. Cases of angle closure glaucoma, traumatic glaucoma and other secondary glaucoma were excluded from study.

All patients were subjected to comprehensive ophthalmic evaluation which included visual acuity, anterior segment examination including pupillary reaction, anterior chamber angle assessment by gonioscopy, intra-ocular pressure measurement by Goldmann applanation tonometry, slit lamp

biomicroscopy, pachymetry, visual field testing (Oculus twin field) and Optical Coherence Tomography (Zeiss cirrus HD OCT). Visual field testing was done within three months of OCT. Second reliable field was analysed as per Anderson's criteria for all patients.

Single well trained examiner did OCT after pupillary dilatation. Optic nerve head scans were acquired with 4 mm concentric maps. The GCC maps were based on macular protocol centred on fovea with a cube of 512 \times 128 with automated measurement of GCC and internal limiting membrane. OCT scans with signal strength more than 6 were included for analysis.

ONH parameters on cirrus HD OCT like CD ratio, para papillary RNFL thickness and GCC were calculated for each case. One eye of each patient was randomized to be part of analysis.

The results were collected on Microsoft excel (Office 2013) and analysed using SPSS (Version 17 IBM NY) software using ANOVA. The means were compared and a p value < 0.05 was considered significant. Area under the curve for GCC was also calculated.

Results

Results: A total 108 patients were enrolled in the study. The mean age of glaucoma patients was 58.54 (SD 13.78) years, pre-perimetric glaucoma was 53.71 (SD 17.553) years and controls was 52 (SD 14.88). In the glaucoma group, 54.54% were males and 45.45% were females. In the pre-perimetric glaucoma group, 46.66% were males and 53.33% were females and in control group 53.33% were males and 46.66% were females.

The data showed a normal distribution. The difference of mean for RNFL and GCC by ANOVA was statistically significant for controls, glaucoma patients and pre-perimetric glaucoma group RNFL ($p < 0.001$) and GCC ($p < 0.001$). When comparing the difference of means of GCC and RNFL with normal values, it was found to be statistically significant ($p < 0.001$) and GCC ($p < 0.001$) (Table 1).

Pearson correlation coefficient for ganglion cell layer had a statistically significant correlation ($r = 0.455$) with the RNFL thickness ($p < 0.05$).

Receiver operating characteristics curves for GCC were drawn as shown in Fig 1. The area under the curve (AUC) 0.835 with 95% Confidence interval 0.751–0.899 which was statistically significant ($p < 0.0001$)

Table 1. Comparative OCT data for study group.

Diag		CDR	RNFL	GCC
Control	Mean	.4907	99.6000	85.1667
	N	30	30	30
	Std. deviation	.11380	10.32773	9.07092
	95% CONF interval	0.44–0.53	95.74–103.45	81.77–88.55
Poag	Mean	.6876	71.6364	69.1970
	N	33	33	33
	Std. deviation	.12150	20.15110	12.72113
	95% CONF Interval	0.64–0.73	64.49–78.78	64.68–73.70
Pre-perimetric	Mean	.6716	77.3111	71.8556
	N	45	45	45
	Std. deviation	.091870	17.07313	10.70929
	95% CONF Interval	.64–0.69	72.18–82.44	68.63–75.07

Comparative data.




Figure 1. Receiver operating characteristic curve.

Discussion

Glaucoma is a progressive optic nerve disorder characterized by loss of retinal axons. Early diagnosis of this disorder is a challenge because the visual field changes which are considered a gold standard for diagnosis do not show changes until significant neuronal loss occurs.

Due to advancement in technology, various non-perimetric modalities for diagnosis of glaucoma are evolving. Optical coherence tomography performs high resolution, cross sectional imaging of optic nerve head, RNFL and macula. A retinal thickness analysis can be obtained from this data. This has led to better understanding and early detection of glaucomatous changes.² The testing is rapid, non-invasive, patient and examiner friendly.⁴ In the present study we found that values of RNFL and GCC were lower both in glaucoma and pre-perimetric glaucoma patients indicating its utility in early diagnosis of glaucoma. Experimental animal models have shown that GCC loss correlates well with visual field defects.³ Studies have mentioned utility of electroretinogram and ganglion cell axons imaging in early diagnosis of glaucoma.⁵ In addition GCC can be used as a supplementary tool in picking up cases of pre-perimetric glaucoma as Ganglion Cell Complex loss is significant in pre-perimetric glaucoma as well.⁶ In glaucoma GCC is first to be affected so evaluation of GCC is early indicator of damage. Central macular thickness has been used by some authors for early diagnosis of glaucoma but with a limited role.⁷ Central macular thickness, though very commonly performed OCT test may not have a role in early glaucoma diagnosis. In some studies, RNFL was found to have good discrimination power in detection of glaucoma. GCC had also shown to be equally good predictor of damage due to glaucoma.⁸ Some of the studies have tried to differentiate the type of glaucoma based on the nature of GCC loss but other studies have failed to establish such a difference.⁹ Repeated RNFL measurements have been used for detection of glaucoma progression over a period of time.¹⁰ Similar periodic GCC measurements can be utilized to detect progression of glaucoma in very nascent stage.¹¹

The limitations of this method include lack of wide normative database across different ethnic groups. Large population based studies are required to be done. Limited availability of equipment in African and Asian countries could be another limitation.

Some authors evaluating role of GCC have found the role of outer retinal thickness in GCC measurement.¹² They proposed GCC to outer retinal ratio as a better parameter to compensate for individual variation in outer retinal thickness. This study concluded that GCC to total retinal thickness (G/T) ratio is a complimentary indicator for diagnosing glaucoma in early stages and was found to have better sensitivity than RNFL thickness alone.

Mapping of GCC based on residual cells without averaging the total GCC thickness used by some authors has shown to be another early indicator of glaucoma. Minimum GCC on 360° spoke showed a better ability to differentiate between healthy and glaucomatous eyes.¹³

The ability of GCC as compared to RNFL to detect early glaucoma is found to be slightly better in some studies.¹⁴ Addition of GCC data to RNFL data has been found to enhance the detection of glaucoma in both pre-perimetric and perimetric groups.¹⁵

There are few limitations of the present study. Firstly this study did not include long term follow up of patients so further studies are required to study the role of serial GCC scans in glaucoma progression. Secondly, we also could not compare the visual field loss severity indicators like mean deviation (MD) with GCC or RNFL loss which could have been a better parameter to study in advanced cases of glaucoma because this was beyond the scope of this study, as our main aim was to study role of GCC in early diagnosis in glaucoma.

In conclusion, our study has shown that measurement of GCC is a newer tool in early diagnosis of glaucoma. It is more useful in pre-perimetric glaucoma where in conjunction with RNFL analysis it can pick-up pre-perimetric glaucoma cases.

Conflict of interest

The authors declared that there is no conflict of interest.

References

- Pagliara MM, Lepore D, Balestrazzi E. The role of OCT in glaucoma management. *Prog Brain Res* 2008;**173**:139–48.
- Kader M, Ramakrishnan R, Mittal S, Ambatkar S. Retinal nerve fibre layer thickness measurements in normal Indian population by optical coherence tomography. *Indian J Ophthalmol* 2006;**54**:11–5.
- Harwerth RSC-DL, Shen F, Smith 3rd EL, Crawford ML. Ganglion cell losses underlying visual field defects from experimental glaucoma. *Invest Ophthalmol Vis Sci* 1999;**40**:2242–50.
- Mwanza JCOJ, Budenz DL, Chang RT, Knight OJ, Feuer WJ. Macular ganglion cell-inner plexiform layer: automated detection and thickness reproducibility with spectral domain-optical coherence tomography in glaucoma. *Invest Ophthalmol Vis Sci* 2011;**52**:8323–9.
- Falsini B, Marangoni D, Salgarello T. Structure-function relationship in ocular hypertension and glaucoma: interindividual and interocular analysis by OCT and pattern ERG. *Graefes Arch Clin Exp Ophthalmol* 2008;**246**:1153–62.
- le Pv TO, Chopra V, Francis BA, Ragab O, Varma R, Huang D. Regional correlation among ganglion cell complex, nerve fiber layer, and visual field loss in glaucoma. *Invest Ophthalmol Vis Sci* 2013;**54**:4287–95.
- Barisic F, Sicaja AJ, Ravlic MM, Novak-Laus K, Ivekovic R, Mandic Z. Macular thickness and volume parameters measured using optical coherence tomography (OCT) for evaluation of glaucoma patients. *Coll Antropol* 2012;**36**:441–5.
- Firat PG, Doganay S, Demirel EE, Colak C. Comparison of ganglion cell and retinal nerve fiber layer thickness in primary open-angle glaucoma and normal tension glaucoma with spectral-domain OCT. *Graefes Arch Clin Exp Ophthalmol* 2013;**251**:831–8.
- Kim NR, Hong S, Kim JH, Rho SS, Seong GJ, Kim CY. Comparison of macular ganglion cell complex thickness by fourier-domain oct in

- normal tension glaucoma and primary open-angle glaucoma. *J Glaucoma* 2013;**22**:133–9.
10. Lee EJ, Kim TW, Park KH, Seong M, Kim H, Kim DM. Ability of stratus OCT to detect progressive retinal nerve fiber layer atrophy in glaucoma. *Invest Ophthalmol Vis Sci* 2009;**50**:662–8.
 11. Moura AL, Raza AS, Lazow MA, De Moraes CG, Hood DC. Retinal ganglion cell and inner plexiform layer thickness measurements in regions of severe visual field sensitivity loss in patients with glaucoma. *Eye (Lond)* 2012;**26**:1188–93.
 12. Kita Y, Kita R, Takeyama A, Takagi S, Nishimura C, Tomita G. Ability of optical coherence tomography-determined ganglion cell complex thickness to total retinal thickness ratio to diagnose glaucoma. *J Glaucoma* 2012 E pub ahead of print.
 13. Kohei Takayama MH, Mary Durbin, Noriko Nakano, Satoshi Morooka, Tadamichi Akagi, Hanako Ohashi Ikeda, et al. A novel method to detect local ganglion cell loss in early glaucoma using spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci* 2012;**53**:6904–13.
 14. Azusa Akashi AK, Makoto Nakamura, Masashi Fujihara, Yuko Negi, Akira Negi. Comparative assessment for the ability of cirrus, RTVue and 3D OCT to diagnose glaucoma. *Invest Ophthalmol Vis Sci* 2013;**54**:4478–84.
 15. Sung MS, Yoon JH, Park SW. Diagnostic validity of macular ganglion cell-inner plexiform layer thickness deviation map algorithm using cirrus HD-OCT in preperimetric and early glaucoma. *J Glaucoma* 2014;**23**(8):e144–51.