

SCIENTIFIC REPORTS

OPEN

Author Correction: A rapid and nondestructive protocol for whole-mount bone staining of small fish and *Xenopus*

Hiromi Sakata-Haga¹, Maimi Uchishiba², Hiroki Shimada¹, Tsuyoshi Tsukada¹, Mayumi Mitani¹, Tomohiro Arikawa³, Hiroki Shoji³ & Toshihisa Hatta¹

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-018-25836-4>, published online 10 May 2018

In this Article, the legend of Figure 3 is incorrect:

“A flowchart of the rapid (RAP) system for bone staining (RAP-B). The new procedure for bone staining which is rapid, nondestructive, and allowed to obtain high-definition, fine bone-stained specimens, was based on our RAP system. Cartilage staining (RAP-C) can be applied as single staining or optionally added before the bone staining for double staining of bone and cartilage (RAP-B/C, Supplementary Fig. S8). FIXATIVE: 5% formalin, 5% Triton X-100, 1% potassium hydroxide (KOH); B-STAINING SOLUTION: 0.05% alizarin red S, 20% ethylene glycol, 1% KOH; C-STAINING SOLUTION: 50–70% ethanol, 20% acetate, 0.015–0.02% alcian blue; CLEARING SOLUTION: 20% Tween 20, 1% KOH; ENHANCEMENT SOLUTION: 20% ethylene glycol, 5% Triton X-100, 1% KOH.”

should read:

“A flowchart of the rapid (RAP) system for bone staining (RAP-B). The new procedure for bone staining which is rapid, nondestructive, and allowed to obtain high-definition, fine bone-stained specimens, was based on our RAP system. Cartilage staining (RAP-C) can be applied as single staining or optionally added before the bone staining for double staining of bone and cartilage (RAP-B/C, Supplementary Fig. S8). FIXATIVE: 5% formalin, 5% Triton X-100, 1% potassium hydroxide (KOH); B-STAINING SOLUTION: 0.05% alizarin red S, 20% ethylene glycol, 1% KOH; C-STAINING SOLUTION: 50–70% ethanol, 20% acetic acid, 0.015–0.02% alcian blue; CLEARING SOLUTION: 20% Tween 20, 1% KOH; ENHANCEMENT SOLUTION: 20% ethylene glycol, 5% Triton X-100, 1% KOH.”



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018

¹Department of Anatomy, Kanazawa Medical University, Ishikawa, Japan. ²Department of Obstetrics and Gynecology, Wakayama Medical University, Wakayama, Japan. ³Department of Biology, Kanazawa Medical University, Ishikawa, Japan. Correspondence and requests for materials should be addressed to T.H. (email: thatta@kanazawa-med.ac.jp)