



# Recent progress in radioactive seed implantation brachytherapy of non-small cell lung cancer: a narrative review

Xinyu Wang<sup>1#</sup>, Sen Tian<sup>1,2#</sup>, Hui Shi<sup>1</sup>, Hao Qin<sup>1</sup>, Wei Zhang<sup>1</sup>, Yuchao Dong<sup>1</sup>, Chong Bai<sup>1</sup>

<sup>1</sup>Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Naval Medical University, Shanghai, China; <sup>2</sup>Department of Respiratory and Critical Care Medicine, No. 906 Hospital of the Chinese People's Liberation Army Joint Logistic Support Force, Ningbo, China

**Contributions:** (I) Conception and design: C Bai; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: C Bai; (V) Data analysis and interpretation: S Tian, X Wang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

<sup>#</sup>These authors contributed equally to this work.

**Correspondence to:** Chong Bai, MD, FCCP. Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Naval Medical University, No. 168 Changhai Road, Yangpu District, Shanghai 200433, China. Email: Chongbai@smmu.edu.cn.

**Background and Objective:** Brachytherapy, a new form of radiation therapy, has been used to treat lung cancer and consists of two main forms of treatment: endobronchial brachytherapy and radioactive seed implantation brachytherapy (RSI-BT), the latter of which is used to treat non-small cell lung cancer (NSCLC). The use of RSI-BT in the treatment of NSCLC at our centre has yielded some positive results.

**Methods:** To more fully consider the context of this application, we conducted a search of PubMed from 2018 to March 5, 2023. The search included a combination of the MeSH terms: “brachytherapy” and “lung neoplasm”.

**Key Content and Findings:** The majority of NSCLC patients who received RSI-BT achieved positive benefits. Most patients had a progression-free survival (PFS) of between 12 and 18 months. Additionally, radioactive particle stent implantation as a specific RSI-BT has shown therapeutic potential in the treatment of malignant airway obstruction. With the application of new technologies, RSI-BT will become more precise, efficient and inexpensive.

**Conclusions:** This review demonstrates that RSI-BT can be therapeutic in the treatment of both early and advanced NSCLC with manageable complications. There have also been reports on the combination of RSI-BT with other therapies, but more research is needed on the combination of RSI-BT with them.

**Keywords:** Brachytherapy; non-small cell lung cancer (NSCLC); radioactive seed implantation; combination; intervention

Submitted Oct 17, 2023. Accepted for publication Jan 18, 2024. Published online Mar 25, 2024.

doi: 10.21037/jtd-23-1600

**View this article at:** <https://dx.doi.org/10.21037/jtd-23-1600>

## Introduction

Lung cancer is the malignancy with the highest mortality rate in the male population, and approximately 85% of lung cancer patients have non-small cell lung cancer (NSCLC) (1). For such patients, radiotherapy is usually used as an important alternative treatment method (2). External beam radiotherapy (EBRT), as a conventional radiation therapy, has been one of the optional therapies

for the treatment of NSCLC. Although EBRT has some advantages, such as mature technology, and more convenient. However, due to the limitation of therapeutic mechanism, EBRT needs to pass the accelerated photon or electron beam through normal tissues before it can reach the tumour tissues. On the one hand, this causes additional damage to normal tissues; on the other hand, in order to achieve the therapeutic dose, EBRT needs to increase the irradiation dose, which in turn exacerbates the

**Table 1** Clinical studies related to radioactive seed implantation brachytherapy in the last 5 years

Therapy	NSCLC stages	Year	Content	Outcome (positive & negative)
RSI-BT alone	Early-stage	2020	Safety and efficiency	11 CR, 9 PR, 2 PD cases, OS of 11.7±7.6 months (7) 3 CR, 1 PR and 1 PD cases (8)
		2019	Retrospective clinical study	Get better ORR, slight fever (9). Inhibit local tumour growth 4 cases of pneumothorax, 3 cases of hemoptysis, and 2 cases of particle displacement (10)
		2020	RSI-BT vs. microwave ablation	Longer OS and DFS (11)
		2021	Malignant airway compression	Relieves compression, include radiation-related pulmonary reactions 13.33%; skin reactions 13.33%; radiation-related esophageal reactions 6.67%, and leukopenia 10% (12)
		2022	Safety and efficiency	Better LCR, acute grade 1–2 toxicity was identified in 4 patients (13)
		2023	A randomized trial, a phase 2 clinical trial	Safe and effective, distant disease, corresponding to 40% of patients and 65.2% of all failures. 2.7% patients developed grade 1 acute pneumonitis. Grade 2 and 3 acute esophagitis occurred in 14.7% and 5.3% patients, respectively (14,15)
RSI-BT + EBRT	Advanced	2018	RSI-BT vs. EBRT	Higher quality of life scores, no significant differences were observed for complications (16)
	Early-stage	2020	RSI-BT vs. SART	Higher LCR and 1-year survival rate, the indicator of radioactive pneumonia V20 and V30 in Groups A and B was 6.06% and 4.207%, and 11.32% and 7.111%, respectively (17)
RSI-BT + chemotherapy	Advanced	2020	RSI-BT vs. second-line chemotherapy	Higher DCR, the incidence of hematologic toxicity in group A was significantly lower than that in group B (18)
		2021	RSI-BT + BACE	Better LRR, no serious complications occurred in the two groups (19)
		2022		Longer PFS and OS, the main adverse events included fever (100%), pain (65.2%), liver function impairment (65.2%), fatigue (56.5%), and nausea and vomiting (52.2%) (20)
RSI-BT + targeted therapy	Advanced	2021	RSI-BT + EGFR-TKIs	Higher ORR, DCR and 2-year OS rate longer PFS (21)

NSCLC, non-small cell lung cancer; RSI-BT, radioactive seed implantation brachytherapy; CR, complete response; PR, partial response; PD, progressive disease; OS, overall survival; ORR, objective response rate; DFS, disease-free survival; LCR, local control rate; EBRT, external beam radiotherapy; SART, stereotactic ablative radiotherapy; DCR, disease control rate; BACE, bronchial artery chemoembolization; LRR, local response rate; PFS, progression-free survival; EGFR, epidermal growth factor receptor; TKI, tyrosine kinase inhibitor.

damage to the tissues in the irradiation path (3). However, brachytherapy has some advantages in the above problems. Since the irradiation comes from inside the tumour, the irradiation to the surrounding tissues can be effectively reduced. The efficacy of brachytherapy depends on the direct irradiation of tumours with high doses of radiation from a radiation source. Brachytherapy has a considerable dose-dependent advantage of radiation dose gradients over conventional external irradiation techniques. In fact, if the tumour moves during radiation, the radiation source also

moves, because the radiation source is inside the tumour. This property differs from conventional EBRT, which requires an additional margin to account for setup and organ motion uncertainties (4–6). In this review, we describe a list based on the current literature and present promising clinical data regarding radioactive seed implantation brachytherapy (RSI-BT) in NSCLC treatment (*Table 1*). We present this article in accordance with the Narrative Review reporting checklist (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1600/rc>).

**Table 2** Brachytherapy and lung neoplasms literature research

Items	Specification
Date of search	03/02/2023 to 03/05/2023
Databases searched	PubMed
Search terms used	Search terms: “Brachytherapy [MeSH]” and “Lung Neoplasms [MeSH]”. Filters: English
Timeframe	2018 to 2023
Inclusion and exclusion criteria	Inclusion criteria: experimental design; discussed brachytherapy (mainly radioactive seed implantation brachytherapy) and its application in lung neoplasms (mainly non-small cell lung cancer) treatment; mentioned the effects of therapeutics on early-stage and advanced non-small cell lung cancer  Exclusion criteria: literature reviews; case studies; only mentioned brachytherapy and lung neoplasms without description of specific therapeutic effects
Selection process	The selection process was conducted independently by the authors and was approved by a board-certified pulmonologist. The search produced 53 articles that were examined according to the inclusion and exclusion criteria; this reduced the total to 15 articles. Then, we narrowed the list to those that discussed the specific radioactive seed implantation brachytherapy

MeSH, Medical Subject Headings.

## Methods

The PubMed database was initially searched on March 02, 2023. *Table 2* highlights the search conducted using the combination of the Medical Subject Headings (MeSH) terms: “Brachytherapy”, and “Lung Neoplasms”. And we integrated clinical studies including “radioactive seed implantation brachytherapy” to make it more accurate.

### Technical background

The mechanisms of RSI-BT for cancer lesions include direct irradiation to promote apoptosis of cancer cells and inhibition of vascular growth in cancerous tissues (22-24). DNA strands are the primary target of therapeutic irradiation, and radiation-induced DNA damage can lead directly to cell death, cell cycle redistribution and microenvironmental changes without reparations. The therapeutic index of radiation reflects the differential response between tumour and normal tissues because tumour cells have a lower capacity for DNA repair than normal tissue cells. The ideal technique should be able to deliver a therapeutic dose to the tumour while minimizing the dose to the organs at risk (25).

Iridium-192 (Ir-192), cobalt-60 (Co60), iodine-125 (I-125) and palladium-103 (Pd-103) are most commonly used in modern brachytherapy (26,27). I-125, whose efficacy and safety in RSI-BT have been established, is the principal particle utilized in clinical practice (4). Especially for

elderly NSCLC patients who have lost the chance for surgery and are in poor physical condition, brachytherapy can control tumour progression and improve quality of life.

The total dose can be obtained by (I) continuous low-dose-rate (LDR) irradiation; (II) repeating low-intensity pulses hourly for several days (pulsed-dose-rate irradiation); or (III) delivering several portions of a high dose at a time [high-dose-rate (HDR) irradiation] (4,26). Radioactive seed implantation brachytherapy is commonly used in LDR brachytherapy (LDR-BT), while HDR brachytherapy (HDR-BT) and pulsed-dose-rate brachytherapy (PDR-BT) have shown advantages in the treatment of internal bronchial lesions (4). As an example, Ir-192, which is commonly used as a source for HDR-BT, typically delivers at a rate of  $\geq 12$  Gy/h, compared to 0.4–2.0 Gy/h of LDR-BT. The linear quadratic (LQ) equation can be used to calculate the total biologically effective dose (BED), which can be used to assess the frequency that HDR-BT/LDR-BT is received (28). At the same time, more precise treatment planning systems (TPSs) and the use of high-resolution computed tomography (CT) have made dose control easier.

The standard treatment planning procedure for 3D volumetric image-based brachytherapy involves three main steps: (I) anatomic contouring, (II) applicator digitization, and (III) dosimetric planning and optimization, using a dedicated TPS (29). Steps (I) and (II) would be accomplished by using TPS, but step (III) usually needs



**Figure 1** A 38-year-old female with right middle lung adenocarcinoma T1bN0M0 stage IA who was treated with I-125 radioactive particle placement in the First Affiliated Hospital of Naval Medical University. (A) Before brachytherapy; (B) six months after brachytherapy; (C) two years after brachytherapy.

more BT teams' effort and time. Accurate delineation of the clinical target volume (CTV) and organ at risk (OAR) is a crucial step in radiotherapy treatment planning (12,30). However, it is still a labor-intensive process to manually contour the CTV and OARs in a 3D image volume with some guidelines exist to help define the contours. Deep-learning-based image processing has demonstrated tremendous potential to help this (31). Additionally, dose calculations based on the AAPM Task Group 43 formalism are currently the mainstream (32). Some model-based calculation methods are increasingly being used to improve efficiency, such as Monte Carlo simulation or solving the Boltzmann transport equation (33).

In recent years, 3D noncoplanar templates, polylactic acid templates, and automatic needles have gradually been applied in clinical treatment (34-36). They enhance RSI-BT in numerous ways, including particle selection, needle insertion, template selection and design, making brachytherapy more precise and less expensive. A study found that particle implantation using a 3D noncoplanar template requires fewer probes, less rib injury, and a higher dose in the target organ region than a conventional 3D coplanar template to achieve the same therapeutic dose, making RSI-BT for NSCLC more individualized and precise (34). Additionally, polylactic acid-based templates can reduce the cost of 3D noncoplanar templates, making RSI-BT cheaper (35). Automated probes have effectively posed a challenge to conventional manual probes in terms of probe implantation, and investigations have revealed that the average misses in probe angle and depth are  $2^\circ$  and 1 mm, respectively, and the average segmentation time per needle is 0.238 s, which is already useful (36).

## Clinical applications

### RSI-BT alone for NSCLC

#### Early-stage NSCLC

The location of the tumour lesion, the tumour's clinical stage, and other factors influence the specific strategy used for RSI-BT of NSCLC. RSI-BT can be effective under a variety of circumstances.

Zhao *et al.* demonstrated that CT-guided brachytherapy with I-125 implantation improved local control rates, decreased side effects, and improved patient quality of life. Of the 26 patients, 11 patients had complete response (CR) and 9 patients had partial response (PR), with an overall survival (OS) of  $11.7 \pm 7.6$  months (7). A study including six patients also demonstrated the safety and efficacy of CT-guided I-125 implantation for BT, with only one patient progressing in the 12-month postoperative follow-up (8). For patients with early-stage NSCLC who refuse to undergo surgery, the disease can be completely controlled or even eradicated by RSI-BT (Figure 1). However, RSI-BT was not linked to a lower risk of local or distant metastases or better outcomes in patients with clinical stage I NSCLC, according to a meta-analysis by Chen *et al.* (37). Regardless, it may reduce local recurrence (LR) when the BED is  $>100$  Gy. In addition, a study comparing the effectiveness of brachytherapy and external beam therapy in 543 patients with limited resection for NSCLC found that while survival rates for patients with lung cancer with T1-T4N0M0 were comparable in the groups receiving RSI-BT and EBRT (compared to EBRT, brachytherapy was associated with a decreased risk of death for both OS and disease-specific survival (DSS) [hazard ratio (HR) 0.604; 95% confidence interval (CI): 0.380-0.961 and



**Figure 2** Implantation of radioactive particles by bronchoscope for central lung cancer in the First Affiliated Hospital of Naval Medical University. (A) Before brachytherapy; (B) one month after brachytherapy; (C) three months after brachytherapy.

HR 0.524; 95% CI: 0.303–0.908, respectively), patients with T1 had higher survival rates after receiving brachytherapy (38). In conclusion, RSI-BT for early-stage NSCLC has been shown to be safe and therapeutically effective.

### Advanced NSCLC

In advanced lung cancer, the therapeutic effect of RSI-BT is more limited and is more about relieving cancer symptoms and improving patients' quality of life. The first topic covered is the function of CT-guided RSI-BT in the management of advanced NSCLC. I-125 RSI-BT alone significantly improved clinical outcomes and decreased the incidence of myelosuppression compared to chemotherapy, according to a meta-analysis by Zhang *et al.* (39). RSI-BT, however, might cause lung damage. To support the combined findings of complications, larger samples and higher quality randomized controlled studies are needed. The majority of studies on RSI-BT have demonstrated that the method is safe and reliable with controllable side effects and less harm to nearby organs while having the same or better local control rate, lower resource cost, and far lower overall radiation dose compared to conventional local treatment (10,13,24,40–45). However, the manageability of the difficulties associated with RSI-BT and issues with dose management have also been brought up by scholars in several studies (13,41,42). Moreover, advanced lung cancer-related airway constriction can be treated using bronchoscope-guided I-125 RSI-BT (Figure 2), and this therapy may also benefit patients with metastatic lung cancer (46,47).

### Radioactive stent implantation

A novel kind of radioactive particle stent implantation is being used in clinical practice in addition to the currently

available BT procedure. A meta-analysis of radioactive and plain stent insertion for the treatment of malignant airway stenosis was performed by Meng *et al.* (48). It revealed that radioactive stent insertion was sufficient to reduce the rate of stent restenosis and extend the patient's OS. New stents incorporating radioactive particles have also demonstrated therapeutic value in the treatment of inoperable malignant airway obstruction. Wang *et al.* discovered that from the second month after implantation, the group with radiographic stents had lower tracheal restenosis than the group with ordinary stents and a greater median survival (170 *vs.* 123 days,  $P < 0.05$ ). This study involved 66 patients with malignant airway obstruction (49). Moreover, radioactive stents can aid in increasing stent patency and OS and relieve superior vena cava stenosis caused by lung cancer (50). Although the safety has only been partially tested, a growing body of research points to the therapeutic potential of radioactive stents in the treatment of cavernous organs (51). However, additional reliable clinical trials are still needed.

### Combining RSI-BT with other treatments

#### RSI-BT combined with EBRT

RSI-BT can be used not only alone in clinical treatment but also in combination with other existing therapies to improve treatment outcomes. One study found that brachytherapy can improve tissue radiosensitivity by increasing  $pO_2$  in the tumour microenvironment, providing evidence for the combination of RSI-brachytherapy and EBRT (52). Retrospective research by Li *et al.* further demonstrated the efficacy of repeated I-125 seed implantation in conjunction with EBRT in the management of LR and metastasis in stage III/IV NSCLC [the overall (complete + partial) response rate was 87.4%]. The local control rates after

the first, second, and third years were 94.1%, 58.8% and 41.2%, respectively (53). Dual I-125 therapy is efficient in boosting the therapeutic dose to tumour lesions following EBRT failure. With an objective response rate (ORR) of 76%, 10 patients had CR, and 9 patients had PR after I-125 implantation in a retrospective study comprising 25 patients with HNSTS who underwent surgery and experienced recurrence after EBRT from 2006 to 2018 (54).

### RSI-BT combined with chemotherapy

RSI-BT has been used in combination with chemotherapy in a relatively large number of studies. Addressing the mechanism behind their combination, a study by Rong *et al.* found that lobaplatin (LBP) inhibits the AKT/mTOR pathway, promoting the apoptotic and antiproliferative effects of I-125 in NSCLC cells (55). I-125 particle implantation in conjunction with chemotherapy can improve the short-term local control rate [risk ratio (RR) 1.34; 95% CI: 1.09–1.65%,  $P=0.005$ ] and efficiency in early-stage lung cancer, but more clinical evidence is required (56). To control the progression of advanced lung cancer with less harmful side effects, which is crucial for elderly patients, RSI-BT can also be used in combination with second-line chemotherapeutic drugs, particularly after first-line chemotherapy fails (18,19,37,57). Moreover, bronchial artery embolization chemotherapy, a particular type of chemotherapy, may be used with RSI-BT. The combination can significantly slow tumour growth and enhance NSCLC prognosis (20,58–60). In a randomized controlled trial, patients who underwent bronchial artery chemoembolization (BACE) combined with I-125 RSI-BT achieved a better disease remission rate than those who underwent BACE alone (62.75% *vs.* 41.18%) (58). A pilot study, however, found no discernible difference in the effects of injectable medications used alone and in combination (61). Moreover, RSI-BT coupled with chemotherapy can ease severe airway obstruction caused by NSCLC. Jiang *et al.* effectively combined chemotherapy with the use of conventional transbronchial needle aspiration (C-TBNA) to implant radioactive particles into a bronchial lesion in a 51-year-old patient (62). Even though the majority of the currently used regimens of RSI-BT combined with chemotherapy have produced superior outcomes, further high-quality randomized controlled studies are still required to prove the combination therapy's safety and effectiveness.

### RSI-BT combined with targeted therapy

RSI-BT combined with targeted therapy has been applied in the treatment of NSCLC (63–66). According to Feng *et al.*, overall effectiveness in the apatinib combined with RSI-BT group was higher than that in the control group, and no fatal consequences occurred, demonstrating the therapy's safety and effectiveness (63). The elderly cases of combined apatinib and RSI-BT described by Zhang *et al.* can also be used to corroborate the findings of Feng *et al.* (64). In addition to clinical use, according to He *et al.* on A549 mice, RSI-BT combined with anlotinib prevented bone loss in tumour-bearing mice, but there was no significant difference between single and multiple particle implantations at the same total dose (65).

### RSI-BT combined with immunotherapy

Clinical reports of immunotherapy combined with RSI-BT in recent years are less frequent, and some studies have focused more on the mechanistic aspect. A safety investigation of oncologic therapies, which included RSI-BT, revealed no unanticipated or unmanageable adverse effects when RSI-BT was combined with immune checkpoint inhibition therapy (67). The pembrolizumab and I-125 RSI-BT combination demonstrated promising outcomes in NSCLC, and PD-L1 can be employed as a prognostic biomarker, according to a study by Wang *et al.* (68). Furthermore, immunotherapy can alter the shape of radioactive particles that enter tumour cells, improving the effectiveness and precision of RSI-BT (69). Although the use of immunotherapy and RSI-BT in combination has shown some positive clinical results, further research on the underlying concepts is still needed. The delivery method for RSI-BT can even be changed entirely.

### Limitations

There are still several side effects that limit RSI-BT in clinical treatment, such as radiation pneumonia, pneumothorax, and haemorrhage (70). The values of the incidence of haemorrhage, pneumothorax, and radiographic lung injury have been reported in the literature to be 0.14% (95% CI: 0.07–0.21%), 0.19% (95% CI: 0.11–0.28%), and 0.00% (95% CI: 0.00–0.03%), respectively (56). Meanwhile, RSI-BT can temporarily halt the progression of the disease. The lack of a unified standard industry practice and potential complications brought on by operator technical issues may impact the effectiveness of the treatment.

According to research conducted in Belgium, there is a need for advancements in RSI-BT-related education, awareness, implementation, collaboration, and cooperation (71). Consequently, additional data support and standardized training of staff are still needed.

## Conclusions

As technology advances, RSI-BT is an effective local oncologic intervention. It can be used alone or in conjunction with surgery, radiation, chemotherapy, immunotherapy, and targeted therapy to play a more complex role in clinical care. Its applications are promising, and RSI-BT plays a significant role in the treatment of digestive illnesses, prostate cancer, breast cancer, and lung cancer (72,73). To ensure treatment safety and efficacy, standardization of RSI-BT is needed. Meanwhile, multicentre, randomized clinical controlled trials are needed to validate the efficacy of minimally invasive interventions for tumours, including RSI-BT.

## Acknowledgments

The authors thank Shanghai Municipal Hospital Respiratory and Critical Care Medicine Specialist Alliance.

*Funding:* This work was supported by the Collaborative Innovation Cluster Project of Shanghai Municipal Health Commission (No. 2020CXJQ03), the Clinical Research Plan of SHDC (No. SHDC2020CR1021B-005) and the National Natural Science Foundation of China (No. 82000102).

## Footnote

*Reporting Checklist:* The authors have completed the Narrative Review reporting checklist. Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1600/rc>

*Peer Review File:* Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1600/prf>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1600/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related

to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021;71:209-49.
2. Araújo A, Barroso A, Parente B, et al. Unresectable stage III non-small cell lung cancer: Insights from a Portuguese expert panel. *Pulmonology* 2023;S2531-0437(22)00284-7.
3. Tanderup K, Ménard C, Polgar C, et al. Advancements in brachytherapy. *Adv Drug Deliv Rev* 2017;109:15-25.
4. Stewart A, Parashar B, Patel M, et al. American Brachytherapy Society consensus guidelines for thoracic brachytherapy for lung cancer. *Brachytherapy* 2016;15:1-11.
5. Casutt A, Kinj R, Ozsahin EM, et al. Fiducial markers for stereotactic lung radiation therapy: review of the transthoracic, endovascular and endobronchial approaches. *Eur Respir Rev* 2022;31:210149.
6. Costa R, Aires F, Rodrigues D, et al. Results of surgery versus stereotactic body radiotherapy for lung cancer. *Pulmonology* 2023;29:342-4.
7. Zhao J, Zhi Z, Zhang H, et al. Efficacy and safety of CT-guided (125)I brachytherapy in elderly patients with non-small cell lung cancer. *Oncol Lett* 2020;20:183-92.
8. Yang DY, Lin YP, Xue C, et al. CT-guided percutaneous implantation of (125)I particles in treatment of early lung cancer. *J Thorac Dis* 2020;12:5996-6009.
9. Dai F, Wang J, An H, et al. Therapy of 125I particles implantation inhibited the local growth of advanced non-small cell lung cancer: a retrospective clinical study. *Am J Transl Res* 2019;11:3737-49.
10. Kou F, Gao S, Liu S, et al. Preliminary clinical efficacy of iodine-125 seed implantation for the treatment of advanced malignant lung tumors. *J Cancer Res Ther*

- 2019;15:1567-73.
11. Liu B, Wang Y, Tian S, et al. Salvage treatment of NSCLC recurrence after first-line chemotherapy failure: Iodine-125 seed brachytherapy or microwave ablation? *Thorac Cancer* 2020;11:697-703.
  12. Yoon SM, Suh R, Abtin F, et al. Outcomes with multi-disciplinary management of central lung tumors with CT-guided percutaneous high dose rate brachyablation. *Radiat Oncol* 2021;16:99.
  13. Xiang L, Ren PR, Li HX, et al. Effect of 3-Dimensional Interstitial High-Dose-Rate Brachytherapy With Regional Metastatic Lymph Node Intensity-Modulated Radiation Therapy in Locally Advanced Peripheral Non-Small Cell Lung Cancer: 5-Year Follow-up of a Phase 2 Clinical Trial. *Int J Radiat Oncol Biol Phys* 2023;115:347-55.
  14. Sur R, Pond G, Falkson C, et al. BRACHY: A Randomized Trial to Evaluate Symptom Improvement in Advanced Non-Small Cell Lung Cancer Treated With External Beam Radiation With or Without High-Dose-Rate Intraluminal Brachytherapy. *Int J Radiat Oncol Biol Phys* 2023;116:601-10.
  15. Xiang Z, Wang L, Yan H, et al. (125)I seed brachytherapy versus external beam radiation therapy for the palliation of painful bone metastases of lung cancer after one cycle of chemotherapy progression. *Onco Targets Ther* 2018;11:5183-93.
  16. Zhang Z, Mao H, Wang X, et al. Comparison of I(125) seed brachytherapy (radioactive seed brachytherapy) joint three-dimensional conformal radiotherapy and stereotactic ablative radiotherapy on early nonsmall cell lung cancer. *J Cancer Res Ther* 2020;16:1560-8.
  17. Wang H, Lu J, Zheng XT, et al. Oligorecurrence Non-small Cell Lung Cancer After Failure of First-Line Chemotherapy: Computed Tomography-Guided (125)I Seed Implantation vs. Second-Line Chemotherapy. *Front Oncol* 2020;10:470.
  18. Xiang Z, Zhong Z, Mu L, et al. The Clinical Value of Computed Tomography (CT)-Guided (125)I Brachytherapy for Locally Advanced Non-Small Cell Lung Cancer After Progression of Concurrent Radiochemotherapy. *Cancer Manag Res* 2021;13:5297-307.
  19. Zhao G, Liu S, Liu Y, et al. CalliSpheres® microsphere transarterial chemoembolization combined with (125)I brachytherapy for patients with non-small-cell lung cancer liver metastases. *Front Oncol* 2022;12:882061.
  20. Wang X, Wang D. Clinical analysis of 125I seed implantation combined with epidermal growth factor receptor-tyrosine kinase inhibitors in advanced non-small cell lung cancer. *J BUON* 2021;26:1879-86.
  21. Zhang T, Mo Z, Duan G, et al. (125)I Seed Promotes Apoptosis in Non-small Lung Cancer Cells via the p38 MAPK-MDM2-p53 Signaling Pathway. *Front Oncol* 2021;11:582511.
  22. Xiang GL, Zhu XH, Lin CZ, et al. 125I seed irradiation induces apoptosis and inhibits angiogenesis by decreasing HIF-1alpha and VEGF expression in lung carcinoma xenografts. *Oncol Rep* 2017;37:3075-83.
  23. Cheng J, Ma S, Yang G, et al. The Mechanism of Computed Tomography-Guided 125I Particle in Treating Lung Cancer. *Med Sci Monit* 2017;23:292-9.
  24. Chargari C, Magne N, Guy JB, et al. Optimize and refine therapeutic index in radiation therapy: Overview of a century. *Cancer Treat Rev* 2016;45:58-67.
  25. Chargari C, Deutsch E, Blanchard P, et al. Brachytherapy: An overview for clinicians. *CA Cancer J Clin* 2019;69:386-401.
  26. Parashar B, Port J, Arora S, et al. Analysis of stereotactic radiation vs. wedge resection vs. wedge resection plus Cesium-131 brachytherapy in early stage lung cancer. *Brachytherapy* 2015;14:648-54.
  27. Zaorsky NG, Horwitz EM. Chapter 44 - Brachytherapy for Prostate Cancer: An Overview. In: Mydlo JH, Godec CJ, editors. *Prostate Cancer (Second Edition)*. San Diego: Academic Press; 2016:399-411.
  28. Mayadev J, Qi L, Lentz S, et al. Implant time and process efficiency for CT-guided high-dose-rate brachytherapy for cervical cancer. *Brachytherapy* 2014;13:233-9.
  29. Wu B, Ricchetti F, Sanguineti G, et al. Patient geometry-driven information retrieval for IMRT treatment plan quality control. *Med Phys* 2009;36:5497-505.
  30. Georg D, Hopfgartner J, Gora J, et al. Dosimetric considerations to determine the optimal technique for localized prostate cancer among external photon, proton, or carbon-ion therapy and high-dose-rate or low-dose-rate brachytherapy. *Int J Radiat Oncol Biol Phys* 2014;88:715-22.
  31. LeCun Y, Bengio Y, Hinton G. Deep learning. *Nature* 2015;521:436-44.
  32. Rivard MJ, Coursey BM, DeWerd LA, et al. Update of AAPM Task Group No. 43 Report: A revised AAPM protocol for brachytherapy dose calculations. *Med Phys* 2004;31:633-74.
  33. Beaulieu L, Carlsson Tedgren A, Carrier JF, et al. Report of the Task Group 186 on model-based dose calculation methods in brachytherapy beyond the TG-43 formalism: current status and recommendations for clinical



- implementation. *Med Phys* 2012;39:6208-36.
34. Ji Z, Sun H, Jiang Y, et al. Comparative study for CT-guided (125)I seed implantation assisted by 3D printing coplanar and non-coplanar template in peripheral lung cancer. *J Contemp Brachytherapy* 2019;11:169-73.
  35. Han X, Fang S, Sheng R, et al. Dosimetry verification of three-dimensional printed polylactic acid template-guided precision (125) I seed implantation for lung cancer using a desktop three-dimensional printer. *J Appl Clin Med Phys* 2021;22:202-9.
  36. Zheng Y, Jiang S, Yang Z, et al. Automatic needle detection using improved random sample consensus in CT image-guided lung interstitial brachytherapy. *J Appl Clin Med Phys* 2021;22:121-31.
  37. Chen ZK, Fan J, Li FQ, et al. I-125 seeds with chemotherapy for progressive non-small-cell lung cancer after first-line treatment: a meta-analysis. *J Cardiothorac Surg* 2022;17:75.
  38. Patel MA, Fazli Y, Sivakumar S, et al. Brachytherapy vs external beam therapy among NSCLC patients undergoing limited surgical resection. *J Cancer Res Clin Oncol* 2021;147:853-61.
  39. Zhang W, Li J, Li R, et al. Efficacy and safety of iodine-125 radioactive seeds brachytherapy for advanced non-small cell lung cancer-A meta-analysis. *Brachytherapy* 2018;17:439-48.
  40. Doggett SW, Chino S, Lempert T. A novel approach for salvage treatment of non-small-cell lung cancer: percutaneous CT fluoroscopy-guided permanent seed brachytherapy for salvage treatment of lung cancer: long-term results of a case series. *J Contemp Brachytherapy* 2019;11:174-9.
  41. Wang Y, Zhu L, Lin X, et al. Therapeutic Effect of CT-guided <sup>125</sup>I Seed Implantation on Advanced Lung Cancer and Pulmonary Metastatic Carcinoma. *Zhongguo Fei Ai Za Zhi* 2020;23:424-8.
  42. Wang Z, Chen S, Su M, et al. Analysis of the clinical efficacy and safety of computerized tomography-guided 125 I seed implantation in the treatment of non-small cell lung cancer that relapsed after chemoradiotherapy. *J Cancer Res Ther* 2022;18:426-31.
  43. Chatzikonstantinou G, Zamboglou N, Baltas D, et al. Image-guided interstitial high-dose-rate brachytherapy for dose escalation in the radiotherapy treatment of locally advanced lung cancer: A single-institute experience. *Brachytherapy* 2019;18:829-34.
  44. Rashid A, Pinkawa M, Haddad H, et al. Interstitial single fraction brachytherapy for malignant pulmonary tumours. *Strahlenther Onkol* 2021;197:416-22.
  45. Ji Z, Jiang Y, Guo F, et al. Radiation-related Adverse Effects of CT-guided Implantation of (125)I Seeds for Thoracic Recurrent and/or Metastatic Malignancy. *Sci Rep* 2019;9:14803.
  46. Ji Z, Jiang Y, Guo F, et al. Safety and efficacy of CT-guided radioactive iodine-125 seed implantation assisted by a 3D printing template for the treatment of thoracic malignancies. *J Cancer Res Clin Oncol* 2020;146:229-36.
  47. Zhao Z, Yao W, Zhang T, et al. Computed tomography-guided implantation of (125)I radioactive seeds in patients with malignant airway compression induced by advanced lung cancer: effectiveness and safety in 40 patients. *J Contemp Brachytherapy* 2020;12:343-50.
  48. Meng QK, Yu XY, Li W, et al. Radioactive and normal stent insertion for the treatment of malignant airway stenosis: A meta-analysis. *Brachytherapy* 2021;20:883-91.
  49. Wang Y, Lu J, Guo JH, et al. A Novel Tracheobronchial Stent Loaded with (125)I Seeds in Patients with Malignant Airway Obstruction Compared to a Conventional Stent: A Prospective Randomized Controlled Study. *EBioMedicine* 2018;33:269-75.
  50. Yuan FK, Fu YF, Shi YB, et al. Radioactive Seed Strand Efficacy in Superior Vena Cava Stenting Due to Non-small-cell Lung Cancer Obstruction. *Vasc Endovascular Surg* 2022;56:173-9.
  51. Zhu HD, Guo JH, Mao AW, et al. Conventional stents versus stents loaded with (125)iodine seeds for the treatment of unresectable oesophageal cancer: a multicentre, randomised phase 3 trial. *Lancet Oncol* 2014;15:612-9.
  52. Cron GO, Beghein N, Crokart N, et al. Changes in the tumor microenvironment during low-dose-rate permanent seed implantation iodine-125 brachytherapy. *Int J Radiat Oncol Biol Phys* 2005;63:1245-51.
  53. Li F, Wang L, Zhang Y, et al. A Retrospective Study on Using a Novel Single Needle Cone Puncture Approach for the Iodine-125 Seed Brachytherapy in Treating Patients With Thoracic Malignancy. *Front Oncol* 2021;11:640131.
  54. Chen Y, Jiang Y, Ji Z, et al. Efficacy and safety of CT-guided (125)I seed implantation as a salvage treatment for locally recurrent head and neck soft tissue sarcoma after surgery and external beam radiotherapy: A 12-year study at a single institution. *Brachytherapy* 2020;19:81-9.
  55. Rong JH, Li D, Li YL. Lobaplatin Enhances Radioactive (125)I Seed-Induced Apoptosis and Anti-Proliferative Effect in Non-Small Cell Lung Cancer by Suppressing the AKT/mTOR Pathway. *Onco Targets Ther* 2021;14:289-300.

56. Chen E, Wang J, Zhang H, et al. Analysis of the efficacy and safety of iodine-125 seeds implantation in the treatment of patients with inoperable early-stage non-small cell lung cancer. *J Contemp Brachytherapy* 2021;13:347-57.
57. Yue TH, Xing W. (125)I Seed Brachytherapy Combined with Single-Agent Chemotherapy in the Treatment of Non-Small-Cell Lung Cancer in the Elderly: A Valuable Solution. *Onco Targets Ther* 2020;13:10581-91.
58. Xie P, He L, Zhang Y. Analysis on the Efficacy of Bronchial Artery Chemoembolization Combined with 125I Seed Implantation in the Therapy of Advanced Non-Small-Cell Lung Cancer Based on the Medical Database. *Biomed Res Int* 2022;2022:7376844.
59. Xie P, Zhang Y, He L. Prognostic Evaluation of CT Imaging Big Data-Assisted Arterial Chemoembolization Combined with (125)I Seed Implantation for Non-Small-Cell Lung Cancer. *Comput Math Methods Med* 2022;2022:3472982.
60. Chen C, Wang W, Yu Z, et al. Combination of computed tomography-guided iodine-125 brachytherapy and bronchial arterial chemoembolization for locally advanced stage III non-small cell lung cancer after failure of concurrent chemoradiotherapy. *Lung Cancer* 2020;146:290-6.
61. Zhao Z, Tu J, Fan X, et al. Drug-eluting beads-transcatheter arterial chemoembolization with or without iodine-125 treatment is effective and tolerable in treating advanced non-small cell lung cancer patients: a pilot study. *Transl Cancer Res* 2020;9:3191-202.
62. Jiang AG, Lu HY, Ding ZQ. Implantation of (125)I radioactive seeds via c-TBNA combined with chemotherapy in an advanced non-small-cell lung carcinoma patient. *BMC Pulm Med* 2019;19:205.
63. Feng Y, Yang B, Li X. Clinical analysis of 125I seed implantation combined with Apatinib in the treatment of locally advanced lung cancer: A case series. *J Pak Med Assoc* 2021;71:1025-7.
64. Zhang Y, Yin S, Jia Y, et al. Safety and efficacy of apatinib combined with iodine-125 in chemotherapy-refractory advanced lung cancer: A case report. *Medicine (Baltimore)* 2020;99:e21600.
65. He F, Bao Q, Bai J, et al. Effects of I-125 seeds combined with anlotinib on tumor growth and bone metabolism in A549 tumor-bearing mice. *Int J Radiat Biol* 2021;97:1578-88.
66. Xiang L, Zhang JW, Lin S, et al. Computed Tomography-Guided Interstitial High-Dose-Rate Brachytherapy in Combination With Regional Positive Lymph Node Intensity-Modulated Radiation Therapy in Locally Advanced Peripheral Non-Small Cell Lung Cancer: A Phase 1 Clinical Trial. *Int J Radiat Oncol Biol Phys* 2015;92:1027-34.
67. Leppelmann KS, Mooradian MJ, Ganguli S, et al. Thermal Ablation, Embolization, and Selective Internal Radiation Therapy Combined with Checkpoint Inhibitor Cancer Immunotherapy: Safety Analysis. *J Vasc Interv Radiol* 2021;32:187-95.
68. Wang S, Zhang J, Meng FJ, et al. Combination of pembrolizumab and (125)I attenuates the aggressiveness of non-small cell lung cancer. *Oncol Lett* 2020;19:4142-50.
69. Kelly G, Milligan JJ, Mastria EM, et al. Intratumoral delivery of brachytherapy and immunotherapy by a thermally triggered polypeptide depot. *J Control Release* 2022;343:267-76.
70. Zhang F, Wang J, Guo J, et al. Chinese expert consensus workshop report: Guideline for permanent iodine-125 seeds implantation of primary and metastatic lung tumors (2020 edition). *J Cancer Res Ther* 2020;16:1549-54.
71. Salembier C, De Hertogh O, Daisne JF, et al. Brachytherapy in Belgium in 2018. A national survey of the brachytherapy study group of the Belgian Society for Radiotherapy and Oncology (BeSTRO). *Radiother Oncol* 2020;150:245-52.
72. Terheyden MM, Melchert C, Kovács G. External beam boost versus interstitial high-dose-rate brachytherapy boost in the adjuvant radiotherapy following breast-conserving therapy in early-stage breast cancer: a dosimetric comparison. *J Contemp Brachytherapy* 2016;8:294-300.
73. Otake S, Goto T. Stereotactic Radiotherapy for Oligometastasis. *Cancers (Basel)* 2019;11:133.

**Cite this article as:** Wang X, Tian S, Shi H, Qin H, Zhang W, Dong Y, Bai C. Recent progress in radioactive seed implantation brachytherapy of non-small cell lung cancer: a narrative review. *J Thorac Dis* 2024;16(3):2167-2176. doi: 10.21037/jtd-23-1600