

## APPLIED ECOLOGY

# Inconsistencies undermine the conclusion that agriculture is a dominant source of NO<sub>x</sub> in California

Tai McClellan Maa<sup>1\*</sup>, Sarah Waldo<sup>2\*</sup>, Tom Bruulsema<sup>1\*</sup>, Rob Mikkelsen<sup>1\*</sup>

**Almaraz *et al.* reported that agricultural soils are a dominant source of NO<sub>x</sub> pollution in California (20 to 32% of total statewide NO<sub>x</sub> emissions). However, this conclusion may be undermined by the lack of agreement between their modeled estimates and previously reported empirical measurements, the extrapolation of NO<sub>x</sub> fluxes during hot moments to derive annual estimates, and the overestimation of nitrogen fertilizer consumption in California.**

Almaraz *et al.* (1) estimated that the mean NO<sub>x</sub> emission flux from California's agricultural soils was 19.8 kg of N ha<sup>-1</sup> year<sup>-1</sup>, in comparison to 1 kg of N ha<sup>-1</sup> year<sup>-1</sup> from natural ecosystems. The study estimated that 161,100 metric tons of NO<sub>x</sub>-N year<sup>-1</sup> was emitted from California's soils, and that 127,000 metric tons of this was derived from agricultural land. A bottom-up model was scaled up on the basis of the N surplus and theoretical NO, N<sub>2</sub>O, and N<sub>2</sub> partitioning based on soil conditions, and verified using top-down airborne NO<sub>x</sub> measurements. Using these two approaches, the study identified agricultural regions in California that are potential hot spots for NO<sub>x</sub> emissions due to the spatial distribution of soil environmental conditions and N fertilizer inputs. We agree that accurate accounting of all sources of NO<sub>x</sub> is important, particularly for the management and mitigation of direct emissions from agricultural soils.

The magnitude of NO<sub>x</sub> emitted from agricultural soils presented in this work, however, is considerably greater than the magnitudes reported in previously published work. Given that background NO<sub>x</sub> fluxes were estimated at 1 kg of N ha<sup>-1</sup> year<sup>-1</sup> and fertilizer inputs averaged 132 kg ha<sup>-1</sup>, attributing 19.8 kg of NO<sub>x</sub>-N ha<sup>-1</sup> year<sup>-1</sup> to agricultural land implies a fertilizer-induced emission factor in excess of 14%. A recent global meta-analysis reported a fertilizer-induced NO emission factor of 1.2% (2). Another earlier global assessment concluded that the emission factor for NO is similar to that for N<sub>2</sub>O (3). While measurements in California were not included in previous global assessments, measurements from similar agroecosystems were included. The assertion that the state of California has twice the NO flux and an emission factor eightfold greater than the previous global assessment for vegetable croplands (2) warrants a high degree of scrutiny. Emission factors can vary regionally owing to soil texture, climate, and other factors, but Almaraz *et al.* did not discuss their findings within the context of these global assessments nor provide an analysis of uncertainty needed to interpret their estimate.

The authors concluded that their estimates of total NO<sub>x</sub> flux were "slightly higher than, although comparable with, the few number of empirical [chamber-based] measurements of NO<sub>x</sub> emissions from the San Joaquin Valley's cropland soils (made between July and September 1995)." This discrepancy was attributed to greater fertilizer use and population growth in the last 20 years.

<sup>1</sup>International Plant Nutrition Institute, 5550 Triangle Parkway, Suite 300, Peachtree Corners, GA 30092, USA. <sup>2</sup>National Risk Management Research Laboratory, U.S. Environmental Protection Agency, Office of Research and Development, 26 West Martin Luther King Drive, Cincinnati, OH 45268, USA.

\*Corresponding author. Email: tom.bruulsema@ipni.net (T.B.); tmaaz@ipni.net (T.M.M.); waldo.sarah@epa.gov (S.W.); rmikkelsen@ipni.net (R.M.)

Copyright © 2018  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. No claim to  
original U.S. Government  
Works. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

However, three potential issues arise from this argument. First, 45% of the modeled values were more than twofold greater than these averaged empirical measurements, and 31% were at least an order of magnitude larger. The paper cites a study that included 2850 NO<sub>x</sub> measurements at 26 sites in 10 different cropping systems and reported average peak (noon-time) and diel (adjusted for lower night-time temperature) fluxes of 4.0 and 1.9 kg of N ha<sup>-1</sup> year<sup>-1</sup>, respectively, if hourly summer fluxes are extrapolated to annual emissions. Even the agreement between the top-down and bottom-up estimates reported in Almaraz *et al.* is difficult to assess without an analysis of model uncertainty; and even when canopy exchange is considered, the bottom-up modeled flux for July and August is 32 to 47% greater than the top-down estimate based on airborne measurements collected in July and August.

Second, too much weight has been given to the hot moments of measured emissions when scaling up temporally. The best agreement between modeled and observed high-magnitude NO fluxes reported in Almaraz *et al.* occurs in the Imperial Valley, CA. The fluxes modeled by Almaraz *et al.* appear to agree with chamber-based measurements published in Oikawa *et al.* (4), which was reported by Almaraz *et al.* as 21 kg of N ha<sup>-1</sup> year<sup>-1</sup> in table 1. Almaraz *et al.* derived this estimate by extrapolating an average flux of 66 ng of NO<sub>x</sub>-N m<sup>-2</sup> s<sup>-1</sup> to represent annual emissions. This extrapolation is inappropriate because the NO measurements by Oikawa *et al.* were largely designed to characterize emission pulses following summer fertilization. Because of the aims of that research, the measurements are weighted toward high-emission periods. Furthermore, Oikawa *et al.* reported integrated emission factors of 1.8 to 6.6%, less than the emission factor if the annual average emission rate were 21 kg of N ha<sup>-1</sup> year<sup>-1</sup> (~11%, if averaged across both small and large field measurements). Excluding or down-adjusting the Oikawa *et al.* "Observed NO" from table 1 limits the range of the empirical data and further challenges the quantitative coherence between modeled and observed values.

Third, the perception that state-wide N fertilizer consumption in agriculture has increased since the 1990s is not supported by the available data. California fertilizer N sales plateaued in the early 2000s (5). According to the Association of American Plant Food Control Officials, annualized mineral N fertilizer sales have declined to pre-2000 levels (6). The California Department of Food and Agriculture reports the sales of Home and Garden fertilizer blends and compost for organic agriculture under code 0 ("Identified by Grade"). Following reporting changes in 2012, 44% of total fertilizer sales have fallen in this category, resulting in a tonnage increase by more than 300,000 metric tons (7). However, Almaraz *et al.*

did not assess nonagricultural fertilizer inputs, such as for Home and Garden fertilizer use, in urban areas. Future nitrogen modeling groups using these data should be aware of these reporting differences, as well as the need to differentiate among the uses and sources of nitrogen fertilizers (agricultural versus nonagricultural uses) to properly interpret the model outputs.

The impact of these inconsistencies is difficult to assess because Almaraz *et al.* did not present their N budget for California. The N rate and total emission data estimated in this study infer an agricultural area of 6,423,737 ha and a natural ecosystem area of 33,810,000 ha. The total annual N fertilizer inputs to agriculture would thereby amount to 848,000 metric tons. This estimate exceeds fertilizer N sales in California from 1980 to 2007, a period for which Almaraz *et al.* used crop-specific data to generate N inputs, by almost 50% (6). It also well exceeds total nitrogen fertilizer sales in 2016, even when additional sources of nitrogen fertilizers are included (7). It is unclear whether this estimate includes other sources of reactive N (that is, N from biological fixation and deposition), although the authors state that manure N was not included. Is it possible that the authors overestimated fertilizer N inputs? Or are the authors using the term “fertilizer” when they actually meant “reactive” N?

We ask that the authors upload their complete N budget and assumptions for water-filled pore space, along with an uncertainty analysis, for California as supplemental data. This will help the reader properly compare and otherwise address the apparent inconsistencies in emission factors and allow for an improved assessment of the role of agriculture as a source of NO<sub>x</sub> in California.

*EPA Disclaimer: This document has been reviewed in accordance with U.S. Environmental Protection Agency policy and approved for external publication. The views expressed in this article are those of the authors and do not necessarily represent the opinions or policies of the U.S. Environmental Protection Agency. Any mention of trade*

*names or commercial products does not constitute endorsement or recommendation for use.*

## REFERENCES AND NOTES

1. M. Almaraz, E. Bai, C. Wang, J. Trousdell, S. Conley, I. Faloon, B. Z. Houlton, Agriculture is a major source of NO<sub>x</sub> pollution in California. *Sci. Adv.* **4**, eaao3477 (2018).
2. S. Liu, F. Lin, S. Wu, C. Ji, Y. Sun, Y. Jin, S. Li, Z. Li, J. Zou, A meta-analysis of fertilizer-induced soil NO and combined NO+N<sub>2</sub>O emissions. *Glob. Change Biol.* **23**, 2520–2532 (2017).
3. A. F. Bouwman, L. J. M. Boumans, N. H. Batjes, Modeling global annual N<sub>2</sub>O and NO emissions from fertilized fields. *Glob. Biogeochem. Cycles* **16**, 1080 (2002).
4. P. Y. Oikawa, C. Ge, J. Wang, J. R. Eberwein, L. L. Liang, L. A. Allsman, D. A. Grantz, G. D. Jenerette, Unusually high soil nitrogen oxide emissions influence air quality in a high-temperature agricultural region. *Nat. Commun.* **6**, 8753 (2015).
5. T. S. Rosenstock, D. Liptzin, J. Six, T. P. Tomich, Nitrogen fertilizer use in California: Assessing the data, trends and a way forward. *Calif. Agric.* **67**, 68–79 (2013).
6. J. Slater, B. Kirby, *Commercial Fertilizers* (The Association of American Plant Food Control Officials in partnership with The Fertilizer Institute, Fertilizer/Ag Lime Control Service, University of Missouri, 1985–2014).
7. California Department of Food and Agriculture, *Fertilizer Tonnage Report by Year* (Feed, Fertilizer, and Livestock Drugs Regulatory Services, Division of Inspection Services, 2018); [https://www.cdfa.ca.gov/is/ffldrs/Fertilizer\\_Tonnage.html](https://www.cdfa.ca.gov/is/ffldrs/Fertilizer_Tonnage.html).

## Acknowledgments

**Funding:** No funding was involved in this work. **Author contributions:** All authors conceived of and reviewed the technical comment. T.M.M. and S.W. wrote the manuscript. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and in the references cited. Additional data related to this paper may be requested from the authors.

Submitted 1 March 2018

Accepted 31 July 2018

Published 12 September 2018

10.1126/sciadv.aat4706

**Citation:** T. M. Maaz, S. Waldo, T. Bruulsema, R. Mikkelsen, Inconsistencies undermine the conclusion that agriculture is a dominant source of NO<sub>x</sub> in California. *Sci. Adv.* **4**, eaat4706 (2018).