



Toward Systems Models for Obesity Prevention: A Big Role for Big Data

Adele R Tufford,¹ Christos Diou,² Desiree A Lucassen,¹ Ioannis Ioakimidis,³ Grace O'Malley,^{4,5} Leonidas Alagialoglou,⁶ Evangelia Charmandari,^{7,8} Gerardine Doyle,^{9,10} Konstantinos Filis,¹¹ Penio Kassari,^{7,8} Tahar Kechadi,¹² Vassilis Kilintzis,¹³ Esther Kok,¹ Irini Lekka,¹³ Nicos Maglaveras,¹³ Ioannis Pagkalos,¹⁴ Vasileios Papapanagiotou,⁶ Ioannis Sarafis,⁶ Arsalan Shahid,¹² Pieter van 't Veer,¹ Anastasios Delopoulos,⁶ and Monica Mars¹

¹Division of Human Nutrition and Health, Wageningen University and Research, Wageningen, Netherlands; ²Department of Informatics and Telematics, Harokopio University of Athens, Athens, Greece; ³Department of Biosciences and Nutrition, Karolinska Institute, Stockholm, Sweden; ⁴W82GO Child and Adolescent Weight Management Service, Children's Health Ireland at Temple Street, Dublin, Ireland; ⁵Division of Population Health Sciences, School of Physiotherapy, Royal College of Surgeons in Ireland University for Medicine and Health Sciences, Dublin, Ireland; ⁶Department of Electrical and Computer Engineering, Aristotle University of Thessaloniki, Thessaloniki, Greece; ⁷Division of Endocrinology, Metabolism, and Diabetes, First Department of Pediatrics, National and Kapodistrian University of Athens Medical School, "Aghia Sophia" Children's Hospital, Athens, Greece; ⁸Division of Endocrinology and Metabolism, Center for Clinical, Experimental Surgery and Translational Research, Biomedical Research Foundation of the Academy of Athens, Athens, Greece; ⁹College of Business, University College Dublin, Dublin, Ireland; ¹⁰Geary Institute for Public Policy, University College Dublin, Dublin, Ireland; ¹¹COSMOTE Mobile Telecommunications, Athens, Greece; ¹²CeADAR: Ireland's Centre for Applied AI, University College Dublin, Dublin 4, Ireland; ¹³Lab of Computing, Medical Informatics, and Biomedical Imaging Technologies, Department of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece; and ¹⁴Department of Nutritional Sciences and Dietetics, School of Health Sciences, International Hellenic University, Thessaloniki, Greece

ABSTRACT

The relation among the various causal factors of obesity is not well understood, and there remains a lack of viable data to advance integrated, systems models of its etiology. The collection of big data has begun to allow the exploration of causal associations between behavior, built environment, and obesity-relevant health outcomes. Here, the traditional epidemiologic and emerging big data approaches used in obesity research are compared, describing the research questions, needs, and outcomes of 3 broad research domains: eating behavior, social food environments, and the built environment. Taking tangible steps at the intersection of these domains, the recent European Union project "BigO: Big data against childhood obesity" used a mobile health tool to link objective measurements of health, physical activity, and the built environment. BigO provided learning on the limitations of big data, such as privacy concerns, study sampling, and the balancing of epidemiologic domain expertise with the required technical expertise. Adopting big data approaches will facilitate the exploitation of data concerning obesity-relevant behaviors of a greater variety, which are also processed at speed, facilitated by mobile-based data collection and monitoring systems, citizen science, and artificial intelligence. These approaches will allow the field to expand from causal inference to more complex, systems-level predictive models, stimulating ambitious and effective policy interventions. *Curr Dev Nutr* 2022;6:nzac123.

Keywords: behavior, built environment, big data, childhood obesity, monitoring systems, physical activity, systems models

© The Author(s) 2022. Published by Oxford University Press on behalf of the American Society for Nutrition. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Manuscript received November 8, 2021. Initial review completed March 24, 2022. Revision accepted July 28, 2022. Published online July 30, 2022. doi: <https://doi.org/10.1093/cdn/nzac123>

Supported by Horizon2020 Framework Programme project grant agreement no. 727688: "BigO: Big data against childhood obesity."

Author disclosures: the authors report no conflicts of interest.

Address correspondence to ART (e-mail: a.r.tufford@uu.nl).

Abbreviations used: FAIR, findable, accessible, interoperable, reusable; GPS, Global Positioning System; mHealth, mobile health.

Introduction

Obesity is one of the most important global health issues, owing to its association with significant morbidity and mortality, as well as increased public health costs. Overweight and obesity—as defined by abnormal or excessive fat accumulation that may impair health—has risen dramatically in prevalence during the last 4 decades in every global region (1–3). As of 2016, 13% of all adults had obesity and 39% had overweight; whereas, 28% of children had overweight or obesity as of 2017 (4). Overweight and obesity are risk factors for a broad set of non-communicable diseases. Individuals with obesity were also recently estimated to be 1.5 times more likely to die of COVID-19 (5), because

the associated immune system compromise, itself influenced by associated comorbidities, complicates treatment, recovery, and vaccination efficacy (6). Consequently, excess weight contributes to a substantial loss in disease-free years and avoidable deaths (3), which makes the identification of the major causes and drivers of obesity a primary concern. Although the "obesogenic environment" is a key predictor, specific causal factors related to the built and social environments remain underexplored. The global and uniform rise in obesity prevalence points to complex, multifactorial, and systemic causal factors (e.g., income, education, food quality), in contrast to the assumption that obesity is an individual's responsibility stemming from poor lifestyle choices (7, 8).

Although current approaches to explain obesity development consider many etiologic factors—from genetic to molecular to environmental levels—there is still an urgent need for tools and methods able to capture the complexity of the factors and systems contributing to obesity. Nearly 20 y ago, the Ecological Systems Theory was applied to obesity, leading to the ecological model of obesity, wherein an individual's weight status is heavily influenced by their social, economic, cultural, and environmental context (9). This model was further elaborated by the Foresight project, which identified 108 key variables relevant to energy balance, grouped into 7 themes: food production, food consumption, social psychology, individual psychology, physiology, individual physical activity, and the physical activity environment (10). These systems models, as well as others in obesity and nutrition (11–15), have offered tangible public health intervention points around which to shape policies and model scenarios aimed at obesity prevention and management, as well as a framework for modeling the concentric spheres of agents (e.g., individual, family, community, government, industry) that influence body weight gain (9, 16, 17). In addition, the *Lancet* Commission on Obesity has made critical leaps in systems thinking with landmark articles such as “The Global Syndemic” (11), delivering a transdisciplinary framework for addressing the interconnections between obesity, undernutrition, and climate change. Despite this, a holistic, data-driven understanding of *how* the complex milieu of the built and dietary environments and socioeconomic position (income, education, and health inequalities) affects individual behavior (nutrition and physical activity level) is still lacking. Although recent studies have made advances in capturing a wider range of systemic factors using mixed-methods approaches (18), static or traditional survey-based approaches fail to appropriately capture the complexity of factors and dynamic interactions between the individual and their environment (19, 20).

The adoption of *big data* creates the potential to generate synergies between biological and ecological approaches toward a better understanding of the etiology of obesity (21). Big data have the potential to identify intervention points for obesity prevention or management *in vivo*, across different environmental settings, as well as create data-driven causal maps of obesity determinants. A recent Delphi study sought to define big data in the context of obesity research, concluding that big data:

...is always digital, has a large sample size, and a large volume or variety or velocity of variables that require additional computing power. It can include quantitative, qualitative, observational, or interventional data from a wide range of sources (e.g., government, commercial, cohorts) that have been collected for research or other purposes, and may include one or several datasets. (22)

In this framework, volume concerns sample size, number of variables, or measurement occasions; variety concerns the types of data and numbers of variables; and velocity means that data are generated and processed in real-time or at speed—translating them into interfaces to make them readily usable (22). Moreover, the study noted that the variety component of big data in obesity research is more nuanced than its traditional definition within the big data domain, involving quantitative, qualitative, and observational data from a broad range of sources, e.g., clinical cohorts, census data, Geographic Information Systems (GIS) data, social media, advertising, and commercial sales data. To date, big data are under-utilized and under-exploited in obesity research. There

are few widely available data sets concerning obesity that conform to the foregoing definition; few studies have used a wide variety of relevant variables, permitting multivariate investigations, and seemingly none have used velocity (22–24). Furthermore, it is not yet clear how obesity research can ethically use big data methods, although ethical frameworks are emerging from the literature (25, 26).

In this report, we investigate the potential for big data to support and expand current methodologies aimed toward systems approaches in the domain of obesity etiology. The complementarity and novel additions of big data to current obesity research approaches are discussed, as well as important risks and considerations when collecting such data, and the roles for domain and technological expertise.

Traditional Compared with Emergent Approaches to Study Causal Relations in the Etiology of Obesity

The Ecological Systems Theory of obesity proposed 20 y ago added structure and form to our understanding of obesity as a chronic disease arising from biological, social, and environmental interactions (9). Combined with the arrival of the genomic and information ages, complex health (and biological) approaches to obesity etiology have emerged, including the Foresight model. In what follows, we discuss the contributions and pitfalls of current epidemiologic approaches to obesity research and highlight the ways in which big data may contribute.

Epidemiologic: diet, socioeconomic, and food-environment drivers of obesity

Epidemiologic research in obesity has led to critical insights into obesity's noncommunicable disease sequelae and its disproportionate prevalence in socioeconomically disadvantaged groups in high-income countries. This line of research has also motivated hypothesis-driven research on the links between obesity and living environments; particularly as it relates to the drivers of food consumption, physical activity, and social psychology (27). Epidemiologic surveillance over time has also revealed the changing relation between these aforementioned factors and weight status over time; such as the associations between weight status and urban compared with rural living environments, or history of breastfeeding (28, 29). These insights have informed important public health interventions aimed at lowering inequalities in obesity, for example, programs targeting breastfeeding rates, school nutrition, or the reduction of sedentary behavior (30).

The large-scale WHO European Childhood Obesity Surveillance Initiative (COSI) has collected data from >300,000 children in Europe, leading to the development of a standardized monitoring system, and a variety of insights on, e.g., diet and physical activity patterns by country (31). The Healthy Behavior in School-Aged Children study, examining data from >130,000 adolescents from 29 European countries, was able to show that individual differences in physical activity could largely be explained by between-country differences in environmental variables, such as temperature, safety, household income, and public health policy (32). In the United States, the National Collaborative on Childhood Obesity Research has delivered critical research frameworks for childhood obesity, such as compendiums of physical activity codes relevant to youth, and measurement and surveillance needs for childhood obesity prevention from a variety of entry points (i.e., nutrition,

physical activity, and food environments) (33–35). Despite these advances, these traditional data assessment methods lack the variety and granularity of data needed to explore a wider array of determinants—both group and individual—relevant to obesity. Classical epidemiologic approaches, such as dietary and exposure assessments followed by linkage to BMI, are strongly focused on the outcome of body size, rather than on complex associations between exposures, environments, and risk of obesity-related health complications.

In addition, BMI is often used as a metric for weight status; however, this is a proxy measure of adiposity and therefore an incomplete assessment of individual health status. Some individuals within the normal-weight range may display all characteristics of obesity, and those outside the normal range, not (36, 37). Importantly, the use of BMI and BMI cutoffs as a measure for weight status is problematic when comparing people with a different racial background, because body posture and build differ widely between these groups (36, 37). Moreover, these approaches also tend to orient (methodologically) from a biomedical lens. This means genes, nutrients, and foods, rather than social and environmental determinants (e.g., socioeconomic factors and the built environment), have taken the center stage (38, 39). Epidemiology also tends to side on inference of (health) outcomes based on behaviors and exposure, rather than data-driven prediction models of outcomes under differing or altered circumstances (40).

The more complex associations between exposures, environments, and outcomes are difficult to capture using traditional “offline” methods (food-frequency surveys, dietary intake recall) (41, 42). A systematic review of cohort studies that included both individual and ecological data purporting to examine the influence of food environment on obesity concluded that all available studies ($n = 71$) were of low quality and error-prone; and associations were predominantly null, failing to deliver meaningful conclusions or evidence (43). An additional scoping review on advanced methods linking dietary behaviors and the food environment uncovered only 5 studies with >100 participants that incorporated both Global Positioning System (GPS) and food intake data (44). None were transnational, and the period of data collection was a maximum of 7 d. The authors concluded that this paucity of data hinders the development of advanced modeling of food behaviors, such as agent-based modeling or complex-adaptive systems modeling. Incorporating environmental attributes in a manner that still enables the identification of causal relations, with attention to data veracity and reproducibility of variables, is a critical challenge for 21st-century epidemiologists. This is particularly the case for factors relevant to the social determinants of health as pertaining to obesity—for example, quality and accessibility of healthy foods or availability of athletics and recreation facilities.

Big data: toward a more complex systems etiology of obesity

Big data are new and underexploited in obesity research. The most relevant contributions so far are from “found/real-world data” scraped for other purposes, such as social media, retail, and environmental sensing, rather than generated for specific obesity-related objectives or hypotheses (22). An agreed-upon scope and definition of big data for obesity research is both nuanced and still emerging (22). Moreover, an overview of case studies purposing to use big data in obesity research did not identify any studies using advanced sensors/apps for use by individuals or participants (45). The studies relied on post hoc analysis of 1-time

administered surveys or electronic health records, in combination with data from external sources—e.g., land-value and census records (24). On the other hand, very recent advances have been made in using accelerometers to measure physical activity and its link to overweight and obesity risk, such as the International Children’s Accelerometry Database Collaboration study (46). Gathering big data on obesity via mobile surveillance/monitoring has the promise to link relevant variables at the level of individuals and populations—allowing synergy between the biomedical and epidemiologic approaches. The emergence of new assessment methods means that individual or group-level data of large populations can be collected in real-time, allowing researchers to move away from average or self-reported values of food intake, mood, physical activity, or geolocation. This brings enormous potential to examine dietary behaviors more accurately over time and across contexts, which can lead to advanced predictive and causal models for planned interventions. The aforementioned Delphi study highlighted the need for the adoption of more advanced statistical, machine learning, artificial intelligence, and database methods to address complex hypotheses (22). These research avenues could include the influence of social media on food intake, or the prediction of weight status based on distinct environmental features—associations that are challenging to determine using traditional methods. Next, we outline 3 emerging areas where big data can be used in the obesity research context.

Mobile sensing.

Mobile sensing and mobile health (mHealth) tools have the potential to continuously harvest behavioral and environmental data in real-time. This field is rapidly expanding (47), but remains underused in research contexts owing to the reliance on mainly commercially developed tools, and the lack of approved devices and systems for broad use in clinical trials. These approaches have the potential, although still challenging in a research context, to measure both physical activity and food intake (both self-reported and sensor-based), as well as collect data related to the built and food environments. These tools avoid the drawbacks of certain self-reported measures, and may also feed-forward either specific interventions or behavioral suggestions to the user, i.e., the use of “nudging” approaches to increase physical activity behavior (48). These strategies could be adopted not only using mobile phone-based sensors and apps but also via alternative means of data collection such as smartwatches, food-intake sensors, and other wearables (45, 49, 50). To date, a few pilot mHealth tools have been made available to gather information on food environments in a participatory way (51, 52). The European study DEDIPAC (DEterminants of DIet and Physical ACTivity) made steps toward capacity building for big data in the field, making progress, for example, in the adoption of physical activity sensor-derived data (53, 54). Measurement of physical activity over time among diverse groups is therefore feasible; however, broader efforts to incorporate these measures in real-time, in combination with social/environmental determinants of physical activity, have not yet been achieved.

Citizen science.

The approach of citizen science uses large and engaged populations as data providers, involving the participant in the whole scientific process. This approach not only facilitates “feeding-forward” of health information to participants, but also exploits and/or enhances their knowledge

base of science and health (55). Citizen science tools for health monitoring in the public domain are generally developed in controlled populations, followed by dissemination to the public; or involve crowd-sourced data contributions or scraping of existing databases. In the health care domain, projects include SCAMPI (the Smart City Active Mobile Phone Intervention), a mobile intervention tool to promote physical activity among sedentary adults (56), and Big data against childhood Obesity (BigO), discussed in detail in what follows. The Our Voice project, a mobile community engagement tool, was deployed in 5 countries and collects photos and voice memos from individuals on the theme of physical activity and food access (57). A major drawback to a citizen science approach from a research perspective, particularly as linked to health-related behaviors, is that it is accompanied by an uncontrolled sample population. Therefore, uncontrolled data collection protocols and reliance on self-reported measures of health make it challenging to derive valid associations (58). The use of traditional statistical approaches on this type of data can be erroneous. New statistical approaches need to be developed and used that consider selectivity biases, the sparseness of data, and noise. These aforementioned issues have been recognized by Eurostat and all European Union national statistical authorities, but can take inspiration from success in other domains, for example in environmental monitoring of biodiversity (59, 60).

Moreover, the use of citizen science tools may be biased to urban, relatively wealthy, highly educated individuals, although this bias is also a known pitfall in food intake and nutrition survey responses. The steady increase in smartphone use in all global regions, particularly among younger users, could overcome this with appropriate dissemination efforts (61).

Artificial intelligence.

The rapid evolution in the technological development of artificial intelligence, including machine and deep learning, is moving research beyond data collection and toward knowledge extraction and prediction. Artificial intelligence promises to mine purchase, Internet, and social media data in an aggregated manner, allowing analysis, visualization, and prediction from individual and group-level big data. In the context of obesity, this technique has the promising potential outcome of deep learning-derived prediction of population behavior and weight status, based on environments or proposed interventions. Artificial intelligence has the potential to overcome some of the aforementioned sampling biases in citizen science-based approaches, eventually leading to the automatic fitting of prediction models and the identification of modifiers and confounders—allowing the prediction of the outcomes of health interventions (62, 63). So far, this type of prediction has been largely the domain of tech giants such as Facebook, IBM Watson, Google, and Amazon. For example, the power of Google's data access and intelligent computing capacity is being combined in "Google Health," which aims at predicting and treating a variety of health issues. Critics are concerned about this form of industry research, including a lack of standards for ethical accountability and informed consent (63). In the public domain, the predominant use of artificial intelligence as it pertains to diet and eating behavior has been in computational neuroscience, with emergent applications aimed at automating eating behavior analyses (64–67). The development of complex adaptive systems and agent-based models of complex drivers and determinants of eating behavior within varied food environments have the po-

tential to generate rich and accurate predictions of behavior (68–70). However, the volume and variety of data required to build reliable and accurate models so far remain largely unavailable in publicly available data sets owing to data-sharing challenges (44). The next step for computational researchers concerned with dietary behavior will be to apply artificial intelligence techniques to big (rich) data comprising food accessibility, preference and choice (i.e., dietary patterns), eating behavior (i.e., eating rate, satiety, satisfaction and fulfillment, setting), health profile, psychology, socioeconomic status, and various aspects of the built environment—leading to insights into obesity etiology and guiding targeted prevention measures. As until very recently these techniques and data have largely remained the property of industry, there must be increased attention paid to responsible knowledge transfer, in part via public-private partnerships, and the required ethical frameworks (71).

How Can Big Data Complement Traditional Research Approaches?

The advanced means of data acquisition described already are only beginning to be used in research and public health contexts. Although most researchers are no doubt aware of their promise, epidemiologists concerned with obesity and public health are at risk of lagging behind big tech or profit-driven data acquisition pursuits. The European Centre for Disease Prevention and Control has promoted a policy on the use of big data in epidemiology (72, 73). This approach will not only allow the pursuit of cross-cutting hypotheses but improve data quality and make resource-intensive surveillance efforts more efficient (74). Here, we discuss 3 thematic areas where the big data tools already described can complement traditional approaches in obesity research: 1) food intake and eating behavior, 2) social food environments, and 3) the built environment and physical activity. All 3 themes have the potential to utilize mobile sensing, citizen science, and artificial intelligence approaches described in the previous section, complementing and aiding traditional methods. Representative examples of specific research questions and a subset of data needs per theme are proposed in **Table 1**. The wide variety of big data types are not captured here exhaustively; this is intended to illustrate potential research avenues and stimulate already-existing disciplinary framing to adopt aspects of big data. Many variables sit across themes and their exploration will only serve to further benefit systems thinking on obesity.

1. Food intake and eating behavior—the nutritional theme

Implementing effective nutrition interventions for obesity prevention requires a more complete understanding of individual (biological) and food (nutritional, chemical, and physical) factors that influence eating behavior and appetite (75–77). These factors include both food characteristics (taste, texture, food matrix, energy density, food form, satiation and satiety, satisfaction, and fulfillment) and individual physiology (metabolic and endocrine responses, body composition, and level of physical activity). Big data have the potential to complement single-study-derived insights into food intake, eating rate, and their link to satiety. These can be linked to relevant bio-social individual characteristics such as history of breastfeeding, health comorbidities, and subjectively reported satiety and motivations for food choice. For example, mobile-supported objective measurement of eating rate was recently found to be

TABLE 1 Examples of big data contributions to obesity research, by research dimension¹

Research dimension	Research question	Data needed	Potential data sources	Methods needed	Potential outcomes
Food/nutrient intake and eating behavior—nutritional theme	How do food characteristics (taste, texture) interact with social setting to control satiety?	<ul style="list-style-type: none"> Rich food composition and features database [volume/variety] Objective measures of social setting: eating regularity and duration, family, school, and peer eating activity [volume/variety] Timing of food intake and objective/subjective satiety measures in real-time, not, e.g., daily average [velocity] 	<ul style="list-style-type: none"> National food consumption surveys EFSA food composition database (97) Publicly available nutrition and food composition applications—as documented (98, 99) 	<ul style="list-style-type: none"> Automated or semiautomated food profiling/food composition spectrometry in FAIR databases Automated meal sensing (wearable sensors/AI-driven meal picture analysis, observational restaurants) Mobile-based surveys 	<ul style="list-style-type: none"> Research: integrated biological, economic, and social sciences understanding of the determinants of food intake Innovations: novel food types with high satiety index Policies: health policy promoting specific eating practices in schools, families, and workplaces, according to healthy and sustainable FB DGs; tailored to context
Social food environments—psycho-social theme	What role does food and diet portrayal in digital media (advertising and social media) play in food and beverage intake?	<ul style="list-style-type: none"> Real-time food-intake assessments and subjective/objective determinants of satiety [volume, velocity] Social media and real-world advertising exposure logs [variety] Source of digital media advertising/marketing (e.g., via industry or from users themselves) 	<ul style="list-style-type: none"> National food consumption surveys Social media content, behavior and network data via APIs/web scraping Regional/local census and statistics bureaus 	<ul style="list-style-type: none"> Mobile/sensor-based food intake measurements Citizen/crowd-sourced mapping tools Monitoring of digital advertisements (e.g., via Web browser/application plugins) 	<ul style="list-style-type: none"> Research: role of social media, influence, and social networks in obesogenic behaviors Innovations: effective healthy diet-targeted advertisement campaigns/nudging Policies: food advertisement policies for more targeted restriction/promotion of certain foods

(Continued)

TABLE 1 (Continued)

Research dimension	Research question	Data needed	Potential data sources	Methods needed	Potential outcomes
Built environment—environmental theme	What combinations of features of the built environment influence physical activity levels and weight status?	<ul style="list-style-type: none"> • GIS-derived points of interest per region [volume/variety] • Socioeconomic, education, ability, and health characteristics [volume/variety] • Real-time, GPS-correlated physical activity measurement [volume/velocity] 	<ul style="list-style-type: none"> • National food consumption surveys • GIS/Google/Foursquare • USDA Food Environment Atlas • Regional/local census and statistics bureaus (e.g., Eurostat) • Map the Meal Gap (US) (100) • Food Environment Atlas (US) (101) • Global Physical Activity Observatory (102) 	<ul style="list-style-type: none"> • Computational tools for scraping environment characteristics, linked in real-time to user activity • Mobile-based tracking of anthropometry and activity levels 	<ul style="list-style-type: none"> • Research: relation between built environment, physical activity, health-related behaviors, and obesity prevalence • Innovations: interactive intervention/policy design tools; tools for real-time monitoring of policies and interventions • Policies: toward healthier, physical-activity-promoting, and accessible urban and suburban environments

¹ AI, artificial intelligence; API, Application Programming Interface; EFSA, European Food Safety Authority; FAIR, findable, accessible, interoperable, reusable; FBDG, food-based dietary guideline; GIS, Geographic Information Systems; GPS, Global Positioning System.

associated with higher BMI among adolescents (66, 67, 78). Libraries of food characteristics can be linked to genomics and metabolomics data sets, and/or advanced food-intake sensors, as well as data relevant to satiety. The timing of food consumption and the time occupied with food ingestion could also be linked to the level of intake and physical activity. Such integrated big data approaches would greatly enhance our understanding of how food intake is influenced or can be modified and can also help to overcome some of the issues inherent in relying on the metric of BMI as a measure of weight status, moving toward a more holistic assessment of health risk on both an individual and a population level.

2. Social food environments—the psychological theme

Branches of nutrition science and consumer behavior are typically concerned with the social determinants of food intake and eating behavior, including the social environment (e.g., eating alone compared with eating in a group) (79), in addition to availability, accessibility, affordability, and exposure to food advertising (80). Big data allow investigation and modeling of both the real-world and the social-media/digital food environments as determinants of food intake, taking into account cultural, racial, and socioeconomic background and the surrounding food environment, building on food intake data described in theme 1. Research must address the social drivers and motives that influence individuals in the mixed “real” and digital food environment, and how these drivers influence food preference, intake, satiety, and dietary profiles. Monitoring social dietary behaviors via methods more advanced than traditional surveys or observational studies will become possible, such as observational purchase data from supermarkets and restaurants or surveillance of food- and diet-related individual interactions with social media. This dimension also concerns the influence of national or regional agricultural policy and subsidies on food availability and affordability and the generation of data infrastructures able to categorize and compare these factors in relation to epidemiologic outcomes. The INFORMAS healthy food environment policy index is a critical step toward a framework within which big data on food environments can be monitored (81).

3. Built environment—the environmental theme

The design and implementation of effective public health policy addressing obesity require an understanding of what features, or combinations of features, in the built environment influence food availability and obesogenic behaviors—both food intake, as outlined in the first 2 themes, as well as physical activity. Studies of these associations typically use limited cohorts or data of insufficient granularity and variety (e.g., census data), although recent studies have made progress in linking neighborhood composition to body composition and movement behaviors (82). Big data will allow for data-driven testing of the ecological model of obesity, linking objective aspects of behavior (step and accelerometry data, GPS coordinates of locations visited) to weight status, demographic data, and detailed characteristics of the built environment such as densities of food-related points of interest and rich data on their offerings and affordability (via geo-sensing and digital mapping).

Summary

Table 1 outlines potential research questions in each of these dimensions, the required data, the big data and traditional complementary

methods needed, and potential outcomes in the realm of innovation or public health policy. These questions may serve as a conceptual starting point for epidemiologists, health professionals, and social scientists seeking to use big data in obesity research.

How Can Big Data Contribute to Evidence-Informed Policy Making? Case Study of the BigO Project

Given that many forms of data known to be highly relevant to obesity are currently out of reach of traditional epidemiologic research (e.g., mobile/GPS data; social media and digital advertisement data concerning the social food environment; rich mapping of food environments), immediate steps must involve improved interdisciplinary collaboration of computer and data scientists and engineers, led by nutrition and medical domain-experts. Big data in obesity have the potential to examine intervention outcomes or observational studies and expand causal inference of ecological associations to more complex, systems-level multivariate models, disentangling individual-level and environmental determinants in specific groups and contexts. Big data collection tools have the potential to test associations using an unbiased representation of multiple variables of interest, such as aspects of the built environment and physical activity. More excitingly, the use of big data also expands the possibility for much-needed predictive models, predicting, for example, how changes to the built environment could change physical activity and diet-related behaviors, or how changes in behavioral risk factors for obesity affect prevalence.

Responding to a lack of effective data-driven policies for the prevention of childhood obesity, BigO: Big data against childhood obesity, a Horizon2020 project, was launched in 2016. The project took action to address the lack of behavioral data processed in real-time (velocity), as well as the need for comprehensive tools for stakeholders seeking to design and implement policies. Using an mHealth and citizen science approach, BigO built a technology platform to capture environmental features and link them to individual behavior and weight status of children. The project is intended as a starting point for assessing the link between weight status and the built environment using data gathered from the platform, as well as a blueprint for future projects aiming to gather multivariate big data on diverse behaviors related to individual and environmental characteristics. To our knowledge, BigO is the only program to have collected data on physical activity and the built environment as related to weight status (BMI *z* score) within the same individual, as well as subjective/qualitative measures of health and well-being, in multiple countries and socioeconomic groups and across all BMI strata (83, 84). Figure 1 shows the scope of the data collected about individual factors, behaviors, and environmental characteristics.

The project involved >5500 children who acted as “citizen scientists,” submitting data via a smartwatch or smartphone application. The children were spread across 4 European countries, recruited via schools and obesity clinics, with weight status being both self-reported and clinic/school-measured reports. The BigO program developed a user-friendly mHealth app (MyBigO), because it was intended for use by children from 9 y of age onwards. To achieve this, a rigorous testing procedure was adopted consisting of 6 levels: code quality testing, unit testing, black box testing, integration testing, load testing, and beta testing (assimilation of both system and acceptance testing).




FIGURE 1 Variables in a mobile citizen-science approach for obesity. An example data map, modeled after the BigO system, of the types of variables pertinent to obesity able to be collected via mobile health approaches. These variables pertain to individual characteristics (demographic, anthropometric), individual behavior (physical activity, food consumption, mood), and the built environment (points of interest visited, food advertisement exposure).

This testing procedure proved effective because the app was widely accepted by both children and obesity clinicians, and successful at collecting the intended breadth and volume of data (85–87). The collected data are big in *variety*: GPS data, accelerometry, mood questionnaires, meal photos, and food advertisements encountered in their offline environment are recorded alongside anthropometry. They amount to big *volume*: the >5500 children submitted >100,000 meal photos to the platform between 2018 and 2020, >31,000 mood questionnaire responses, and >130,000 d of GPS and accelerometry recordings. The data were also processed at *velocity*: data were processed near real-time into dashboards to monitor, for example, the average activity counts per neighborhood of different municipalities, in relation to features of the built environment such as food outlets and physical activity locations. Publicly available e-dashboards were constructed, which can be found at <http://bigo.med.auth.gr:3838/>. Password-protected dashboards were available for clinic and school use, and are intended as health monitoring tools for the medical staff within schools and clinics. The public health dashboard is publicly available and is intended for eventual public health use as a policy planning tool. This technical platform fills a major gap in data-driven policy planning for obesity prevention.

Rather than starting from biological causal hypotheses of physical activity and weight status, data in BigO were gathered among a large and diverse population, which will be followed by hypothesis-generating models linking environmental characteristics, physical activity, and weight status. Making objective measurements of physical activity, linked to aspects of the built environment and weight status, all within the same individual, and taking into account the granular temporality of food intake and physical activity, promises to reveal dynamic behavior of individuals and populations in real-time (83, 88, 89). The system also has the potential to function as an intervention tool, feeding-forward rewards to incentivize the use of the app, which may have the added effect of incentivizing healthier behaviors. In this line, BigO was also conceived and implemented for use in obesity clinics to

track, manage, and eventually explain varying dimensions of weight loss on an individual level (85). Further exploitation of these data in the clinical context will provide insights into individual factors relevant to weight loss rate, maintenance, and recidivism in response to specific interventions.

Research projects and initiatives such as BigO build on the interplay between individual, social, and environmental determinants of behavior to help guide the development, monitoring, and evaluation of preventative strategies. To make such initiatives maximally relevant to public health policy, there is a need to exploit the data generated to feed advanced predictive models of individual and population determinants of BMI and eating patterns. The end goal for these types of initiatives is the development of contextually effective, ambitious data-driven policies targeting a large segment of the obesity map: individual food intake and physical activity, the school environment, and the built environment.

Limitations and Considerations

Despite its promises, several important caveats and limitations surround the use of big data, particularly when tracking multiple individual characteristics or behaviors across time and space. These limitations are discussed and supplemented with experience from the BigO project and presented further in Table 2.

Privacy concerns are significant when using mobile tools, especially among children

Although the General Data Protection Regulation, which has brought about a guarantee of location privacy, provides procedures for obtaining informed consent for the use of GPS in the age of smartphones, this is an emerging area still subject to varying degrees of stringency depending on country and institution (90). The potential involvement of young,

TABLE 2 Overcoming challenges to the use of big data in obesity research¹

Challenge	Potential solutions
Privacy	<ul style="list-style-type: none"> • Automatic local/offline prechecks built into apps for identity-violating content • Process location locally to geohashes or other aggregated values • Assign tiered data access to investigators • Involve domain experts/researchers, educational and citizens' groups (end users) in privacy co-design
User bias	<ul style="list-style-type: none"> • Align with city-lab and health equity and health promotion initiatives • Specific targeting of communities of lower socioeconomic position via schools, clinics, and community organizations • Control for access to health care
Veracity	<ul style="list-style-type: none"> • Domain-specific standards for specific types of noisy, error-prone data and volume reduction • Open-access, user-friendly tools to facilitate implementation of data standards
Legacy and FAIR data	<ul style="list-style-type: none"> • Integrate big data in nutrition and public health with emerging research infrastructures • Align emerging data and projects to the European Open Science Cloud and surveillance efforts [e.g., INFORMAS (81, 103)]
Industry vs. research interest	<ul style="list-style-type: none"> • Ethical forums, conferences, and guidelines for big data industry–academic partnerships • Research training in big data techniques and communication with technical experts

¹FAIR, findable, accessible, interoperable, reusable.

vulnerable data contributors (i.e., children and adolescents), as well as centralized, cross-border data collection schemes, further complicates matters. A powerful and innovative approach was developed within BigO to support privacy based on spatial aggregation of behavioral data, described in Diou et al. (83). Issues related to the location were resolved by processing location data locally in each mobile device before sending them to the central server and grouping them into larger geographic zones, or “geohashes.” However, all data collected by children, such as pictures related to their food intake, were at risk of violating individual privacy. For instance, some children sent photos of themselves to the server. These data were preprocessed manually, and all images deemed irrelevant or identity-revealing were deleted from the system. However, this procedure was time-consuming, and it is not a scalable solution; future efforts to incorporate automatic image recognition should help to overcome this challenge.

Big data warehouse architecture should implement privacy-aware protocols for monitoring and storing personal data related to the behaviors of a potentially vulnerable population

Individual health status and behavioral data sets must be protected and kept private. On the other hand, these data must remain analyzable and ideally reusable to extract useful knowledge that advances research and practices. The aim is to create an environment where private and sensitive data can be analyzed without revealing the identity of individuals. In BigO, an n-tier architecture to deal with the data analysis was proposed, with various defined investigator roles, where each role had a very limited view of the data, together with anonymity procedures. There also exists the challenge of acting on anonymized data collected from vul-

nerable populations, where, e.g., child protection or welfare is a concern. The development of data analysis pipelines for either encrypted or anonymized data remains a complex task.

In terms of sampling, without attention to recruitment and dissemination, citizen science tools risk including only the already healthy and wealthy, groups which are the least affected by overweight and obesity

Managing bias in sample populations in big data approaches which incorporate large populations is not something that has yet been systematically addressed in the big data approaches toward obesity to date. This issue exacerbates the already present challenge of determining causality from uncontrolled data collection. Sampling bias was a particular concern for data gathered in the BigO project, where many of the participating schools were fee-paying and richly resourced, whereas the clinical population represented children from a much wider range of socioeconomic backgrounds, with varying access to health care depending on the country.

Concerning the scope of data extracted, it is not always the case that more volume means more quality

Filtering out the appropriate level of data from big data collection methods (e.g., physical activity data sampled by the second or minute) is critical to reaching reliable conclusions. Approachable or standardized methods for both data reduction and managing data veracity are needed (e.g., low GPS accuracy, missing data from an individual not carrying their device, or errors in translation from signal to behavior). When considering large populations, engaging individuals over long periods of time to contribute data poses many logistical and tactical challenges,

as does coping with the ensuing mismatched or missing data. In the BigO project, domain expertise, gathered via a Delphi study, was essential to prioritize the most relevant variables to childhood obesity, and considerable effort was made to maximize user engagement (91). Despite the high volume of data that can be captured, many critical factors relevant to obesity remain unexplored. This is evidenced by the many remaining unmapped systems-level domains and nodes (as mapped by the Foresight report) (92) relevant to obesity, and they are not yet incorporated into big data frameworks. For example, physiologic factors (satiety, extent of digestion and metabolism, predisposition to reduced physical activity level), societal influence (peer pressure, parental control), and individual physical activity (degree of physical activity education, parental modeling of activity, physical barriers to movement) (92).

Ensuring a legacy for big data collected in public-funded research projects is also an emerging concern

Considerable amounts of quantitative, qualitative, and observational or intervention data already exist, collected for research or other purposes. Ensuring that these data are findable, accessible, interoperable, reusable (FAIR) and using them to feed existing models should therefore be a priority. Of particular concern is ensuring the interoperability and reuse (FAIR-ification) of data while maintaining individual privacy rights. The legacy of data, particularly when gathered in time-limited research projects, can also be threatened. Alignment with existing and emerging health and digital science research infrastructures, as well as conforming to the European Open Science Cloud is a first step to overcoming some of these legacy concerns. Alignment and FAIR-ification can also ensure the reuse of big data collected in the framework of health and obesity research for other domains to which it is of tremendous value, e.g., urban design, consumer behavior, and educational sciences. Ensuring the interoperability of big data in obesity is of particular interest given the emerging linkages between obesity and COVID-19. Big data mobile surveillance tools could pivot to assess the impact of lockdowns on diet, physical activity, and well-being, or ongoing efforts linking the built environment and obesity can be aligned to infection mapping (93).

Balancing obesity domain expertise with the required technical and computational expertise—academic or not—remains a key challenge in academic and policy pursuits using big data

The data analytics industry holds a monopoly on tool development, which risks overtaking nutrition and obesity research. If academic nutrition and epidemiology experts do not undertake action to shape big data use in obesity research through the lens of their domain expertise, critical health policy insights may risk being biased by profit-driven motives (25, 26). Moreover, domain experts in nutrition and epidemiology require training in the incorporation of big data, and hypothesis-generating rather than hypothesis-testing approaches. The development of agri-food policy over the past half-century has been fostered by economically driven industry motives and has (unintentionally) contributed to the overweight and obesity epidemic and its associated health costs (94–96). Now there is, therefore, a critical opportunity to change course to ensure the health of present and future generations, leveraging scientific and policy interest in the value of public goods to increase public health, well-being, and equity.

Conclusion

The collection of big and rich data on diet-related behaviors and physical activity, concerning the built environment, is critical and timely to arrive at a more advanced and integrated understanding of the inherently complex etiology of obesity. Adoption of big data approaches has not been rapid in the field of obesity, as there remains ambiguity regarding its integration into traditional approaches. The Foresight model has provided a critical first framework to conceptualize and organize the multitude of factors determining energy balance; the data-driven collection and exploration of these have the potential to lead to new hypotheses for traditional biomedical approaches (e.g., more advanced understanding of food properties' effects on metabolism) or epidemiologic approaches (e.g., multivariate, longitudinal models of environmental determinants of physical activity levels and food intake). The multicausal nature of the obesity epidemic makes it challenging to disentangle and quantify the influence of biology, behavior, and built environment, calling for an approach wider in scope than traditional epidemiologic methods alone. This can lead to advanced predictive models of the effects of public health interventions, and the ability to prioritize the most effective interventions for a given context. Critical attention and consideration must be paid to data privacy, management, reusability of data, and balancing industry with research interests to facilitate the use of big data as a public research resource. Investing in and adopting tools to allow the mapping of behavior about the social and built environment will provide a much-needed evidence base for the design and implementation of effective and contextually tailored public health policies and strategies aimed at the prevention of overweight and obesity.

Acknowledgments

The authors' responsibilities were as follows—ART: conceived of the structure and content and wrote the manuscript; MM: contributed to writing, made editorial contributions to the paper, and had primary responsibility for the final content; CD, DAL, EK, II, GOM, LA, EC, GD, KF, PK, TK, VK, IL, NM, IP, VP, IS, AS, PvV, and AD: provided input and edits on content; and all authors: read and approved the final manuscript.

References

1. Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, Imamura F, et al. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol* 2019;7(3):231–40.
2. The GBD 2015 Obesity Collaborators, Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K The GBD 2015 Obesity Collaborators, et al. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med* 2017;377(1):13–27.
3. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;390(10113):2627–42.
4. Spinelli A, Buoncristiano M, Nardone P, Starc G, Hejgaard T, Júlíusson PB, et al. Thinness, overweight, and obesity in 6- to 9-year-old children from 36 countries: the World Health Organization European Childhood Obesity Surveillance Initiative—COSI 2015–2017. *Obes Rev* 2021;22(Suppl 6):e13214.

5. Popkin BM, Du S, Green WD, Beck MA, Algaith T, Herbst CH, et al. Individuals with obesity and COVID-19: a global perspective on the epidemiology and biological relationships. *Obes Rev* 2020;21(11):e13128.
6. Frasca D, Blomberg BB. The impact of obesity and metabolic syndrome on vaccination success. *Interdiscip Top Gerontol Geriatr* 2020;43:86–97.
7. Lee A, Cardel M, Donahoo WT. Social and environmental factors influencing obesity. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K al. et editors. South Dartmouth (MA): Endotext; 2019.
8. Puhl RM, Heuer CA. Obesity stigma: important considerations for public health. *Am J Public Health* 2010;100(6):1019–28.
9. Davison KK, Birch LL. Childhood overweight: a contextual model and recommendations for future research. *Obes Rev* 2001;2(3):159–71.
10. Jebb SA, Kopelman P, Butland B. Executive summary: FORESIGHT ‘Tackling Obesity: Future Choices’ project. *Obes Rev* 2007;8(s1):vi–ix.
11. Swinburn BA, Kraak VI, Allender S, Atkins VJ, Baker PI, Bogard JR, et al. The global syndemic of obesity, undernutrition, and climate change: *The Lancet* Commission report. *Lancet* 2019;393(10173):791–846.
12. Xue H, Slivka L, Igusa T, Huang TT, Wang Y. Applications of systems modelling in obesity research. *Obes Rev* 2018;19(9):1293–308.
13. Allender S, Brown AD, Bolton KA, Fraser P, Lowe J, Hovmand P. Translating systems thinking into practice for community action on childhood obesity. *Obes Rev* 2019;20(S2):179–84.
14. Wells JCK, Marphatia AA, Amable G, Siervo M, Friis H, Miranda JJ, et al. The future of human malnutrition: rebalancing agency for better nutritional health. *Glob Health* 2021;17(1):119.
15. Walker AE, Wattick RA, Olfert MD. The application of systems science in nutrition-related behaviors and outcomes implementation research: a scoping review. *Curr Dev Nutr* 2021;5(9):nzab105.
16. Marmot M. Commentary: social determinants and the health gap: creating a social movement. *Int J Epidemiol* 2017;46(4):1335–9.
17. Ohri-Vachaspati P, DeLia D, DeWeese RS, Crespo NC, Todd M, Yedidia MJ. The relative contribution of layers of the Social Ecological Model to childhood obesity. *Public Health Nutr* 2015;18(11):2055–66.
18. Gadsby EW, Hotham S, Eida T, Lawrence C, Merritt R. Impact of a community-based pilot intervention to tackle childhood obesity: a ‘whole-system approach’ case study. *BMC Public Health* 2020;20(1):1818.
19. Kukulu K, Sarvan S, Muslu L, Yirmibesoglu SG. Dietary habits, economic status, academic performance and body mass index in school children: a comparative study. *J Child Health Care* 2010;14(4):355–66.
20. Andrade VMB, de Santana MLP, Fukutani KF, Queiroz ATL, Arriaga MB, Conceicao-Machado MEP, et al. Multidimensional analysis of food consumption reveals a unique dietary profile associated with overweight and obesity in adolescents. *Nutrients* 2019;11(8):1946.
21. Mooney SJ, Westreich DJ, El-Sayed AM. Commentary: epidemiology in the era of big data. *Epidemiology* 2015;26(3):390–4.
22. Vogel C, Zwolinsky S, Griffiths C, Hobbs M, Henderson E, Wilkins E. A Delphi study to build consensus on the definition and use of big data in obesity research. *Int J Obes* 2019;43(12):2573–86.
23. Fuller D, Buote R, Stanley K. A glossary for big data in population and public health: discussion and commentary on terminology and research methods. *J Epidemiol Community Health* 2017;71(11):1113–7.
24. Wilkins E, Aravani A, Downing A, Drewnowski A, Griffiths C, Zwolinsky S, et al. Evidence from big data in obesity research: international case studies. *Int J Obes* 2020;44(5):1028–40.
25. Mastroianni AC, Kahn JP, Kass NE. *The Oxford handbook of public health ethics*. New York (NY): Oxford University Press; 2019.
26. Xafis V, Schaefer GO, Labude MK, Brassington I, Ballantyne A, Lim HY, et al. An ethics framework for big data in health and research. *Asian Bioeth* 2019;11(3):227–54.
27. Cauchi D, Glonti K, Petticrew M, Knai C. Environmental components of childhood obesity prevention interventions: an overview of systematic reviews. *Obes Rev* 2016;17(11):1116–30.
28. Rito AI, Buoncristiano M, Spinelli A, Salanave B, Kunešová M, Hejgaard T, et al. Association between characteristics at birth, breastfeeding and obesity in 22 countries: the WHO European Childhood Obesity Surveillance Initiative—COSI 2015/2017. *Obes Facts* 2019;12(2):226–43.
29. Bixby H, Bentham J, Zhou B, Di Cesare M, Paciorek CJ, Bennett JE, et al. Rising rural body-mass index is the main driver of the global obesity epidemic in adults. *Nature* 2019;569:260–4.
30. Bamba CL, Hillier FC, Cairns JM, Kasim A, Moore HJ, Summerbell CD. How effective are interventions at reducing socioeconomic inequalities in obesity among children and adults? Two systematic reviews. *Public Health Res* 2015;3(1):Southampton (UK): NIHR Journals Library. pp. 1–446.
31. Williams J, Buoncristiano M, Nardone P, Rito AI, Spinelli A, Hejgaard T, et al. A snapshot of European children’s eating habits: results from the fourth round of the WHO European Childhood Obesity Surveillance Initiative (COSI). *Nutrients* 2020;12(8):2481.
32. Weinberg D, Stevens G, Bucksch J, Inchley J, de Looze M. Do country-level environmental factors explain cross-national variation in adolescent physical activity? A multilevel study in 29 European countries. *BMC Public Health* 2019;19(1):680.
33. Foti KE, Perez CL, Knapp EA, Kharmats AY, Sharfman AS, Arteaga SS, et al. Identification of measurement needs to prevent childhood obesity in high-risk populations and environments. *Am J Prev Med* 2020;59(5):746–54.
34. National Collaborative on Childhood Obesity Research (NCCOR). *Health, behavioral design, and the built environment White Paper*. Washington (DC): NCCOR; 2017.
35. Kirkpatrick S, Reedy J, Samuels A, Lytle L. Towards improved measurement of individual diet behaviors and food environment exposures: resources from the National Collaborative on Childhood Obesity Research (P16–026–19). *Curr Dev Nutr* 2019;3(Supplement_1):nzz050.P16–026–19.
36. Gallagher D, Visser M, Sepúlveda D, Pierson RN, Harris T, Heymsfield SB. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups? *Am J Epidemiol* 1996;143(3):228–39.
37. Liu B, Du Y, Wu Y, Snetselaar LG, Wallace RB, Bao W. Trends in obesity and adiposity measures by race or ethnicity among adults in the United States 2011–18: population based study. *BMJ* 2021;372:n365.
38. Swinburn BA, Caterson I, Seidell JC, James WP. Diet, nutrition and the prevention of excess weight gain and obesity. *Public Health Nutr* 2004;7(1A):123–46.
39. Kiraly C, Turk MT, Kalarchian MA, Shaffer C. Applying ecological frameworks in obesity intervention studies in Hispanic/Latino youth: a systematic review. *Hisp Health Care Int* 2017;15(3):130–42.
40. Broadbent A. Causation and prediction in epidemiology: a guide to the “methodological revolution”. *Stud Hist Philos Biol Biomed Sci* 2015;54:72–80.
41. Archer E, Hand GA, Blair SN. Validity of U.S. nutritional surveillance: National Health and Nutrition Examination Survey caloric energy intake data, 1971–2010. *PLoS One* 2013;8(10):e76632.
42. Brouwer-Brolsma E, Lucassen D, de Rijk M, Slotegraaf A, Perenboom C, Borgonjen K, et al. Dietary intake assessment: from traditional paper-pencil questionnaires to technology-based tools. In: Athanasiadis I, Frysinger S, Schimak G, Knibbe W, editors. *Environmental software systems. Data science in action. ISESS 2020. IFIP Advances in Information and Communication Technology*, vol. 554. Cham (Switzerland): Springer; 2020. pp. 7–23.
43. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CA. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Obesity (Silver Spring)* 2015;23(7):1331–44.
44. Giabbanelli PJ, Crutzen R. Using agent-based models to develop public policy about food behaviours: future directions and recommendations. *Comput Math Methods Med* 2017;5742629.
45. Vu T, Lin F, Alshurafa N, Xu W. Wearable food intake monitoring technologies: a comprehensive review. *Computers* 2017;6(1):4.
46. Werneck AO, Silva DR, Oyeyemi AL, Fernandes RA, Steene-Johannesen J, Andersen LB, et al. Physical activity attenuates metabolic risk of adolescents with overweight or obesity: the ICAD multi-country study. *Int J Obes* 2020;44(4):823–9.
47. Tully L, Burls A, Sorensen J, El-Moslemany R, O’Malley G. Mobile health for pediatric weight management: systematic scoping review. *JMIR Mhealth Uhealth* 2020;8(6):e16214.

48. O'Malley GC, Baker PRA, Francis DP, Perry I, Foster C. Incentive-based interventions for increasing physical activity and fitness. *Cochrane Database Syst Rev* 2012;12(1):CD009598.
49. Lucassen DA, Lasschuijt MP, Camps G, Van Loo EJ, Fischer ARH, de Vries RAJ, et al. Short and long-term innovations on dietary behavior assessment and coaching: present efforts and vision of the Pride and Prejudice consortium. *Int J Environ Res Public Health* 2021;18(15):7877.
50. He S, Li S, Nag A, Feng S, Han T, Mukhopadhyay SC, et al. A comprehensive review of the use of sensors for food intake detection. *Sens Actuators A* 2020;315:112318.
51. Dunford E, Trevena H, Goodsell C, Ng KH, Webster J, Millis A, et al. FoodSwitch: a mobile phone app to enable consumers to make healthier food choices and crowdsourcing of national food composition data. *JMIR Mhealth Uhealth* 2014;2(3):e37.
52. Vandevijvere S, Williams R, Tawfiq E, Swinburn B. A food environments feedback system (FoodBack) for empowering citizens and change agents to create healthier community food places. *Health Promot Int* 2019;34(2):277–90.
53. Lakerveld J, van der Ploeg HP, Kroeze W, Ahrens W, Allais O, Andersen LF, et al. Towards the integration and development of a cross-European research network and infrastructure: the DEterminants of DIet and Physical ACTivity (DEDIPAC) knowledge hub. *Int J Behav Nutr Phys Act* 2014;11(1):143.
54. Stok FM, Hoffmann S, Volkert D, Boeing H, Enseñauer R, Stelmach-Mardas M, et al. The DONE framework: creation, evaluation, and updating of an interdisciplinary, dynamic framework 2.0 of determinants of nutrition and eating. *PLoS One* 2017;12(2):e0171077.
55. Perelló J, Cigarini A, Vicens J, Bonhoure I, Rojas-Rueda D, Nieuwenhuijsen MJ, et al. Large-scale citizen science provides high-resolution nitrogen dioxide values and health impact while enhancing community knowledge and collective action. *Sci Total Environ* 2021;789:147750.
56. Ek A, Alexandrou C, Delisle Nyström C, Direito A, Eriksson U, Hammar U, et al. The Smart City Active Mobile Phone Intervention (SCAMPI) study to promote physical activity through active transportation in healthy adults: a study protocol for a randomised controlled trial. *BMC Public Health* 2018;18(1):880.
57. King AC, Winter SJ, Sheats JL, Rosas LG, Buman MP, Salvo D, et al. Leveraging citizen science and information technology for population physical activity promotion. *Transl J Am Coll Sports Med* 2016;1(4):30–44.
58. Rowbotham S, McKinnon M, Leach J, Lamberts R, Hawe P. Does citizen science have the capacity to transform population health science? *Crit Public Health* 2019;29(1):118–28.
59. European Statistical System Committee (ESSC). Bucharest Memorandum on official statistics in a datafied society (trusted smart statistics). Luxembourg City (Luxembourg): Eurostat; 2018.
60. Pocock MJO, Chapman DS, Sheppard LJ, Roy HE. Choosing and using citizen science: a guide to when and how to use citizen science to monitor biodiversity and the environment. Wallingford (United Kingdom): NERC/Centre for Ecology & Hydrology; 2014.
61. Pew Research Center. Smartphone ownership is growing rapidly around the world, but not always equally. Washington (DC): Pew Research Center; 2019.
62. Callaghan CT, Rowley JLL, Cornwell WK, Poore AGB, Major RE. Improving big citizen science data: moving beyond haphazard sampling. *PLoS Biol* 2019;17(6):e3000357.
63. Wiggins A, Wilbanks J. The rise of citizen science in health and biomedical research. *Am J Bioeth* 2019;19(8):3–14.
64. Kyritsis K, Fagerberg P, Ioakimidis I, Chaudhuri KR, Reichmann H, Klingelhofer L, et al. Assessment of real life eating difficulties in Parkinson's disease patients by measuring plate to mouth movement elongation with inertial sensors. *Sci Rep* 2021;11(1):1632.
65. Konstantinidis D, Dimitropoulos K, Langlet B, Daras P, Ioakimidis I. Validation of a deep learning system for the full automation of bite and meal duration analysis of experimental meal videos. *Nutrients* 2020;12:209.
66. Papapanagiotou V, Diou C, Ioakimidis I, Sodersten P, Delopoulos A. Automatic analysis of food intake and meal microstructure based on continuous weight measurements. *IEEE J Biomed Health Inform* 2019;23(2):893–902.
67. Kyritsis K, Diou C, Delopoulos A. A data driven end-to-end approach for in-the-wild monitoring of eating behavior using smartwatches. *IEEE J Biomed Health Inform* 2021;25(1):22–34.
68. Schill C, Anderies JM, Lindahl T, Folke C, Polasky S, Cárdenas JC, et al. A more dynamic understanding of human behaviour for the Anthropocene. *Nat Sustain* 2019;2(12):1075–82.
69. Sardi S, Vardi R, Meir Y, Tugendhaft Y, Hodassman S, Goldental A, et al. Brain experiments imply adaptation mechanisms which outperform common AI learning algorithms. *Sci Rep* 2020;10(1):6923.
70. Fellous JM, Sapiro G, Rossi A, Mayberg H, Ferrante M. Explainable artificial intelligence for neuroscience: behavioral neurostimulation. *Front Neurosci* 2019;13:1346.
71. Drewnowski A, Caballero B, Das JK, French J, Prentice AM, Fries LR, et al. Novel public-private partnerships to address the double burden of malnutrition. *Nutr Rev* 2018;76(11):805–21.
72. About Digital Health. Big data for epidemiology [Internet]. About Digital Health; 2019 [cited 2021 Oct 1]. Available from: <https://aboutdigitalhealth.com/2019/10/02/big-data-for-epidemiology/>.
73. European Centre for Disease Prevention and Control (ECDC). Consultation on digital technologies for public health functions. Stockholm (Sweden): ECDC; 2019.
74. Pentland A, Reid TG, Heibeck T. Big data and health: revolutionizing medicine and public health. Report of the Big Data and Health Working Group 2013. Qatar: World Innovation Summit for Health (WISH); 2013.
75. Forde CG, Mars M, de Graaf K. Ultra-processing or oral processing? A role for energy density and eating rate in moderating energy intake from processed foods. *Curr Dev Nutr* 2020;4(3):nzaa019.
76. Forde CG, Almiron-Roig E, Brunstrom JM. Expected satiety: application to weight management and understanding energy selection in humans. *Curr Obes Rep* 2015;4(1):131–40.
77. Forde CG, Fogel A, McCrickerd K. Children's eating behaviors and energy intake: overlapping influences and opportunities for intervention. *Nestle Nutr Inst Workshop Ser* 2019;91:55–67.
78. Fagerberg P, Charmandari E, Diou C, Heimeier R, Karavidopoulou Y, Kassari P, et al. Fast eating is associated with increased BMI among high-school students. *Nutrients* 2021;13(3):880.
79. Higgs S, Thomas J. Social influences on eating. *Curr Opin Behav Sci* 2016;9:1–6.
80. Boyland EJ, Nolan S, Kelly B, Tudur-Smith C, Jones A, Halford JC, et al. Advertising as a cue to consume: a systematic review and meta-analysis of the effects of acute exposure to unhealthy food and nonalcoholic beverage advertising on intake in children and adults. *Am J Clin Nutr* 2016;103(2):519–33.
81. Vandevijvere S, Barquera S, Caceres G, Corvalan C, Karupaiah T, Kroker-Lobos MF, et al. An 11-country study to benchmark the implementation of recommended nutrition policies by national governments using the Healthy Food Environment Policy Index, 2015–2018. *Obes Rev* 2019;20(S2):57–66.
82. Caetano IT, Miranda VPN, Dos Santos FK, Dos Santos Amorim PR. Adolescent's movement behaviors and built environment: a latent class analysis. *BMC Public Health* 2021;21(1):1937.
83. Diou C, Sarafis I, Papapanagiotou V, Ioakimidis I, Delopoulos A. A methodology for obtaining objective measurements of population obesogenic behaviors in relation to the environment. *Stat J IAOS* 2019;35(4):677–90.
84. Diou C, Sarafis I, Papapanagiotou V, Alagialoglou L, Lekka I, Filos D, et al. BigO: a public health decision support system for measuring obesogenic behaviors of children in relation to their local environment. *Annu Int Conf IEEE Eng Med Biol Soc* 2020:5864–7.
85. Browne S, Kechadi MT, O'Donnell S, Dow M, Tully L, Doyle G, et al. Mobile health apps in pediatric obesity treatment: process outcomes from a feasibility study of a multicomponent intervention. *JMIR Mhealth Uhealth* 2020;8(7):e16925.
86. Arthurs N, Browne S, Boardman R, O'Donnell S, Doyle G, Kechadi T, et al. The BigO application: usability and engagement among adolescents and children with obesity. European and International Congress on Obesity Online; EP-432 [abstr]. *Obes Rev* 2020;21(S1):e13118. p. 251.

87. Browne S, Boardman R, Arthurs N, O'Donnell S, Doyle G, Kechadi T, et al. A clinical portal for childhood obesity management: acceptability and usability among healthcare professionals. *European and International Congress on Obesity Online; EP-494 [abstr]*. *Obes Rev* 2020;21(S1):e13118. p. 286.
88. Sarafis I, Diou C, Delopoulos A. Behaviour profiles for evidence-based policies against obesity. *Annu Int Conf IEEE Eng Med Biol Soc* 2019:3596–9.
89. Sarafis I, Diou C, Papapanagiotou V, Alagioglou L, Delopoulos A. Inferring the spatial distribution of physical activity in children population from characteristics of the environment. *Annu Int Conf IEEE Eng Med Biol Soc* 2020:5876–9.
90. Bu-Pasha S, Alén-Savikko A, Mäkinen J, Guinness R, Korpisaari P. EU law perspectives on location data privacy in smartphones and informed consent for transparency. *Eur Data Prot Law Rev* 2016;2(3):312–23.
91. O'Donnell S, Doyle G, O'Malley G, Browne S, O'Connor J, Mars M, et al. Establishing consensus on key public health indicators for the monitoring and evaluating childhood obesity interventions: a Delphi panel study. *BMC Public Health* 2020;20:1733.
92. Morris MA, Wilkins E, Timmins KA, Bryant M, Birkin M, Griffiths C. Can big data solve a big problem? Reporting the obesity data landscape in line with the Foresight obesity system map. *Int J Obes* 2018;42(12):1963–76.
93. Filos D, Lekka I, Kilintzis V, Stefanopoulos L, Karavidopoulou Y, Maramis C, et al. Exploring associations between children's obesogenic behaviors and the local environment using big data: development and evaluation of the Obesity Prevention dashboard. *JMIR Mhealth Uhealth* 2021;9(7):e26290.
94. Elinder LS. Obesity, hunger, and agriculture: the damaging role of subsidies. *BMJ* 2005;331(7528):1333–6.
95. Hall KD. Did the food environment cause the obesity epidemic? *Obesity (Silver Spring)* 2018;26(1):11–3.
96. James WP. The fundamental drivers of the obesity epidemic. *Obes Rev* 2008;9(s1):6–13.
97. European Food Safety Authority (EFSA). EFSA Food Composition Database [Internet]. Parma (Italy): EFSA; 2021 [cited 2021 Oct 1]. Available from: <https://www.efsa.europa.eu/en/data-report/food-composition-data>.
98. Onwezen MC, Bouwman EP, van Trijp HCM. Participatory methods in food behaviour research: a framework showing advantages and disadvantages of various methods. *Foods* 2021;10(2):470.
99. Maringer M, van't Veer P, Klepacz N, Verain MCD, Normann A, Ekman S, et al. User-documented food consumption data from publicly available apps: an analysis of opportunities and challenges for nutrition research. *Nutr J* 2018;17(1):59.
100. Feeding America. Map the meal gap [Internet]. Chicago (IL): Feeding America; 2021 [cited 2021 Oct 1]. Available from: <https://map.feedingamerica.org/>.
101. USDA Economic Research Service (ERS). Food Environment Atlas [Internet]. Washington (DC): USDA ERS; 2020 [cited 2021 Oct 1]. Available from: <https://www.ers.usda.gov/data-products/food-environment-atlas/>.
102. University of California San Diego (UCSD). Global Observatory for Physical Activity [Internet]. San Diego (CA): UCSD; 2021 [cited 2021 Oct 1]. Available from: <https://new.globalphysicalactivityobservatory.com/>.
103. Swinburn B, Sacks G, Vandevijvere S, Kumanyika S, Lobstein T, Neal B, et al. INFORMAS (International Network for Food and Obesity/non-communicable diseases Research, Monitoring and Action Support): overview and key principles. *Obes Rev* 2013;14(S1):1–12.