

CrossMark
click for updatesCite this: *Chem. Sci.*, 2015, 6, 2419

Two-photon sensitive protecting groups operating via intramolecular electron transfer: uncaging of GABA and tryptophan†

Karolina A. Korzycka,^{‡a} Philip M. Bennett,^{‡a} Eduardo Jose Cueto-Diaz,^b Geoffrey Wicks,^c Mikhail Drobizhev,^c Mireille Blanchard-Desce,^b Aleksander Rebane^{cd} and Harry L. Anderson^{*a}

Improved photo-labile protecting groups, with high sensitivity to two-photon excitation, are needed for the controlled release of drugs, as tools in neuroscience and physiology. Here we present a new modular approach to the design of caging groups based on photoinduced electron transfer from an electron-rich two-photon dye to an electron acceptor, followed by scission of an ester to release a carboxylic acid. Three different electron acceptors were tested: nitrobenzyl, phenacyl and pyridinium. The nitrobenzyl system was ineffective, giving only photochemical decomposition and no release of the carboxylic acid. The phenacyl system also performed poorly, liberating the carboxylic acid in 20% chemical yield and 0.2% photochemical yield. The pyridinium system was most successful, and was tested for the release of two carboxylic acids: γ -amino butyric acid (GABA) and tryptophan. The caged GABA undergoes photochemical cleavage with a chemical yield of >95% and a photochemical yield of 1%; it exhibits a two-photon absorption cross section of 1100 GM at 700 nm, corresponding to a two-photon uncaging cross section of 10 ± 3 GM.

Received 6th December 2014

Accepted 2nd February 2015

DOI: 10.1039/c4sc03775h

www.rsc.org/chemicalscience

Introduction

Light-sensitive protecting groups have continued to gain importance as tools for investigating the role of physiologically active compounds, ever since they were first applied in biological system in 1978.¹ A number of UV- and visible-light cleavable cages have been developed² to allow rapid spatially and temporally controlled photo-release of various biomolecules within living cells.³ In the 1990s, the concept of uncaging was extended to take advantage of two-photon absorption (TPA),^{4,5} a nonlinear optical phenomenon in which excitation occurs by the simultaneous absorption of two photons, each having half the energy of the corresponding one-photon process. The main advantage of TPA is that excitation is effectively restricted to the focal volume, giving tight spatial control. Furthermore, TPA

allows the use of near-IR wavelengths which diminishes photo-damage and improves tissue penetration by reducing scattering and avoiding absorption by natural pigments. Unfortunately, most protecting groups optimized for one-photon uncaging display low efficiency under two-photon excitation, because their chromophores lack the specific features required for efficient TPA. The process of uncaging can be divided into two steps: light absorption followed by bond scission. The efficiency of TPA is quantified by the TPA cross-section (δ_a), while the quantum yield of uncaging (ϕ_u) measures the efficiency of photo-induced bond scission. The product of these two parameters, the two-photon uncaging cross-section ($\delta_u = \delta_a \phi_u$) is a figure of merit reflecting the overall sensitivity of a protecting group to two-photon uncaging. Typically, δ_u values reported to date lie between 0.05 and 2.5 GM (ref. 2c, 6, 7) (1 GM = 10^{50} cm⁴ s per photon) whereas $\delta_u > 3$ GM is desired for efficient uncaging in living cells.⁸ A common strategy for enhancing the TPA cross-section of a pre-existing protecting platform is to extend the π -system and modulate the strength of electron donating/accepting substituents. Dipolar, quadrupolar and octupolar architectures have been explored for functionalizing established caging groups such as coumarin,⁹ *o*-nitrobenzyl,^{10,11c} 2-(*o*-nitrophenyl)propyl,¹¹ phenacyl^{11c} and quinoline¹² with vinyl, phenyl, styryl, dihydronaphthalenyl, thienyl, fluorenyl and triphenylamine groups. The strategy of incorporating an existing protecting group into a conjugated donor- π -acceptor system led to a dipolar protecting group within the

^aOxford University, Department of Chemistry, Chemistry Research Laboratory, 12 Mansfield Road, Oxford, OX1 3TA, UK. E-mail: harry.anderson@chem.ox.ac.uk; Fax: +44 (0)1865 285002; Tel: +44 (0)1865 275704

^bUniversité de Bordeaux, Institut des Sciences Moléculaires, CNRS UMR 5255, 33400 Bordeaux, France

^cDepartment of Physics, Montana State University, Bozeman, MT 59717, USA

^dNational Institute of Chemical Physics and Biophysics, Tallinn 12618, Estonia

† Electronic supplementary information (ESI) available: One- and two-photon absorption spectra, emission spectra, synthetic procedures and characterization, square-wave and cyclic voltammetry, photolysis protocols. See DOI: 10.1039/c4sc03775h

‡ K.A.K. and P.M.B. made equal contribution to this work.



2-(*o*-nitrophenyl)propyl series with a record δ_u of 11 GM at 800 nm.^{11a} However, as absorption and bond scission are inherently related, any alteration in the caging group influences both δ_a and ϕ_u . Therefore some structurally modified cages, such as BNSF (2,7-bis-{4-nitro-8-[3-(2-propyl)-styryl]}-9,9-bis-[1-(3,6-dioxahexyl)]-fluorene, δ_u of 5 GM at 800 nm, 65% yield of uncaging),^{11b} suffer from light-induced side reactions or decreased yield of release, compared to their parent protecting units. Since it proves extremely difficult to enhance δ_a while preserving a high value of δ_u , several attempts have been made to explore an alternative approach, in which the absorption and release steps are decoupled and occur in different parts of the caging platform. A modular design allows each process to be optimized independently. This concept was first demonstrated for one-photon photolysis, where two spatially separated steps of uncaging were linked by intramolecular photoinduced electron transfer (PeT)¹³ or triplet sensitization.¹⁴ Recently this strategy has been implemented in two-photon uncaging systems: the fluorenyl-nitroindolyl derived protecting group (δ_u of 0.5 GM at 730 nm), where the absorption step is followed by intramolecular energy transfer-mediated release¹⁵ and 2-(*o*-nitrophenyl)propyl-thioxanthone with intermolecular FRET (δ_u of 0.86 GM at 766 nm).¹⁶ Here, we report a study of PeT-mediated uncaging in a two-photon excitable system.¹⁷ Drawing upon previously reported designs,¹³ we devised a protecting group, the removal of which operates *via* intramolecular PeT between a photoexcited electron-donor (a TPA dye with high δ_a) and an electron-acceptor (pre-existing release unit) to achieve efficient release of physiologically active compounds (Fig. 1a).

Our studies began with choosing a suitable electron-donor with high δ_a . We selected a symmetric banana-shaped bis-thiynyl fluorene (BEF) dye, in which the core is extended with substituted anilines *via* acetylene bridges.¹⁸ A pyridinium salt




Fig. 1 Concept of a photolabile protecting group operating *via* photoinduced electron transfer. Upon two-photon excitation (i) the dye unit donates an electron to the release unit (ii), which undergoes photochemical reaction and liberates the drug (iii).

was chosen as a potential electron-acceptor, since it has been demonstrated to release carboxylic acids upon PeT.^{13b,c,19} Fig. 1b shows the proposed mechanism of photo-deprotection. Light absorption generates a photosensitizer-based singlet state, which is quenched by electron transfer to the release group. The resulting charge-shifted state decays by σ -bond cleavage to liberate the physiologically active carboxylic acid. The symmetrical design was chosen to simplify the synthesis, while the aniline unit was substituted with heptaethyleneglycol chains to promote solubility in aqueous media. A model dye-unit **BEF-OH** was synthesized for the purpose of photophysical and electrochemical studies (Fig. 2). We used caged tryptophan, **BEF-Pyr-Trp**, for testing the intramolecular PeT mediated uncaging mechanism. L-Tryptophan (Trp) was selected as a model amino acid due to the presence of an indole chromophore that allows release to be quantified by HPLC, with UV detection. We also prepared caged γ -amino butyric acid (GABA), **BEF-Pyr-GABA**, to explore the utility of our protecting group for release of an inhibitory neurotransmitter.

Results and discussion

Synthesis

The reference dye **BEF-OH** was prepared as shown in Scheme 1. The synthesis started from *N*-(4-iodophenyl)diethanolamine²⁰ (**1**) by coupling with TIPS-acetylene to give intermediate **2**, which was subsequently alkylated with hexaethyleneglycol monomethyl ether tosylate. Removal of TIPS group from **3** resulted in the key aniline intermediate **4**, which was coupled with the central fluorene diiodide **5** (see ESI†) under Sonogashira conditions to give **BEF-OH**. In a convergent synthesis towards GABA and tryptophan derivatives, the fluorene core was appended with the pyridinium esters before conjugation with the substituted aniline units **4** using Sonogashira cross-coupling (Scheme 2). The final step involved removal of the Boc protecting groups to yield the trifluoroacetate salts **BEF-Pyr-GABA** and **BEF-Pyr-Trp**.

One- and two-photon absorption and fluorescence spectra

The one-photon absorption and emission spectra of **BEF-OH** in water and polar organic solvents (EtOH and THF) are shown in Fig. 3. The fluorene-based dye exhibits a strong absorption at






Fig. 2 Structure of model electron-donor **BEF-OH**, caged tryptophan, **BEF-Pyr-Trp**, and caged GABA, **BEF-Pyr-GABA**; R = (CH₂CH₂O)₇CH₃.





Scheme 1 Reagents: (i) TIPS-acetylene, Pd(OAc)₂, PPh₃, CuI, DIPA, 50 °C, 16 h, (ii) CH₃(OCH₂CH₂)₆OTs, NaH, THF, reflux, 48 h, (iii) TBAF, THF, 20 °C, 12 h, (iv) **4**, Pd(OAc)₂, PPh₃, CuI, DIPA, MeCN, 20 °C, 3 h; Ts = *p*-toluenesulfonate, TIPS = triisopropylsilyl, THF = tetrahydrofuran, DIPA = diisopropylamine, TBAF = tetra-*n*-butylammonium fluoride, R = (CH₂CH₂O)₇CH₃.



Scheme 2 Reagents: (i) MeCN, reflux, 36 h, (ii) **4**, Pd(OAc)₂, PPh₃, CuI, DIPA, 20 °C, 1 h, brine wash, (iii) BF₃·OEt₂, DCM, 0 °C → 20 °C, 5 h, (iv) MeCN, 100 °C, 24 h, (v) **4**, Pd(OAc)₂, PPh₃, CuI, DIPA, DCM, MeCN, 20 °C, 2 h, then purification in TFA-buffered solvent, (vi) HCO₂H, TIPS-H, 20 °C, 1 h.




Fig. 3 Absorption (solid) and emission (dashed) spectra of **BEF-OH** in THF (black), EtOH (red) and water (blue). Excitation wavelength: 366 nm.

300–400 nm (maximum: 380 nm; $\epsilon_{380} = 9.0 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$). Comparison of the absorption spectra in water and organic solvents reveals that the one-photon absorption properties of **BEF-OH** are independent of the solvent polarity. However, the fluorescence spectrum is sensitive to the environment, being red-shifted and broadened in polar solvents. The Stokes shifts are 78, 83 and 128 nm in THF, EtOH and water, respectively. Bathochromic shifts in the luminescence spectra of banana-shaped fluorene-dyes have previously been attributed to symmetry breaking and the formation of a polar excited state, which is stabilized in polar solvents.²¹ The fluorescence quantum yield (ϕ_f) of **BEF-OH** is about 0.42 in organic solvents (0.43 in THF, 0.41 in EtOH, referenced to quinine in 0.5 M H₂SO₄), but it falls to 0.1 in water. The TPA spectra of **BEF-OH** and its *t*-butyldimethylsilyl ether derivative, **BEF-OTBDMS** (Fig. S28[†]), in water and EtOH, respectively, are compared in Fig. 4. The TPA maxima are 1150 GM at 700 nm (for **BEF-OTBDMS** in EtOH) and 1100 GM at 715 nm (for **BEF-OH** in water). These cross-sections are similar to those reported previously for closely related dyes.^{18,22} The spectrum is slightly broader and red-shifted in water, but the spectra are similar, revealing that the TPA is insensitive to the solvent environment. In both solvents, there is a shoulder in the TPA spectrum at twice the wavelength of the one-photon allowed S₀ → S₁ transition. However, the TPA spectra are dominated by peaks corresponding to the one-photon forbidden, two-photon allowed higher-energy electronic or vibronic transitions. This behavior is similar to that reported for slightly non-centrosymmetric D- π -D quadrupolar chromophores.¹⁸

The model electron acceptor, *N*-methyl pyridinium hexafluorophosphate (**Pyr**) (Fig. 5), displays weak absorption in the UV region ($\lambda_{\text{max}} = 270 \text{ nm}$, $\epsilon_{270} = 0.5 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ in THF, Fig. S26[†]), with no significant absorption at wavelengths greater than 300 nm. The difference between the absorption spectra of the dye unit **BEF-OH** and release platform **Pyr**, mean that the fluorene dye is the only absorbing species at wavelengths longer than 300 nm.




Fig. 4 TPA spectra (circles) overlaid with double-wavelength one-photon absorption spectra (lines); blue – **BEF-OH** recorded in water, black – **BEF-OTBDMS** (*t*-butyldimethylsilyl ether derivative) recorded in EtOH.



of caged compounds under conditions identical to those of the final target application. We have not investigated how bicarbonate catalyzes this hydrolysis reaction, but the hydrolysis of α -amino acid esters under similar conditions has been attributed to CO_2 -mediated carbamate formation and intramolecular cyclization.²⁵ The low hydrolytic stability of GABA-pyridinium esters in standard aCSF buffer poses a limitation for the use of these compounds under strictly physiological conditions that will need to be addressed in future molecular designs.

The hydrolytic stability of **BEF-Pyr-Trp** was monitored only by HPLC and is different from that of **BEF-Pyr-GABA**. **BEF-Pyr-Trp** has a half-life of only about 2 h at pH 7.4 (in both PBS and NaHCO_3 based aCSF buffers). The hydrolysis proceeds more slowly at pH 3.0 (citric acid/citrate buffer), with a half-life of 3 h (Fig. S36†). The close proximity of the protonated amino group to the ester functionality probably increases the electrophilicity of the carbonyl center, enhancing the hydrolytic instability of the tryptophan derivative. **BEF-Pyr-Trp** also undergoes decomposition at low pH, in a reaction that appears to be associated with the fluorene dye rather than ester hydrolysis (Fig. S37†).

Uncaging studies

We evaluated the uncaging properties of **BEF-Pyr-Trp** and **BEF-Pyr-GABA** in a series of one-photon irradiation experiments. The method used to monitor release of the amino acid was dictated by properties of caged species. The high aqueous solubility of **BEF-Pyr-GABA** facilitated photolysis at mM concentration and we followed the reaction by ^1H NMR spectroscopy. It was not possible to monitor the release of GABA by HPLC because this amino acid lacks a UV chromophore. In contrast, **BEF-Pyr-Trp** gave broad ^1H NMR spectra at mM concentrations, presumably due to aggregation. Photolysis studies were conducted at μM concentration and the indole motif in the side chain of tryptophan enabled quantification of the released amino acid by HPLC.

One-photon uncaging of the pyridinium-based protecting group was initially investigated by irradiating a 1 mM D_2O solution of **BEF-Pyr-GABA** with a broad UV-A source (300–400 nm, peak 350 nm) in an NMR tube. At this concentration, the transmittance of the solution is negligible across the entire wavelength range of the light source. ^1H NMR spectroscopy (with *t*-butanol as an internal reference) demonstrated the release of GABA with a chemical yield of >95% (Fig. 7 and 8). Two GABA molecules are released from each molecule of **BEF-Pyr-GABA**, which demonstrates that the BEF chromophore is able to undergo two cycles of photoreduction. We were unable to identify the chemical products generated by photolysis of the caging group.

The photolysis of **BEF-Pyr-Trp** (20 μM in water) with 300–400 nm light (350 nm peak) was monitored by HPLC. Complete consumption of the starting material resulted in release of tryptophan with 83% chemical yield. No photochemically generated byproducts were detected but decomposition of the chromophore unit was observed after the uncaging events.

The quantum yield of uncaging of **BEF-Pyr-GABA** was determined by comparison with the commercially available DPNI-GABA, which has a known ϕ_u of 0.085.²⁶ Solutions of **BEF-Pyr-GABA** and DPNI-GABA (1 mM, D_2O) were irradiated




Fig. 7 ^1H -NMR spectra from a representative uncaging experiment, which shows the disappearance of caged-GABA signals with a concurrent increase of a new set of multiplets later found to be free GABA (signals "a-c"). Decrease in intensity of signal "d" (5.29 ppm) is not shown.

simultaneously (300–400 nm, peak 350 nm), and their respective rates of uncaging (k_u) were determined by ^1H -NMR (see ESI†). Under these concentrated conditions, the rate of uncaging depends only on the light intensity and the quantum yield, but not on the molar absorption coefficient. The uncaging quantum yield of **BEF-Pyr-GABA** was calculated using eqn (4) to give $\phi_u = 0.009 \pm 0.003$.

$$\phi_u(\text{BEF-Pyr-GABA}) = 0.085 \left(\frac{k_u(\text{BEF-Pyr-GABA})}{k_u(\text{DPNI-GABA})} \right) \quad (4)$$

This determination of the uncaging quantum yield of **BEF-Pyr-GABA** was verified using ferric oxalate actinometry,²⁷ which




Fig. 8 The change in concentration of **BEF-Pyr-GABA** (black squares) and free GABA (red triangles) over time. Concentration of free GABA and **BEF-Pyr-GABA** were determined by integration of signal "c" and "d" respectively relative to *t*-butanol. The presence of free GABA was confirmed by doping an irradiated solution with an authentic sample, which resulted in no new signals and an increase in the intensity of the suspected GABA signals (Fig. S30†).



gave an uncaging quantum yield of $\phi_u = 0.088 \pm 0.004$ for DPNI-GABA (in close accord with the published value) and an uncaging quantum yield of $\phi_u = 0.009 \pm 0.004$ for **BEF-Pyr-GABA**.

The fluorescence quenching experiments showed that 98% of absorption events lead to charge transfer, while the overall uncaging quantum yield is only about 1%, which suggests that decay of the excited state is dominated by back electron transfer to the ground state. Nevertheless, due to the high value of the TPA cross section (1100 GM at 700 nm), the calculated two-photon uncaging cross section for **BEF-Pyr-GABA** is $\delta_u = \delta_a \phi_u = 10 \pm 3$ GM at 700 nm, which is comparable to the highest reported value (11 GM for the 2-(*o*-nitrophenyl)propyl caged GABA).^{11a}

The uncaging quantum yield for **BEF-Pyr-Trp** was also determined by ferric oxalate actinometry. A solution of **BEF-Pyr-Trp** (~1.5 μ M, pH 3.0) was irradiated at 360 nm with a fluorimeter and photorelease of tryptophan was monitored by HPLC (Fig. 9), using a protocol designed to take account of competing background hydrolysis (see ESI†). The quantum yield of uncaging was measured for **BEF-Pyr-Trp** as $\phi_u = 0.0025$ (at 360 nm), so the calculated two-photon uncaging cross-section for **BEF-Pyr-Trp** at 720 nm is only $\delta_u = 2.5$ GM. The reasons for the difference in uncaging quantum yield between **BEF-Pyr-Trp** and **BEF-Pyr-GABA** are unclear and will require further investigation.

To prove that photo-cleavage of our new protecting group is sensitized by the absorption of the fluorene-based dye, rather than by direct excitation of pyridinium unit, we investigated the efficiency of uncaging as a function of irradiation wavelength. For this purpose, solutions of **BEF-Pyr-Trp** (1.5 μ M, pH 3.0) were irradiated at 340, 360, 380, 400 and 420 nm for 3800 s, and dark control experiments were carried out, to account for background hydrolysis (Fig. 9). The extent of uncaging correlates closely with the absorption spectrum of the BEF chromophore (Fig. 10), confirming the active role of the fluorene-dye in uncaging and PeT mediated release of tryptophan from **BEF-Pyr-Trp**.




Fig. 9 The change in concentration of tryptophan upon photolysis of **BEF-Pyr-Trp** at pH 3.0 and 360 nm; red circles – photolyzed sample, blue squares – control samples stored in the dark, black triangles – net concentration of tryptophan released upon uncaging. The concentration of tryptophan was determined by HPLC.

Preliminary experiments were carried out to test the two-photon excited release of GABA in the proximity of cultured neurons using **BEF-Pyr-GABA**. Upon two-photon excitation at 720 nm (300 fs, Ti:sapphire laser; 2–5 ms duration), we observed changes in membrane potential with kinetics consistent with activation of GABA-A receptors, indicating GABA release. Further experiments are needed to test the effectiveness of this caged compound, and to quantify side-effects, such as the biological activity of the caged drug.

Non-pyridinium designs

The pyridinium group is an excellent electron-acceptor, and the results presented above show that it gives very efficient PeT, however the efficiency of cleavage of the charge-separated state is disappointing ($\phi_u \approx 1\%$). The pyridinium group may also exhibit undesirable reactivity towards nucleophiles, so we decided to explore other electron acceptors. We chose to investigate phenacyl, for which uncaging *via* PeT has been previously reported^{13b} and nitrobenzyl esters (Fig. 11). The reduction potentials of both methyl 4-acetylbenzoate (**Phen**) and methyl 4-nitrobenzoate (**NB**) were measured by cyclic and square wave voltammetry, giving E_{RED} of **Phen**: -2.18 V; E_{RED} of **NB**: -1.47 V (vs. Fc/Fc^+ in THF with 0.1 M Bu_4PF_6). We estimated the values of the Coulombic term, eqn (2), in **BEF-Phen-Ind** and **BEF-NB-Trp** from the distance of photoinduced charge separation (a) using molecular mechanics calculations (see ESI†). This gave $w(\text{D}^+\text{A}^-) = -0.21$ eV for both compounds. Calculation of Gibbs energy of the photoinduced electron transfer (ΔG_{ET}), according to eqn (1), revealed that it is energetically favorable in both systems, giving ΔG_{ET} of -0.54 eV and -1.25 eV for **Phen** and **NB** respectively. To evaluate the photo-release properties of phenacyl and nitrobenzyl derived groups, we synthesized their tryptophan analogues for HPLC-monitored uncaging experiments. In the case of the phenacyl group, attack of the free amino group of tryptophan on the ketone group to form a 6-membered ring posed a limitation for protection of α -amino acids. This issue was overcome by use of the alternative




Fig. 10 Normalized absorption spectrum of **BEF-Pyr-Trp** in citric acid/sodium citrate buffer (pH 3.0) (black line) plotted in the scale with extent of photolysis, expressed as yield of released tryptophan upon 3800 s of irradiation of **BEF-Pyr-Trp** at 340–420 nm (blue circles).






Fig. 11 The structures of alternative electron acceptor units: methyl 4-acetylbenzoate (**Phen**) and methyl 4-nitrobenzoate (**NB**) and corresponding protecting groups based on the banana-shaped fluorene dye **BEF-Phen-Ind** and **BEF-NB-Trp**; R = $(\text{CH}_2\text{CH}_2\text{O})_7\text{CH}_3$.

structure: 3-indolepropionic acid (**Ind**), which possesses an indole chromophore but lacks the amino group, allowing us to preserve the absorption properties of tryptophan while avoiding cyclization. Structures of nitrobenzyl tryptophan (**BEF-NB-Trp**) and phenacyl protected 3-indolepropionic acid (**BEF-Phen-Ind**) are shown in Fig. 11 (for synthesis see ESI†).

Photolysis of **BEF-Phen-Ind** was investigated by irradiating a solution in water (17 μM) with a broad UV-A light source (300–400 nm; peak: 350 nm), monitoring the progress of uncaging by HPLC. 3-Indolepropionic acid was liberated with a modest 20% chemical yield, despite full consumption of the starting material. HPLC analysis did not reveal formation of any other products of photolysis and we were unable to define the fate of the remaining 80% of starting material. The quantum yield of uncaging was determined by use of ferric oxalate actinometry, giving $\phi_u = 0.0022$, corresponding to a two-photon uncaging cross section of $\delta_u = 2.4 \text{ GM}$ at 700 nm. The limited scope of substrates that can be caged (due to the reactivity of the ketone group), the low chemical yield of photorelease and the susceptibility to hydrolysis under physiological conditions make ester-linked phenacyl platforms unattractive release units.

Photolysis of **BEF-NB-Trp** was tested in a range of solvents (5 μM concentration in water, NaHCO_3 -aCSF, acetonitrile, methanol and THF) using a broad UV-A light source (300–400 nm; peak: 350 nm), but no release of tryptophan was observed. Photochemical decomposition of **BEF-NB-Trp** occurred, but it did not result in information of free tryptophan. The same result was observed when irradiation was carried out at 280 nm. Tryptophan was liberated cleanly in 75% yield by hydrolysis of **BEF-NB-Trp** in the dark over 40 h in NaHCO_3 -based aCSF buffer.

Conclusions

In this study, we have investigated an approach to the rational design of two-photon photo-labile protecting groups. A protecting group has been developed which operates by PeT between an electron-rich fluorene-based dye and a pyridinium electron-acceptor. The fluorescence of the dye is quenched by

electron transfer to the pyridinium; charge-transfer leads to bond scission, to liberate a carboxylic acid. Our protecting group has been demonstrated to release the neurotransmitter GABA and amino acid L-tryptophan upon irradiation with light of wavelength 340–420 nm, in aqueous solution in nearly quantitative chemical yields. This group exhibits a high TPA cross-section (1100 GM at 700 nm) and highly efficient charge-transfer between the electron donor and acceptor was observed (98%). The fast back electron transfer from the charge-shifted state reduces the overall quantum efficiency of uncaging to around 1% which, when combined with the TPA cross-section, results in a two-photon uncaging cross-section of approximately $\delta_u = 10 \text{ GM}$ (700 nm) for **BEF-Pyr-GABA**. Wavelength-dependent uncaging experiments confirmed electron-transfer mediated release of caged tryptophan with efficiency of release proportional to the extinction coefficient of the fluorene dye within 340–420 nm.

A key objective for future research will be to apply the modular design strategy demonstrated in this study to create an electron donor-acceptor pair for which the back electron transfer is suppressed, so that bond-scission becomes the main decay pathway. The susceptibility of pyridinium esters towards hydrolysis can lead to practical difficulties for uncaging studies in aqueous media, and it would be useful to extend these systems to non-ester linking unit that is more stable to aqueous hydrolysis, such as carbamates.

Acknowledgements

We thank Prof. Seth R. Marder for valuable discussion, Prof. Ole Paulsen, Dr Yu Zhang and Dr Golnaz Borghei for carrying out preliminary electrophysiology experiments with **BEF-Pyr-GABA**. This work was funded by the European Commission Seventh Framework Programme (TOPBIO ITN, project-grant agreement no. 264362). We thank the EPSRC Mass Spectrometry Service (Swansea) for mass spectra.

Notes and references

- J. H. Kaplan, B. Forbush III and J. F. Hoffman, *Biochemistry*, 1978, **17**, 1929.
- (a) A. P. Pelliccioli and J. Wirz, *Photochem. Photobiol. Sci.*, 2002, **1**, 441; (b) C. G. Bochet, *J. Chem. Soc., Perkin Trans. 1*, 2002, 125; (c) P. Klán, T. Šolomek, C. G. Bochet, A. Blanc, R. Givens, M. Rubina, V. Popik, A. Kostikov and J. Wirz, *Chem. Rev.*, 2013, **113**, 119; (d) H. Yu, J. Li, D. Wu, Z. Qiu and Y. Zhang, *Chem. Soc. Rev.*, 2010, **39**, 464.
- G. Mayer and A. Heckel, *Angew. Chem., Int. Ed.*, 2006, **45**, 4900.
- M. Pawlicki, H. A. Collins, R. G. Denning and H. L. Anderson, *Angew. Chem., Int. Ed.*, 2009, **48**, 3244.
- W. Denk, *Proc. Natl. Acad. Sci. U. S. A.*, 1994, **91**, 6629.
- (a) G. Bort, T. Gallavardin, D. Ogden and P. I. Dalko, *Angew. Chem., Int. Ed.*, 2013, **52**, 4526; (b) A. Specht, F. Bolze, Z. Omran, J.-F. Nicoud and M. Goeldner, *HFSP J.*, 2009, **3**, 255; (c) C. Brieke, F. Rohrbach, A. Gottschalk, G. Mayer and A. Heckel, *Angew. Chem., Int. Ed.*, 2012, **51**, 8446; (d)



- D. Warther, S. Gug, A. Specht, F. Bolze, J.-F. Nicoud, A. Mourot and M. Goeldner, *Bioorg. Med. Chem.*, 2010, **18**, 7753.
- 7 (a) O. D. Fedoryak, J.-Y. Sul, P. G. Haydon and G. C. R. Ellis-Davies, *Chem. Commun.*, 2005, 3664; (b) G. C. R. Ellis-Davies, M. Matsuzaki, M. Paukert, H. Kasai and D. E. Bergles, *J. Neurosci.*, 2007, **27**, 6601; (c) N. Gagey, P. Neveu and L. Jullien, *Angew. Chem., Int. Ed.*, 2007, **46**, 2467; (d) N. Gagey, P. Neveu, C. Benbrahim, B. Goetz, I. Aujard, J.-B. Baudin and L. Jullien, *J. Am. Chem. Soc.*, 2007, **129**, 9986; (e) T. Furuta, S. S. Wang, J. L. Dantzker, T. M. Dore, J. W. Bybee, E. M. Callaway, W. Denk and R. Y. Tsien, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 1193; (f) V. Nikolenko, R. Yuste, L. Zayat, L. M. Baraldo and R. Etchenique, *Chem. Commun.*, 2005, 1752; (g) M. Salierno, E. Marceca, D. S. Peterka, R. Yuste and R. Etchenique, *J. Inorg. Biochem.*, 2010, **104**, 418; (h) E. Fino, R. Araya, D. S. Peterka, M. Salierno, R. Etchenique and R. Yuste, *Front. Neural Circuits*, 2009, **3**, 1; (i) A. Specht, J.-S. Thomann, K. Alarcon, W. Wittayanan, D. Ogden, T. Furuta, Y. Kurakawa and M. Goeldner, *ChemBioChem*, 2006, **7**, 1690; (j) Y. Zhu, C. M. Pavlos, J. P. Toscano and T. M. Dore, *J. Am. Chem. Soc.*, 2006, **128**, 4267; (k) M. J. Davis, C. H. Kragor, K. G. Reddie, H. C. Wilson, Y. Zhu and T. M. Dore, *J. Org. Chem.*, 2009, **74**, 1721; (l) M. Petit, C. Tran, T. Roger, T. Gallavardin, H. Dhimane, F. Palma-Cerda, M. Blanchard-Desce, F. C. Acher, D. Ogden and P. I. Dalko, *Org. Lett.*, 2012, **14**, 6366.
- 8 N. I. Kiskin, R. Chillingworth, J. A. McCray, D. Piston and D. Ogden, *Eur. Biophys. J.*, 2002, **30**, 588.
- 9 (a) J. P. Olson, H.-B. Kwon, K. T. Takasaki, C. Q. Chiu, M. J. Higley, B. L. Sabatini and G. C. R. Ellis-Davies, *J. Am. Chem. Soc.*, 2013, **135**, 5954; (b) J. M. Amatrudo, J. P. Olson, G. Lur, C. Q. Chiu, M. J. Higley and G. C. R. Ellis-Davies, *ACS Chem. Neurosci.*, 2014, **5**, 64; (c) Y. Sakamoto, S. Boinapally, C. Katan and M. Abe, *Tetrahedron Lett.*, 2013, **54**, 7171; (d) C. Bao, G. Fan, Q. Lin, B. Li, S. Cheng, Q. Huang and L. Zhu, *Org. Lett.*, 2012, **14**, 572.
- 10 (a) I. Aujard, C. Benbrahim, M. Gouget, O. Ruel, J.-B. Baudin, P. Neveu and L. Jullien, *Chem.-Eur. J.*, 2006, **12**, 6865; (b) S. Boinapally, B. Huang, M. Abe, C. Katan, J. Noguchi, S. Watanabe, H. Kasai, B. Xue and T. Kobayashi, *J. Org. Chem.*, 2014, **79**, 7822.
- 11 (a) L. Donato, A. Mourot, C. M. Davenport, C. Herbivo, D. Warther, J. Léonard, F. Bolze, J.-F. Nicoud, R. H. Kramer, M. Goeldner and A. Specht, *Angew. Chem., Int. Ed.*, 2012, **51**, 1840; (b) S. Gug, F. Bolze, A. Specht, C. Bourgogne, M. Goeldner and J.-F. Nicoud, *Angew. Chem., Int. Ed.*, 2008, **47**, 9525; (c) A. Specht, F. Bolze, L. Donato, C. Herbivo, S. Charon, D. Warther, S. Gug, J.-F. Nicoud and M. Goeldner, *Photochem. Photobiol. Sci.*, 2012, **11**, 578; (d) S. Gug, S. Charon, A. Specht, K. Alarcon, D. Ogden, B. Zietz, J. Léonard, S. Haacke, F. Bolze, J.-F. Nicoud and M. Goeldner, *ChemBioChem*, 2008, **9**, 1303; (e) F. Bolze, J.-F. Nicoud, C. Bourgogne, S. Gug, X. H. Sun, M. Goeldner, A. Specht, L. Donato, D. Warther, G. F. Turi and A. Losonczy, *Opt. Mater.*, 2012, **34**, 1664.
- 12 S. Picard, E. Genin, G. Clermont, V. Hugues, O. Mongin and M. Blanchard-Desce, *New J. Chem.*, 2013, **37**, 3899.
- 13 (a) D. E. Falvey and C. Sundararajan, *Photochem. Photobiol. Sci.*, 2004, **3**, 831; (b) K. Lee and D. E. Falvey, *J. Am. Chem. Soc.*, 2000, **122**, 9361; (c) C. Sundararajan and D. E. Falvey, *Org. Lett.*, 2005, **7**, 2631.
- 14 (a) D. Wöll, J. Smirnova, W. Pfeleiderer and U. E. Steiner, *Angew. Chem., Int. Ed.*, 2006, **45**, 2975; (b) G. Papageorgiou, D. Ogden and J. E. T. Corrie, *J. Org. Chem.*, 2004, **69**, 7228; (c) G. Papageorgiou, D. Ogden and J. E. T. Corrie, *Photochem. Photobiol. Sci.*, 2008, **7**, 423.
- 15 (a) S. Picard, E. J. Cueto-Diaz, E. Genin, G. Clermont, F. Acher, D. Ogden and M. Blanchard-Desce, *Chem. Commun.*, 2013, **49**, 10805; (b) E. J. Cueto Diaz, S. Picard, V. Chevasson, J. Daniel, V. Hugues, O. Mongin, E. Genin and M. Blanchard-Desce, *Org. Lett.*, 2015, **17**, 102.
- 16 M. C. Pirrung, T. M. Dore, Y. Zhu and V. S. Rana, *Chem. Commun.*, 2010, **46**, 5313.
- 17 Related studies were reported in a PhD thesis by Jing Wang, University of Arizona, 2007; thesis advisor: S. R. Marder.
- 18 O. Mongin, L. Porrès, M. Charlot, C. Katan and M. Blanchard-Desce, *Chem.-Eur. J.*, 2007, **13**, 1481.
- 19 (a) C. Sundararajan and D. E. Falvey, *J. Org. Chem.*, 2004, **69**, 5547; (b) C. Sundararajan and D. E. Falvey, *J. Am. Chem. Soc.*, 2005, **127**, 8000; (c) C. Sundararajan and D. E. Falvey, *Photochem. Photobiol. Sci.*, 2006, **5**, 116.
- 20 C. Monnereau, E. Blart and F. Odobel, *Tetrahedron Lett.*, 2005, **46**, 5421.
- 21 F. Terenziani, A. Painelli, C. Katan, M. Charlot and M. Blanchard-Desce, *J. Am. Chem. Soc.*, 2006, **128**, 15742.
- 22 C. Katan, M. Charlot, O. Mongin, C. Le Droumaguet, V. Jouikov, F. Terenziani, E. Badaeva, S. Tretiak and M. Blanchard-Desce, *J. Phys. Chem. B*, 2010, **114**, 3152.
- 23 IUPAC. Compendium of Chemical Terminology, 2nd ed. (the "Gold Book"). Compiled by A. D. McNaught and A. Wilkinson. Blackwell Scientific Publications, Oxford (1997). XML on-line corrected version: <http://www.goldbook.iupac.org> (2006-) created by M. Nic, J. Jirat, B. Kosata; updates compiled by A. Jenkins. ISBN 0-9678550-9-8.
- 24 A. I. Ciuciu, K. A. Korzycka, W. J. Lewis, P. M. Bennett, H. L. Anderson and L. Flamigni, *Phys. Chem. Chem. Phys.*, 2015, **17**.
- 25 (a) I. M. Kovach, I. H. Pitman and T. Higuchi, *J. Pharm. Sci.*, 1981, **70**, 881; (b) R. W. Hay and L. Main, *Aust. J. Chem.*, 1968, **21**, 155.
- 26 G. Papageorgiou and J. E. T. Corrie, *Tetrahedron*, 2007, **63**, 9668.
- 27 (a) J. Goedhart and T. W. J. Gadella, *Biochemistry*, 2004, **43**, 4263; (b) H. J. Kuhn, S. E. Braslavsky and R. Schmidt, *Pure Appl. Chem.*, 1989, **61**, 187; (c) M. Montalti, A. Credi, L. Prodi and M. T. Gandolfi, *Handbook of Photochemistry*, Taylor & Francis Group, Florida, 2006, pp. 601–604.

