

Effects of Ovarian Cycle on Hemodynamic Responses during Dynamic Exercise in Sedentary Women

Hyun-Min Choi¹, Charles L. Stebbins², Hosung Nho¹, Mi-Song Kim¹, Myoung-Jei Chang¹, and Jong-Kyung Kim¹

¹Graduate School of Physical Education, Kyung Hee University, Yongin 446-701, Korea, ²Department of Internal Medicine, University of California at Davis, California 95616, USA

This study tested the hypothesis that effects of the menstrual cycle on resting blood pressure carry over to dynamic exercise. Eleven healthy females were studied during the early (EP; low estrogen, low progesterone) and late follicular (LP; high estrogen, low progesterone) menstrual phases. Stroke volume (SV), heart rate (HR), cardiac output (CO), systolic blood pressure (SBP), diastolic blood pressure (DBP), and total vascular conductance (TVC) were assessed at rest and in response to mild and moderate cycling exercise during EP and LP. During EP, compared to LP, baseline SBP (111±1 vs. 103±2 mmHg), DBP (71±2 vs. 65±2 mmHg) and mean arterial pressure (MAP) (84±2 vs. 78±1 mmHg) were higher and TVC (47.0±1.5 vs. 54.9±4.2 ml/min/mmHg) was lower ($p < 0.05$). During exercise, absolute values of SBP (Mild: 142±4 vs. 127±5 mmHg; Moderate: 157±4 vs. 144±5 mmHg) and MAP (Mild: 100±3 vs. 91±3 mmHg; Moderate: 110±3 vs. 101±3 mmHg) were also higher, while TVC was lower (Mild: 90.9±5.1 vs. 105.4±5.2 ml/min/mmHg; Moderate: 105.4±5.3 vs. 123.9±8.1 ml/min/mmHg) during EP ($p < 0.05$). However, exercise-induced increases in SBP, MAP and TVC at both work intensities were similar between the two menstrual phases, even though norepinephrine concentrations were higher during LP. Results indicate that blood pressure during dynamic exercise fluctuates during the menstrual cycle. It is higher during EP than LP and appears to be due to additive effects of simultaneous increases in baseline blood pressure and reductions in baseline TVC.

Key Words: Blood pressure, Dynamic exercise, Estrogen, Menstrual cycle, Vascular conductance

INTRODUCTION


Several studies have shown that blood pressure fluctuates during the phases of the menstrual cycle and tends to be highest at the onset of menstruation (e.g., early follicular phase) [1-4]; Moreover, these peak values are associated with time reductions in peripheral vascular reactivity [1,5]. These observations raise the possibility that the blood pressure response to exercise may also fluctuate during the menstrual cycle. In this regard, administration of estrogen can attenuate contraction-induced increases in blood pressure in cats [6] and forearm vasoconstriction in postmenopausal women [7]; suggesting that reduced levels of estrogen during menstruation may lead to abnormal blood pressure responses to exercise.

These previous studies focused on effects of isometric exercise. Effects of dynamic exercise on healthy premenopausal women may be of interest because there are differences in the pattern of the cardiovascular response to the two modes of exercise. For example, at a given level of oxygen consumption, blood pressure increases more during static compared to dynamic exercise [8]. Also, differences in muscle mass, tension production and metabolism during dynamic and static contraction may lead to differential effects on the magnitude of the corresponding blood pressure responses [9,10]. Thus, an understanding of the extent to which low levels of estrogen affect the pressor response to dynamic exercise and the hemodynamic mechanisms responsible may be clinically important as this type of activity is performed in clinical exercise testing and cardiac rehabilitation programs and has been shown to lower both resting and exercise blood pressure in postmenopausal women [11].

Results of a small cohort of previous investigations have revealed no menstrual cycle-induced differences in blood pressure during dynamic exercise. However, differences in

Received July 2, 2013, Revised September 23, 2013,
Accepted October 8, 2013

Corresponding to: Jong-Kyung Kim, Graduate School of Physical Education, Kyung Hee University, 1, Seocheon-dong, Giheung-gu, Yongin 446-701, Korea. (Tel) 82-31-201-2778, (Fax) 82-31-201-3743, (E-mail) kyung19692002@khu.ac.kr

 This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABBREVIATIONS: EP, early follicular phase; LP, late follicular phase; VO_{2max} , volume of maximal oxygen uptake; HR, heart rate; SV, stroke volume; CO, cardiac output; TVC, total vascular conductance; SBP, systolic blood pressure; DBP, diastolic blood pressure; MAP, mean arterial blood pressure; SVi, SV index; BSA, body surface area; HPLC, high-performance liquid chromatography.

methodology may have had intervening effects on the results. They include selection of time points for exercise testing where estrogen concentrations were not different [12] or where menstrual cycle effects on resting blood pressure were not seen [13,14]. Based on the totality of these observations, we tested the hypotheses that, when resting blood pressure is greater in the early versus the late follicular phase of the menstrual cycle: 1) blood pressure during dynamic exercise is also greater and 2) this greater blood pressure is associated with augmented vasoconstriction.

METHODS

Study population

Data were collected from Kyung Hee University in Yongin, Korea. All procedures were reviewed and approved by the Kyung Hee University Institutional Review Board (KHU 2011-002). All subjects signed informed consent. We studied 11 healthy women (age 25 ± 1 yr) during two phases of their ovarian cycle: days 1~4 (early follicular phase, EP) and days 10~12 (late follicular phase, LP) [15]. These time points were chosen because estrogen levels are near their lowest concentrations during days 1~4 and are near their maximal levels during days 10~12 [16]. Moreover, progesterone levels are low at both of these time points [16]. This was an important consideration, because progesterone can antagonize beneficial effects of estrogen on vascular reactivity [17].

All subjects were nonsmokers in good health who were not taking cardiovascular medications and had regular menstrual cycles as assessed by medical history. All subjects were sedentary. The order in which the cycle phase was studied was the late follicular phase followed by the early follicular phase.

Exercise protocols

Resting blood pressure was evaluated in a seated position. We obtained a minimum of 2 blood pressure measurements, which were taken 2 min apart using a sphygmomanometer. To determine the relative exercise intensities of the two workloads used in the study (40% and 60% of maximal oxygen uptake [VO_{2max}]), a maximal exercise test was performed using a cycle ergometer (Manark 828, Sweden). All subjects cycled at 0 watts for 2 min followed by increases of 30 watts every min until they could no longer maintain a pedal cadence of 60 rpm. Respiratory gases were measured using an Ultima CPX Metabolic Measurement Cart (Medgraphic, USA). The VO_{2max} obtained from this test was used as an index of functional capacity. All subjects, in turn, completed a progressive exercise test consisting of 5 min of cycling at each of the two work intensities (mild: 40% of VO_{2max} and moderate: 60% of VO_{2max}). This test was performed during both the early and late follicular phases of the ovarian cycle.

Measurement of hemodynamic variables

Stroke volume (SV) and HR were measured continuously via impedance cardiography (Physio Flow, Manatec Biomedical, France). This noninvasive device provides real-time CO data and measures hemodynamic parameters in heal-

thy subjects [8,18,19]. In brief, SV was measured by the bioimpedance method based on changes in transthoracic impedance during the cardiac cycle [20]. One set of two electrodes (one transmitting and one sensing) was placed above the supraclavicular fossa at the base of the left side of the neck. A second set was placed at the xiphoid process. A third set was used for recording the ECG. CO was determined according to the following formula: $CO = HR \times SV_i \times BSA$, where HR (heart rate) was measured from the R-R interval obtained from the first derivative of the ECG. SV_i is the SV index (i.e., $SV \div BSA$). BSA (body surface area) (m^2) was determined based on the Haycock formula: $BSA = 0.024265 \times BM^{0.5378} \times H^{0.3964}$, where BM is the body mass in kilograms and H is height in centimeters.

Brachial artery blood pressure was measured by a sphygmomanometer during mild and moderate exercise in steady-state conditions. Mean arterial blood pressure (MAP) was calculated using the formula: $MAP = [(SBP - DBP) \times 1/3] + DBP$. Total vascular conductance (TVC) was calculated as $CO \div MAP$. Blood pressure was measured by the same investigator in each subject for all interventions.

Measurement of catecholamine

To obtain blood samples for measurement of norepinephrine and epinephrine concentrations, a catheter with a 20-gauge needle was inserted into a brachial vein. Five ml of venous blood were obtained at rest and during the last min of each exercise workload. Samples were then centrifuged at 3,000 rpm for 10 min. Subsequently, plasma concentrations were isolated and stored at 5°C until analysis.

Catecholamine concentrations were determined by reverse-phase high-performance liquid chromatography (HPLC). Extraction was performed by selective absorption from aluminum oxide (Plasma-Catecholamine-Kit, BIO-RAD, Korea). Samples were then added to a 200 μ l internal standard solution + 1 ml Tris Buffer. The solution was shaken for 12 min followed by centrifugation at 2,500 rpm for 5 min. Subsequently, the supernatant (minus the alumina) was aspirated using a vacuum aspirator. The samples were then centrifuged again at 2,500 rpm for 2 min. Subsequently, 200 μ l of the supernatant was injected onto the HPLC column and eluted by mobile phase. The intra-assay coefficient of variation was 4.0% for norepinephrine and 4.0% for epinephrine.

Data analysis

Cardiac output measurements using impedance cardiography (i.e., averaging procedures in steady-state conditions) have been found to be reliable and valid in healthy subjects [8,18,19]. Accordingly, this technique accurately assesses the absolute values of CO at rest and during exercise. Effects of dynamic exercise on MAP, SBP, DBP, HR, SV, CO and TVC have been expressed as absolute values and as changes from baseline.

The average values of HR, SV, and CO at each 30 sec interval at rest and during mild and moderate exercise were used for comparison purposes. Blood pressure was assessed between the 4th and 5th min of each workload, just before collection of blood samples. Mean values of all variables were compared between EP and LP via the Student's paired t test. Means were considered to be statistically different at $p < 0.05$.

RESULTS

Subject characteristics are presented in Table 1, Fig. 1 depicts comparisons of the absolute values of the hemodynamic variables between the two menstrual phases at rest and during exercise. Resting SBP, DBP, and MAP were significantly higher in the early compared to the late follicular phase ($p < 0.05$). TVC was also lower in the early follicular phase ($p < 0.05$). Conversely, no statistically significant differences in resting HR, SV, or CO were found between these two phases. During exercise, absolute values of SBP and MAP were greater while those of TVC were lower during the early versus the late follicular phase of the menstrual cycle ($p < 0.05$). This was the case for both exercise intensities (40% and 60% of VO_{2max}). On the other hand, no differences between phases were found for DBP, HR, SV or CO during either work intensity. When the magnitude of the exercise-induced cardiovascular responses were compared (i.e., changes from rest to exercise) (Fig. 2), no differences between the two menstrual phases were found for

any of the variables during either work load. Menstrual cycle phase had no effect on plasma concentrations of epinephrine at rest or during exercise (Fig. 3). Resting levels of norepinephrine also were not statistically different between the early and late follicular phases. However, 7 of the 9 subjects tested had higher levels during the late fol-

Table 1. Physical characteristics of subjects

Variables	Subjects (N=11)
Age (yrs)	25±1.0
Height (cm)	167.3±1.6
Weight (kg)	57.5±1.3
BMI (kg/m ²)	20.5±0.4
VO _{2max} (ml/kg/min)	32±1.1

Values are mean±standard error; BMI, body mass index; HR, heart rate; SBP, systolic blood pressure; DBP, diastolic blood pressure; MAP, mean arterial pressure.




Fig. 1. Effects of the early (EP) and late (LP) follicular phases of the menstrual cycle on systolic blood pressure (SBP), diastolic blood pressure (DBP), mean arterial blood pressure (MAP), heart rate (HR), stroke volume (SV), cardiac output (CO), and total vascular conductance (TVC) at rest and during cycling exercise at 40% and 60% of maximal oxygen uptake. Values are mean±standard error. * $p < 0.05$ vs. EP.




Fig. 2. Effects of the early (EP) and late (LP) follicular phases of the menstrual cycle on changes (Δ) in systolic blood pressure (SBP), diastolic blood pressure (DBP), mean arterial blood pressure (MAP), heart rate (HR), stroke volume (SV), cardiac output (CO), and total vascular conductance (TVC) in response to cycling exercise at 40% and 60% of maximal oxygen uptake. Values are mean±standard error. * $p < 0.05$ vs. EP.




Fig. 3. Effects of the early (EP) and late (LP) follicular (LP) phases of the menstrual cycle on plasma concentrations of epinephrine and norepinephrine at rest and during cycling exercise at 40% and 60% of maximal oxygen uptake. Values are mean±standard error. * $p < 0.05$ vs. EP.

licular phase, suggesting a trend for an increase in this catecholamine. During cycling exercise, norepinephrine values were higher during the late follicular phase in response to both cycling workloads ($p < 0.05$) (Fig. 3). The magnitudes of the exercise-evoked increases in norepinephrine were also greater during the late follicular than the early follicular phases at both work intensities (40% VO_{2max} : $\Delta 175 \pm 49$ vs. $\Delta 48 \pm 9$ pg/ml; 60% VO_{2max} : $\Delta 128 \pm 49$ vs. $\Delta 611 \pm 193$ pg/ml) ($p < 0.05$).

DISCUSSION

The most interesting new finding of this study is that absolute MAP during dynamic exercise in healthy premenopausal women is greater in the early compared to the late follicular phase of the menstrual cycle. The enhanced blood pressure during cycling in the early follicular phase was most likely due to an additive effect of a concomitant elevation in resting blood pressure because the magnitude of the changes in the exercise-induced pressor responses were not different between the early and late follicular phases during either workload.

TVC at rest and during dynamic exercise was also lower in the early follicular phase, suggesting that peripheral vasoconstriction contributed to the higher blood pressure response seen during this phase of the menstrual cycle. It is not immediately clear why these apparent changes in peripheral vasoconstriction occurred. There is evidence that reactivity of peripheral arteries and microvascular arterioles improves significantly during the late follicular phase and then returns to baseline levels in early follicular phase [1,5]. This effect appears to be mediated by estrogen-induced increases in NO bioavailability and vascular reactivity that are associated with concurrent reductions in blood pressure [1,21]. Consequently, reductions in endothelial function during the early follicular phase may explain, in part, the concomitant attenuation of TVC at rest and during exercise.

Our observations differ from results of Lewandowski et al. [12], who found no differences in the blood pressure in response to progressive treadmill exercise between the pre-ovulatory follicular, (i.e., 2 days before ovulation) and the luteal (i.e., 2~3 days before menstruation) phases. However, no differences in plasma estrogen concentrations

were seen and progesterone levels were over 20 fold greater during the luteal phase.

Kim et al. [14] and Hartwich et al. [13] also reported no menstrual effects on blood pressure during cycling when the early (1~5 days) and late (10~15 days) follicular phases were compared. Unlike our study and others [1-4], they saw no menstrual cycle effects on resting blood pressure. It is not clear why effects on resting blood pressure were not seen, especially since measurements were taken at the same time points during the menstrual cycle as ours were. One possible explanation is that our subjects were sedentary and had a relative low functional capacity (i.e., low maximal O_2 uptake [VO_{2max} : 32 ± 1 ml·kg⁻¹·min⁻¹]) whereas the two previous studies enrolled subjects that had a higher functional capacity (VO_{2max} : 37 ± 2 ml·kg⁻¹·min⁻¹) and/or engaged in physical activity 2~3 days a week [13,14]. This is of note because even moderate levels of chronic physical activity can improve vascular compliance [22,23]. Consequently, the differential effects of the menstrual cycle on vascular conductance, which we observed in our sedentary subjects, may have been minimized in the more active subjects in the Kim et al. [14] and Havlik et al. [22] studies. This outcome may have offset or reduced differences in resting blood pressure between the early and late follicular phases in these subjects compared to ours.

Menstrual cycle related changes in the sympathetic nervous system also may have contributed to the differences in blood pressure and conductance during exercise that we found in our subjects. Estrogen can reduce blood pressure, and there is evidence that this effect is related to decreases in sympathetic nerve activity [24]. Unexpectedly, we found that concentrations of plasma norepinephrine during exercise were higher during the late than the early follicular phases and that a trend for this phenomenon was also seen at rest (i.e. 7 out of 9 subjects testing displayed higher norepinephrine concentrations during the LP phase). This finding suggests that sympathetic nerve activity was also augmented during this phase. While these outcomes seem counterintuitive, higher resting sympathetic nerve activity and concentrations of norepinephrine have previously been reported during the luteal versus the early follicular phases, even though luteal estrogen concentrations were also higher [25]. Since estrogen enhances the basal release of the nitric oxide [26], parallel increases in sympathetic outflow may indicate opposition to vasodilation evoked by estrogen-induced increases in nitric oxide.

Understanding differential effects of the menstrual cycle on the blood pressure response to dynamic exercise and the hemodynamic mechanisms underlying these differences may be beneficial for maximizing the effectiveness and safety of exercise interventions in premenopausal women. In this regard, results of maximal exercise testing for diagnosis of cardiovascular disease or determination of functional capacity will likely be less reliable if the phase of the menstrual cycle is not held constant. Moreover, blood pressure may reach unsafe levels in premenopausal women with preexisting hypertension or heart failure that undergo exercise testing during the late luteal or early follicular phases of the menstrual cycle. Thus, understanding intervening effects of the menstrual cycle on dynamic exercise may lead to safer and more effective exercise interventions that avoid abnormal cardiovascular responses.

Results of this study indicate that menstrual cycle effects on resting blood pressure carry over to dynamic exercise. The fact that the magnitudes of the exercised-induced in-

crease in blood pressure were not affected by the menstrual cycle suggests that elevated absolute blood pressure during the early follicular phase is an additive effect of concomitant elevations in resting blood pressure. Elevated resting and exercise blood pressure in the early follicular phase occurred in concert with lower levels of vascular conductance, which suggests that low levels of estrogen also lead to increased peripheral vasoconstriction. The mechanism responsible for the reduced TVC (i.e., enhanced vasoconstriction) may be due to decreases in endothelial function, vascular compliance and nitric oxide availability by low levels of estrogen [1,17,27].

ACKNOWLEDGEMENTS

We sincerely thank the subjects for volunteering to participate in this study.

REFERENCES

1. Adkisson EJ, Casey DP, Beck DT, Gurovich AN, Martin JS, Braith RW. Central, peripheral and resistance arterial reactivity: fluctuates during the phases of the menstrual cycle. *Exp Biol Med (Maywood)*. 2010;235:111-118.
2. Dunne FP, Barry DG, Ferriss JB, Grealy G, Murphy D. Changes in blood pressure during the normal menstrual cycle. *Clin Sci (Lond)*. 1991;81:515-518.
3. Kavitha C, Jamuna BL, Vijayakumar GS. Pressor response during normal menstrual cycle. *Int J Biol Med Res*. 2012;3:1975-1981.
4. Kelleher C, Joyce C, Kelly G, Ferriss JB. Blood pressure alters during the normal menstrual cycle. *Br J Obstet Gynaecol*. 1986;93:523-526.
5. Williams MR, Westerman RA, Kingwell BA, Paige J, Blombery PA, Sudhir K, Komesaroff PA. Variations in endothelial function and arterial compliance during the menstrual cycle. *J Clin Endocrinol Metab*. 2001;86:5389-5395.
6. Schmitt PM, Kaufman MP. Estrogen attenuates the exercise pressor reflex in female cats. *J Appl Physiol (1985)*. 2003;95:1418-1424.
7. Fadel PJ, Wang Z, Watanabe H, Arbique D, Vongpatanasin W, Thomas GD. Augmented sympathetic vasoconstriction in exercising forearms of postmenopausal women is reversed by oestrogen therapy. *J Physiol*. 2004;561:893-901.
8. Kaplan V, Bucklar GB, Bloch KE. Noninvasive monitoring of cardiac output during exercise by inductance cardiography. *Med Sci Sports Exerc*. 2003;35:747-752.
9. Hietanen E. Cardiovascular responses to static exercise. *Scand J Work Environ Health*. 1984;10:397-402.
10. Kaufman MA, Forster HV. Reflexes controlling circulatory, ventilatory, and airway responses to exercise. In: Handbook of physiology Sect. 12. Exercise: regulation and integration of multiple systems. Bethesda, 1996.
11. Swift DL, Earnest CP, Katzmarzyk PT, Rankinen T, Blair SN, Church TS. The effect of different doses of aerobic exercise training on exercise blood pressure in overweight and obese postmenopausal women. *Menopause*. 2012;19:503-509.
12. Lewandowski J, Pruszczyk P, Elaffi M, Chodakowska J, Wocial B, Switalska H, Januszewicz W, Zukowska-Grojec Z. Blood pressure, plasma NPY and catecholamines during physical exercise in relation to menstrual cycle, ovariectomy, and estrogen replacement. *Regul Pept*. 1998;75-76:239-245.
13. Hartwich D, Aldred S, Fisher JP. Influence of menstrual cycle phase on muscle metaboreflex control of cardiac baroreflex sensitivity, heart rate and blood pressure in humans. *Exp Physiol*. 2013;98:220-232.
14. Kim A, Deo SH, Fisher JP, Fadel PJ. Effect of sex and ovarian hormones on carotid baroreflex resetting and function during dynamic exercise in humans. *J Appl Physiol (1985)*. 2012;112:1361-1371.
15. Ettinger SM, Silber DH, Gray KS, Smith MB, Yang QX, Kunselman AR, Sinoway LI. Effects of the ovarian cycle on sympathetic neural outflow during static exercise. *J Appl Physiol (1985)*. 1998;85:2075-2081.
16. Boron WF, Boulpaep EL. Medical physiology: a cellular and molecular approach. Philadelphia: Elsevier Saunders; 2005.
17. English JL, Jacobs LO, Green G, Andrews TC. Effect of the menstrual cycle on endothelium-dependent vasodilation of the brachial artery in normal young women. *Am J Cardiol*. 1998;82:256-258.
18. Richard R, Lonsdorfer-Wolf E, Charloux A, Doutreleau S, Buchheit M, Oswald-Mammosser M, Lampert E, Mettauer B, Geny B, Lonsdorfer J. Non-invasive cardiac output evaluation during a maximal progressive exercise test, using a new impedance cardiograph device. *Eur J Appl Physiol*. 2001;85:202-207.
19. Tordi N, Mourot L, Matusheski B, Hughson RL. Measurements of cardiac output during constant exercises: comparison of two non-invasive techniques. *Int J Sports Med*. 2004;25:145-149.
20. Bougault V, Lonsdorfer-Wolf E, Charloux A, Richard R, Geny B, Oswald-Mammosser M. Does thoracic bioimpedance accurately determine cardiac output in COPD patients during maximal or intermittent exercise? *Chest*. 2005;127:1122-1131.
21. Choi OH, Lee SH, Kim EJ, Kim KH, Rhim BY. Expression of nitric oxide synthase and endothelin-1 in human uterine artery from full-term pregnancies. *Korean J Physiol Pharmacol*. 2005;9:165-172.
22. Havlik RJ, Phillips CL, Brock DB, Lohman K, Haskell W, Snell P, O'Toole M, Ribisl P, Vaitkevicius P, Spurgeon HA, Lakatta EG, Pullen P. Walking may be related to less vascular stiffness in the Activity Counseling Trial (ACT). *Am Heart J*. 2005;150:270-275.
23. Tanaka H, Safar ME. Influence of lifestyle modification on arterial stiffness and wave reflections. *Am J Hypertens*. 2005;18:137-144.
24. Wyss JM, Carlson SH. Effects of hormone replacement therapy on the sympathetic nervous system and blood pressure. *Curr Hypertens Rep*. 2003;5:241-246.
25. Minson CT, Halliwill JR, Young TM, Joyner MJ. Influence of the menstrual cycle on sympathetic activity, baroreflex sensitivity, and vascular transduction in young women. *Circulation*. 2000;101:862-868.
26. Sudhir K, Jennings GL, Funder JW, Komesaroff PA. Estrogen enhances basal nitric oxide release in the forearm vasculature in perimenopausal women. *Hypertension*. 1996;28:330-334.
27. Westendorp IC, Bots ML, Grobbee DE, Reneman RS, Hoeks AP, Van Popele NM, Hofman A, Witteman JC. Menopausal status and distensibility of the common carotid artery. *Arterioscler Thromb Vasc Biol*. 1999;19:713-717.