

SCIENTIFIC REPORTS



OPEN

Fitness costs of increased cataract frequency and cumulative radiation dose in natural mammalian populations from Chernobyl

Philipp Lehmann^{1,2}, Zbyszek Boratyński³, Tapio Mappes¹, Timothy A. Mousseau⁴ & Anders P. Møller⁵

Received: 11 June 2015
Accepted: 21 December 2015
Published: 27 January 2016

A cataract is a clouding of the lens that reduces light transmission to the retina, and it decreases the visual acuity of the bearer. The prevalence of cataracts in natural populations of mammals, and their potential ecological significance, is poorly known. Cataracts have been reported to arise from high levels of oxidative stress and a major cause of oxidative stress is ionizing radiation. We investigated whether elevated frequencies of cataracts are found in eyes of bank voles *Myodes glareolus* collected from natural populations in areas with varying levels of background radiation in Chernobyl. We found high frequencies of cataracts in voles collected from different areas in Chernobyl. The frequency of cataracts was positively correlated with age, and in females also with the accumulated radiation dose. Furthermore, the number of offspring in female voles was negatively correlated with cataract severity. The results suggest that cataracts primarily develop as a function of ionizing background radiation, most likely as a plastic response to high levels of oxidative stress. It is therefore possible that the elevated levels of background radiation in Chernobyl affect the ecology and fitness of local mammals both directly through, for instance, reduced fertility and indirectly, through increased cataractogenesis.

A cataract is a clouding of the lens that affects vision by increasing opacity and reducing light transmission to the retina¹. Thus cataracts reduce visual acuity of the bearer, up to the point of total blindness. In humans, cataracts generally develop as a consequence of ageing, as old lens fibres cease to be removed at the same rate as new lens fibres are being added^{2,3}. There are, however several extrinsic factors which can lead to the formation of presenile cataracts, such as smoking⁴, ultraviolet radiation⁵ or dehydration⁶. A common denominator is oxidative stress, which is related to several of the aforementioned extrinsic factors and to age, as antioxidant levels decrease with age^{7–9}. While relatively well studied in humans, the prevalence of cataracts in wild mammals, and their potential ecological significance (i.e. potential fitness costs) remains unknown. In birds, for which visual clues are key components of their life history strategies, cataracts have a significant negative effect on survival and fitness^{10,11}.

A major cause of oxidative stress in animals is ionizing radiation¹². Ionizing radiation produces free radicals that consume antioxidant stores and, consequently, lead to oxidative stress⁸. The lenses have been suggested to be particularly vulnerable to the effects of ionizing radiation^{12,13}, due to the effect of oxidative stress on the high amount of protein that makes up the bulk of the lens⁸. While acute doses of between 0.2 and 0.5 Gy have been shown to lead to detectable changes in human lenses¹⁴, there is still significant uncertainty of the relationship between acute and chronic radiation dose and cataract formation in humans^{13,15} and other mammals. In the present study we investigate if elevated frequencies of cataracts are found in eyes of bank voles (*Myodes* [= *Clethrionomys*] *glareolus*) (Schreiber) collected from natural populations in areas with varying levels of background radiation in Chernobyl Exclusion Zone¹⁶. In previous studies no major detrimental effects of ionizing

¹Centre of Excellence in Biological Interactions Research, Department of Biological and Environmental Science, P.O. Box 35, FI-40014 University of Jyväskylä, Finland. ²Department of Zoology, SE-106 91 University of Stockholm, Sweden. ³CIBIO/InBIO, Research Center in Biodiversity and Genetic Resources, University of Porto, 4485–661 Vairão, Portugal. ⁴Department of Biological Sciences, University of South Carolina, Columbia, SC 29208, USA. ⁵Laboratoire d'Ecologie, Systématique et Evolution, CNRS UMR 8079, Université Paris-Sud, Bâtiment 362, F-91405 Orsay Cedex, France. Correspondence and requests for materials should be addressed to P.L. (email: philipp.lehmann@zoologi.su.se)

	Total N	Length (mm)	Weight (g)	Molar length (mm)	Cataracts (score*)	Radiation ($\mu\text{Sv}/\text{hour}$)	Accumulated radiation (μSv)
Females	34	92.9 \pm 1.2	22.5 \pm 0.8	1.2 \pm 0.2	2.0 \pm 0.3	2.5 \pm 0.8	8020 \pm 2924
Males	54	90.8 \pm 1.1	22.2 \pm 0.6	0.9 \pm 0.1	2.1 \pm 0.2	1.4 \pm 0.3	6909 \pm 2317

Table 1. Descriptive data on morphological traits, cataract score and average background radiation levels in the collection areas and lifetime accumulated radiation dose. Years and locations were pooled. Numbers are expressed as mean \pm standard error. *0 = no cataract in either eye, 4 = large cataract in each eye.

Effect	df _(n,d)	F	P
a) Full model			
Corrected model	5, 69	14.947	<0.001
Sex	1, 69	7.437	0.008
Age	1, 69	7.387	0.008
ARD	1, 69	1.323	0.254
Sex* ARD	1, 69	5.371	0.023
Age*ARD	1, 69	0.270	0.605
b) Female			
Corrected model	2, 24	20.094	<0.001
Age	1, 24	18.761	<0.001
ARD	1, 24	5.545	0.027
c) Male			
Corrected model	2, 45	15.566	<0.001
Age	1, 45	26.723	<0.001
ARD	1, 45	0.600	0.443

Table 2. Generalized linear mixed models testing the effect of age, sex and level of accumulated radiation dose (ARD) on the presence and extent of cataracts in lenses of bank voles from Chernobyl. (a) includes all data, while (b) and (c) show results for the data split by sex. Three parameters and their interactions were estimated using restricted maximum-likelihood procedures. Sampling year and location were included as random factors in each of the models. For a) estimate \pm residual: 1.313 \pm 0.253, $z = 5.187$, $P < 0.001$, b) 1.240 \pm 0.385, $z = 3.221$, $P = 0.001$ c), 1.540 \pm 0.331, $z = 4.654$, $P < 0.001$. df_(n,d), degrees of freedom from numerator (n) and denominator (d); F, test statistic, P, probability, significant values ($P < 0.05$) are shown in bold.

radiation have been seen in abundance patterns, life history traits and biopsies of bank voles from Chernobyl^{17,18} even though these animals carry significant body burdens of radionuclides¹⁹ and variation in dorsal fur coloration has been attributed to background radiation¹⁶. In the present study we tested detrimental effects of ionizing radiation on an important, and susceptible tissue, the lens, and if the possible injuries of lenses are related to the fitness components (e.g. number of offspring) in natural mammalian populations.

Results

Cataracts were found in 57 out of 80 voles from Chernobyl (71%). Females and males had cataracts at roughly equal frequency (males: 73%, females 67%, Table 1). While the cataract score increased with age in both sexes, the influence of accumulated radiation dose on cataract score differed between female and male voles (Table 2a). Therefore the data were split by sex and re-analysed. The interaction between age and accumulated radiation dose was not significant in either sex (females: $F_{1,23} = 0.092$, $P = 0.764$; males: $F_{1,44} = 0.396$, $P = 0.533$) and removed from the final model. The frequency of cataracts was positively correlated with age in both females and males (Fig. 1A,B), and age explained a significant amount of the variance in cataract scores in both sexes (Table 2b,c). In addition, the frequency of cataracts was also positively correlated with accumulated radiation dose. However, this effect was only significant for female voles, but not for the male voles (Table 2b,c, Fig. 1C,D). Finally, in the analysis of female litter size the interaction between cataract score and age was non-significant and removed ($F_{1,10} = 4.078$, $P = 0.071$) after which the analysis was redone with only the main effects. Number of offspring in pregnant female voles was positively correlated with age and strongly negatively correlated with cataract score (Full model: $F_{2,11} = 14.916$, $P = 0.001$; Age: $F_{1,11} = 23.194$, $P = 0.001$; Cataract score: $F_{1,11} = 25.901$, $P < 0.001$; Random factors: estimate \pm residual: 0.105 \pm 0.111, $z = 0.941$, $P = 0.347$, Fig. 2). Overall all results stay unchanged if only animals from 2011 (the majority of samples) were analysed (data not shown).

Discussion

Significant frequencies of cataracts were found in voles collected from different areas in Chernobyl. The frequency of cataracts increased as a function of age, which either could be due to cumulative exposure to background radiation with age, or an inherent effect of ageing, such as, for instance, decreased antioxidant production⁷⁻⁹. A complicating factor is that the relationship between age and radiation sensitivity might be non-linear, as has




Figure 1. The upper panels show cataract scores regressed against the age estimate for (A) female and (B) male bank voles collected from Chernobyl. The lower panel shows cataract scores corrected for age (standardized residuals from a generalized linear regression on the data split by sex), regressed against the logarithm of lifetime accumulated radiation dose, for (C) female and (D) male bank voles. The symbol after the R^2 value denotes the significance level of the regression (n.s. = $P > 0.05$; * $P < 0.05$; ** $P < 0.005$; *** $P < 0.001$). Dashed lines refer to the 95% confidence interval limits of the regression (solid) line.

been shown in rats²⁰. The results nevertheless suggest that even though no major detrimental effects of ionizing radiation are seen in population (or geographic) genetic structure or life-history traits of bank voles^{17,18,19}, but see²⁰, ionizing radiation does have a significant negative effect on bank voles, seen through the induction of severe cataract formation.

While voles of both sexes had significant cataract frequencies the sexes differed in their responses to local radiation. The cataract score in female voles was positively correlated with the accumulated radiation dose, which further supports the notion that cataracts primarily are gained as a function of accumulated ionizing background radiation^{7,21}. That the correlation with accumulated radiation dose was only significant for females could be due to several reasons, such as higher baseline oxidative stress levels associated with reproductive investment^{22,23}, but see²⁴. Also female voles that were not pregnant at the time of dissection, in general had mating scars indicating previous mating efforts. In humans it has been suggested that females in general have higher cataract prevalence than males^{25–27}. It is also possible that male voles suffer more strongly from cataracts, due to higher movement activity and increased predation risk²⁸, than females and therefore, due to high mortality, were under-represented in the current dataset. However, since females show a strong negative relationship between cataract score and fecundity (Fig. 2), the first hypothesis, stronger negative effects on females as a function of increased reproductive effort, seems more plausible. It must be stated that increased radiation might also directly lower fertility^{29,30} and indeed, also lifetime accumulated radiation was negatively correlated with fecundity (results not shown). Thus




Figure 2. Litter size in female bank voles from Chernobyl was significantly negatively correlated with age and cataract score. The figure shows a regression between litter size and the Pearson residual from a regression between age and cataract score. The symbol after the R^2 value denotes the significance level of the regression ($***P < 0.001$). Dashed lines refer to the 95% confidence interval limits of the regression (solid) line.

there might be multiple effects on fertility due to radiation. Finally it should be noted that while the effect of radiation was significant only in females, lens deformities were discovered in the majority of male voles (73%). The average background radiation level in Europe varies, but is around $0.27 \mu\text{Sv/h}^{12}$. Compared to that, most areas, even relatively clean ones, added significant background radiation to the average (the median value of $0.274 \mu\text{Sv/h}$ amounts to a 50% increase in the level of ionizing background radiation). Thus, even though no correlation was seen in male voles between age corrected cataract score and local background radiation, the overall high frequency of cataracts does support the notion that also male voles in Chernobyl suffer from increased prevalence of lens deformities. It must finally be stated that short-lived animals such as voles with high general mortality, especially in males, makes partitioning out radiation-specific mortality challenging.

The lifetime accumulated chronic doses of ionizing radiation estimated in the present study for voles, which had lens deformities (cataract score ≥ 1), were on average $0.01 \pm 0.003 \text{ Sv}$ (min: 0.0002, max: 0.08). In a study on human children exposed to comparably small cumulative doses of radiation from Chernobyl as in the current study (between 0.029 to 0.086 Sv) a small (3.6%) but significantly increased proportion showed lens changes when compared to non-exposed children³¹, see also¹⁵. In humans the threshold dose currently estimated for radiation cataractogenesis is 0.5 Sv for acute exposures³². Thus, the present study suggests that voles might develop cataracts at lower accumulated radiation doses than humans as a consequence of chronic exposure to ionizing radiation (Table 3). This could be due to several reasons. Firstly, the method used to assess eye deformities in the current study, by microscopically examining dissected lenses, could be more sensitive in detecting small eye lesions than in many large scale human studies where eyes are investigated using non-invasive clinically ascertained methods e.g.²⁷ or slit-lamp exams e.g.¹⁵. Furthermore, since radiation induced cataract formation depends on the rate at which damaged lens epithelial cells divide, differentiate and migrate³³, small mammals with high metabolic rate and short lifespan might have lower response thresholds than humans. Also in other animal models, such as the mouse and rat, significantly lower response thresholds for cataract formation have been documented than for humans³⁴. A second, methodological issue relates to how lifetime accumulated radiation dose was calculated. In the current study the background radiation rate (in $\mu\text{Sv/h}$) was multiplied with estimated lifespan in hours. This yields a dose reflecting accumulated background radiation on a very local scale, since background radiation reflects trap location. While not strongly dispersing, voles can move significant distances in search of mates, while foraging or to find better habitats^{28,35}. Thus the dose measure might not reflect individual variation to the extent needed for accurate calculations on dose-response thresholds. This remains an important future task. Also more careful partitioning of sex-specific effects is important, since male and female voles differ in several behavioural and life-history characters, including dispersal, home range size and predation risk^{28,35,36} which could influence how they are exposed to, and accumulate radiation as well as how cataracts influence survival.

The ecological significance of the high frequency of cataracts in voles from Chernobyl is challenging to deduce, since vision is less important than other senses in bank voles. Indeed, sensory dependence differences and its consequences for individual fitness might be a potential explanation for why higher frequencies of cataracts are recorded in bank voles than in live birds from Chernobyl, which rely more on vision¹¹. However, the negative correlation between litter size and cataract seen in breeding females suggests that cataracts may have a negative impact on fitness in wild bank voles (Fig. 2) although at this stage we cannot rule out the possibility that

Reference	Species	Time since exposure	Numbers in cohort	Dose range (Sv)	Odds ratio Sv ⁻¹ (lower, upper 95% CI)
Hsieh et al. 2010 ¹⁵	<i>Homo sapiens</i>	7 years	73	0.005–0.06	Increased risk, but OR not calculated.
Day et al. 1995 ³¹	<i>Homo sapiens</i>	5–7 years	991	0.030	Increased risk, but OR not calculated.
Chodick et al. 2008 ⁴⁰	<i>Homo sapiens</i>	~19 years	35705	0.005–0.06	1.98 (-0.69, 4.6) ^a
Neriishi et al. 2007 ⁴¹	<i>Homo sapiens</i>	55–57 years	3761	0–3	1.39 (1.24, 1.55)
Worgul et al. 2007 ⁴²	<i>Homo sapiens</i>	12–14 years	8607	0–1	1.65 (1.18, 2.30)
Rafnsson et al. 2005 ⁴³	<i>Homo sapiens</i>	Working life	445	0–0.48	3.02 (1.44, 6.35)
Markiewicz et al. 2015 ⁴⁴	<i>Mus musculus</i>	1–10 months	144	0–2	Dose response peak at 0.5 Sv ^b , but OR not calculated.
Current study	<i>Myodes glareolus</i>	Chronic exposure	88	0.0002–0.08	1.69 (0.81, 3.53) ^c

Table 3. Summary of results from studies measuring eye deformities in response to low doses of radiation compared to current study. ^aExcess relative risk. ^bAt 500 mGy. ^cThe logarithm (log₁₀) of the predictor, life-time accumulated background radiation in μSv, was used to generate the odds ratio.

radiation exposure affects both traits independently. Given the results presented here for a rodent and past related studies on birds, it is likely that elevated frequencies of cataracts have a profound effect on the ecology of animal populations affected by nuclear accidents.

Methods

Study animals, study areas, general morphology and fecundity. Study animals were captured with Ugglan multiple-capture live traps (Grahnb, Sweden) baited with sunflower seeds and potato, during the summers of 2011, 2013 and 2014, from 41 locations in the Chernobyl region of Ukraine¹⁶. In 2011, 64 voles (26 females, 38 males), in 2013, 12 voles (5 females, 7 males) and in 2014, 12 voles (3 females, 9 males) were captured. Animals were killed by cervical dislocation and stored at –20 °C on site. All procedures were performed in accordance with relevant guidelines and regulations. The study was approved by the Finnish Ethical Committee (license numbers: ESLH-2008-04660/Ym-23 and ESLH-2009-09663/Ym-23). Vole length was measured as the ventral distance between the tip of the snout to the anus of the straightened animal (to the nearest 0.1 mm), while weight of the animal was measured after removal of embryos if present (Mettler Toledo XS204, to the nearest 0.1 g). Number of offspring (embryos) was counted from all pregnant females (14 in total).

Measurements of background radiation. Soil radiation level at the exact trap location was measured with a hand-held dosimeter (Inspector, SE International, Inc, Summertown, TN, USA) calibrated to measure Sieverts (Sv). The average radiation varied between trapping sites from 0.05 to 59.70 μSv/hour. The actual dose in individuals is likely to be strongly correlated to these “residential” measurements although the effects of ingestion of foods of varying activity levels is likely to increase the variance of doses considerably³⁷.

Detection and quantification of cataracts. Cataracts were identified through visual inspection of lenses dissected from eyes of voles under a preparation microscope (Olympus SZ-CTV). Dissections were carefully performed in distilled water in 2013 (animals collected 2011) and 2014 (animals collected 2013 and 2014) after thawing animals on ice. After removal, the lens was placed on millimetre paper on top of a line. The presence and size of cataracts were divided into three categories: 0 = no cataracts, line on paper clearly visible, 1 = small cataract, line partially visible, 2 = large cataract, line not visible¹⁰. Eye scores were combined, and thus each animal got a cataract score from 0 to 4, with 0 representing no cataracts and 4 representing large cataracts in both eyes. Uncertain cases (13 animals with eye or head damage) were removed from analysis. No bias between the right and left eye was seen (paired sample t-test: $t_{74} = -0.363$, $P = 0.718$).

Estimation of age and lifetime accumulated radiation. For age determinations the first molar of the right and left mandibles was removed and the length of the oral root from the upper part of the neck of the tooth measured under a calibrated stereomicroscope (Olympus SZ61) and converted to mm at a resolution of 0.01 mm. Since teeth wear out with time, this measure is strongly positively correlated with age in bank voles^{36,38}. We estimated the age of voles in hours with the following equation, from³⁸:

$$y = 0.186x - 0.027 \quad (1)$$

where y is root length in mm and x the age in months (30 day periods). The lifespan was converted from months to hours and multiplied with radiation in μS/h to produce an estimate of lifetime accumulated radiation.

Statistical analyses. The data was first analysed with a generalized linear mixed model with cataract score as dependent variable, sex as factor and age (in hours) and the logarithm of accumulated radiation dose (log₁₀μSv) as

covariates. The main effects as well as the interactions age*radiation and sex*radiation were included. Collection year and location were added as random (block) factors. Following this analysis the data were split by sex, and the cataract score then re-analysed as described above. The non-significant interaction between age*accumulated radiation was removed from the final models³⁹. A generalized linear mixed model was employed since the cataract score data was slightly overdispersed due to zero-inflation. The influence of cataracts on number of offspring in reproducing females was tested with a generalized linear mixed effect model. Number of offspring was included as target variable, while age and cataract score were added as explanatory variables. Collection year and location were added as random (block) factors. All statistical tests were performed with the IBM SPSS 20.0 software (IBM SPSS Inc., Chicago, IL, USA).

References

- Riordan-Eva, P. & Cunningham, E. T. Jr. (eds) *Vaughan & Asbury's General Ophthalmology*. (The McGraw-Hill Companies, Inc. USA, 2011).
- Taylor, H. R. Epidemiology of age-related cataract. *Eye*. **13**, 445–448 (1999).
- Allen, D. & Vasavada, A. Cataract and surgery for cataract. *BMJ*. **33**, 128–132 (2006).
- Kelly, S. P., Thornton, J., Edwards, R., Sahu, A. & Harrison, R. Smoking and cataract: review of causal association. *J. Cataract Refract. Surg.* **31**, 2395–2404 (2005).
- Taylor, H. R. *et al.* Effect of ultraviolet radiation on cataract formation. *N. Engl. J. Med.* **319**, 1429–1433 (1988).
- Minassian, D. C., Mehra, V. & Verrey, J. D. Dehydrational crises: a major risk factor in blinding cataract. *Br. J. Ophthalmol.* **73**, 100–105 (1989).
- Collman, G. W., Shore, D. L., Shy, C. M., Checkoway, H. & Luria, A. S. Sunlight and other risk factors for cataracts: an epidemiologic study. *Am. J. Public Health*. **78**, 1459–1462 (1988).
- Taylor, A., Jacques, P. F. & Epstein, E. M. Relations among aging, antioxidant status, and cataract. *Am. J. Clin. Nutr.* **62**, 1439S–1447S (1995).
- Costantini, D. (ed.) *Oxidative stress and hormesis in evolutionary ecology and physiology*. (Springer-Verlag, Berlin Heidelberg, 2014).
- Galvan, I., Erritzøe, J., Wakamatsu, K. & Møller, A. P. High prevalence of cataracts in birds with pheomelanin-based colouration. *Comp. Biochem. Physiol. A*. **162**, 259–264 (2012).
- Mousseau, T. A. & Møller, A. P. Elevated frequency of cataracts in birds from Chernobyl. *PLoS ONE*. **8**, e66939 (2013).
- UNSCEAR. *Sources and effects of ionizing radiation*. In: *Effects*. Vol. 2. (United Nations, New York, NY, 2008)
- International Commission on Radiological Protection. ICRP statement on tissue reactions and early and late effects of radiation in normal tissues and organs—threshold doses for tissue reactions in a radiation protection context. ICRP publication 118. *Ann. ICRP* **41**, 1–322 (2012).
- Ainsbury, E. A. *et al.* Radiation cataractogenesis: a review of recent studies. *Radiat. Res.* **172**, 1–9 (2009).
- Hsieh, W. A. *et al.* Lens Opacities in Young individuals Long after Exposure to Protracted Low-Dose-Rate γ Radiation in 60Co-Contaminated buildings in Taiwan. *Rad. Research*. **173**, 197–204 (2010).
- Boratynski, Z., Lehmann, P., Mappes, T., Mousseau, T. A. & Møller, A. P. Increased radiation from Chernobyl decreases the expression of red colouration in natural populations of bank voles (*Myodes glareolus*). *Sci. Rep.* **4**, 7141 (2014).
- Baker, R. J. *et al.* Consequences of polluted environments on population structure: the bank vole (*Clethrionomys glareolus*) at Chernobyl. *Ecotoxicology*. **10**, 211–216 (2001).
- Rodgers, B. E., Wickliffe, J. K., Phillips, C. J., Chesser, R. K. & Baker, R. J. Experimental exposure of naïve bank voles (*Clethrionomys glareolus*) to the Chernobyl, Ukraine, environment: a test of radioresistance. *Environ. Toxicol. Chem.* **20**, 1936–1941 (2001).
- Chesser, R. K. *et al.* 2000 Concentrations and dose rate estimates of 124, 137cesium and 90strontium in small mammals at Chernobyl, Ukraine. *Environ. Toxicol. Chem.* **19**, 305–312 (2000).
- Merriam, G. R. & Szechter, A. The relative radiosensitivity of rat lenses as a function of age. *Rad. Research*. **62**, 488–497 (1975).
- Møller, A. P. & Mousseau, T. The effect of natural variation in background radioactivity on humans, animals and other organisms. *Biol. Rev.* **88**, 226–254 (2013).
- Casanueva, E. & Viteri, F. E. Iron and oxidative stress in pregnancy. *J. Nutr.* **133**, 1700S–1708S (2003).
- Mutinati, M. *et al.* Oxidative stress during pregnancy in the sheep. *Repr. Dom. Anim.* **48**, 353–357 (2013).
- Oldakowski, L. *et al.* Is Reproduction costly? No increase of oxidative damage in breeding bank voles. *J. Exp. Biol.* **215**, 1799–1805 (2012).
- The Eye Diseases Prevalence Research Group. Prevalence of cataract and pseudophakia/aphakia among adults in the United States. *Epidemiology*. **122**, 487–494 (2004).
- Klein, B. E. K., Klein, R., Lee, K. E. & Gangnon, R. E. Incidence of age-related cataract over a 15-year interval. *Ophthalmology*. **155**, 477–482 (2008).
- Neriishi, K. *et al.* Radiation dose and cataract surgery incidence in atomic bomb survivors, 1986–2005. *Radiology*. **265**, 167–174 (2012).
- Mappes, T., Halonen, M., Suhonen, J. & Ylönen, H. Selective avian predation on population of field vole *Microtus agrestis*: greater vulnerability of males and subordinates. *Ethol. Ecol. & Evol.* **5**, 519–527 (1993).
- International Commission on Radiological Protection. Biological effects after prenatal irradiation (Embryo and Fetus). ICRP publication 90. *Ann. ICRP* **41**, 1–322 (2012).
- Little, M. P. A review of non-cancer effects, especially circulatory and ocular diseases. *Radiat. Environ. Biophys.* **52**, 435–449 (2013).
- Day, R., Gorin, M. B. & Eller, A. W. Prevalence of lens changes in Ukrainian children residing around Chernobyl. *Health. Phys.* **68**, 632–642 (1995).
- Kleiman, N. J. Radiation cataract. *ICRP Proceedings*. **41**, 80–97 (2012).
- Worgul, B. V. & Rothstein, H. Radiation cataract and mitosis. *Ophthalm. Res.* **7**, 21–32 (1975).
- Worgul, B. V., Smilenov, L. & Brenner, D. J. Mice heterozygous for the ATM gene are more sensitive to both X-ray and heavy ion exposure than are wildtypes. *Adv. Space. Res.* **35**, 254–259 (2005).
- Sandell, M., Agrell, J., Erlinge, S. & Nelson, J. Natal dispersal in relation to population density and sex ratio in the field vole, *Microtus agrestis*. *Oecologia*. **83**, 145–149 (1990).
- Meri, T., Halonen, M., Mappes, T. & Suhonen, J. Younger bank voles are more vulnerable to avian predation. *Can. J. Zool.* **86**, 1074–1078 (2008).
- Garnier-Laplace, J. *et al.* Are radiosensitivity data derived from natural field conditions consistent with data from controlled exposures? A case study of Chernobyl wildlife chronically exposed to low dose rates. *J. Environ. Rad.* **121**, 12–21 (2013).
- Gustafsson, T. O., Andersson, C. B. & Westlin, L. M. Determining the age of Bank voles - a laboratory study. *Acta. Theriol.* **27**, 275–282 (1982).
- Sokal, R. R. & Rohlf, F. J. (eds) *Biometry, the principles and practice of statistics in biological research*. (W.H. Freeman and Company, New York, 2003).
- Chodick, G. *et al.* Risk of cataract after exposure to low doses of ionizing radiation: a 20-year prospective cohort study among US radiologic technologists. *Am. J. Epidemiol.* **168**, 620–631 (2008).

41. Neriishi, K. *et al.* Postoperative cataract cases among atomic bomb survivors: radiation dose response and threshold. *Radiat. Res.* **168**, 404–408 (2007).
42. Worgul, B. V. *et al.* Cataracts among Chernobyl clear-up workers: implications regarding permissible eye exposures. *Radiat. Res.* **167**, 233–243 (2007).
43. Rafnsson, V. *et al.* Cosmic radiation increases the risk of nuclear cataract in airline pilots: a population-based case-control study. *Arch. Ophthalmol.* **123**, 1102–1105 (2005).
44. Markiewicz, E. *et al.* Nonlinear ionizing radiation-induced changes in eye lens cell proliferation, cyclin D1 expression and lens shape. *Open Biol.* **5**, 150011 (2015).

Acknowledgements

The authors wish to thank Igor Chizhevsky and Gennadi Milinevski for logistic support and help. The project was funded by the Academy of Finland to T. Mappes (project number 268670). Z. Boratyński is a postdoctoral grantee from the Portuguese Foundation for Science and Technology (RH/BPD/84822/2012).

Author Contributions

All authors planned the study. Z.B., T.M., T.A.M. and A.P.M. participated in field work. P.L. and T.M. took phenotypic measurements, prepared the dataset and conducted initial analyses. P.L. conducted final statistical analyses and prepared the first version of the manuscript. All authors prepared the final version of the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Lehmann, P. *et al.* Fitness costs of increased cataract frequency and cumulative radiation dose in natural mammalian populations from Chernobyl. *Sci. Rep.* **6**, 19974; doi: 10.1038/srep19974 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>