

SURVEY

Prevalence of Malocclusion among 10–12-year-old Schoolchildren in Kozhikode District, Kerala: An Epidemiological Study

¹Retna Kumari Narayanan, ²MT Jeseem, ³TV Anupam Kumar

ABSTRACT

Background: A malocclusion is an irregularity of the teeth or a malrelationship of the dental arches beyond the range of what is accepted as normal.

Objectives: To determine the prevalence of malocclusion in children aged 10–12 years in Kozhikode district of Kerala, South India.

Materials and methods: A descriptive cross-sectional study was conducted among schoolchildren aged 10–12 years in six schools in Kozhikode district of Kerala, South India. A total of 2,366 children satisfied the inclusion criteria. Occlusal characteristics like crossbite, open bite, deep bite, protrusion of teeth, midline deviations, midline diastema and tooth rotation were recorded. The data were tabulated and analyzed using Chi-square test.

Results: The results revealed that the overall prevalence of malocclusion was 83.3%. Of this, 69.8% of the children had Angle's class I malocclusion, 9.3% had class II malocclusion (division 1 = 8.85%, division 2 = 0.5%) and 4.1% had class III malocclusion; 23.2% showed an increased overjet (>3 mm), 0.4% reverse overjet, 35.6% increased overbite (>3 mm), 0.29% open bite, 7.2% crossbite with 4.6% crossbite of complete anterior teeth, 63.3% deviation of midline, 0.76% midline diastema and 3.25% rotated tooth. No significant differences in gender distributions of malocclusions were noted except for increased overjet and overbite.

Conclusion: There is high prevalence of malocclusion among schoolchildren in Kozhikode district of Kerala. Early interception and early correction of these malocclusions will eliminate the potential irregularities and malpositions in the developing dentofacial complex.

Keywords: Crossbite, Malocclusion, Prevalence.

How to cite this article: Narayanan RK, Jeseem MT, Kumar TVA. Prevalence of Malocclusion among 10–12-year-old Schoolchildren in Kozhikode District, Kerala: An Epidemiological Study. *Int J Clin Pediatr Dent* 2016;9(1):50-55.

Source of support: Nil

Conflict of interest: None

¹Professor and Head, ²Postgraduate Student, ³Associate Professor

¹⁻³Department of Pedodontics and Preventive Dentistry Government Dental College, Kozhikode, Kerala, India

Corresponding Author: Retna Kumari Narayanan, Professor and Head, Department of Pedodontics and Preventive Dentistry Government Dental College, Kozhikode, Kerala, India, Phone: 04952350598, e-mail: dr.retnakumari@gmail.com

INTRODUCTION

Malocclusion is a continuum ranging from an ideal occlusion to considerable deviation from normal.¹ It has large impact on individual and society in terms of discomfort, quality of life and social and functional limitations. The etiology of malocclusion may be genetic, environmental or more commonly a combination of them. In addition, local factors such as adverse oral habits, anomalies in number, form and developmental position of teeth can also cause malocclusion.² Early interception and early correction of these malocclusions will prevent their progression to its full form and will exclude factors interfering with the regular development of the dental arches.³

In India, the prevalence of malocclusion varies from 20–43%.⁴ Presently, there is insufficient literature regarding the prevalence of malocclusion in Kerala state of India. Therefore, the present study was carried out to determine the prevalence of malocclusion and associated variables among schoolchildren in Kozhikode district of Kerala.

AIMS AND OBJECTIVES

- To determine the prevalence of malocclusion among 10 to 12-year-old schoolchildren.
- To identify the proportion of various types of malocclusion.
- To determine the proportion of different variables of malocclusions like crossbite, open bite, protrusion of teeth, deep bite, midline diastema, midline deviations and rotation of teeth.

MATERIALS AND METHODS

A descriptive cross-sectional study was conducted to assess the prevalence of malocclusion among schoolchildren aged 10–12 years in six schools in Kozhikode district of Kerala. The study was undertaken by the Department of Pedodontics and Preventive Dentistry, Government Dental College, Kozhikode and was conducted for a period of 1 year.

The permission for conducting the study was taken from the District Education Officer, Kozhikode. The

necessary information, such as names of all schools in Kozhikode district, their addresses and total number of students studying in each division in each school was obtained from the Education Council for the construction of a sample frame. Considering an average of 400 students from each school, six schools were randomly selected using cluster sampling method. To make up the estimated sample size, a total of 2,424 students were examined, among which 2,366 children satisfied the inclusion criteria.

Ethical clearance was obtained from the Ethical Committee of Government Dental College. Permission from the school authorities and consent from parents of children examined were obtained before the commencement of the study.

Inclusion Criteria

- The children of the age group of 10–12 years who were present on the day of examination with the informed consent of their parents
- Children who had all the permanent first molars erupted.

Exclusion Criteria

- Previous history of orthodontic treatment
- Craniofacial anomalies
- Uncooperative child
- Medically compromised child.

Examination of the Children

The oral examination was conducted by a single trained examiner using disposable gloves, standard mouth mirror and probe. The children were examined while seated on chair with good natural light/artificial illumination during class hours in a predetermined order. Each child was examined using the World Health Organization (WHO) criteria for oral health assessment. The assessment of the dental occlusion was carried out using disposable gloves, sterilized standard mouth mirrors and probes and calipers. All occlusal relationships were evaluated at a centric occlusion position which was achieved by asking the child to swallow and then to bite in his or her most posterior teeth. Class of malocclusion in Angle's system of classification, presence of variables like crossbite, open bite, deep bite, protrusion of teeth, tooth rotation, midline deviations and midline diastema were recorded. Children with class I molar relationship, normal overbite and overjet, proper alignment and no gross irregularities of tooth were categorized in normal occlusion group.

Personal data and previous history of orthodontic treatment were obtained directly from the children.

Statistical Analysis

The prevalence of malocclusion was represented in proportions. Differences in proportion among the group were analyzed using Chi-square test and data were analyzed using statistical software Statistical Packages for the Social Sciences (SPSS); p values less than 0.05 were considered as statistically significant.

RESULTS

Among the 2,366 children examined for the prevalence of malocclusion, 54.1% were boys and 45.9% were girls. The age and gender distribution of the children examined are shown in Table 1.

Among the children examined, 83.3% presented with malocclusions (Graph 1). To categorize the malocclusion in the examined children, 69.8% had class I malocclusion, 9.3% had class II malocclusion and 4.1% had class III malocclusion. Within class II malocclusion, 8.85% was division 1 type and 0.5% was division 2 type. The distribution of subjects based on gender and Angle's class of malocclusion was not statistically significant (Table 2).


Of the 2,366 children who were examined for overjet, 64.1% had normal overjet (<3 mm) and 23.2% had an increased overjet (>3 mm). A small percentage (4.2%) had reverse overjet (Table 3).

Among the total children examined, 64.1% had normal overbite (<3 mm), 35.6% had an increased overbite (>3 mm) and a small percentage (0.29%) had open bite (Table 4). The gender distribution was statistically significant in this group. Among the total of seven children (0.29%) with open bite (Table 5), six were purely skeletal with a male:female ratio of 1:5 and only one male child presented with dental open bite.

To present the crossbite, among the total children examined, 7.2% had teeth in crossbite, of which 1.2% presented

Table 1: Descriptive statistics of the children examined

Age (in years)	Male n (%)	Female n (%)	Total n (%)
10	358 (52.6)	322 (47.4)	680 (28.7)
11	334 (53.7)	288 (46.3)	622 (26.3)
12	589 (55.4)	475 (44.6)	1064 (45)
Total	1281 (54.1)	1085 (45.9)	2366 (100)



Graph 1: Prevalence of malocclusion

Table 2: The classification of the subjects based on Angle's class of malocclusion

Occlusion		Male n (%)	Female n (%)	Total n (%)	
Normal occlusion		187 (47.1)	210 (52.9)	397 (16.7)	
Class I malocclusion		911 (55.1)	741 (44.9)	1652 (69.8)	
Class II malocclusion	Div. 1	127 (60.8)	82 (39.2)	209 (8.85)	221 (9.3)
	Div. 2	3 (25)	9 (75)	12 (0.5)	
Class III malocclusion	Class III	53 (55.2)	43 (44.8)	96 (4.1)	
Total		1281 (54.1)	1085 (45.9)	2366 (100)	

Chi-square = 2.249; p = 0.325

Table 3: Distribution of overjet among the subjects

Overjet	Male n (%)	Female n (%)	Total n (%)
Normal (<3 mm)	929 (51.4)	878 (48.65)	1807 (76.4)
Increased (>3 mm)	346 (63.0)	203 (37.0)	549 (23.2)
Reverse overjet	6 (60)	4 (40)	10 (0.4)

Chi-square = 22.713; p = 0.000

Table 4: Distribution of overbite among the subjects

Overbite	Male n (%)	Female n (%)	Total n (%)
Normal (<3 mm)	775 (51.0)	742 (49)	1517 (64.1)
Increased (>3 mm)	504 (59.9)	338 (40.1)	842 (35.6)
Open bite	2 (28.5)	5 (71.5)	7 (0.29)

Chi-square = 17.199; p = 0.000

Table 5: Distribution of open bite among the subjects

Type of openbite	Male	Female	Total
Skeletal	1 (16.7%)	5 (83.3%)	6 (85.7)
Dental	1 (100%)	-	1 (14.3%)
Total	2 (28.5%)	5 (71.5%)	7 (100%)

with complete crossbite, 4.6% had crossbite of complete anterior teeth, 66.3% were with single or multiple anterior teeth in crossbite, 18.6% had unilateral posterior teeth in crossbite and 9.3% had both anterior and unilateral posterior teeth in crossbite (Table 6). Among the single-tooth crossbite, the most frequently noted tooth was maxillary right lateral incisor (61.4%) followed by maxillary left lateral incisor (38.5%). Gender distribution was also significant as only male children had complete crossbite.

In the examination of the midline with respect to the maxillary arch, 36.6% had no deviation, 30.23%

Table 6: Distribution of crossbite among the subjects

Type of crossbite	Male	Female	Total	
Complete (anterior + bilateral posterior)	2 (100%)	-	2 (1.2%)	
Anterior	Complete	4 (50%)	4 (50%)	8 (4.6%)
	Single tooth	62 (54.3%)	52 (45.6%)	114 (66.3%)
Unilateral posterior	17 (53.1%)	15 (46.8%)	32 (18.6%)	
Anterior + unilateral posterior	10 (62.5%)	6 (37.5%)	16 (9.3%)	
Total	95 (55.23%)	77 (44.77%)	172 (100%)	

Table 7: Distribution of midline deviation

Midline	Male	Female	Total
No deviation	457 (52.7%)	409 (47.2%)	866 (36.6%)
Deviated to right	382 (53.4%)	333 (46.6%)	715 (30.2%)
Deviated to left	442 (56.3%)	343 (43.7%)	785 (33.2%)

Chi-square = 2.283; p = 0.319

Table 8: Distribution of midline diastema and rotation


Variable	Male	Female	Total
Midline diastema	9 (50%)	9 (50%)	18 (0.76%)
Tooth rotation	34 (44.2%)	43 (55.8%)	77 (3.25%)

had deviation to right and 33.17% had deviation to left (Table 7).

Among the children examined for midline diastema, 0.76% had midline diastema (Table 8) and it was completely in the maxillary arch.

Rotation of tooth was found to be the most common individual tooth irregularities in our study group. In the total group of children examined, 3.25% had rotation of tooth (Table 8). Lateral incisors (64.9%) were found to be the more frequently rotated type of teeth. The most rotated teeth were maxillary right lateral incisors (19.4%).

Overall distribution of the variables of malocclusion among subjects examined (Graph 2) revealed that deep bite (35.6%) was the most prevalent, followed by increased overjet (23.2%) and crossbite (7.2%). The least noted characteristics were open bite (0.29%) and midline diastema (0.76%).



Graph 2: Prevalence and gender distribution of different variables of malocclusion

DISCUSSION

Malocclusion is one of the most common dental problems in mankind. Maloccluded teeth can cause psychosocial problems related to impaired dentofacial aesthetics, disturbances of oral function, such as mastication, swallowing and speech, and greater susceptibility to trauma and periodontal disease.

Numerous studies have been published regarding the prevalence of malocclusion in various populations. The results have shown wide variations. Differences in the age ranges of the populations studied, the number of subjects examined and differences in the registration methods are probably the most important factors explaining these variations.⁵

In the last decade, a number of studies have attempted to examine the malocclusion problem on a population basis using cross-sectional examinations of groups claimed to be representative of the Indian nation. But very few studies were known to be reported based on population of Kerala state of India. The present study was conducted among 2,366 schoolchildren aged 10–12 years to determine the prevalence of malocclusion in Kozhikode district of Kerala, India.

The present study evaluated the occlusion status of the subjects using Angle's classification of malocclusion. Of the children examined in the present study, 16.7% of subjects reported with normal occlusion and 83.3% with malocclusions. The prevalence of malocclusion in our study is almost similar to the study done by Kaur et al⁶ in Karnataka, India (87.79%) and Ajayi⁷ in Nigeria (84.1%). When compared with our study, a higher prevalence of malocclusion was reported by Abu Alhaija et al⁸ in Jordan (92%) and Rwakatemala⁹ in Tanzania (97.6%). But the studies conducted by Hemapriya et al¹⁰ in Kancheepuram (75%) and Trehan et al¹¹ in Jaipur (66.7%) reported a lower prevalence than the present study.

The prevalence of class I malocclusion seen in the present study (69.8%) is almost similar to the findings by Trehan et al¹¹ in Jaipur (57.9%) and Das and Reddy¹² in Bengaluru (61.6%), India. This is found to be higher according to studies of Phaphe et al¹³ in Bagalkot (17.8%) and Vibhute et al¹⁴ in Mumbai (49.1%), India and lower when compared to the findings of Mtaya et al⁵ in Tanzania (93.6%) and Ajayi⁷ in Nigeria (80.7%). Class II malocclusion seen in the present study (9.03%) is similar with the findings of Sridharan et al¹⁵ in Tumkur (10%) and Muppa et al¹⁶ in Andhra Pradesh (9.95), India. This is found to be higher than the results of Mtaya et al⁵ in Tanzania (4.4%) and Shrestha et al¹⁷ in Kathmandu (2.5%) and was lower when compared with the studies of Abu Alhaija et al⁸ in Jordan (18.8%) and Phaphe et al¹³ in Bagalkot, India

(30.1%). Class III malocclusion (4.1%) found in the present study was almost similar to that reported by Vibhute et al¹⁴ in Mumbai (5.7%), India and Thilander et al¹⁸ in Columbia (3.7%). This is found to be higher according to Abu Alhaija et al⁸ in Jordan (1.4%) and Das et al¹² in Bengaluru, India (0.6%) and lower when compared with the findings of Farahani et al¹ in Iran (7.8%) and Cedikoglu et al¹⁹ in Turkey (16.7%). There was no significant difference in gender distribution for the prevalence of different classes of malocclusion in our study.

In the present study, overjet less than 3 mm was categorized as normal and it was found that 23.2% of the subjects were with increased overjet, which is similar to the findings of Abu Alhaija et al⁸ in Jordan (24.7%) and Gelgor et al²⁰ in Central Anatolia (25.1%). The studies by Farahani et al¹ in Iran (31.7%) and Hemapriya et al¹⁰ in Kancheepuram, India (61.4%) reported a higher prevalence of increased overjet, while lower prevalence was reported in the studies by Siddegowda and Rani²¹ in Karnataka (6.3%), India and Poeung et al²² in Cambodia (8.1%).

In the present study, overbite less than 3 mm was categorized as normal and it was found that 38.0% of the subjects were with increased overbite, which is similar to the findings of Cedikoglu et al¹⁹ in Turkey (36.6%) and Nainani and Sugandh²³ in Nagpur, India (38.0%). But the studies by Tausche³ in Dredsen (46.2%) and Siddegowda and Rani²¹ in Karnataka, India (51.75%) reported a higher prevalence of increased overbite, and studies by Rwakatemala⁹ in Tanzania (20%) and Phaphe et al¹³ in Bagalkot, India (9.2%) reported a lower prevalence than that in the present study.

In our study, only 0.29% of the subjects reported with anterior open bite. The studies by Farahani et al¹ in Iran (1.6%), Mtaya et al⁵ in Tanzania (1.8%) and Ciuffolo et al²⁴ in Italy (1.7%) also reported a very low prevalence of open bite. When compared with our study, the findings of the studies conducted by Nainani and Sugandh²³ in Nagpur (2.98%), India, Ajayi⁷ in Nigeria (4.1%) and Poeung et al²² in Cambodia (16.4%) were much higher. In the present study, it was found that 7.1% children had crossbite which corroborates with the studies by Nainani and Sugandh²³ in Nagpur, India (5.5%) and Abu Alhaija et al⁸ in Jordan (6.7%). The studies by Siddegowda and Rani²¹ in Karnataka (18%), India and Poeung et al²² in Cambodia (14.7%) reported with a higher prevalence of crossbite. In our study, 4.2% reported with anterior crossbite, which is similar to the findings of Muppa et al¹⁶ in Andhra Pradesh (4.98), India and Bittencourt et al²⁵ in Brazil (5%). But the studies by Phaphe et al¹³ in Bagalkot (7.2%), India and Cedikoglu et al¹⁹ in Turkey (14.1%) showed a higher value of crossbite.

A midline diastema is considered to be present when there is a space of at least 2 mm between the maxillary central incisors. Among the total subjects evaluated, only 0.76% were with maxillary midline diastema. This finding is very lower when compared with the studies by Phaphe et al¹³ in Bagalkot (18%), Hemapriya et al¹⁰ in Kancheepuram (35.2%), India and Ajayi⁷ in Nigeria (19.5%). The children reported with rotated tooth in the present study was 3.25%, which was lower when compared with the findings of Vibhute et al¹⁴ in Mumbai (13.1%) and Nainani and Sugandh²³ in Nagpur (15.3%).

It is essential to identify and localize the wide range of deviations from occlusal development that may arise and that must be intercepted before the end of the active growth stage. Problems of a functional nature that arise from these morphological changes may become more complex skeletal problems in the future with serious psychosocial consequences for the developing individual.⁴

The findings of the present study will be very useful for the early interceptive measures as well as early correction of the malocclusion, thus reducing its severity in the permanent dentition.

CONCLUSION

The following conclusions are drawn from the present study:

- Prevalence of malocclusion was found to be 83.3%. Class I malocclusion was the most prevalent type (69.8%), followed by class II malocclusion (9.3%), with 8.85% of division 1 type and 0.5% of division 2 type, and class III malocclusion (4.1%).
- Prevalence of normal occlusion was seen more in females, whereas prevalence of malocclusion was more in males. This difference was not statistically significant.
- The prevalence of crossbite and openbite was 7.2 and 0.29% respectively. Maxillary right lateral incisor was the most common tooth in crossbite.
- The prevalence of excessive overjet (protrusion of maxillary teeth) and excessive overbite (deep bite) was 23.2 and 35.6%, respectively.
- The prevalence of midline diastema was 0.76%.
- The prevalence of rotation was 3.25% and the most commonly rotated tooth was maxillary right lateral incisor followed by mandibular right lateral incisor.
- There was no statistically significant difference in midline deviation.

To conclude, the results of the present study confirmed that there is increased prevalence of malocclusion among children in the 10–12 years age group. The finding of this study will provide baseline data for implementing

early interceptive treatment for the elimination of factors inhibiting dental arch development as well as skeletal jaw growth.

REFERENCES

1. Farahani AB, Eslamipour F. Malocclusion and occlusal traits in an urban Iranian population. An epidemiological study of 11- to 14-year-old children. *Eur J Orthod* 2009 Oct;31(5):477-484.
2. Kumar DA, Varghese RK, Chaturvedi SS, Agrawal C, Fating C, Makkad RS. Prevalence of malocclusion among children and adolescents residing in orphanages of Bilaspur, Chhattisgarh, India. *J Adv Oral Res* 2012 Sep-Dec;3(3):21-28.
3. Tausche E. Prevalence of malocclusion in the early mixed dentition and orthodontic treatment need. *Eur J Orthod* 2004 Jun 1;26(3):237-244.
4. Agarwal SS, Chopra SS, Jayan B, Verma M. Epidemiology in orthodontics—a literature review. *Orthod Cyber J* 2013 Sep.
5. Mtaya M, Brudvik P, Astrom AN. Prevalence of malocclusion and its relationship with sociodemographic factors, dental caries, and oral hygiene in 12- to 14-year-old Tanzanian schoolchildren. *Eur J Orthod* 2009 Oct;31(5):467-476.
6. Kaur H, Pavithra US, Abraham R. Prevalence of malocclusion among adolescents in South Indian population. *J Int Soc Prev Community Dent* 2013 Jul;3(2):97-102.
7. Ajayi EO. Prevalence of malocclusion among school children in Benin City, Nigeria. *J Med Biomed Res* 2008;7(1 and 2):5-11.
8. Abu Alhajja ES, Al-Khateeb SN, Al-Nimri KS. Prevalence of malocclusion in 13-15 year-old North Jordanian school children. *Community Dent Health* 2005 Dec;22(4):266-271.
9. Rwakatema S. Prevalence of malocclusion among 12-15-year-olds in Moshi, Tanzania, using Bjork's criteria. *East Afr Med J* 2006 Jul;83(7):372-380.
10. Hemapriya S, Ingle NA, Chaly PE, Reddy VC. Prevalence of malocclusion and orthodontic treatment needs among 12 and 15 years old rural school children in Kancheepuram district. *J Oral Health Community Dent* 2013 May;7(2):84-90.
11. Trehan M, Chugh VK, Sharma S. Prevalence of malocclusion in Jaipur, India. *Int J Clin Pediatr Dent* 2009 Jan;2(1):23-25.
12. Das UM, Reddy VD. Prevalence of malocclusion among school children in Bangalore, India. *Int J Clin Pediatr Dent* 2008 Sep;1(1):10-12.
13. Phaphe S, Kallur R, Vaz A, Gajapurada J, Sugaraddy, Mattigatti S. To determine the prevalence rate of malocclusion among 12 to 14-year-old schoolchildren of urban Indian population (Bagalkot). *J Contemp Dent Pract* 2012 May 1;13(3):316-321.
14. Vibhute AH, Vibhute NA, Daule R. Prevalence of malocclusion characteristics and chief motivational factor for treatment in orthodontic patients from Maharashtra, India. *J Orthod Res* 2013;1(2):62-65.
15. Sridharan K, Udupa V, Srinivas H, Kumar S, Sandbhor S. Prevalence of Class II malocclusion in Tumkur population. *J Dent Sci Res* 2011;2(2):1-5.
16. Muppa R, Bhupathiraju P, Duddu MK, Dandempally A, Karre DL. Prevalence and determinant factors of malocclusion in population with special needs in South India. *J Indian Soc Pedod Prev Dent* 2013 Apr-Jun;31(2):87-90.
17. Shrestha BK, Yadav R, Basel P. Prevalence of malocclusion among high school students in Kathmandu Valley. *Orthod J Nepal* 2012;2(1):1-5.

18. Thilander B, Pena L, Infante C, Parada SS, Mayorga CD. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod* 2001 Apr 1;23(2):153-167.
19. Cedikoglu M, Akpinar S, Yavuz I. The pattern of malocclusion in a sample of orthodontic patients from Turkey. *Med Oral Patol Oral Cir Bucal* 2010 Sep 1;15(5):e791-e796.
20. Gelgor IE, Karaman AI, Ercan ER. Prevalence of malocclusion among adolescents in Central Anatolia. *Eur J Dent* 2007 Jul;1(3):125-131.
21. Siddegowda R, Rani MS. A cross-sectional epidemiological survey on prevalence of malocclusion in government, aided and private school children of Karnataka. *Univ J Public Health* 2013;1(3):124-130.
22. Poeung P, Kruger E, Tennant M. The prevalence of malocclusion, dental irregularities and orthodontic treatment need in 13-15 year olds in Teuk Klaing, Cambodia. *J Int Oral Health* 2011 Oct;3(5):19-28.
23. Nainani JT, Relan Sugandh. Prevalence of malocclusion in school children of Nagpur rural region—an epidemiological study. *J Ind Dent Assoc* 2011;5(8):865-867.
24. Ciuffolo F, Manzoli L, D'Attilio M, Tecco S, Muratore F, Festa F, Romano F. Prevalence and distribution by gender of occlusal characteristics in a sample of Italian secondary school students: a cross-sectional study. *Eur J Orthod* 2005 Dec;27 (6):601-606.
25. Bittencourt MAV, Machado AW. An overview of the prevalence of malocclusion in 6 to 10-year-old children in Brazil. *Dental Press J Orthod* 2010 Nov-Dec;15(6):113-122.