



Genome Sequence of a Gammaherpesvirus from a Common Bottlenose Dolphin (*Tursiops truncatus*)

Andrew J. Davison,^a  Kuttichantran Subramaniam,^b Karen Kerr,^a
Jessica M. Jacob,^b Nelmarie Landrau-Giovanetti,^b Michael T. Walsh,^c
Randall S. Wells,^{c,d} Thomas B. Waltzek^b

MRC-University of Glasgow Centre for Virus Research, Glasgow, United Kingdom^a; Department of Infectious Diseases and Pathology, College of Veterinary Medicine, University of Florida, Gainesville, Florida, USA^b; Department of Large Animal Clinical Sciences, College of Veterinary Medicine, University of Florida, Gainesville, Florida, USA^c; Chicago Zoological Society's Sarasota Dolphin Research Program, c/o Mote Marine Laboratory, Sarasota, Florida, USA^d

ABSTRACT A herpesvirus genome was sequenced directly from a biopsy specimen of a rectal lesion from a female common bottlenose dolphin. This genome sequence comprises a unique region (161,235 bp) flanked by multiple copies of a terminal repeat (4,431 bp) and contains 72 putative genes. The virus was named common bottlenose dolphin gammaherpesvirus 1.

Health assessments of common bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, FL, were carried out in 2015 (1). Physical examination of a 13-year-old long-term resident female on 15 May determined that the animal weighed 142.2 kg, was lactating, had mild lung disease, and exhibited a raised proliferative cutaneous lesion at the border of the rectum. A biopsy specimen of the lesion was taken and stored at -80°C . DNA was extracted by using a Qiagen DNeasy blood and tissue kit. A sequencing library was prepared by using an Illumina Nextera XT DNA library preparation kit and sequenced by using a 600-cycle version 3 cartridge on an Illumina MiSeq.

The data set of 8,994,640 quality-trimmed reads was assembled *de novo* by using SPAdes (2). A large herpesvirus-related contig was extended and joined to other contigs manually, essentially forming a circular sequence consisting of a unique region (U) linked to a complete and a partial copy of a terminal repeat (TR). The right genome terminus was mapped to a nucleotide in TR from the presence of multiple reads commencing at the same position, and the adjacent nucleotide was assigned as the left genome terminus. The sequence contained seven large tandem repeats that were possibly heterogeneous and could not be solved by additional sequencing of PCR products. Representative solutions consistent with the size of PCR products and the available sequence data were adopted. Coverage analysis indicated that the linear viral genome consists of U flanked on each side by multiple copies of TR. The final sequence was represented as U followed by a complete and a partial copy of TR. The size of this sequence was 167,212 bp, with the sizes of U and TR being 161,235 bp and 4,431 bp, respectively. The integrity of the sequence was assessed by inspection of a read assembly generated by using Bowtie 2 (3). A total of 626,924 reads (7.0% of the total) aligned at an average coverage of 1,092 reads per nucleotide.

Predicted amino acid sequences from open reading frames (ORFs) of >50 codons were screened for relatives in other organisms (especially herpesviruses) and to locate probable initiation codons and splicing patterns. Substantially overlapping ORFs and ORFs of <150 codons that lacked relatives and other features (e.g., hydrophobic domains) were ruled out. Among the final set of 72 genes, 4 (De4, De7, De9, and De10) lacked relatives and other features, and were considered marginal. Phylogenetic anal-

Received 21 June 2017 Accepted 23 June 2017 Published 3 August 2017

Citation Davison AJ, Subramaniam K, Kerr K, Jacob JM, Landrau-Giovanetti N, Walsh MT, Wells RS, Waltzek TB. 2017. Genome sequence of a gammaherpesvirus from a common bottlenose dolphin (*Tursiops truncatus*). *Genome Announc* 5:e00777-17. <https://doi.org/10.1128/genomeA.00777-17>.

Copyright © 2017 Davison et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Thomas B. Waltzek, tbwaltzek@ufl.edu.

yses indicated that the virus from which the genome presumably originated is a deeply rooted member of subfamily *Gammaherpesvirinae* of family *Herpesviridae*.

The virus was named common bottlenose dolphin gammaherpesvirus 1 strain Sarasota (proposed species *Delphinid gammaherpesvirus 1*) and is the first herpesvirus from a marine mammal for which a genome sequence has been determined. Alpha-herpesviruses and gammaherpesviruses have been reported previously in the cetacean families Balaenopteridae, Delphinidae, Kogiidae, Monodontidae, Phocoenidae, Physeteridae, and Ziphiidae (4–8).

Accession number(s). The common bottlenose dolphin gammaherpesvirus 1 strain Sarasota genome sequence has been deposited in GenBank under accession no. [KY965444](https://doi.org/10.1093/genbank/KY965444).

ACKNOWLEDGMENTS

Bottlenose dolphin samples were collected under National Marine Fisheries Service Scientific Research Permit no. 15543 (issued to R. S. Wells), and with Mote Marine Laboratory IACUC approval. Health assessments were conducted through the support of Dolphin Quest, Inc. Sequencing was supported by the Florida Fish and Wildlife Conservation Commission (grant no. 15145); the views expressed herein are those of the authors and not necessarily those of the Commission. Sequence analyses were supported by the UK Medical Research Council (grant no. MC_UU_12014/3).

REFERENCES

1. Wells RS, Rhinehart HL, Hansen LJ, Sweeney JC, Townsend FI, Stone R, Casper DR, Scott MD, Hohn AA, Rowles TK. 2004. Bottlenose dolphins as marine ecosystem sentinels: developing a health monitoring system. *Ecohealth* 1:246–254. <https://doi.org/10.1007/s10393-004-0094-6>.
2. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
3. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
4. Maness HTD, Nollens HH, Jensen ED, Goldstein T, LaMere S, Childress A, Sykes J, St Leger J, Lacave G, Latson FE, Wellehan JFX. 2011. Phylogenetic analysis of marine mammal herpesviruses. *Vet Microbiol* 149:23–29. <https://doi.org/10.1016/j.vetmic.2010.09.035>.
5. Miyoshi K, Nishida S, Sone E, Tajima Y, Makara M, Yoshioka M, Nakamura M, Yamada TK, Koike H. 2011. Molecular identification of novel alpha- and gammaherpesviruses from cetaceans stranded on Japanese coasts. *Zool Sci* 28:126–133. <https://doi.org/10.2108/ZSJ.28.126>.
6. Bellehumeur C, Lair S, Romero CH, Provost C, Nielsen O, Gagnon CA. 2015. Identification of a novel herpesvirus associated with a penile proliferative lesion in a beluga (*Delphinapterus leucas*). *J Wildl Dis* 51:244–249. <https://doi.org/10.7589/2013-11-293>.
7. Melero M, Crespo-Picazo JL, Rubio-Guerri C, García-Párraga D, Sánchez-Vizcaíno JM. 2015. First molecular determination of herpesvirus from two mysticete species stranded in the Mediterranean Sea. *BMC Vet Res* 11:283. <https://doi.org/10.1186/s12917-015-0596-1>.
8. van Elk C, van de Bildt M, van Run P, de Jong A, Getu S, Verjans G, Osterhaus A, Kuiken T. 2016. Central nervous system disease and genital disease in harbor porpoises (*Phocoena phocoena*) are associated with different herpesviruses. *Vet Res* 47:28. <https://doi.org/10.1186/s13567-016-0310-8>.