

9-(3-Bromo-5-chloro-2-hydroxyphenyl)-10-(2-hydroxyethyl)-1,2,3,4,5,6,7,8,9,10-decahydroacridine-1,8-dione

Shaaban K. Mohamed,^{a,b} Mehmet Akkurt,^{c*} Peter N. Horton,^d Antar A. Abdelhamid^e and Mahmoud A. A. El Remaily^e

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^bChemistry Department, Faculty of Science, Minia University, El-Minia, Egypt, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^dSchool of Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, England, and ^eDepartment of Chemistry, Sohag University, 82524 Sohag, Egypt

Correspondence e-mail: akkurt@erciyes.edu.tr


Received 3 December 2012; accepted 9 December 2012

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.028; wR factor = 0.077; data-to-parameter ratio = 16.9.

In the title compound, $C_{21}H_{21}\text{BrClNO}_4$, the dihydropyridine ring adopts a flattened boat conformation. The 3-bromo-5-chloro-2-hydroxyphenyl ring forms a dihedral angle of $84.44(7)^\circ$ with the dihydropyridine mean plane. The molecular conformation is stabilized by an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond, with an $S(8)$ ring motif. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming a three-dimensional network.

Related literature

For the synthesis and bioactivity of acridines, see, for example: Karolak-Wojciechowska *et al.* (1996). For related structures, see: Abdelhamid *et al.* (2011a,b); Mohamed *et al.* (2012); Guo *et al.* (2004); Suganya & Sureshbabu (2012); Yogavel *et al.* (2005). For ring puckering parameters, see: Cremer & Pople (1975). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$C_{21}H_{21}\text{BrClNO}_4$
 $M_r = 466.74$
Monoclinic, $P2_1/n$
 $a = 8.810(2)\text{ \AA}$
 $b = 13.809(3)\text{ \AA}$
 $c = 15.797(4)\text{ \AA}$
 $\beta = 100.026(4)^\circ$

$V = 1892.5(8)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 2.34\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.22 \times 0.14 \times 0.03\text{ mm}$

Data collection

Rigaku AFC12 (Right)
diffractometer
Absorption correction: multi-scan
(*CrystalClear-SM Expert*;
Rigaku, 2012)
 $T_{\min} = 0.627$, $T_{\max} = 0.933$

14959 measured reflections
4299 independent reflections
4126 reflections with $I > 2\sigma(i)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.077$
 $S = 1.03$
4299 reflections

255 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.86\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.38\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A \cdots O2	0.84	1.88	2.6749 (19)	158
O4—H4 \cdots O2 ⁱ	0.84	1.94	2.782 (2)	176
C6—H6B \cdots O3 ⁱⁱ	0.99	2.33	3.051 (2)	129
C20—H20A \cdots O3 ⁱⁱ	0.99	2.53	3.486 (2)	163
C20—H20B \cdots O1 ⁱ	0.99	2.57	3.492 (2)	154

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *CrystalClear-SM Expert* (Rigaku, 2012); cell refinement: *CrystalClear-SM Expert*; data reduction: *CrystalClear-SM Expert* Expert; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

The EPSRC National Crystallography Service is gratefully acknowledged for the data collection. The authors are thankful to Manchester Metropolitan University, Sohag University and Erciyes University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5278).

References

- Abdelhamid, A. A., Mohamed, S. K., Allahverdiyev, M. A., Gurbanov, A. V. & Ng, S. W. (2011b). *Acta Cryst. E67*, o785.
- Abdelhamid, A. A., Mohamed, S. K., Khalilov, A. N., Gurbanov, A. V. & Ng, S. W. (2011a). *Acta Cryst. E67*, o744.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Guo, C., Tu, S., Li, T. & Zhu, S. (2004). *Acta Cryst. E60*, o2035–o2037.
- Karolak-Wojciechowska, J., Mrozek, A., Amiel, P., Brouant, P. & Barbe, J. (1996). *Acta Cryst. C52*, 2939–2941.

- Mohamed, S. K., Abdelhamid, A. A., Maharramov, A. M., Khalilov, A. N., Gurbanov, A. V. & Allahverdiyev, M. A. (2012). *J. Chem. Pharm. Res.* **4**, 955–965.
- Rigaku (2012). *CrystalClear-SM Expert*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Sughanya, V. & Sureshbabu, N. (2012). *Acta Cryst. E* **68**, o2755.
- Yogavel, M., Velmurugan, D., Murugan, P., Shanmuga Sundara Raj, S. & Fun, H.-K. (2005). *Acta Cryst. E* **61**, o2761–o2763.

supplementary materials

Acta Cryst. (2013). E69, o85–o86 [doi:10.1107/S1600536812050222]

9-(3-Bromo-5-chloro-2-hydroxyphenyl)-10-(2-hydroxyethyl)-1,2,3,4,5,6,7,8,9,10-decahydroacridine-1,8-dione

Shaaban K. Mohamed, Mehmet Akkurt, Peter N. Horton, Antar A. Abdelhamid and Mahmoud A. A. El Remaily

Comment

Acridine derivatives are one of the oldest and most successful classes of bioactive agents (Karolak-Wojciechowska *et al.*, 1996). Further to our on-going study on the synthesis and biological assessment of acridines (Mohamed *et al.*, 2012; Abdelhamid *et al.*, 2011*a,b*), we report herein the synthesis and crystal structure determination of the title compound (I).

In the title compound (I), (Fig. 1), the dihydropyridine ring (N1/C1/C2/C7/C8/C13) is almost planar with a maximum deviation of 0.160 (2) Å for C1. The C14–C19 phenyl ring forms a dihedral angle of 84.44 (7)° with the dihydropyridine mean plane. In the 1,2,3,4,5,6,7,8,9,10-decahydroacridine ring system, the puckering parameters (Cremer & Pople, 1975) for the A(C2–C7), B(N1/C1/C2/C7/C8/C13) and C(C8–C13) rings are $Q_T = 0.4695$ (18) Å, $\theta = 121.6$ (2) °, $\varphi = 341.3$ (2) ° (for A); $Q_T = 0.2607$ (16) Å, $\theta = 77.7$ (4) °, $\varphi = 167.6$ (4) ° (for B) and $Q_T = 0.4511$ (19) Å, $\theta = 126.1$ (2) °, $\varphi = 351.9$ (3) ° (for C), respectively. The cyclohexenone rings A and C adopt sofa conformations, whereas the central ring B adopts flattened boat conformation. In (I), the bond lengths and angles are within normal ranges and comparable with those in related similar compounds (Sughanya & Sureshbabu, 2012; Yogavel *et al.*, 2005; Guo *et al.*, 2004). The ethanol group is not coplanar with the attached 1,4-dihydropyridine ring, with a N1—C20—C21—O4 torsion angle of -174.31 (14)°.

The molecular conformation is stabilized by an intramolecular O—H···O hydrogen bond (Table 1), which forms a pseudo-eight-membered ring with graph set S(8) (Bernstein *et al.*, 1995).

In the crystal, molecules are linked by O—H···O and C—H···O hydrogen bonds, forming three dimensional network (Fig. 2, Table 1).

Experimental


A mixture of 112 mg (0.001 mol) cyclohexane-1,3-dione, 236 mg (0.001 mol) 3-bromo-5-chloro-2-hydroxybenzaldehyde and 61 mg (0.001 mol) 2-aminoethanol in 50 ml ethanol was refluxed at 350 K and monitored by TLC till completion after 2 h. A mass solid precipitate was deposited on cooling, filtered and dried under vacuum then washed with cold ethanol and dried again.. The raw product was recrystallized from dimethyl formamide then triturated with ether to afford a good yield (73%) of high quality yellow plats (m.p. 483 K) that were suitable for X-ray difraction.

Refinement


All H-atoms were placed in calculated positions with O—H = 0.84 Å, and C—H = 0.95 for aromatic, 0.99 for methylene and 1.00 Å for methine C—H = 0.97 Å for methylene $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$. They were refined using a riding model approximation with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$ for hydroxyl and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ for the other H atoms.

Computing details

Data collection: *CrystalClear-SM Expert* (Rigaku, 2012); cell refinement: *CrystalClear-SM Expert* (Rigaku, 2012); data reduction: *CrystalClear-SM Expert Expert* (Rigaku, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

**Figure 1**

The title compound (I) with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.

**Figure 2**

View of the packing and hydrogen bonding of (I) down the a axis. The hydrogen atoms not involved in the hydrogen bonds have been omitted for clarity.

9-(3-Bromo-5-chloro-2-hydroxyphenyl)-10-(2-hydroxyethyl)-1,2,3,4,5,6,7,8,9,10-decahydroacridine-1,8-dione

Crystal data



$M_r = 466.74$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 8.810 (2)$ Å

$b = 13.809 (3)$ Å

$c = 15.797 (4)$ Å

$\beta = 100.026 (4)^\circ$

$V = 1892.5 (8)$ Å³

$Z = 4$

$F(000) = 952$

$D_x = 1.638 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71075$ Å

Cell parameters from 4812 reflections

$\theta = 2.5-27.5^\circ$

$\mu = 2.34 \text{ mm}^{-1}$

$T = 100$ K

Plate, yellow

$0.22 \times 0.14 \times 0.03$ mm

Data collection

Rigaku AFC12 (Right)
diffractometer

Radiation source: Rotating Anode

Detector resolution: 28.5714 pixels mm⁻¹

profile data from ω -scans

Absorption correction: multi-scan

(*CrystalClear-SM Expert*; Rigaku, 2012)
 $T_{\min} = 0.627$, $T_{\max} = 0.933$

14959 measured reflections

4299 independent reflections

4126 reflections with $I > 2\sigma(i)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -10 \rightarrow 11$

$k = -17 \rightarrow 14$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.077$

$S = 1.03$

4299 reflections

255 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0485P)^2 + 1.1263P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$
 $\Delta\rho_{\max} = 0.86 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.38 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Rigaku *CrystalClear-SM* Expert 3.1 b5

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.02039 (2)	1.02608 (1)	0.11100 (1)	0.0178 (1)
Cl2	0.89028 (5)	1.13026 (3)	0.42922 (3)	0.0206 (1)
O1	0.85205 (14)	0.84648 (9)	0.14843 (8)	0.0171 (3)
O2	0.91879 (13)	0.68133 (9)	0.23546 (7)	0.0155 (3)
O3	0.44222 (13)	0.87300 (9)	0.18903 (8)	0.0175 (3)
O4	0.69992 (16)	0.85677 (12)	0.68490 (9)	0.0294 (4)
N1	0.65599 (15)	0.75304 (10)	0.46456 (8)	0.0125 (4)
C1	0.69843 (17)	0.79777 (11)	0.29224 (10)	0.0113 (4)
C2	0.79574 (17)	0.72563 (11)	0.34966 (10)	0.0114 (4)
C3	0.90436 (17)	0.66869 (11)	0.31212 (10)	0.0124 (4)
C4	0.99634 (19)	0.59201 (12)	0.36547 (11)	0.0169 (4)
C5	1.03428 (19)	0.62450 (13)	0.45893 (11)	0.0165 (4)
C6	0.88912 (18)	0.65080 (12)	0.49409 (11)	0.0152 (4)
C7	0.77954 (17)	0.71276 (11)	0.43370 (10)	0.0121 (4)
C8	0.54946 (17)	0.81352 (11)	0.32480 (10)	0.0114 (4)
C9	0.42314 (18)	0.85498 (12)	0.26225 (10)	0.0134 (4)
C10	0.27404 (19)	0.87759 (14)	0.29195 (11)	0.0202 (5)
C11	0.24551 (19)	0.80920 (14)	0.36209 (11)	0.0203 (5)
C12	0.38110 (18)	0.80674 (14)	0.43714 (11)	0.0165 (4)
C13	0.53272 (17)	0.79239 (11)	0.40620 (10)	0.0122 (4)
C14	0.78575 (17)	0.89267 (11)	0.28563 (10)	0.0113 (4)
C15	0.85590 (17)	0.91091 (12)	0.21354 (10)	0.0128 (4)
C16	0.93144 (18)	0.99934 (13)	0.20933 (10)	0.0135 (4)
C17	0.94463 (17)	1.06742 (12)	0.27456 (11)	0.0143 (4)
C18	0.87751 (19)	1.04633 (12)	0.34544 (11)	0.0146 (4)
C19	0.79829 (18)	0.96063 (12)	0.35129 (10)	0.0131 (4)
C20	0.65018 (19)	0.75423 (14)	0.55771 (10)	0.0168 (4)
C21	0.6898 (2)	0.85513 (15)	0.59499 (12)	0.0242 (5)
H1	0.67320	0.76910	0.23330	0.0140*
H1A	0.85100	0.78990	0.16790	0.0260*
H4	0.61310	0.84430	0.69740	0.0440*
H4A	1.09300	0.57970	0.34330	0.0200*

H4B	0.93660	0.53100	0.36150	0.0200*
H5A	1.10350	0.68140	0.46340	0.0200*
H5B	1.08930	0.57180	0.49410	0.0200*
H6A	0.83570	0.59050	0.50590	0.0180*
H6B	0.91900	0.68570	0.54920	0.0180*
H10A	0.18800	0.87260	0.24270	0.0240*
H10B	0.27710	0.94490	0.31370	0.0240*
H11A	0.22750	0.74320	0.33800	0.0240*
H11B	0.15140	0.82970	0.38350	0.0240*
H12A	0.38450	0.86830	0.46960	0.0200*
H12B	0.36580	0.75330	0.47660	0.0200*
H17	0.99800	1.12660	0.27080	0.0170*
H19	0.75240	0.94830	0.40040	0.0160*
H20A	0.72430	0.70640	0.58780	0.0200*
H20B	0.54570	0.73570	0.56680	0.0200*
H21A	0.78940	0.87620	0.58030	0.0290*
H21B	0.60970	0.90150	0.56860	0.0290*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0204 (1)	0.0193 (1)	0.0158 (1)	0.0017 (1)	0.0089 (1)	0.0053 (1)
Cl2	0.0244 (2)	0.0190 (2)	0.0201 (2)	-0.0061 (2)	0.0086 (2)	-0.0083 (2)
O1	0.0224 (6)	0.0168 (6)	0.0136 (5)	-0.0002 (5)	0.0072 (5)	-0.0009 (5)
O2	0.0148 (5)	0.0180 (6)	0.0145 (5)	0.0010 (4)	0.0048 (4)	-0.0014 (5)
O3	0.0163 (6)	0.0228 (6)	0.0126 (5)	0.0012 (5)	0.0006 (4)	0.0024 (5)
O4	0.0222 (7)	0.0506 (9)	0.0163 (6)	-0.0094 (6)	0.0056 (5)	-0.0090 (6)
N1	0.0129 (6)	0.0163 (7)	0.0084 (6)	0.0000 (5)	0.0021 (5)	-0.0004 (5)
C1	0.0106 (7)	0.0136 (7)	0.0096 (7)	0.0000 (5)	0.0016 (5)	-0.0002 (6)
C2	0.0101 (7)	0.0114 (7)	0.0123 (7)	-0.0003 (5)	0.0011 (5)	-0.0008 (6)
C3	0.0107 (7)	0.0114 (7)	0.0150 (7)	-0.0021 (5)	0.0020 (5)	-0.0024 (6)
C4	0.0164 (7)	0.0155 (8)	0.0192 (8)	0.0035 (6)	0.0045 (6)	0.0019 (7)
C5	0.0138 (7)	0.0182 (8)	0.0168 (8)	0.0026 (6)	0.0005 (6)	0.0028 (6)
C6	0.0154 (7)	0.0158 (8)	0.0142 (7)	0.0012 (6)	0.0020 (6)	0.0026 (6)
C7	0.0106 (7)	0.0116 (7)	0.0137 (7)	-0.0021 (5)	0.0013 (5)	-0.0012 (6)
C8	0.0108 (7)	0.0117 (7)	0.0116 (7)	-0.0006 (5)	0.0020 (5)	-0.0017 (6)
C9	0.0126 (7)	0.0134 (7)	0.0136 (7)	-0.0010 (6)	0.0009 (6)	-0.0012 (6)
C10	0.0130 (7)	0.0288 (9)	0.0183 (8)	0.0048 (7)	0.0017 (6)	0.0013 (7)
C11	0.0134 (7)	0.0293 (9)	0.0182 (8)	-0.0011 (7)	0.0027 (6)	-0.0009 (7)
C12	0.0116 (7)	0.0262 (9)	0.0127 (7)	-0.0004 (6)	0.0047 (6)	-0.0018 (7)
C13	0.0110 (7)	0.0120 (7)	0.0132 (7)	-0.0010 (5)	0.0011 (6)	-0.0020 (6)
C14	0.0085 (6)	0.0134 (7)	0.0115 (7)	0.0013 (5)	0.0005 (5)	0.0018 (6)
C15	0.0105 (7)	0.0166 (8)	0.0110 (7)	0.0027 (6)	0.0012 (5)	-0.0001 (6)
C16	0.0114 (7)	0.0170 (7)	0.0126 (7)	0.0031 (6)	0.0038 (6)	0.0059 (6)
C17	0.0109 (7)	0.0141 (7)	0.0176 (8)	0.0011 (6)	0.0015 (6)	0.0025 (6)
C18	0.0138 (7)	0.0155 (7)	0.0142 (7)	0.0012 (6)	0.0015 (6)	-0.0029 (6)
C19	0.0114 (7)	0.0168 (8)	0.0114 (7)	0.0007 (6)	0.0029 (6)	0.0008 (6)
C20	0.0149 (7)	0.0271 (9)	0.0083 (7)	0.0013 (6)	0.0018 (6)	0.0002 (6)
C21	0.0209 (9)	0.0358 (11)	0.0163 (8)	-0.0070 (7)	0.0044 (7)	-0.0073 (8)

Geometric parameters (\AA , $^{\circ}$)

Br1—C16	1.8939 (17)	C14—C19	1.389 (2)
Cl2—C18	1.7480 (18)	C14—C15	1.410 (2)
O1—C15	1.356 (2)	C15—C16	1.398 (2)
O2—C3	1.2517 (19)	C16—C17	1.385 (2)
O3—C9	1.223 (2)	C17—C18	1.385 (2)
O4—C21	1.408 (2)	C18—C19	1.385 (2)
O1—H1A	0.8400	C20—C21	1.529 (3)
O4—H4	0.8400	C1—H1	1.0000
N1—C20	1.481 (2)	C4—H4A	0.9900
N1—C7	1.384 (2)	C4—H4B	0.9900
N1—C13	1.406 (2)	C5—H5A	0.9900
C1—C14	1.532 (2)	C5—H5B	0.9900
C1—C2	1.510 (2)	C6—H6A	0.9900
C1—C8	1.506 (2)	C6—H6B	0.9900
C2—C3	1.443 (2)	C10—H10A	0.9900
C2—C7	1.371 (2)	C10—H10B	0.9900
C3—C4	1.500 (2)	C11—H11A	0.9900
C4—C5	1.524 (2)	C11—H11B	0.9900
C5—C6	1.524 (2)	C12—H12A	0.9900
C6—C7	1.501 (2)	C12—H12B	0.9900
C8—C13	1.351 (2)	C17—H17	0.9500
C8—C9	1.470 (2)	C19—H19	0.9500
C9—C10	1.503 (2)	C20—H20A	0.9900
C10—C11	1.510 (3)	C20—H20B	0.9900
C11—C12	1.530 (2)	C21—H21A	0.9900
C12—C13	1.513 (2)	C21—H21B	0.9900
C15—O1—H1A	109.00	C2—C1—H1	108.00
C21—O4—H4	109.00	C8—C1—H1	108.00
C13—N1—C20	119.40 (13)	C14—C1—H1	108.00
C7—N1—C20	121.30 (13)	C3—C4—H4A	110.00
C7—N1—C13	119.30 (13)	C3—C4—H4B	110.00
C2—C1—C14	111.47 (13)	C5—C4—H4A	110.00
C2—C1—C8	109.12 (13)	C5—C4—H4B	110.00
C8—C1—C14	112.17 (13)	H4A—C4—H4B	108.00
C1—C2—C7	121.55 (14)	C4—C5—H5A	109.00
C1—C2—C3	117.41 (13)	C4—C5—H5B	109.00
C3—C2—C7	121.02 (14)	C6—C5—H5A	109.00
O2—C3—C4	120.40 (14)	C6—C5—H5B	109.00
C2—C3—C4	119.06 (14)	H5A—C5—H5B	108.00
O2—C3—C2	120.52 (14)	C5—C6—H6A	109.00
C3—C4—C5	110.03 (14)	C5—C6—H6B	109.00
C4—C5—C6	111.52 (14)	C7—C6—H6A	109.00
C5—C6—C7	113.07 (14)	C7—C6—H6B	109.00
C2—C7—C6	121.69 (14)	H6A—C6—H6B	108.00
N1—C7—C6	117.60 (13)	C9—C10—H10A	109.00
N1—C7—C2	120.64 (14)	C9—C10—H10B	109.00
C1—C8—C13	122.65 (14)	C11—C10—H10A	109.00

C9—C8—C13	121.91 (14)	C11—C10—H10B	109.00
C1—C8—C9	115.43 (13)	H10A—C10—H10B	108.00
C8—C9—C10	118.16 (14)	C10—C11—H11A	109.00
O3—C9—C8	120.16 (15)	C10—C11—H11B	109.00
O3—C9—C10	121.64 (15)	C12—C11—H11A	109.00
C9—C10—C11	111.44 (15)	C12—C11—H11B	109.00
C10—C11—C12	112.04 (14)	H11A—C11—H11B	108.00
C11—C12—C13	111.51 (14)	C11—C12—H12A	109.00
N1—C13—C8	120.51 (14)	C11—C12—H12B	109.00
N1—C13—C12	117.61 (13)	C13—C12—H12A	109.00
C8—C13—C12	121.85 (14)	C13—C12—H12B	109.00
C1—C14—C15	120.25 (14)	H12A—C12—H12B	108.00
C15—C14—C19	119.57 (14)	C16—C17—H17	121.00
C1—C14—C19	120.17 (14)	C18—C17—H17	121.00
O1—C15—C16	119.10 (14)	C14—C19—H19	120.00
C14—C15—C16	118.28 (14)	C18—C19—H19	120.00
O1—C15—C14	122.61 (14)	N1—C20—H20A	110.00
C15—C16—C17	122.46 (15)	N1—C20—H20B	110.00
Br1—C16—C15	118.54 (12)	C21—C20—H20A	110.00
Br1—C16—C17	119.00 (13)	C21—C20—H20B	110.00
C16—C17—C18	117.80 (15)	H20A—C20—H20B	108.00
Cl2—C18—C19	118.86 (13)	O4—C21—H21A	109.00
C17—C18—C19	121.64 (15)	O4—C21—H21B	109.00
Cl2—C18—C17	119.50 (13)	C20—C21—H21A	109.00
C14—C19—C18	120.20 (15)	C20—C21—H21B	109.00
N1—C20—C21	110.34 (14)	H21A—C21—H21B	108.00
O4—C21—C20	111.92 (16)		
C7—N1—C13—C12	161.29 (14)	C5—C6—C7—C2	12.9 (2)
C7—N1—C20—C21	103.01 (17)	C5—C6—C7—N1	-170.23 (14)
C13—N1—C20—C21	-77.86 (18)	C1—C8—C13—C12	-178.67 (15)
C7—N1—C13—C8	-16.8 (2)	C9—C8—C13—N1	-179.81 (14)
C13—N1—C7—C2	11.4 (2)	C1—C8—C9—O3	1.0 (2)
C20—N1—C7—C2	-169.52 (15)	C1—C8—C9—C10	-176.73 (14)
C20—N1—C13—C12	-17.9 (2)	C13—C8—C9—O3	-179.82 (15)
C20—N1—C13—C8	164.11 (15)	C9—C8—C13—C12	2.2 (2)
C20—N1—C7—C6	13.6 (2)	C13—C8—C9—C10	2.4 (2)
C13—N1—C7—C6	-165.53 (14)	C1—C8—C13—N1	-0.7 (2)
C14—C1—C2—C7	98.95 (17)	O3—C9—C10—C11	151.65 (16)
C8—C1—C14—C15	-137.57 (15)	C8—C9—C10—C11	-30.6 (2)
C2—C1—C8—C13	20.3 (2)	C9—C10—C11—C12	54.0 (2)
C14—C1—C8—C9	75.41 (17)	C10—C11—C12—C13	-49.4 (2)
C2—C1—C8—C9	-160.57 (13)	C11—C12—C13—N1	-156.62 (15)
C2—C1—C14—C15	99.74 (17)	C11—C12—C13—C8	21.4 (2)
C2—C1—C14—C19	-79.27 (18)	C1—C14—C15—O1	-1.0 (2)
C8—C1—C14—C19	43.4 (2)	C19—C14—C15—C16	-2.4 (2)
C8—C1—C2—C7	-25.5 (2)	C1—C14—C19—C18	179.85 (15)
C14—C1—C2—C3	-82.55 (17)	C15—C14—C19—C18	0.8 (2)
C14—C1—C8—C13	-103.74 (17)	C1—C14—C15—C16	178.62 (14)

C8—C1—C2—C3	153.02 (13)	C19—C14—C15—O1	177.99 (15)
C1—C2—C7—C6	-172.02 (14)	C14—C15—C16—Br1	-178.05 (12)
C1—C2—C3—O2	1.8 (2)	O1—C15—C16—Br1	1.6 (2)
C7—C2—C3—C4	2.2 (2)	O1—C15—C16—C17	-177.82 (15)
C1—C2—C7—N1	11.2 (2)	C14—C15—C16—C17	2.5 (2)
C7—C2—C3—O2	-179.65 (15)	C15—C16—C17—C18	-1.1 (2)
C3—C2—C7—C6	9.5 (2)	Br1—C16—C17—C18	179.52 (12)
C1—C2—C3—C4	-176.32 (14)	C16—C17—C18—Cl2	-179.70 (13)
C3—C2—C7—N1	-167.22 (14)	C16—C17—C18—C19	-0.6 (2)
C2—C3—C4—C5	-34.7 (2)	C17—C18—C19—C14	0.7 (3)
O2—C3—C4—C5	147.20 (15)	Cl2—C18—C19—C14	179.80 (12)
C3—C4—C5—C6	55.54 (18)	N1—C20—C21—O4	-174.31 (14)
C4—C5—C6—C7	-45.49 (19)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1A \cdots O2	0.84	1.88	2.6749 (19)	158
O4—H4 \cdots O2 ⁱ	0.84	1.94	2.782 (2)	176
C1—H1 \cdots O1	1.00	2.48	2.918 (2)	106
C6—H6B \cdots O3 ⁱⁱ	0.99	2.33	3.051 (2)	129
C20—H20A \cdots O3 ⁱⁱ	0.99	2.53	3.486 (2)	163
C20—H20B \cdots O1 ⁱ	0.99	2.57	3.492 (2)	154

Symmetry codes: (i) $x-1/2, -y+3/2, z+1/2$; (ii) $x+1/2, -y+3/2, z+1/2$.