

Outcomes in patients with non-ST-elevation acute coronary syndrome randomly assigned to invasive *versus* conservative treatment strategies: A meta-analysis

Ying-Qing Li,^{1*#} Na Liu,^{1#} Jian-Hua Lu^{1#}

¹Guangzhou Medical University, Guangzhou First People's Hospital, Emergency Department 1 Panfu Road, Guangzhou, People's Republic of China. [#]Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Guangzhou, People's Republic of China.

OBJECTIVE: The goal of the present study was to compare the prognoses of patients with non-ST-elevation acute coronary syndromes who were treated with invasive or conservative treatment strategies.

METHODS: We performed a meta-analysis of studies of patients with non-ST-elevation acute coronary syndromes to assess the benefits of an invasive strategy vs. a conservative strategy for short- and long-term survival. We searched PubMed for studies published from 1990 to November 2012 that investigated the effects of an invasive vs. conservative strategy in patients with non-ST-elevation acute coronary syndromes. The following search terms were used: "non-ST-elevation myocardial infarction", "unstable angina", "acute coronary syndromes", "invasive strategy", and "conservative strategy". The primary endpoints were all-cause mortality at 30 days and 1 year.

RESULTS: Seven published studies were included in the present meta-analysis. The pooled analyses show that an invasive strategy decreased the risk of death (risk ratio [0.839] [95% confidence interval {0.648-1.086}; I^2 , 86.46%]) compared to a conservative strategy over a 30-day-period. Furthermore, invasive treatment also decreased patient mortality (risk ratio [0.276] [95% confidence interval {0.259-0.294}; I^2 , 94.58%]) compared to a conservative strategy for one year.

CONCLUSION: In non-ST-elevation acute coronary syndromes, an invasive strategy is comparable to a conservative strategy for decreasing short- and long-term mortality rates.

KEYWORDS: Non-ST-elevation Acute Coronary Syndrome; Invasive Strategy; Conservative Strategy; Meta-analysis; Mortality.

Li YQ, Liu N, Lu JH. Outcomes in patients with non-ST-elevation acute coronary syndrome randomly assigned to invasive *versus* conservative treatment strategies: A meta-analysis. *Clinics*. 2014;69(6):398-404.

Received for publication on October 1, 2013; First review completed on November 7, 2013; Accepted for publication on November 27, 2013

E-mail: hare0914@aliyun.com

*corresponding author

#contributed equally to this study

Tel.: +86 20-81048888

INTRODUCTION

Non-ST-elevation acute coronary syndrome (NSTEMI) includes non-ST-elevation myocardial infarction (NSTEMI) and unstable angina (UA), which accounted for more than two thirds of all cases of acute coronary syndrome (ACS) in the United States in 2009 (1). Furthermore, cardiovascular disease, including NSTEMI, still remains the leading

cause of mortality worldwide (2). NSTEMI represents an increasingly frequent cause of hospital admission, as it is the most frequent presentation of coronary instability in patients with prior cardiac events or coronary revascularizations (3). Compared with ST-elevation myocardial infarction (STEMI), NSTEMI is usually associated with a partial or transient coronary obstruction that results in a temporary reduction in coronary blood flow, and microemboli have also been implicated in the pathophysiology of NSTEMI (4). Recognizing the difference between NSTEMI and STEMI is critical when developing therapeutic strategies for NSTEMI. For instance, thrombolytic management is not indicated in NSTEMI management (5). Experimental data from randomized clinical trials have clearly demonstrated the efficacy of conservative or invasive approaches in the above mentioned patients (6-8). In clinical

Copyright © 2014 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

DOI: 10.6061/clinics/2014(06)06



Table 1 - Characteristics of the included NSTEMI-ACS studies.

Sources	Study Design	Number of Deaths/ Patients who Underwent Invasive Treatment for 30 days	Number of Deaths/ Patients who Underwent Conservative Treatment for 30 days	Number of Deaths/ Patients who Underwent Invasive Treatment for 1 year	Number of Deaths/ Patients who Underwent Conservative Treatment for 1 year
OCS					
Leslie et al., (11) 2003* [△]	Retrospective	100/4535	97/3362	181/4535	309/3362
Timm et al., (12) 2007 [△]	Prospective			101/945	227/815
Cindy et al., (13) 2008 [△]	Prospective			55/1382	82/754
Rogério et al., (14) 2009* [△]	Prospective	1/386	13/334	16/386	46/334
Kouraki et al., (15) 2011 [△]	Prospective			98/1215	185/771
Joakim et al., (16) 2011 [△]	Prospective			814/28112	2381/18343
RCTs					
William et al., (17) 1998* [△]	RCT	33/442	7/256	52/442	24/256

NSTEMI-ACS: non-ST-elevation acute coronary syndrome; RCT: randomized controlled trials. * Included in the meta-analysis of all-cause mortality within 30 days for patients with NSTEMI-ACS. [△] The seven studies included in the meta-analysis of all-cause mortality within 1 year for NSTEMI-ACS patients.

practice, physicians caring for NSTEMI-ACS patients are faced with immediate decisions about substantially different therapeutic strategies that may significantly affect short- and long-term outcomes (9).

An increasing number of studies have demonstrated that an invasive treatment strategy rather than a conservative approach improves the outcomes of NSTEMI-ACS patients (6-8,10-16). However, some studies have shown that a conservative strategy provides the same benefit as an invasive strategy for NSTEMI-ACS patients, and clearly shown that the efficacy of invasive treatment was not definite (17,18).

According to the 2012 American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines (19), medical therapy remains a cornerstone in managing patients with NSTEMI-ACS, and we found that in the above mentioned studies that demonstrated the benefit of an invasive strategy, the administration of antithrombotics (antiplatelets or oral anticoagulants) statins was the primary step prior to interventional therapy. However, in the conservative treatment group, the NSTEMI-ACS patients also received interventional therapy during the index hospitalization when medical therapy failed or if substantial residual ischemia was documented in some of the randomized control trials (RCTs) (6,8,9,18,20,21).

It is necessary to clearly distinguish between the different efficacies for invasive and conservative strategies for NSTEMI-ACS patients. Therefore, we performed the present meta-analysis to analyze the benefits between invasive and conservative strategies for NSTEMI-ACS patients.

METHODS

We followed the Preferred Reporting Items for Systematic reviews and Meta-Analysis (PRISMA) guidelines (22) for randomized clinical trials and the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) guidelines (23) for observational cohort studies in the present meta-analysis.

Study search strategy

Accessing the PubMed database, we conducted a literature search for research papers published from 1990 to November 2012. The following search terms and keywords were used: “non-ST-elevation myocardial infarction”, “unstable angina”, “acute coronary syndromes”, “invasive

strategy”, and “conservative strategy”. In addition, we also manually checked the reference list of each article. The main focus of the study was RCTs and observational cohort studies (OCS). We excluded those research papers with the following keywords: “review”, “review literature”, “review of reported cases”, “review, academic”, “review, multi-case”, “review, tutorial”, “case reports”, “congresses”, “interview”, “overall”, “comment”, “letter”, “practice guideline”, “scientific integrity review”, “news”, “news-paper article”, and “address”. The search was limited by the search words of “>19 years”, “Publication Date from 1990/01/01 to 2012/11/01”, “English”, and “Human”.

Study Selection

Eligible patients had experienced NSTEMI-ACS (i.e., NSTEMI or UA) that was treated with invasive or conservative treatment during the admission. The NSTEMI-ACS diagnosis was based on the American College of Cardiology Foundation/American Heart Association (ACCF/AHA) Guidelines 2012 for NSTEMI-ACS (19). In our present meta-analysis, the invasive group contained patients who underwent cardiac catheterization, diagnostic angiography, coronary artery bypass graft (CABG) surgery, and intra-aortic balloon pump (IABP) performed during the index hospitalization with or without revascularization; the conservative group contained patients who did not undergo catheterization, angiography, CABG, or IABP during the index hospitalization. Conservative therapy consists of vigorous anti-ischemic therapy with nitrates, statins, β -blockers, and (sometimes) calcium channel blockers, as well as antithrombotic therapy with aspirin, clopidogrel, or ticagrelor, and low molecular weight heparin (LMWH) while the patients are in the hospital. In ACS patients, antithrombotics can effectively inhibit thrombus formation and coagulation, and statins can play important roles in stabilizing plaque, inhibiting vascular endothelial inflammation, improving endothelial function, and reducing cholesterol levels. The main endpoints were death from all causes at 30 days and 1 year. Patients who received both conservative and interventional therapy in the hospital were excluded.

We originally planned to perform the meta-analysis in two parts: RCTs and OCS analyzed in two separate meta-analyses; however, there was only one RCT included in the present work (17). For this reason, we retrospectively



analyzed all included studies at two different time points (30 days and 1 year) by meta-analysis.

Data Extraction

In addition to information about the study design, characteristics, and sample size, we extracted the following data from the research papers: the actual number of survivors and non-survivors and corresponding cohort sizes. The main outcome variables were 30-day and 1-year survival rates.

Statistical Analysis

We performed the meta-analysis using Comprehensive Meta-analysis software, version 2.2.064 (Biostat, Englewood, NJ, USA). Risk ratios (RR) and 95% confidence intervals (95% CI) were calculated for each study and pooled with a random effects model because the heterogeneity of all meta-analyses could not be ignored, as indicated by an I^2 statistics above 50% and a $p < 0.10$. Heterogeneity among studies was formally assessed by the Q and I^2 statistics. Publication bias was assessed by calculating the fail-safe number (Nfs) (24) according to the formula $Nfs_{.05} = (\Sigma Z/1.645)^2 - N$. A fail-safe number indicates the number of non-significant,

unpublished studies that would need to be added to a meta-analysis to reduce the overall statistically significant results to non-significance. If the number is large relative to the number of observed studies, the summary conclusions are fairly reliable. All analyses were performed using the SPSS 17.0 software package (SPSS, Inc., Chicago, IL, USA), and p -values < 0.05 were considered to be statistically significant.

RESULTS

Seven studies were included in the meta-analysis; one study was an RCT, and the other six were OCS; the total number of patients in all seven studies was 61,652 (11-17) (Figure 1) (Table 1).

There were three studies in the first meta-analysis (11,14,17); we used all-cause mortality at 30 days as the primary endpoint, and 9,315 patients were included in the systematic review. After the data were analyzed, we found that an invasive strategy was associated with a statistically significant 15.6% decreased mortality (RR, 0.839 [95% CI, 0.648-1.086]) compared to conservative strategy during the 30 days after hospital discharge (Figure 2). In this meta-analysis,




Figure 1 - Study selection process for the studies included in the Meta-Analysis (based on the PRIMA guidelines).



Meta-analysis




Figure 2 - Primary analysis of the OCS in NSTEMI-ACS within 30 days.

we used a random-effects model to consider the effect of heterogeneity among the studies ($Q=14.77, p=0.001, I^2=86.46\%$). The calculation for the Nfs.05 is 20.97, which suggests that there need to be 21 studies with a negative conclusion to reverse the conclusion. The conclusion from this meta-analysis is convincing, with little risk of publication bias.

In the next systematic review, the seven studies were included in the meta-analysis. We compared the mortality of patients treated with an invasive strategy and a conservative strategy for one year after hospital discharge. Of the 37,017 patients with NSTEMI-ACS who received invasive treatments and were discharged, there were 1,317 patients who died within one year. However, 24,635 patients received conservative treatment and were discharged and 3,254 patients died within one year. After the data were pooled, we found that invasive therapy significantly reduced mortality by 73.1% within one year after discharge (RR, 0.276 [95% CI, 0.259-0.294]) compared to conservative therapy (Figure 3). In the second meta-analysis, we also used a random-effects model to consider the effect of heterogeneity among the studies ($Q=110.7, p<0.001, I^2=94.58\%$). The sensitivity analysis reveals that 2,505 negative studies have been missed to reverse the statistical significance (Nfs.05 = 2505). The conclusion from the second meta-analysis is convincing, and the publication bias could be neglected.

DISCUSSION

NSTEMI-ACS is responsible for approximately 1 million admissions to U.S. hospitals and twice as many to European hospitals each year and is associated with an in-hospital mortality rate of approximately 5% (25). Worldwide, more than 4 million people each year are estimated to have a NSTEMI, and long-term mortality is higher in patients with NSTEMI-ACS than in those with ST-elevation acute coronary syndrome (STEMI-ACS) (26,27). In 2012, the ACCF/AHA published an update to their 2011 guidelines for the management of patients with NSTEMI-ACS (19,28). According to the new guidelines, selecting either invasive or conservative strategy is one of the most important decisions that must be made in managing NSTEMI-ACS patients. The ACC/AHA guidelines continue to support an early invasive strategy in patients with high-risk NSTEMI-ACS without serious comorbidities or contraindications to angiography; in contrast, an early conservative approach should be selected for low-risk patients.

The results of our present study were consistent with the new 2012 guidelines, and the two systematic meta-analyses showed that compared to a conservative strategy, an invasive strategy is associated with reduced short- and long-term mortality in patients with NSTEMI-ACS after hospital discharge.

Meta-analysis




Figure 3 - Primary analysis of the NSTEMI-ACS OCS within 1 year.



Meta-analyses are statistical tools for pooling existing evidence; therefore, we considered the strength of the evidence from all seven studies (11-17). We searched and analyzed those articles addressing the treatment strategies for NSTEMI-ACS. Furthermore, we studied the details of the related studies according to the 2012 guidelines (19); these studies include well-known studies such as the FRISC II (Fragmin and Fast Revascularization During Instability in Coronary Artery Disease II) (4), the RITA 3 (Randomized Intervention Trial of Unstable Angina 3) trial (8), TACTICS-TIMI 18 (Treat angina with Aggrastat and Determine Cost of Therapy with an Invasive or Conservative Strategy-Thrombolysis in Myocardial Infarction 18) trial (7), ICTUS (Invasive vs Conservative Treatment in Unstable Coronary Syndromes) trial (29), TIMI IIIB (Thrombolysis in Myocardial Infarction IIIB) (30), MATE (Medicine vs Angiography in Thrombolytic Exclusion) (31), and VINO (Value of First Day Angiography/Angioplasty In Evolving Non-ST Segment Elevation Myocardial Infarction) (32). Based on the above mentioned studies, we found that an invasive strategy could improve the short- or long-term survival rates of NSTEMI-ACS patients compared to a conservative strategy. In addition, a related meta-analysis demonstrated that an invasive strategy had comparable benefits in men and high-risk women for reducing the composite end point of death, MI, or re-hospitalization with NSTEMI-ACS (33).

In our present work, we did not distinguish between gender, age, risk factors, and complications for the two strategies; however, we focused on the all-cause mortality at 30 days and 1 year after hospital discharge. At the same time, we strictly limited the definition of the invasive and conservative strategies. The invasive group contained patients who underwent cardiac catheterization during the index hospitalization with or without revascularization; in contrast, patients who did not receive catheterization in the hospital were included in the conservative group. Catheterization included angiography and percutaneous coronary intervention; moreover, patients who underwent CABG surgery and IABP in the hospital were included in the invasive group. Our purpose was to assess whether a particular strategy decreased all-cause mortality in NSTEMI-ACS patients. Boden et al. (17) demonstrated that some NSTEMI-ACS patients in the conservative group also received interventional therapy, but we analyzed the total details of the study and regrouped the patients into the conservative or invasive management groups; then, we retrospectively analyzed the data and eliminated the interference from the two therapies with one another. Although the William et al. study was a RCT, we included it in the present work after addressing the interfering factors. Thus, seven studies were included in our meta-analysis.

While analyzing our data, we assessed other reviews that compared the effectiveness of both strategies in NSTEMI-ACS patients. These reviews include Núñez et al. (9), Boden (34,35), Neumann et al. (36), Choudhry et al. (37), O'Donoghue et al. (33), Hoenig et al. (38,39), Navarese et al. (40), and Damman P et al. (41) Based on these reviews, we found that over the past 10 years, there is still controversy regarding invasive *versus* conservative strategy in NSTEMI-ACS patients. Compared with a conservative strategy for NSTEMI-ACS patients, some studies reported that an invasive therapy provided comparable benefits and reduced the rates of fatal or nonfatal re-infarction and

hospital readmission but not all-cause mortality (9,33,35,37). However, other studies found that an invasive strategy was not related to improvements in long-term survival and reduced the rate of revascularization compared with conservative management (34,36,40,41).

Furthermore, the above mentioned studies did not strictly distinguish between or define conservative and invasive strategies; for instance, the administration of antithrombotics was the primary step prior to interventional therapy in invasive strategy. However, in some RCTs, patients with NSTEMI-ACS in the conservative treatment group also received interventional therapy during the index hospitalization when medical therapy failed or if substantial residual ischemia was documented. Based on these facts, we defined the concept of conservative and invasive strategies strictly as described in our article and performed the meta-analysis based on these definitions.

Inspired by the above mentioned studies, we strictly redefined the conservative and invasive strategy and retrospectively analyzed seven studies. The NSTEMI-ACS patients who received interventional management in the hospital were included in the invasive group, whereas the other patients were placed in the conservative group. The main end-point events were 30-day and 1-year all-cause mortality. After performing two systematic meta-analyses, we found that invasive therapy improved the prognosis of patients with NSTEMI-ACS (compared with conservative therapy) and reduced the short- and long-term mortality.

We also noted that the results of the Boden et al. study (17) demonstrated that conservative therapies could play more beneficial roles in NSTEMI-ACS patients than did invasive strategies. What are the reasons for the opposite results? We analyzed their findings in detail and further studied its quotation from Ferry et al. (42) and found that the authors excluded 247 patients with high-risk ischemic complications (e.g., unstable angina after infarction, congestive heart failure that did not respond to medical therapy, cardiogenic shock, or symptomatic ventricular arrhythmia) during the first 48 hours after the onset of infarction. Moreover, the invasive theories, techniques and equipment were not then as well established as they are currently. With the development of modern percutaneous coronary intervention technology, invasive strategies show more benefit than conservative medical therapies in NSTEMI-ACS patients.

For patients with NSTEMI-ACS (NSTEMI/UA), an invasive strategy reduces short- and long-term all-cause mortality compared with a conservative strategy. An invasive strategy is preferable to a conservative strategy in treating NSTEMI-ACS. Of course, these results require validation in future RCTs.

■ ACKNOWLEDGMENTS

This study was supported by grants from the National Natural Science Foundation of China (81372023). The funders played no roles in the study design, data collection, analysis, decision to publish, or manuscript preparation. We thank all our colleagues who supported this work.

■ AUTHOR CONTRIBUTIONS

Ying-Qing Li, Na Liu, and Jian-Hua Lu contributed equally to this study. Ying-Qing Li conceived the study, designed the trial and obtained research funding. Liu NA supervised the data collection. Jian-Hua Lu and Ying-Qing Li discussed the study results and implications and commented on the



manuscript at all stages. Ying-Qing Li is the corresponding author and was responsible for the manuscript in its entirety.

REFERENCES

- Giugliano RP, Braunwald E. The Year in Non-ST-Segment Elevation Acute Coronary Syndrome. *J Am Coll Cardiol*. 2012;60(21):2127-39, <http://dx.doi.org/10.1016/j.jacc.2012.08.972>.
- Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, et al. Executive summary: heart disease and stroke statistics – 2012 update: a report from the American Heart Association. *Circulation*. 2012;125(1): 188-97.
- Di Chiara A, Fresco C, Savonitto S, Greco C, Lucci D, Gonzini L, et al. Epidemiology of non-ST elevation acute coronary syndromes in the Italian cardiology network: the BLITZ-2 study. *Eur Heart J* 2006;27(4): 393-405.
- DeWood MA, Stifter WF, Simpson CS, Spores J, Eugster GS, Judge TP, et al. Coronary arteriographic findings soon after non-Q-wave myocardial infarction. *N Engl J Med*. 1986;315(7):417-23.
- Anderson JL, Adams CD, Antman EM, Bridges CR, Califf RM, Casey DE Jr, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with unstable angina/Non-ST-elevation myocardial infarction (updating the 2007 guideline and replacing the 2011 focused update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2011;123(18):e426-e579, <http://dx.doi.org/10.1161/CIR.0b013e318212bb8b>.
- Invasive compared with non-invasive treatment in unstable coronary-artery disease: FRISC II prospective randomised multicentre study. FRagmin and Fast Revascularisation during InStability in Coronary artery disease Investigators. *Lancet*. 1999;354(9180):708-15.
- Cannon CP, Weintraub WS, Demopoulos LA, Vicari R, Frey MJ, Lakkis N, et al. Comparison of early invasive and conservative strategies in patients with unstable coronary syndromes treated with the glycoprotein IIb/IIIa inhibitor tirofiban. *N Engl J Med*. 2001;344(25):1879-87.
- Fox KA, Poole-Wilson PA, Henderson RA, Clayton TC, Chamberlain DA, Shaw TR, et al. Interventional versus conservative treatment for patients with unstable angina or non-ST-elevation myocardial infarction: the British Heart Foundation RITA 3 randomised trial. *Randomized Intervention Trial of Unstable Angina*. *Lancet*. 2002;360(9335):743-51.
- Bach RG, Cannon CP, Weintraub WS, DiBattiste PM, Demopoulos LA, Anderson HV, et al. The effect of routine, early invasive management on outcome for elderly patients with non-ST-segment elevation acute coronary syndromes. *Ann Intern Med*. 2004;141(3):186-95, <http://dx.doi.org/10.7326/0003-4819-141-3-200408030-00007>.
- Diderholm E, Andrén B, Frostfeldt G, Genberg M, Jernberg T, Lagerqvist B, et al. ST depression in ECG at entry indicates severe coronary lesions and large benefits of an early invasive treatment strategy in unstable coronary artery disease; the FRISC II ECG substudy. *The Fast Revascularisation during InStability in Coronary artery disease*. *Eur Heart J*. 2002;23(1):41-9, <http://dx.doi.org/10.1053/euhj.2001.2694>.
- Cho L, Bhatt DL, Marso SP, Brennan D, Holmes DR Jr, Califf RM, et al. An invasive strategy is associated with decreased mortality in patients with unstable angina and non-ST-elevation myocardial infarction: GUSTO IIb trial. *Am J Med*. 2003;114(2):106-11.
- Bauer T, Koeth O, Jünger C, Heer T, Wienbergen H, Gitt A, et al. Effect of an invasive strategy on in-hospital outcome in elderly patients with non-ST-elevation myocardial infarction. *Eur Heart J*. 2007;28(23):2873-8, <http://dx.doi.org/10.1093/eurheartj/ehm464>.
- Lee CH, Tan M, Yan AT, Yan RT, Fitchett D, Grima EA, et al. Use of cardiac catheterization for non-ST-segment elevation acute coronary syndromes according to initial risk: reasons why physicians choose not to refer their patients. *Arch Intern Med*. 2008;168(3):291-6, <http://dx.doi.org/10.1001/archinternmed.2007.78>.
- Teixeira R, Lourenço C, Baptista R, Jorge E, António N, Monteiro S, et al. Invasive versus conservative strategy in non-ST elevation acute coronary syndromes: data from a single Portuguese center. *Rev Port Cardiol*. 2009;28(4):355-73.
- Kleopatra K, Muth K, Zahn R, Bauer T, Koeth O, Jünger C, et al. Effect of an invasive strategy on in-hospital outcome and one-year mortality in women with non-ST-elevation myocardial infarction. *Int J Cardiol*. 2011; 153(3):291-5.
- Alfredsson J, Lindbäck J, Wallentin L, Swahn E. Similar outcome with an invasive strategy in men and women with non-ST-elevation acute coronary syndromes: from the Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). *Eur Heart J* 2011;32(24):3128-36, <http://dx.doi.org/10.1093/eurheartj/ehr349>.
- Boden WE, O'Rourke RA, Crawford MH, Blaustein AS, Deedwania PC, Zoble RG, et al. Outcomes in patients with acute non-Q-wave myocardial infarction randomly assigned to an invasive as compared with a conservative management strategy. Veterans Affairs Non-Q-Wave Infarction Strategies in Hospital (VANQWISH) Trial Investigators. *N Engl J Med*. 1998;338(25):1785-92.
- Savonitto S, Cavallini C, Petronio AS, Murena E, Antonicelli R, Sacco A, et al. Early aggressive versus initially conservative treatment in elderly patients with non-ST-segment elevation acute coronary syndrome: a randomized controlled trial. *JACC Cardiovasc Interv*. 2012;5(9):906-16, <http://dx.doi.org/10.1016/j.jcin.2012.06.008>.
- 2012 Writing Committee Members, Jneid H, Anderson JL, Wright RS, Adams CD, Bridges CR, et al. 2012 ACCF/AHA focused update of the guideline for the management of patients with unstable angina/Non-ST-elevation myocardial infarction (updating the 2007 guideline and replacing the 2011 focused update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2012;126(7):875-910.
- Lagerqvist B, Husted S, Kontny F, Ståhle E, Swahn E, Wallentin L, et al. 5-year outcomes in the FRISC-II randomised trial of an invasive versus a non-invasive strategy in non-ST-elevation acute coronary syndrome: a follow-up study. *Lancet*. 2006;368(9540):998-1004, [http://dx.doi.org/10.1016/S0140-6736\(06\)69416-6](http://dx.doi.org/10.1016/S0140-6736(06)69416-6).
- Núñez J, Sanchis J, Núñez E, Bodi V, Bertomeu-González V, Bosch MJ, et al. Prognostic differences between routine invasive and conservative strategies for the management of high-risk, non-ST segment acute coronary syndromes: Experience from two consecutive periods in a single center. *Eur J Intern Med*. 2007;18(5):409-16.
- Moher D, Liberati A, Tetzlaff J, Altman DG. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2008;149(4):264-9.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of Observational Studies in Epidemiology: A Proposal for Reporting. *JAMA*. 2000;283(15):2008-12, <http://dx.doi.org/10.1001/jama.283.15.2008>.
- Rosenthal R. The file drawer problem and tolerance for null results. *Psychol Bull*. 1979;86:638-41, <http://dx.doi.org/10.1037/0033-2909.86.3.638>.
- Peterson ED, Roe MT, Mulgund J, DeLong ER, Lytle BL, Brindis RG, et al. Association between hospital process performance and outcomes among patients with acute coronary syndromes. *JAMA*. 2006;295(16):1912-20, <http://dx.doi.org/10.1001/jama.295.16.1912>.
- White HD, Chew DP. Acute myocardial infarction. *Lancet*. 2008;372(9638):570-84, [http://dx.doi.org/10.1016/S0140-6736\(08\)61237-4](http://dx.doi.org/10.1016/S0140-6736(08)61237-4).
- Terkelsen CJ, Lassen JF, Nørgaard BL, Gerdes JC, Jensen T, Gotzsche LB, et al. Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. *Eur Heart J*. 2005;26(1):18-26.
- Wright RS, Anderson JL, Adams CD, Bridge CR, Casey DE Jr, Ettinger SM, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with unstable angina/Non-ST-elevation myocardial infarction (updating the 2007 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2011;123(18):2022-60.
- de Winter RJ, Windhausen F, Cornel JH, Dunselman PH, Janus CL, Bendermacher PE, et al. Early invasive versus selectively invasive management for acute coronary syndromes. *N Engl J Med*. 2005;353(11): 1095-104.
- Effects of tissue plasminogen activator and a comparison of early invasive and conservative strategies in unstable angina and non-Q-wave myocardial infarction. Results of the TIMI IIIb Trial. *Thrombolysis in Myocardial Ischemia*. *Circulation*. 1994;89(4):1545-56.
- McCullough PA, O'Neill WW, Graham M, Stomel RJ, Rogers F, David S, et al. A prospective randomized trial of triage angiography in acute coronary syndromes ineligible for thrombolytic therapy. Results of the medicine versus angiography in Thrombolytic Exclusion (MATE) trial. *J Am Coll Cardiol*. 1998;32(3):596-605.
- Spacek R, Widimský P, Straka Z, Jiresová E, Dvůrák J, Polásek R, et al. Value of first day angiography/angioplasty in evolving Non-ST segment elevation myocardial infarction: an open multicenter randomized trial. *The VINO Study*. *Eur Heart J*. 2002;23(3):230-8, <http://dx.doi.org/10.1053/euhj.2001.2735>.
- O'Donoghue M, Boden WE, Braunwald E, Cannon CP, Clayton TC, de Winter RJ, et al. Early invasive vs conservative treatment strategies in women and men with unstable angina and non-ST-segment elevation myocardial infarction: a meta-analysis. *JAMA*. 2008;300(1):71-80, <http://dx.doi.org/10.1001/jama.300.1.71>.
- Boden WE. Avoidance of routine revascularization in the management of patients with non-ST-segment elevation acute coronary syndromes. *Am J Cardiol*. 2000;86(12B):42M-47M, [http://dx.doi.org/10.1016/S0002-9149\(00\)01480-6](http://dx.doi.org/10.1016/S0002-9149(00)01480-6).
- Boden WE. "Routine invasive" versus "selective invasive" approaches to non-ST-segment elevation acute coronary syndromes management in the post-stent/platelet inhibition era. *J Am Coll Cardiol*. 2003;41(4 Suppl S):113S-122S, [http://dx.doi.org/10.1016/S0735-1097\(02\)02963-7](http://dx.doi.org/10.1016/S0735-1097(02)02963-7).
- Neumann FJ, Kastrati A, Pogatsa-Murray G, Mehilli J, Bollwein H, Bestehorn HP, et al. Evaluation of prolonged antithrombotic pretreatment



- ("cooling-off" strategy) before intervention in patients with unstable coronary syndromes: a randomized controlled trial. *JAMA*. 2003; 290(12):1593-9.
37. Choudhry NK, Singh JM, Barolet A, Tomlinson GA, Detsky AS. How should patients with unstable angina and non-ST-segment elevation myocardial infarction be managed? A meta-analysis of randomized trials. *Am J Med* 2005;118(5):465-74.
 38. Hoenig MR, Doust JA, Aroney CN, Scott IA. Early invasive versus conservative strategies for unstable angina & non-ST-elevation myocardial infarction in the stent era. *Cochrane Database Syst Rev*. 2006; 3:CD004815.
 39. Hoenig MR, Aroney CN, Scott IA. Early invasive versus conservative strategies for unstable angina and non-ST-elevation myocardial infarction in the stent era. *Cochrane Database Syst Rev*. 2010;3: CD004815.
 40. Navarese EP, De Servi S, Gibson CM, Buffon A, Castriota F, Kubica J, et al. Early vs. delayed invasive strategy in patients with acute coronary syndromes without ST-segment elevation: a meta-analysis of randomized studies. *QJM*. 2011;104(3):193-200, <http://dx.doi.org/10.1093/qjmed/hcq258>.
 41. Damman P, van Geloven N, Wallentin L, Lagerqvist B, Fox KA, Clayton T, et al. Timing of angiography with a routine invasive strategy and long-term outcomes in non-ST-segment elevation acute coronary syndrome: a collaborative analysis of individual patient data from the FRISC II (Fragmin and Fast Revascularization During Instability in Coronary Artery Disease), ICTUS (Invasive Versus Conservative Treatment in Unstable Coronary Syndromes), and RITA-3 (Intervention Versus Conservative Treatment Strategy in Patients With Unstable Angina or Non-ST Elevation Myocardial Infarction) Trials. *JACC Cardiovasc Interv*. 2012;5(2):191-9, <http://dx.doi.org/10.1016/j.jcin.2011.10.016>.
 42. Ferry DR, O'Rourke RA, Blaustein AS, Crawford MH, Deedwania PC, Carson PE, et al. Design and baseline characteristics of the Veterans Affairs Non-Q-Wave Infarction Strategies In-Hospital (VANQWISH) trial. *J Am Coll Cardiol*. 1998;31(2):312-20, [http://dx.doi.org/10.1016/S0735-1097\(97\)00486-5](http://dx.doi.org/10.1016/S0735-1097(97)00486-5).