



Published in final edited form as:

J Invest Dermatol. 2012 April ; 132(4): 1133–1140. doi:10.1038/jid.2011.415.

Genome-wide meta-analysis of Psoriatic Arthritis Identifies Susceptibility Locus at *REL*

Eva Ellinghaus^{1,‡}, Philip E. Stuart², David Ellinghaus¹, Rajan P. Nair², Sophie Debrus³, John V. Raelson⁴, Majid Belouchi⁵, Trilokraj Tejasvi², Yanming Li⁶, Lam C. Tsoi⁶, Anna T. Onken⁷, Tonu Esko^{8,9,10}, Andres Metspalu^{8,9,10}, Proton Rahman¹¹, Dafna D. Gladman¹², Anne M. Bowcock¹³, Cynthia Helms¹³, Gerald G. Krueger¹⁴, Sulev Koks¹⁵, Külli Kingo¹⁶, Christian Gieger¹⁷, H. Erich Wichmann^{18,19,20}, Ulrich Mrowietz²¹, Stephan Weidinger²¹, Stefan Schreiber^{1,22}, Gonçalo R. Abecasis⁶, James T. Elder^{2,23}, Michael Weichenthal²¹, and Andre Franke¹

¹Institute of Clinical Molecular Biology, Christian-Albrechts-University, 24105 Kiel, Germany

²Department of Dermatology, University of Michigan, Ann Arbor, MI 48109, USA

³Gatineau, QC, Canada J9J 2X6

⁴PGX-Services, Montreal, QC, Canada H2T 1S1

⁵Genizon BioSciences, Inc., St. Laurent, QC, Canada H4T 2C7

⁶Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, MI 48109, USA

⁷Department of Dermatology and Allergy, Technische Universität München, 80802 Munich, Germany

⁸Estonian Genome Center and Center of Translational Genomics, University of Tartu, 50409 Tartu, Estonia

⁹Estonian Biocenter, 50409 Tartu, Estonia

¹⁰Institute of Molecular and Cell Biology, University of Tartu, 50409 Tartu, Estonia

¹¹Department of Medicine, Memorial University, St. John's, Newfoundland A1C 5B8, Canada

¹²Division of Rheumatology University Toronto, Psoriatic Arthritis Program, University Health Network, Toronto, Ontario M5T 2S8, Canada

¹³Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110, USA

¹⁴Department of Dermatology, University of Utah School of Medicine, Salt Lake City, UT 84132, USA

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

[‡]To whom correspondence should be addressed; Eva Ellinghaus (e.ellinghaus@ikmb.uni-kiel.de), Institute of Clinical Molecular Biology, Christian-Albrechts-University Kiel, Schittenhelmstr. 12, D-24105 Kiel, Germany, Tel.: +49-431-597-2746, Fax.: +49-431-597-2196.

Conflict of Interest

The authors declare no conflict of interest

¹⁵Department of Physiology, University of Tartu, 50409 Tartu, Estonia

¹⁶Department of Dermatology and Venerology, University of Tartu, 50409 Tartu, Estonia

¹⁷Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany

¹⁸Institute of Epidemiology I, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany

¹⁹Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-University, 81377 Munich, Germany

²⁰Klinikum Grosshadern, 81377 Munich, Germany

²¹Department of Dermatology, University Hospital Schleswig-Holstein, Christian-Albrechts-University, 24105 Kiel, Germany

²²Popgen Biobank, University Hospital Schleswig-Holstein, Christian-Albrechts-University, 24105 Kiel, Germany

²³Ann Arbor Veterans Affairs Hospital, Ann Arbor, MI 48105, USA

Abstract

Psoriatic arthritis (PsA) is a chronic inflammatory musculoskeletal disease affecting up to 30% of psoriasis vulgaris (PsV) cases and approximately 0.25% to 1% of the general population. To identify common susceptibility loci, we performed a meta-analysis of three imputed genome-wide association studies (GWAS) on psoriasis, stratified for PsA. A total of 1,160,703 SNPs were analyzed in the discovery set consisting of 535 PsA cases and 3,432 controls from Germany, the United States and Canada. We followed up two SNPs in 1,931 PsA cases and 6,785 controls comprising six independent replication panels from Germany, Estonia, the United States and Canada. In the combined analysis, a genome-wide significant association was detected at 2p16 near the *REL* locus encoding c-Rel (rs13017599, $P=1.18 \times 10^{-8}$, OR=1.27, 95% CI=1.18–1.35). The rs13017599 polymorphism is known to associate with rheumatoid arthritis (RA), and another SNP near *REL* (rs702873) was recently implicated in PsV susceptibility. However, conditional analysis indicated that rs13017599, rather than rs702873, accounts for the PsA association at *REL*. We hypothesize that c-Rel, as a member of the Rel/NF- κ B family, is associated with PsA in the context of disease pathways that involve other identified PsA and PsV susceptibility genes including *TNIP1*, *TNFAIP3* and *NF κ BIA*.

INTRODUCTION

Psoriatic arthritis (PsA) is a chronic inflammatory musculoskeletal disease that occurs in the context of psoriasis vulgaris (PsV) affecting up to 30% of psoriasis patients (Gladman *et al.*, 2005). Apart from the skin manifestation of PsV, the clinical phenotype of PsA overlaps with ankylosing spondylitis and rheumatoid arthritis (RA) (Castelino and Barton, 2010). While PsV has a prevalence of up to 3% in populations of European ancestry (Bowcock and Barker, 2003; Griffiths and Barker, 2007), the estimated prevalence of PsA varies from 0.25% to 1% (Gladman *et al.*, 2005). The strong genetic component of PsA is evidenced by

the high sibling recurrence risk (λ_S) of 27–47 (Chandran *et al.*, 2009; Gladman *et al.*, 2003; Myers *et al.*, 2005), which is much higher than the estimated $\lambda_{S=4-11}$ of PsV (Bhalerao and Bowcock, 1998; Elder *et al.*, 1994), suggesting that PsA patients harbor additional susceptibility loci beside the loci that contribute to risk for PsV. In addition to the significant heritable component, environmental risk factors play an important role for both PsV and PsA. The onset of PsA generally occurs in the fourth and fifth decades of life. In nearly 80% of patients, the skin disease (PsV) precedes the arthritis (PsA). The clarification of similarities and differences of the genetic background in PsV and PsA offers chances for well-targeted interventions, i.e. specific therapies for both conditions. It would also provide an opportunity to identify PsV patients with increased risk of PsA and would allow treatment at the earliest signs of joint involvement (Bowes and Barton, 2010). Several susceptibility loci for PsV have been identified with genome-wide levels of statistical significance in populations of European origin, including *HLA-Cw6*, *IL12B*, *IL23R*, *IL4-IL13*, *IL23A*, *TNIP1*, *TNFAIP3*, *LCE3B-LCE3C*, *RNF114*, *TRAF3IP2*, *NFκBIA*, *NOS2*, *FBXL19*, *TYK2*, *IFIH1*, *REL*, *IL28RA* and *ERAP1* (Duffin *et al.*, 2010; Ellinghaus *et al.*, 2010; Liu *et al.*, 2008; Strange *et al.*, 2010; Stuart *et al.*, 2010). Many of the susceptibility loci for PsV tested so far are also genome-wide significantly associated with PsA such as *HLA-Cw6*, *IL12B*, *TNIP1*, *FBXL19* and *TRAF3IP2* or at the level of $P < 0.05$ such as *IL23R*, *IL23A*, *TNFAIP3*, *NFκBIA*, *NOS2* and *IL13* (Bowes and Barton, 2010; Eder *et al.*, 2011; Ellinghaus *et al.*, 2010; Huffmeier *et al.*, 2010; Stuart *et al.*, 2010). However, because the causative allele(s) have not been identified within most of these regions, their actual contribution to disease risk remains unknown.

In order to further our understanding of PsA etiology, we conducted a meta-analysis of the PsA subsets of three independent genome-wide association studies of PsV from Kiel, Germany (Ellinghaus *et al.*, 2010), the Collaborative Association Study of Psoriasis (CASP) (Nair *et al.*, 2009) and Canada (Genizon, unpublished data), consisting of 535 PsA patients and 3,432 healthy controls. Genotype imputation with HapMap phase 3 reference samples considerably increased the genomic coverage of the three GWAS datasets and facilitated the combined analysis. For replication we selected ten SNPs based on their P -value ranking in the meta-analysis and genotyped them in two independent replication sets from Germany and Estonia. The two most strongly associated SNPs were further genotyped in four additional replication cohorts from Michigan, the National Psoriasis Foundation (NPF), the CASP Deep Follow-up (CASP-DFU) and Estonia. Together, this replication set comprised 1,931 PsA cases and 6,785 controls (Table S1). We identified genome-wide significant association for SNP rs13017599, which is located between the genes *REL* and *PUS10* and has been already reported to be associated with RA (Gregersen *et al.*, 2009). Our results indicate that *REL* is an important candidate susceptibility locus not only for PsV and RA but also for PsA.

RESULTS

Before genotype imputation and meta-analysis we applied extensive quality control filters to the three GWAS datasets separately. We excluded all samples with low genotyping rate (<95%) as well as non-European outliers or cryptically related individuals. Further, we excluded SNPs with a minor allele frequency <1%, a genotyping success rate <95% or

deviation of the genotype distribution from Hardy-Weinberg equilibrium in the controls ($P < 10^{-4}$). After applying these stringent quality control filters genotype imputation was performed with Beagle v.3.2.1 (Browning and Browning, 2009) using HapMap phase 3 reference haplotypes (Altshuler *et al.*, 2010). Only SNPs imputed with high confidence (estimated r^2 between imputed and true genotypes > 0.3) in all three GWAS were considered for subsequent statistical analysis, leaving a total of 1,160,703 SNPs. A logistic regression procedure was applied to test both genotyped and imputed SNPs for association. To account for uncertainty in the imputation procedure we used allele dosages from the imputation. After combining the association results of all three scans, a low genomic control value (meta GWAS: $\lambda_{GC}=1.02$) indicated a minimal overall inflation of the test statistics due to population stratification (Figure S1). Our combined discovery panel had 80% power to detect variants conferring odds ratios (OR) of 1.40 or higher at the 5% significance level, assuming a frequency of the disease-associated risk allele of at least 15% in controls (Figure S2). In line with previous studies, we detected the strongest associations with PsA at SNPs in the HLA complex at chromosome 6p21 approximately 35 kb upstream of *HLA-C* (rs12212594, $P=4.98 \times 10^{-38}$, OR=3.60; rs12191877, $P=3.95 \times 10^{-35}$, OR=2.73 and rs10484554, $P=3.03 \times 10^{-34}$, OR=2.72) (Manhattan plot see Figure 1). SNP rs12191877 was previously identified as most strongly associated with PsV in a GWAS of individuals with genuine European ancestry (Nair *et al.*, 2009), while rs10484554 has been described to be most significantly associated in a combined study of PsA and PsV (Liu *et al.*, 2008). In our discovery panel, rs12212594 was in moderate LD with rs12191877 ($r^2=0.57$), indicating that these signals are not independent. Additionally, we found evidence of association at the previously reported susceptibility loci *TNIP1* (Nair *et al.*, 2009) (rs17728338, $P=6.75 \times 10^{-10}$, OR=2.07) and *IL12B* (Liu *et al.*, 2008) (rs3212227, $P=3.82 \times 10^{-7}$, OR=1.64). For selection of follow-up SNPs we excluded all SNPs within the extended MHC region (7,773 SNPs on chromosome 6 at 25–34Mb) as well as SNPs within the genes *TNIP1* and *IL12B* since these loci have already been shown to be associated with PsA at a genome-wide significant level. Furthermore we employed only those SNPs that were available in all three scans, and that showed no heterogeneity between the studies (I^2 statistic=0). We selected ten SNPs for replication (see Manhattan plot, Figure 1) in two independent samples from Germany and Estonia (replication panel 1 and 2) comprising 252 PsA cases and 1,740 controls. All SNPs passed quality control measures which means that they had a high call rate ($>95\%$ in cases or controls), were not monomorphic (minor allele frequency $>1\%$ in cases or controls) and did not deviate from Hardy-Weinberg equilibrium in the control population (HWE $P > 10^{-4}$). Taking into account the minor differing genetic backgrounds of the two replication study groups, we used the Cochran-Mantel-Haenszel test for the combined analysis (Table 1). Of the ten follow-up SNPs, two SNPs were nominally associated with PsA in the replication dataset (rs13017599 and rs2829866 with $P < 0.05$) and were therefore genotyped in four additional replication panels from Michigan, the NPF, the CASP-DFU and Estonia (replication panels 3 through 6), yielding a replication sample consisting of 1,931 PsA cases and 6,785 controls. In the combined replication analysis rs13017599 - located 15 kb downstream of *REL* and 5 kb downstream of *PUS10* - remained significantly associated with PsA, yielding $P=4.24 \times 10^{-5}$ (OR=1.23, 95% CI=1.14–1.33, no heterogeneity: $I^2=0$). In contrast, rs2829866 showed no evidence for association in the extended replication sample ($\alpha=0.005$, calculated as $0.05/10$). The combined analysis of the

discovery and the six replication panels (2,466 PsA cases and 10,217 controls) yielded a genome-wide significant P -value of 1.18×10^{-8} (OR=1.27, 95% CI=1.18–1.35, $r^2=0$) for the lead SNP rs13017599 (Table 2). To fine map a region of 500 kb around the lead SNP rs13017599, we performed imputation based on CEU haplotypes generated by the 1000 Genomes Project for the three GWAS panels used for discovery phase (The 1000 Genomes Project Consortium, 2010). The regional plot of the combined P -values confirmed the results of the HapMap phase 3 imputed meta-analysis, with rs13017599 displaying the strongest association ($P=1.88 \times 10^{-5}$) at this locus (Figure 2). Additionally, we genotyped the SNP rs702873, which was reported to be associated with PsV by Strange *et al.*, in the replication panels 1 through 6 (1,931 PsA cases and 6,785 controls). Meta-analysis of the three GWAS and the six replication study groups (2,466 PsA cases and 10,217 controls) yielded $P=1.93 \times 10^{-7}$ (OR=1.20, 95% CI=1.13–1.28), confirming the newly identified association of this locus with PsA (Table 1 and 2). The lead SNP rs13017599 and the PsV-associated variant rs702873 are about 83 kb apart and are in moderate linkage disequilibrium (LD in 566 CEU haplotypes generated by the 1000 Genomes Project (The 1000 Genomes Project Consortium, 2010); $r^2=0.625$). To test for the independence of these two markers, we performed a logistic regression analysis. When conditioning for the lead SNP rs13017599, rs702873 was not significantly associated with PsA ($P=0.20$), whereas rs13017599 was still associated ($P=2.90 \times 10^{-2}$) in the reverse conditioning, indicating that the two SNPs are not completely independent, but that rs13017599 accounts for the association of PsA at the *REL* gene locus rather than rs702873. In order to investigate whether the *REL* locus was more strongly associated with PsA than with PsV, we additionally analyzed the two *REL* SNPs rs13017599 and rs702873 in PsV cases without PsA from the same three GWAS and six replication panels (5,912 PsV cases without PsA and 10,217 controls). This analysis yielded much weaker association signals compared to the PsA-cohort (rs13017599, $P_{PsV}=3.48 \times 10^{-5}$, OR=1.13, 95% CI=1.08–1.19; rs702873, $P_{PsV}=4.70 \times 10^{-6}$, OR=1.14, 95% CI=1.09–1.19), highlighting that the *REL*-association is stronger in PsA but nevertheless confirming the previously reported association of rs702873 with PsV. For each of the two associations at rs13017599 and rs702873 there is an overlap in confidence intervals for PsA and PsV which further proves that the *REL* gene represents a shared susceptibility locus for PsA and PsV.

DISCUSSION

In addition to the recently reported association of SNP rs13017599 with RA (Gregersen *et al.*, 2009) and SNP rs702873 - located 27kb upstream of the *REL* gene - with PsV (Strange *et al.*, 2010), several other SNPs downstream of *REL* have been identified to be associated with ulcerative colitis (McGovern *et al.*, 2010), Crohn's Disease (Franke *et al.*, 2010), celiac disease (Dubois *et al.*, 2010) and primary sclerosing cholangitis (Janse *et al.*, 2011), demonstrating that *REL* is a shared disease locus of several complex diseases and thereby highlighting its role in inflammatory barrier disease susceptibility. *REL* encodes one of four subunits found in NF- κ B dimers and belongs to the NF- κ B (REL) family of transcription factors which is highly important in the innate immune system. The above mentioned association findings are therefore consistent with the importance of the NF- κ B pathway in these inflammatory diseases. c-Rel contains trans-activation domains that are a prerequisite

for gene transcription, and T cell-mediated immune responses were markedly impaired in c-Rel^{-/-} animals. The confirmed PsA/PsV susceptibility locus *TNFAIP3* and a variety of other genes are regulated by c-Rel in T cells (Bunting *et al.*, 2007). Notably, c-REL mediates differentiation of CD4⁺ FoxP3⁺ regulatory T cells which have the capacity to develop into IL-17 producing cells in psoriasis (Bovenschen *et al.*, 2011). It also promotes IL-12B and IL-23 expression, by which TH-1 and TH-17 types of immune response might be evoked (Mise-Omata *et al.*, 2007; Reinhard *et al.*, 2011). Altogether these findings further support the hypothesis that *REL* is an important integrator of inflammatory signaling pathways observed in the complex emerging landscape of genetic susceptibility for PsA. In conclusion, we were able to provide - to our knowledge previously unreported - evidence at genome-wide significance for involvement of the *REL* locus in PsA susceptibility, and we could confirm the previously reported association with PsV. The current results demand not only a thorough analysis of all genes involved in this pathway to search for additional risk alleles but also overlap analyses between complex disorders sharing a certain genetic background like PsA, PsV and RA. It should be noted however, that because most of the patients develop PsA following the onset of PsV, PsA might be considered as disease within a disease with psoriasis as the parent disease (Eder *et al.*, 2011), so that any association study on PsA is bound to identify PsV susceptibility genes such as the established PsV/PsA risk loci *HLA-Cw6*, *IL12B*, *IL23R*, *TNIP1* and *FBXL19* which do not allow for genetic discrimination between the two conditions. On the other hand, the genetic component of PsA is much higher than that of PsV and only about one third of PsV patients additionally develop PsA, implying that differences in the genetic backgrounds do exist. Genes involved in innate (such as *REL*) and adaptive immunity are expectedly more often shared between different immune-mediated diseases, while genes of skin barrier function are rather restricted to PsV. So far, no PsA-specific genes have been identified with genome-wide significance, but then again, only three GWAS on PsA have been conducted until now, our study included (Huffmeier *et al.*, 2010; Liu *et al.*, 2008). Lack of power to find genetic variants with weaker effects, and rare variants that contribute to disease risk might explain the sparse results, thereby emphasizing the need for larger studies and targeted next-generation sequencing of candidate regions.

MATERIALS & METHODS

Subjects

All German PsA (219) and PsV (733) cases in the GWAS panel and the replication panel 1 were recruited either at the Department of Dermatology of the Christian-Albrechts-University Kiel or the Department of Dermatology and Allergy of the Technical University Munich through local outpatient services. PsA was diagnosed by a clinical finding of joint complaints and radiological and rheumatologic confirmation of criteria according to Moll and Wright (Moll and Wright, 1973), or more recently, to the Classification Criteria for Psoriatic Arthritis (CASPAR) (Taylor *et al.*, 2006). 63 PsA cases from Munich were diagnosed by a physician, with negative diagnosis of RA. Individuals were considered to be affected by PsV if chronic plaque or guttate psoriasis lesions covered more than 1% of the total body surface area or if at least two skin, scalp, nail or joint lesions were clinically diagnosed of psoriasis by a dermatologist.

2,127 German healthy control individuals in the GWAS panel and the replication panel 1 were obtained from the PopGen biobank (Krawczak *et al.*, 2006). 465 German healthy controls were selected from the KORA S4 survey, an independent population-based sample from the general population living in the region of Augsburg, southern Germany (Wichmann *et al.*, 2005).

The CASP GWAS (for details see (Nair *et al.*, 2009; Stuart *et al.*, 2010)) consisted of 335 PsA cases, 915 PsV cases and 1,322 controls after quality control measures. The datasets used for the analyses described in this manuscript were obtained from the database of Genotype and Phenotype (dbGaP, <http://www.ncbi.nlm.nih.gov/gap>).

The Canadian GWAS sample (from Genizon BioSciences) consisted of 139 PsA cases, 614 PsV cases and 987 controls sampled from the Québec founder population (QFP). Membership in the QFP was defined as having four grandparents with French-Canadian family names who were born in the Province of Québec, Canada or in adjacent areas in the provinces of New Brunswick and Ontario or in New England or New York State. This criterion assured that all subjects were descendants of French-Canadians living before the 1960s, after which time admixture with non-French-Canadians became more common. Inclusion criteria were the presence of plaque forming psoriasis determined by a dermatologist, and disease onset between 18 and 40 years of age. Exclusion criteria were other forms of psoriasis, such as pustular, guttate, inverse, erythrodermic and isolated site psoriasis and late onset. The presence of psoriatic arthritis was noted but not an ascertainment criterion.

The samples of the Estonian replication panel 2 were unrelated Estonian Caucasian patients with a clear clinical diagnosis of psoriasis vulgaris, collected at the Department of Dermatology and Venerology and at the Department of Physiology and Centre of Translational Medicine at the University of Tartu. Patients were classified to have psoriatic arthritis if this diagnosis had been established by an experienced rheumatologist. The Estonian replication panel 6 consisted of samples provided by the population-based biobank of the Estonian Genome Center, University of Tartu (EGCUT). Subjects were recruited by general practitioners (GP) and physicians in the hospitals. Physicians in the hospitals were randomly selected from individuals visiting GP offices or hospitals. Diagnosis of PsV and/or PsA based on clinical symptoms was posed by a general practitioner and confirmed by a dermatologist. At the moment of recruitment, the controls did not report diagnosis of osteoarthritis, psoriasis or autoimmune diseases.

Replication panels 3 through 5 (Michigan, NPF and CASP-DFU) consisted of 1,649 PsA cases, 2,349 PsV cases and 4,061 controls of white European ancestry from the United States and Canada. Diagnosis of psoriasis was always confirmed by a dermatologist. All PsA patients were diagnosed by a rheumatologist and/or fulfilled CASPAR criteria for the classification of PsA. All Canadian PsV patients underwent a rheumatological examination to rule out the presence of inflammatory arthritis. Patients with other forms of inflammatory arthritis were excluded. Written, informed consent was obtained from all study participants, and all protocols were approved by the respective institutional ethical review committees of the participating centers. The investigations were conducted according to the Declaration of

Helsinki Principles. The samples were organized in panels which corresponded to the successive steps of the present study and all panels (GWAS panels and replication panels 1 through 6) were independent from each other.

Genotyping

The genotyping for the German GWAS, which was part of the German GWAS initiative funded by the National Genome Research Network (NGFN), was performed by Illumina's service facility using the Illumina HumanHap 550K v1 with 561,466 SNP markers. All experimental steps were carried out according to standard protocols. For genome-wide genotyping of the CASP samples the Perlegen 600k array was used. The genotyping was provided through the Genetic Association Information Network (GAIN). The Illumina Human 1M BeadChip was used for generating the Canadian genome-wide data (Genizon GWAS) as well as for genotypes of SNPs rs13017599 and rs2829866 in replication panel 5 (CASP-DFU). Functionally tested TaqMan SNP Genotyping Assays (Applied Biosystems) were used to genotype variants in replication panels 1 through 6.

Statistical analyses

We used GWAS datasets that passed stringent quality control filters for genotype imputation with Beagle (Browning and Browning, 2009) using HapMap phase 3 reference samples from the CEU, TSI, MEX and GJT collections, as well as for the subsequent statistical analyses. Low genomic control values for all three scans (German GWAS: $\lambda_{GC}=1.01$, CASP GWAS: $\lambda_{GC}=1.02$ and Canadian GWAS: $\lambda_{GC}=1.03$) indicated a minimal overall inflation of the test statistics due to population stratification.

Power calculations were carried out using PS Power and Sample Size v3.0.12 (Dupont and Plummer, 1997) (Figure S2). GWAS data were analyzed using R statistical environment version 2.10.0 and gPLINK v2.049 in combination with PLINK v1.05 (Purcell *et al.*, 2007).

We selected ten SNPs based on their *P*-value ranking for replication in two independent samples from Germany and Estonia comprising 252 PsA cases and 1,740 controls. All SNPs passed quality control measures which means that they had a high call rate (>95% in cases or controls), were not monomorphic (minor allele frequency >1% in cases or controls) and did not deviate from Hardy-Weinberg equilibrium in the control population (exact HWE $P > 10^{-4}$). Two of the ten follow-up SNPs were significantly associated with PsA and were therefore genotyped in four additional replication panels from the United States, Canada and Estonia.

The Cochran-Mantel-Haenszel test was used for combined replication analysis of replication panel 1 and 2 from Germany and Estonia, in order to take the possibility of minor differing genetic backgrounds into account. The initial GWAS meta-analysis was performed using PLINK's meta-analysis function with its standard error of odds ratio weighting option. Association results of replication panel 5 (CASP-DFU) were obtained with the statistical association test "Efficient Mixed-Model Association eXpedited" (EMMAX) to account for sample structure (Kang *et al.*, 2010) and the combined replication analysis of all six replication panels as well as the combined GWAS-replication analysis was performed with

METAL (Willer *et al.*, 2010). Logistic regression analysis to test for the independence of the two *REL* SNPs rs13017599 and rs702873 was performed with PLINK using the conditional analysis commands --logistic and --condition to test one SNP but adding the allelic dosage for the other SNP as a covariate. The conditional analysis was carried out for each cohort separately and results were subsequently combined by means of meta-analysis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank all individuals with psoriasis and/or psoriatic arthritis, their families and physicians for their cooperation. We acknowledge the cooperation of Genizon Biosciences. We wish to thank Tanja Wesse, Tanja Henke, Catharina Fürstenau and Susan Ehlers for expert technical help. This study was supported by the German Ministry of Education and Research (BMBF) through the National Genome Research Network (NGFN) and the Popgen biobank. The project received infrastructure support through the DFG cluster of excellence "Inflammation at Interfaces". This research was also supported by grants R01-AR42742, R01-AR050511, AR050266 and R01-AR054966 from the National Institutes of Health, the Psoriasis and Psoriatic arthritis New Emerging Team grant from the Canadian Institutes of Health Research and the Krembil Foundation. S.W. is supported by a Heisenberg Professorship of the German Research council DFG (WE 2678/4-1). The KORA (Kooperative Gesundheitsforschung in der Region Augsburg) study was initiated and financed by the Helmholtz Zentrum München - National Research Center for Environmental Health, which is funded by the German Federal Ministry of Education, Science, Research and Technology and by the State of Bavaria. Part of this work was financed by the German National Genome Research Network (NGFNPlus: 01GS0823) and was supported within the Munich Center of Health Sciences (MC Health) as part of LMUinnovativ. EGCUT received financing by FP7 grants (201413, 245536), also received targeted financing from Estonian Government SF0180142s08, from University of Tartu in the frame of Center of Translational Genomics and from the European Union through the European Regional Development Fund, in the frame of. We acknowledge EGCUT and Estonian Biocentre personnel, especially Ms. M. Hass and Mr. V. Soo.

Abbreviations used

PsA	psoriatic arthritis
PsV	psoriasis vulgaris
GWAS	genome-wide association study
SNP	single nucleotide polymorphism
CI	confidence interval

References

- Altshuler DM, Gibbs RA, Peltonen L, et al. Integrating common and rare genetic variation in diverse human populations. *Nature*. 2010; 467:52–8. [PubMed: 20811451]
- Bhalerao J, Bowcock AM. The genetics of psoriasis: a complex disorder of the skin and immune system. *Hum Mol Genet*. 1998; 7:1537–45. [PubMed: 9735374]
- Bovenschen HJ, van de Kerkhof PC, van Erp PE, et al. Foxp3+ Regulatory T Cells of Psoriasis Patients Easily Differentiate into IL-17A-Producing Cells and Are Found in Lesional Skin. *J Invest Dermatol*. 2011
- Bowcock AM, Barker JN. Genetics of psoriasis: the potential impact on new therapies. *J Am Acad Dermatol*. 2003; 49:S51–6. [PubMed: 12894126]
- Bowes J, Barton A. The genetics of psoriatic arthritis: lessons from genome-wide association studies. *Discov Med*. 2010; 10:177–83. [PubMed: 20875338]

- Browning BL, Browning SR. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet.* 2009; 84:210–23. [PubMed: 19200528]
- Bunting K, Rao S, Hardy K, et al. Genome-wide analysis of gene expression in T cells to identify targets of the NF-kappa B transcription factor c-Rel. *J Immunol.* 2007; 178:7097–109. [PubMed: 17513759]
- Castelino M, Barton A. Genetic susceptibility factors for psoriatic arthritis. *Curr Opin Rheumatol.* 2010; 22:152–6. [PubMed: 20084005]
- Chandran V, Schentag CT, Brockbank JE, et al. Familial aggregation of psoriatic arthritis. *Ann Rheum Dis.* 2009; 68:664–7. [PubMed: 18524791]
- Dubois PC, Trynka G, Franke L, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet.* 2010; 42:295–302. [PubMed: 20190752]
- Duffin KC, Woodcock J, Krueger GG. Genetic variations associated with psoriasis and psoriatic arthritis found by genome-wide association. *Dermatol Ther.* 2010; 23:101–13. [PubMed: 20415816]
- Dupont WD, Plummer WD. PS power and sample size program available for free on the Internet. *Controlled Clin Trials.* 1997:18.
- Eder L, Chandran V, Pellett F, et al. IL13 gene polymorphism is a marker for psoriatic arthritis among psoriasis patients. *Ann Rheum Dis.* 2011; 70:1594–8. [PubMed: 21613309]
- Elder JT, Nair RP, Guo SW, et al. The genetics of psoriasis. *Arch Dermatol.* 1994; 130:216–24. [PubMed: 8304761]
- Ellinghaus E, Ellinghaus D, Stuart PE, et al. Genome-wide association study identifies a psoriasis susceptibility locus at TRAF3IP2. *Nat Genet.* 2010; 42:991–5. [PubMed: 20953188]
- Franke A, McGovern DP, Barrett JC, et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet.* 2010; 42:1118–25. [PubMed: 21102463]
- Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis.* 2005; 64(Suppl 2):ii14–7. [PubMed: 15708927]
- Gladman DD, Farewell VT, Pellett F, et al. HLA is a candidate region for psoriatic arthritis. evidence for excessive HLA sharing in sibling pairs. *Hum Immunol.* 2003; 64:887–9. [PubMed: 12941544]
- Gregersen PK, Amos CI, Lee AT, et al. REL, encoding a member of the NF-kappaB family of transcription factors, is a newly defined risk locus for rheumatoid arthritis. *Nat Genet.* 2009; 41:820–3. [PubMed: 19503088]
- Griffiths CE, Barker JN. Pathogenesis and clinical features of psoriasis. *Lancet.* 2007; 370:263–71. [PubMed: 17658397]
- Huffmeier U, Uebe S, Ekici AB, et al. Common variants at TRAF3IP2 are associated with susceptibility to psoriatic arthritis and psoriasis. *Nat Genet.* 2010; 42:996–9. [PubMed: 20953186]
- Janse M, Lamberts LE, Franke L, et al. Three ulcerative colitis susceptibility loci are associated with primary sclerosing cholangitis and indicate a role for IL2, REL, and CARD9. *Hepatology.* 2011; 53:1977–85. [PubMed: 21425313]
- Kang HM, Sul JH, Service SK, et al. Variance component model to account for sample structure in genome-wide association studies. *Nat Genet.* 2010; 42:348–54. [PubMed: 20208533]
- Krawczak M, Nikolaus S, von Eberstein H, et al. PopGen: population-based recruitment of patients and controls for the analysis of complex genotype-phenotype relationships. *Community Genet.* 2006; 9:55–61. [PubMed: 16490960]
- Liu Y, Helms C, Liao W, et al. A genome-wide association study of psoriasis and psoriatic arthritis identifies new disease loci. *PLoS Genet.* 2008; 4:e1000041. [PubMed: 18369459]
- McGovern DP, Gardet A, Torkvist L, et al. Genome-wide association identifies multiple ulcerative colitis susceptibility loci. *Nat Genet.* 2010; 42:332–7. [PubMed: 20228799]
- Mise-Omata S, Kuroda E, Niikura J, et al. A proximal kappaB site in the IL-23 p19 promoter is responsible for RelA- and c-Rel-dependent transcription. *J Immunol.* 2007; 179:6596–603. [PubMed: 17982049]
- Moll JM, Wright V. Psoriatic arthritis. *Semin Arthritis Rheum.* 1973; 3:55–78. [PubMed: 4581554]

- Myers A, Kay LJ, Lynch SA, et al. Recurrence risk for psoriasis and psoriatic arthritis within sibships. *Rheumatology (Oxford)*. 2005; 44:773–6. [PubMed: 15757963]
- Nair RP, Duffin KC, Helms C, et al. Genome-wide scan reveals association of psoriasis with IL-23 and NF-kappaB pathways. *Nat Genet*. 2009; 41:199–204. [PubMed: 19169254]
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007; 81:559–75. [PubMed: 17701901]
- Reinhard K, Huber M, Wostl C, et al. c-Rel promotes type 1 and type 17 immune responses during *Leishmania major* infection. *Eur J Immunol*. 2011; 41:1388–98. [PubMed: 21469108]
- Strange A, Capon F, Spencer CC, et al. A genome-wide association study identifies new psoriasis susceptibility loci and an interaction between HLA-C and ERAP1. *Nat Genet*. 2010; 42:985–90. [PubMed: 20953190]
- Stuart PE, Nair RP, Ellinghaus E, et al. Genome-wide association analysis identifies three psoriasis susceptibility loci. *Nat Genet*. 2010; 42:1000–4. [PubMed: 20953189]
- Taylor W, Gladman D, Helliwell P, et al. Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum*. 2006; 54:2665–73. [PubMed: 16871531]
- Team RDC. R: A language and environment for statistical computing. R Foundation for Statistical Computing; Vienna, Austria: 2007.
- The 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature*. 2010; 467:1061–73. [PubMed: 20981092]
- Wichmann HE, Gieger C, Illig T. KORA-gen--resource for population genetics, controls and a broad spectrum of disease phenotypes. *Gesundheitswesen*. 2005; 67(Suppl 1):S26–30. [PubMed: 16032514]
- Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26:2190–1. [PubMed: 20616382]




Figure 1. Results of genome-wide meta-analysis

The negative decadic logarithm of the corresponding P -value from the meta-analysis is shown for each SNP, according to chromosome. All markers that passed quality control criteria, were available in all three GWAS and showed no heterogeneity between the studies were used for plotting. The plot was created with the software environment R version 2.11.1 (Team, 2007). The established PsA loci *HLA-C* and *TNIP1* are highlighted by grey arrows, while the follow-up loci are tagged by black arrows. The novel risk locus at *REL* is highlighted by a red arrow. The *HLA* region stands out clearly from all other loci. For better scaling the Y-axis was limited to a maximum value of $-\log_{10}(P)=10$, thereby truncating the *HLA* signal at this value.




Figure 2. Regional plot of the REL locus
 Regional plot of the negative decadic logarithm of the combined P -values from the imputed meta-analysis of three GWAS panels. A window of 500 kb around the lead SNP rs13017599 (blue filled circle) is shown. The three GWAS panels were imputed with CEU haplotypes generated by the 1000 Genomes Project (August 2010 release) as a reference. The magnitude of linkage disequilibrium (LD) with the central SNP rs13017599, measured by r^2 , is reflected by the color of each SNP symbol (color coding: see upper right corner of the plot).

Table 1
Association results of ten follow-up SNPs from the meta-analysis and of the PsV associated SNP rs702873.

Locus	SNP	Chr. Position (bp)	Alleles	Genome-wide analysis (535 cases, 3,432 controls)		Replication analysis (252 PsA cases, 1,740 controls)				combined analysis (GWAS, repl. 1+2) (787 PsA cases, 5,172 controls)		
				meta P	OR (95% CI)	repl. 1 Germany	repl. 2 Estonia	P _{CMH}	OR (95% CI)	P _{BD}	meta P	OR
<i>STXBP4/HLF</i>	rs171511	17 53,273,910	T/C	9.30×10 ⁻⁶	1.37 (1.19–1.58)	0.39/0.36	0.42/0.44	0.59	1.06 (0.87–1.29)	0.33	8.62×10 ⁻⁵	1.26
<i>RAD50</i>	rs6596086	5 131,952,222	T/C	9.77×10 ⁻⁶	1.51 (1.26–1.81)	0.77/0.75	0.71/0.67	0.26	1.14 (0.91–1.42)	0.77	3.52×10 ⁻⁵	1.35
<i>RELPUF10</i>	rs13017599	2 61,164,331	G/A	1.48×10 ⁻⁵	1.37 (1.19–1.57)	0.69/0.62	0.67/0.63	8.92×10⁻³	1.31 (1.07–1.61)	0.63	4.56×10 ⁻⁷	1.35
<i>PTDSS1</i>	rs7824505	8 97,330,691	A/C	1.49×10 ⁻⁵	1.37 (1.19–1.57)	0.49/0.47	0.39/0.48	0.48	0.93 (0.77–1.13)	0.04	2.00×10 ⁻³	1.20
<i>CCDC65</i>	rs10783293	12 49,308,475	A/G	1.83×10 ⁻⁵	1.34 (1.17–1.53)	0.43/0.44	0.43/0.43	0.68	0.96 (0.79–1.17)	0.80	9.95×10 ⁻⁴	1.20
<i>A2BP1</i>	rs12445208	16 7,325,638	A/C	3.72×10 ⁻⁵	1.48 (1.23–1.78)	0.80/0.81	0.83/0.81	0.99	1.00 (0.78–1.28)	0.51	1.10×10 ⁻³	1.28
<i>IL5RA</i>	rs3792424	3 3,117,700	A/G	3.97×10 ⁻⁵	1.94 (1.42–2.67)	0.91/0.93	0.93/0.91	0.99	1.00 (0.69–1.43)	0.18	2.98×10 ⁻³	1.43
<i>MARCH1</i>	rs7686154	4 164,542,625	T/G	6.56×10 ⁻⁵	2.83 (1.70–4.71)	0.96/0.96	0.92/0.93	0.95	0.99 (0.63–1.54)	0.53	1.08×10 ⁻²	1.55
<i>CCR6/FGFR1OP</i>	rs162295	6 167,454,097	A/C	8.62×10 ⁻⁵	1.36 (1.17–1.59)	0.72/0.73	0.70/0.67	0.88	1.02 (0.82–1.26)	0.37	1.14×10 ⁻³	1.23
<i>JAM2</i>	rs2829866	21 27,059,071	A/T	1.00×10 ⁻⁴	1.32 (1.15–1.52)	0.37/0.30	0.27/0.25	3.75×10⁻²	1.24 (1.01–1.53)	0.39	1.14×10 ⁻⁵	1.30
<i>PAPOLG/REL</i>	rs702873	2 61,081,542	C/T	1.56×10 ⁻⁴	1.29 (1.13–1.48)	0.61/0.56	0.63/0.57	3.38×10⁻²	1.24 (1.02–1.50)	0.77	1.57×10 ⁻⁵	1.28

The top ten SNPs that were selected for follow-up from the meta-analysis are listed. These SNPs were genotyped in the replication case-control panel 1 and 2 from Germany and Estonia, respectively. Additionally, we genotyped the SNP rs702873 that was previously reported to be associated with PsV (Strange *et al.*, 2010). SNPs are ranked according to their *P*-values obtained in the initial meta-analysis. Positions (**Position (bp)**) are in NCBI's build 37. Chromosome: **Chr.** **P_{BD}** lists the asymptotic *P*-value of the Breslow-Day test for heterogeneity. A significant *P*-value indicates a significant heterogeneity between the two replication panels in terms of the disease association. Combined *P*-values (**P_{CMH}**) and combined ORs of the Cochran-Mantel-Haenszel (CMH) test statistic (one degree of freedom) are shown for replication panels 1 and 2. Significant *P*-values (*P*_{CMH}<0.05 [only if *P*_{BD}>0.05]) of the replication panels are highlighted in **bold**.

Table 2

Association results of the two follow-up SNPs and the PsV associated REL SNP in replication panels 1 through 6.

Locus	SNP	Chr. Position (bp)	Alleles	Genome-wide analysis (535 cases, 3,432 controls)		Replication analysis (1,931 cases, 6,785 controls)										meta analysis (2,466 cases, 10,217 controls)			
				meta <i>P</i>	OR (95% CI)	Allele frequencies (cases/controls)						repl. 1+2				repl. 1-6		meta <i>P</i>	OR (95% CI)
						repl. 1 Germany	repl. 2 Estonia 1	repl. 3 Michigan	repl. 4 NPF	repl. 5 CASP-DFU	repl. 6 Estonia 2	<i>P</i> _{CMH^a}	OR (95% CI)	meta <i>P</i>	OR (95% CI)				
REL	rs13017599	2 61,164,331	G/A	1.48×10 ⁻⁵	1.37 (1.19–1.57)	0.69/0.62	0.67/0.63	0.69/0.65	0.71/0.71	0.69/0.64	0.70/0.64	8.92×10 ⁻³	1.31 (1.07–1.61)	4.24×10 ⁻⁵	1.23 (1.14–1.33)	1.18×10 ⁻⁸	1.27 (1.18–1.35)		
REL	rs702873	2 61,081,542	C/T	1.56×10 ⁻⁴	1.29 (1.13–1.48)	0.61/0.56	0.63/0.57	0.62/0.57	0.61/0.62	0.61/0.56	0.64/0.58	3.38×10 ⁻²	1.24 (1.02–1.50)	1.25×10 ⁻⁴	1.17 (1.09–1.26)	1.93×10 ⁻⁷	1.20 (1.13–1.28)		
JAM2	rs2829866	21 27,059,071	A/T	5.73×10 ⁻⁵	1.34 (1.16–1.54)	0.37/0.30	0.27/0.25	0.32/0.31	0.35/0.30	0.32/0.32	0.21/0.26	3.75×10 ⁻²	1.24 (1.01–1.53)	0.11	1.07 (0.98–1.16)	6.33×10 ⁻⁴	1.13 (1.05–1.21)		

We analyzed the top ten SNPs of the GWAS meta analysis (including typed and imputed genotypes) in two independent PsA case-control panels from Germany and Estonia, respectively (repl. 1 and 2). The two SNPs with nominal significant replication results ($P < 0.05$) in these two replication panels as well as the PsV associated SNP rs702873 were genotyped in four additional independent PsA case-control panels (repl. 3 through 6). Results of all ten SNPs are shown in Table 1. Allele frequencies are given for each of the six replication panels separately. Nucleotide positions refer to NCBI build 37. CMH, Cochran-Mantel-Haenszel test; OR, odds ratio; 95% CI, 95% confidence interval; Chr., chromosome.

^a Cochran-Mantel-Haenszel test; Breslow-Day test, $P = 0.63$ for rs13017599, $P = 0.77$ for rs702873 and $P = 0.39$ for rs2829866.