

Dietary nutrient intake and antioxidant status in preeclamptic women

Mahdiye Sheykhi¹, Zamzam Paknahad^{1,2}, Akbar Hasanzadeh¹

¹Food Security Research Center, ²Department of Clinical Nutrition, School of Nutrition and Food Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: Preeclampsia (PE) is the most common cause of maternal death in the world. Some studies showed that inadequate intake of foods rich in antioxidant leads to increase oxidative stress and adverse obstetrical outcomes. The aim of the present study was to investigate the relationship between antioxidant status and dietary nutrient intake in pregnant women with PE.

Materials and Methods: This cross-sectional study was conducted among 55 pregnant women with PE admitted in the Obstetrics and Gynecology department of Shahid Beheshti Hospital in Isfahan, Iran. The subjects were interviewed about demographic data and dietary intakes by using a 168-items semi-quantitative food frequency questionnaire (FFQ). The total antioxidant capacity (TAC) of this serum was measured by using a double-antibody sandwich enzyme-linked immune-sorbent assay (ELISA). Nonparametric correlation statistics were used to meet assumptions of normality and equal variances.

Results: Total antioxidant status was significantly higher in comparison with healthy pregnant women (which measured as pilot). Intake of vitamin E was below the dietary reference intakes, and was positively associated with serum TAC ($r = 0.367$, $P = 0.003$), but this correlation was significantly negative about dietary selenium. There wasn't any significant correlation between intake of vitamin C, β -carotene, riboflavin, copper and serum TAC.

Conclusion: Our findings showed that intake of vitamin E was positively associated with serum TAC. Little support was found on a relationship between dietary intakes of other micronutrients and serum TAC. Further research is required to explore the relationships between maternal nutrient intake and antioxidant status in women with PE.

Key Words: Antioxidant capacity, antioxidant status, diet, nutrition, preeclampsia

Address for correspondence:

Dr. Zamzam Paknahad, Department of Clinical Nutrition, Faculty of Nutrition and Food Sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

E-mail: paknahad@hlth.mui.ac.ir

Received: 05.08.2014, Accepted: 13.09.2014

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.164003

INTRODUCTION

Hypertensive disorders of pregnancy, including gestational hypertension and preeclampsia (PE), may involve multiple organ systems and considered as leading causes of worldwide maternal and fetal mortality.^[1,2]

By 2015, reducing maternal mortality and morbidity is assumed as an objective of the Millennium

Copyright: © 2015 Sheykhi. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Sheykhi M, Paknahad Z, Hasanzadeh A. Dietary nutrient intake and antioxidant status in preeclamptic women. Adv Biomed Res 2015;4:183.

Development Goals (MDGs) of the United Nations Development Program.^[3] PE is the most common cause of maternal death in the world and affecting nearly 10–18% of pregnancies in developing countries.^[4]

The etiology of PE still remains unknown; however, several studies have examined the association of oxidative stress and PE together. Oxidative stress is created by an imbalance among the production of free radicals and antioxidant defense system.^[5-8] Deficiency of both enzymatic and non-enzymatic antioxidants is also observed in women with PE.^[9,10] Some investigators have reported that total antioxidant capacity (TAC) increases in the serum of women with PE.^[11] An inadequate intake of food resources of antioxidants leads to increase oxidative stress and adverse obstetrical outcomes.^[12-16] Nutrients may affect oxidative stress by increasing or decreasing free radicals or antioxidants or by providing substrate for the formation of free radicals.^[17-19] Data suggest that having a very low intake of vitamin C, E, selenium, and zinc in pregnancy is complied with an increased risk of PE.^[20,21] Some studies found that a diet rich in antioxidants, based on antioxidant-rich food consumption, increases the total antioxidant status and promotes better health.^[17] However, little is known about the role of maternal nutritional status on TAC in the PE. Therefore, we investigated the antioxidant status in pregnant women with PE and its relationship with dietary nutrient intake.

MATERIAL AND METHODS

This cross-sectional study was conducted between April and July in 2013 among pregnant women with PE admitted in the Obstetrics and Gynecology department of the Shahid Beheshti Hospital in Isfahan, Iran. It was approved by the Research and Ethics Committee of Isfahan University of Medical Sciences, and written informed consent was obtained from all subjects. Due to the lack of standard serum TAC, a preliminary study was conducted on 25 healthy pregnant women; according to its findings, the sample size was calculated to be 49 pregnant women with PE. Because of probable missing data, we considered the sample size to be 55.

Women were excluded if they did not give consent, were of gestational age under 18 years, were having a history of pre-existing health condition, including diabetes of type 1 or type 2, gestational diabetes, hypertension disorders, renal disease, heart disease, or other conditions that require special diets.

A team of trained gynecologist and dietitians conducted the study under standard protocols. Women

were interviewed about demographic data. Dietary intakes were obtained by using a validated 168-items semi-quantitative food frequency questionnaire (FFQ) Developed in Iran.^[22] The FFQ was a Willet format questionnaire modified for Iranian food items.^[23] This questionnaire was administered to determine frequency of food consumption during pregnancy and ask about the average frequency of intake of during last year. Dietary data were extracted by Nutritionist-4 software (First Databank Inc., Hearst Corp., San Bruno, CA) by a trained dietitian.

Gestational hypertensive patients were considered as subjects, namely with a systolic blood pressure of ≥ 140 mmHg and/or diastolic blood pressure of ≥ 90 mmHg on two occasions more than 4 h apart after 20 weeks in pregnancy. PE was considered to be gestational hypertension associated with proteinuria ≥ 300 mg protein in 24-h urine collection.^[24]

TAC of serum was determined from the sera that was frozen at -70°C and measured by using a double-antibody sandwich enzyme-linked immune-sorbent assay (ELISA) in samples.

Data analysis was performed using Statistical Package for the Social Sciences (SPSS) (version 20:0, SPSS Inc., Chicago, IL) software. Continuous variables are reported as the mean and standard deviation (SD). Non-parametric correlation statistics were used data failed to meet the assumptions of normality and equal variances. Spearman's rho was used to explore the relationships of both maternal dietary antioxidant and TAC. A $P < 0.05$ was considered as statistically significant.

RESULTS

A total of 55 eligible women consented to participate in the study. Table 1 presents data concerning socio-demographic and lifestyle variables. Mean values of age and gestational week were 29 years (SD = 5.5; range = 18–38) and average weeks 34 weeks (SD = 2.7; range = 30–40), respectively. 25.5% of women had a history of PE.

Table 1: General characteristics of women with preeclampsia

Characteristics	SD±Mean (N=55)
Age (years)	29.3±5.5
Education (years)	9.3±4.5
Gestational age (week)	34.1± 2.7
Pre-pregnancy BMI (kg/m ²)	25.7±5.5
Primiparous (%)	10.9
History of preeclampsia (%)	25.5
History of abortion (%)	21.8
History of stillbirth (%)	7.3
History of twin birth (%)	5.5

Based on the results obtained from the FFQ, the mean value of energy intake was 2157.7 ± 670.6 kcal/day. Women had significantly lower intakes of milk ($P = 0.0001$), meat or beans ($P = 0.015$) and higher consumption of grains ($P = 0.0001$) and fruit, vegetables ($P = 0.012$) compared to the recommended minimum number for pregnant women [Table 2]. All women were not taking supplements except iron and folic acid supplementation that is routinely used during pregnancy.

Table 3 compares the mean nutrient intake with Dietary Reference Intakes (DRIs). Women had significantly lower intakes of vitamin E, D, B6, niacin, magnesium, zinc, iron ($P < 0.05$), and higher consumption of vitamins A, C, thiamin, riboflavin, selenium, copper and calcium ($P < 0.05$) in comparing with DRI.

Mean of TAC was 3.86 ± 1.19 $\mu\text{mol/l}$ and significantly higher than healthy pregnant women in pilot study ($P < 0.04$). Because serum TAC failed to meet assumptions of normality distribution, nonparametric Spearman correlation were used. Correlation between serum TAC and dietary antioxidants are shown in Table 4, dietary intake of vitamin E had a significantly positive relationship with serum TAC ($r = 0.367$, $P = 0.003$) and selenium had a significantly negative relationship with serum TAC ($r = -0.298$, $P = 0.014$). However, no relationship was found between intake of vitamin C, β -carotene, riboflavin and copper with serum TAC.

DISCUSSION

In this study intake of vitamin E in women with PE was positively related to serum TAC; however, intake of vitamin E was lower than the dietary reference intakes. Rumbold *et al.*^[20] showed a very low intake of vitamin E in mid to late pregnancy associated with an increased risk of PE and gestational hypertension, even after adjusting for confounding factors. Limberaki *et al.*^[17] found that high consumption of antioxidant-rich foods increased the total antioxidant status and developed better health.

Our findings showed no significant correlation between intake of vitamin C, β -carotene, riboflavin, copper and serum TAC. A possible explanation for this might be concluded as dietary intake of vitamin C was higher than recommended allowance and fruit-vegetable consumption was more than recommended minimum number during pregnancy. Conflicting evidence was shown in the literature as to approve the relationship of intake of vitamins C and E with PE. Zhang *et al.*^[27] demonstrated that women who consumed <85 mg of vitamin C daily (lower than the recommended

Table 2: Comparison of dietary intakes of some different food group in subjects with recommended values

Food group	SD \pm Mean N=55	Recommended minimum number ^[25]	P*
Meat or beans (ounce)	3.7 \pm 2.1	7	0.0001
Milk (cup)	2.4 \pm 1.6	3	0.015
Grains (ounce)	10.7 \pm 5.3	7	0.0001
Fruit, vegetables (cup)	5.7 \pm 1.4	5	0.012

* $P < 0.05$ was considered significant. SD: Standard deviation

Table 3: Dietary nutrient intake and comparing with DRIs among subjects

Nutrient intake	SD \pm Mean N=55	Dietary reference intakes ^[26]	P*
Vitamin A (μg)	1334.3 \pm 807.4	770	0.0001
Beta-carotene (μg)	869.5 \pm 758.9	-	-
Vitamin C (mg)	135.9 \pm 66.6	85	0.0001
Vitamin D (μg)	2 \pm 1.6	15	0.0001
Vitamin E (mg)	13.9 \pm 5.3	15	0.12
Thiamin (mg)	1.6 \pm 0.5	1.4	0.005
Riboflavin (mg)	2.4 \pm 0.8	1.4	0.0001
Niacin (mg)	15.3 \pm 4.6	18	0.0001
Vitamin B6 (mg)	1.4 \pm 0.4	1.9	0.0001
Selenium (μg)	81.5 \pm 40.7	60	0.0001
Zinc (mg)	9.9 \pm 3.1	11	0.011
Iron (mg)	13.5 \pm 4.7	27	0.0001
Copper (mg)	1.6 \pm 0.5	1	0.0001
Magnesium (mg)	317.9 \pm 102.9	350	0.02
Calcium (mg)	1196.2 \pm 538.5	1000	<0.0001

* $P < 0.05$ was considered significant. DRIs: Dietary Reference Intakes

Table 4: Correlation between serum TAC and dietary antioxidants

Variables	r	P*
Vitamin E	0.367	0.003
Vitamin C	-0.041	0.384
β -carotene	0.125	0.181
Riboflavin	0.152	0.134
Selenium	-0.298	0.014
Copper	-0.036	0.398

* $P < 0.05$ was considered significant. TAC: Total antioxidant capacity

dietary allowance), as compared with others, experienced a doubled risk with PE. However, Klemmensen *et al.*^[28] concluded that incidence of PE did not correlate with dietary vitamin C and E intake. There was a decreasing trend in the incidence of severe PE and eclampsia with increasing dietary vitamin C intake.

In the present study dietary intake of selenium had a significant negative relationship with serum TAC. Intake of selenium was higher than dietary reference intakes and grain consumption was more than recommended minimum number during pregnancy.

Wheat is the major dietary sources of selenium and its content in foods depends on the selenium content

of the soil in which plants are grown or animals are raised.^[29-31] But bioavailability and effects on expression of the various selenoproteins depend on the form of selenium-containing products.^[32]

Selenoprotein S is an anti-inflammatory protein that acts primarily to limit the damaging consequences of endoplasmic reticulum stress.^[33] Some studies have suggested that selenium deficiency may be linked to PE.^[34,35] Conversely, others have shown higher serum selenium concentrations have been reported in women with PE.^[36]

Our finding showed women had significantly lower intakes of milk, meat or beans and higher consumption of grains and fruit, vegetables compared to the recommended minimum number for pregnant women, whereas Brantsaeter *et al.*^[37] suggested that a dietary pattern characterized by high intake of vegetables, plant foods, and vegetable oils decreases the risk of PE, whereas a dietary pattern characterized by high consumption of processed meat, sweet drinks, and salty snacks increases the risk.

Little studies were found on investigating issues of the serum TAC in pregnant women with PE and its relationship with dietary nutrient intake, but some research measured total oxidant status (TOS) and TAS in women with PE.

Ozturk *et al.*^[38] findings showed that maternal and cord plasma levels of TAS were significantly correlated with maternal and cord plasma levels of TOS in PE. Women whose pregnancies were complicated with PE and IUGR had elevated levels of TOS and TAS in comparison with healthy pregnant women.^[39] Fenzl *et al.*^[16] found that serum TAC and TOS concentrations were significantly higher in pregnant women with PE compared to pregnant controls. Increase of TOS in all pregnant women points to latent oxidative stress in pregnancy while increase of TAC in the early steps of preeclampsia might represent a defense mechanism of body against stress.

The current study has limitations that should be viewed cautiously. The smaller sample size and observational study design prevent establishing a definitive influence of maternal nutrient intake on antioxidant status. One explanation may be used on conflicting findings in the process of methodology in order to assess dietary intake, where some studies have used 24-h recall compared with a food frequency questionnaire. The study did not provide information regarding serum micronutrient concentrations and limited our ability to assess diet intake and serum TAC.

A major strength of this study is designing a preliminary test for accessing a normal range of TAC among normal pregnant women. However, women were carefully interviewed by dietitian about demographic data and dietary intakes were compared to questionnaires which have been filled by the subjects.

However, more prospective studies with larger sample size are needed to explore the relationships of both maternal nutrient intake and antioxidant status and to confirm the existing findings. These findings highlight that the intake of vitamin E was positively associated with serum TAC. Little support was found in the relationship between dietary intakes of other micronutrients and serum TAC. These data suggest that antioxidant activity could be increased by high dietary intakes of fruit, vegetables and other rich in antioxidant constituents and that increasing consumption of foods rich in antioxidant, may improve pregnancy-induced hypertension and PE. Further research is required to explore the relationships of maternal nutrient intake and antioxidant status in women with PE.

ACKNOWLEDGMENTS

The authors appreciate the Dear mothers whom participated in this study.

REFERENCES

1. Souza JP, Cecatti JG, Faundes A, Morais SS, Villar J, Carroli G, *et al.* Maternal near miss and maternal death in the World Health Organization's 2005 global survey on maternal and perinatal health. *Bull World Health Organ* 2010;88:113-9.
2. Noris M, Perico N, Remuzzi G. Mechanisms of disease: Pre-eclampsia. *Nature clinical practice Nephrology* 2005;1:98-114.
3. MDGR Millennium Development Goals Report 2010. New York, NY, USA: United Nations Department of Economic and Social Affairs; 2010.
4. WHO, "Risking death to give life". Geneva, Switzerland: World Health Organization; 2005.
5. Scholl TO, Leskiw M, Chen X, Sims M, Stein TP. Oxidative stress, diet, and the etiology of preeclampsia. *Am J Clin Nutr* 2005;81:1390-6.
6. Poston L, Chappell LC. Is oxidative stress involved in the aetiology of pre-eclampsia? *Acta Paediatr Suppl* 2001;90:3-5.
7. Rajmakers MT, Dechend R, Poston L. Oxidative stress and preeclampsia: Rationale for antioxidant clinical trials. *Hypertension* 2004;44:374-80.
8. Rice-Evans C, Burdon R. Free radical lipid interactions and their pathological consequences. *Prog Lipid Res* 1993;32:71-110.
9. Huppertz B, Kingdom JC. Apoptosis in the trophoblast-role of apoptosis in placental morphogenesis. *J Soc Gynecol Investig* 2004;11:353-62.
10. Redman CW, Sargent IL. Latest advances in understanding preeclampsia. *Science* 2005;308:1592-4.
11. Mert I, Oruc AS, Yuksel S, Cakar ES, Buyukkagnici U, Karaer A, *et al.* Role of oxidative stress in preeclampsia and intrauterine growth restriction. *J Obstet Gynaecol* 2012;38:658-64.
12. Fowles E, Murphey C, Ruiz RJ. Exploring relationships among psychosocial status, dietary quality and measures of placental development during the 1st trimester in low-income women. *Biol Res Nurs* 2011;13:70-9.
13. Roberts JM, Balk JL, Bodnar LM, Belizan JM, Bergel E, Martinez A. Nutrient involvement in preeclampsia. *J Nutr* 2003;133(Suppl 2):1684S-92.

14. Fiore G, Capasso A. Effects of vitamin E and C on placental oxidative stress: An *in vitro* evidence for the potential therapeutic or prophylactic treatment of preeclampsia. *Med Chem* 2008;4:526-30.
15. Schiff E, Friedman SA, Stampfer M, Kao L, Barrett PH, Sibai BM. Dietary consumption and plasma concentrations of vitamin E in pregnancies complicated by preeclampsia. *Am J Obstet Gynecol* 1996;175:1024-8.
16. Fenzl V, Flegar-Mestric Z, Perkovic S, Andric L, Tatzber F, Zarkovic N, *et al.* Trace elements and oxidative stress in hypertensive disorders of pregnancy. *Arch Gynecol Obstet* 2013;287:19-24.
17. Limberaki E, Eleftheriou P, Vagdatli E, Kostoglou V, Petrou C. Serum antioxidant status among young, middle-aged and elderly people before and after antioxidant rich diet. *Hippokratia* 2012;16:118-23.
18. Winkhofer-Roob BM, Rock E, Ribalta J, Shmerling DH, Roob JM. Effects of vitamin E and carotenoid status on oxidative stress in health and disease. Evidence obtained from human intervention studies. *Mol Aspects Med* 2003;24:391-402.
19. Huang HY, Appel LJ. Supplementation of diets with alpha-tocopherol reduces serum concentrations of gamma- and delta-tocopherol in humans. *J Nutr* 2003;133:3137-40.
20. Rumbold AR, Maats FH, Crowther CA. Dietary intake of vitamin C and vitamin E and the development of hypertensive disorders of pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2005;119:67-71.
21. Negi R, Pande D, Karki K, Kumar A, Khanna RS, Khanna HD. Trace elements and antioxidant enzymes associated with oxidative stress in the pre-eclamptic/eclamptic mothers during fetal circulation. *Clin Nutr* 2012;31:946-50.
22. Mirmiran P, Azadbakht L, Azizi F. Dietary behaviour of Tehranian adolescents does not accord with their nutritional knowledge. *Public Health Nutr* 2007;10:897-901.
23. Willett W. *Nutritional epidemiology*. USA: Oxford University Press; 1998.
24. National High Blood Pressure Education program working group report on high blood pressure in pregnancy. *Am J Obstet Gynecol* 1990;163:1691-712.
25. Mahan LK, Escott-Stump S, Raymond J. *Krause's Food and the nutrition care process*. 13th ed. Philadelphia, PA: Saunders; 2012. p. 361.
26. Mahan LK, Escott-Stump S, Raymond J. *Krause's Food and the nutrition care process*. 13th ed. Philadelphia, PA: Saunders; 2012. p. 1229-30.
27. Zhang C, Williams MA, King IB, Dashow EE, Sorensen TK, Frederick IO, *et al.* Vitamin C and the risk of preeclampsia-results from dietary questionnaire and plasma assay. *Epidemiology* 2002;13:409-16.
28. Klemmensen A, Tabor A, Osterdal ML, Knudsen VK, Halldorsson TI, Mikkelsen TB, *et al.* Intake of vitamin C and E in pregnancy and risk of pre-eclampsia: Prospective study among 57 346 women. *BJOG* 2009;116:964-74.
29. Reilly C. *Selenium in food and health*. 2nd ed, Vol. 26. New York, N.Y.: Springer; 2006. p. 206.
30. Lyons G, Stangoulis J, Graham R. Nutriprevention of disease with high-selenium wheat. *Journal of the Australian College of Nutritional and Environmental Medicine* 2003;22:3-9.
31. Rayman MP. The importance of selenium to human health. *Lancet* 2000;356:233-41.
32. Rayman MP. Food-chain selenium and human health: Emphasis on intake. *Br J Nutr* 2008;100:254-68.
33. Ye Y, Shibata Y, Yun C, Ron D, Rapoport TA. A membrane protein complex mediates retro-translocation from the ER lumen into the cytosol. *Nature* 2004;429:841-7.
34. Mistry HD, Wilson V, Ramsay MM, Symonds ME, Broughton Pipkin F. Reduced selenium concentrations and glutathione peroxidase activity in pre-eclamptic pregnancies. *Hypertension* 2008;52:881-8.
35. Maleki A, Fard MK, Zadeh DH, Mamegani MA, Abasaizadeh S, Mazloomzadeh S. The relationship between plasma level of se and preeclampsia. *Hypertens Pregnancy* 2010;30:180-7.
36. Mahomed K, Williams MA, Woelk GB, Mudzamiri S, Madzime S, King IB, *et al.* Eukocyte selenium, zinc, and copper concentrations in preeclamptic and normotensive pregnant women. *Biol Trace Elem Res* 2000;75:107-18.
37. Brantsaeter AL, Haugen M, Samuelsen SO, Torjusen H, Trogstad L, Alexander J, *et al.* A dietary pattern characterized by high intake of vegetables, fruits, and vegetable oils is associated with reduced risk of preeclampsia in nulliparous pregnant Norwegian women. *J Nutr* 2009;139:1162-8.
38. Ozturk E, Balat O, Acilmis YG, Ozcan C, Pence S, Erel O. Measurement of the placental total antioxidant status in preeclamptic women using a novel automated method. *J Obstetric Gynaecol Res* 2011;37:337-42.
39. Mert I, Oruc AS, Yuksel S, Cakar ES, Buyukkagnici U, Karaer A, *et al.* Role of oxidative stress in preeclampsia and intrauterine growth restriction. *J Obstet Gynaecol Res* 2012;38:658-64.

Source of Support: Isfahan University of Medical Sciences, Food Security and Nutrition Research Center, **Conflict of Interest:** None declared.