

LETTER


Successful treatment of refractory Sneddon-Wilkinson disease (subcorneal pustular dermatosis) with infliximab

Dear Editor,

Sneddon-Wilkinson disease, also known as subcorneal pustular dermatosis (SPD), is a rare neutrophilic dermatosis clinically characterized by vesico-pustular lesions often distributed in an annular or serpiginous pattern, which may evolve in superficial erosions resolving with scaling and residual dyspigmentation.¹ It commonly involves flexural areas and abdomen in middle-aged women. Histologically is characterized by subcorneal pustules with uninvolved epidermis and negative direct and indirect immunofluorescence.¹ Clinical course is relapsing–remitting with frequent flares and, due to its rareness, no treatment guidelines are available. Dapsone represents the first-line therapy whereas corticosteroids, oral retinoids and immunosuppressants (i.e., methotrexate, cyclosporine) represent alternative treatments. In more recent years, anti-tumor necrosis factor (TNF)- α has been successfully administered in multi-drug resistant patients.¹

We present here the case of an 80-years-old woman followed in our department for a SPD diagnosed in 2012, after the appearance of a vesicular eruption located on the abdomen, submammary area, groins and arms. The lesions had a wax and waning course, lasting for few days with periodic relapses (Figure 1A). Laboratory examinations revealed only a slight elevation of C-reactive protein (CRP, 11.2 mg/L) whereas complete blood count, immunoglobulins and biochemistry including liver and renal function were normal. A punch biopsy confirmed the diagnosis of SPD (Figure 1B) while direct and indirect immunofluorescence were negative. At the time of diagnosis the patient was treated with dapsone at 0.8–1 mg/kg/day with clinical improvement, but frequent relapses occurred, which required oral and intravenous corticosteroids administrations. Past therapies in addition to dapsone and corticosteroids included acitretin, withdrawn for a strong pruritus, cyclosporine and methotrexate suspended for

FIGURE 1 (A) erythematous serpiginous patches with central clearing and crusted edge as resolution of previous vesicular lesions located on the flank of our patient during one of her relapses; (B) histology of a vesicular lesion showing subcorneal accumulation of neutrophils with uninvolved epidermis and dermis (hematoxylin and eosin stain, $\times 20$); (C) erythematous and serpiginous active lesions along with residual dyspigmentation on the left flank and the abdomen before the administration of infliximab; (D) only residual hyperpigmentation on the trunk, arms and mammary area after 5 months of therapy with infliximab.



This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Dermatologic Therapy* published by Wiley Periodicals LLC.

inefficacy, and finally colchicine discontinued for severe gastrointestinal side effects.

No other relevant comorbidities were present; in September 2021 the patient experienced another relapse (Figure 1C); blood examination revealed a total white blood cell count (WBC) of $14.31 \times 10^9/L$ with $10.3 \times 10^9/L$ neutrophils (72% of WBC), C-Reactive Protein was 9.7 mg/L and creatinine 1.16 mg/dl while further complete blood count values, liver function, methaemoglobin, electrolytes and immunoglobulins (IgA, IgM, IgG) were all within the normal range.

Following the flare-up of the disease we decided to introduce infliximab (anti-TNF α) at the induction dose of 5 mg/kg at week 0, 2 and 6 followed by maintenance dose of 5 mg/kg every 8 weeks, in combination with dapsone (50 mg/day), while methylprednisolone was gradually tapered. A marked improvement was obtained a week after the first injection; complete clinical remission was achieved after a month and maintained until her last follow-up visit in January 2022 (Figure 1D). No side effects were reported.

SPD belongs to the group of superficial neutrophilic dermatoses characterized by epidermal sterile neutrophil-rich infiltrates.² A significant clinical and histopathological overlap exists between SPD and the annular form of pustular psoriasis and whether these two entities represent different diseases is still a matter of debate; SPD has been classically described in middle-aged woman mainly affecting the intertriginous and flexural areas, as in our patient.³ Its pathophysiology remains elusive but abnormal neutrophil recruitment and activation as well as the evidence of higher level of TNF- α in affected skin and sera of patients suggest that this cytokine could play a crucial role in the etiology of the disease.^{2,4,5} These findings, together with the recalcitrant clinical course, have led to consider anti-TNF- α as a valuable therapy in multi-drug-resistant cases.⁶⁻¹⁰ Voigtländer et al. were the first ones to use infliximab to successfully treat flares of SPD in a patient.⁶ Other reports showed both safety and efficacy of anti TNF- α in patients with SPD associated with monoclonal gammopathy^{7,8} and systemic lupus erythematosus.⁹ Resistance to infliximab is reported in some cases which required dose intensification or switching to a different anti TNF- α to control the disease^{5,7,8,10}; this could be explained by the production of anti-drug antibodies or a reduced adherence to injection schedule.⁸

Basing on previous reports and our experience we suggest to administer infliximab as an early and effective treatment in severe and recalcitrant cases, with a particular attention to its infusion regimen according to the clinical course, either increasing the dose or the frequencies of injections, maintaining it in combination with dapsone or other immunomodulators, and switching to other anti-TNF- α in case of loss of efficacy. Further reports and long-term follow-up of patients are needed to assess the optimal treatment regimen for this rare and challenging disease.

AUTHOR CONTRIBUTIONS

Maurizio Romagnuolo drafted, conceptualized, and designed the manuscript; Simona Muratori, Angelo Cattaneo, Chiara Moltrasio contributed to the acquisition and analysis of the data, Angelo Valerio Marzano revised it. All the authors accepted the final version of the manuscript.

ACKNOWLEDGMENT

Open access funding provided by Università degli Studi di Milano within the CRUI-CARE Agreement.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Maurizio Romagnuolo^{1,2}

Simona Muratori¹

Angelo Cattaneo¹

Angelo Valerio Marzano^{1,2} 

Chiara Moltrasio^{2,3}

¹Dermatology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

²Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Milan, Italy

³Department of Medical Surgical and Health Sciences, University of Trieste, Trieste, Italy

Correspondence

Maurizio Romagnuolo, Dermatology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Via Pace 9, 20122 Milan, Italy.
Email: maurizio.romagnuolo@unimi.it; maurizioromagnuolo@hotmail.it

ORCID

Angelo Valerio Marzano  <https://orcid.org/0000-0002-8160-4169>

REFERENCES

- Cheng S, Edmonds E, Ben-Gashir M, Yu RC. Subcorneal pustular dermatosis: 50 years on. *Clin Exp Dermatol*. 2008;33(3):229-233.
- Marzano AV, Ortega-Loayza AG, Heath M, Morse D, Genovese G, Cugno M. Mechanisms of inflammation in neutrophil-mediated skin diseases. *Front Immunol*. 2019;10:1059.
- Watts PJ, Khachemoune A. Subcorneal pustular dermatosis: a review of 30 years of Progress. *Am J Clin Dermatol*. 2016;17(6):653-671.
- Grob JJ, Mege JL, Capo C, et al. Role of tumor necrosis factor-alpha in Sneddon-Wilkinson subcorneal pustular dermatosis. A model of neutrophil priming in vivo. *J Am Acad Dermatol*. 1991;25(5 Pt 2):944-947.
- Bonifati C, Trento E, Cordiali Fei P, Muscardin L, Amantea A, Carducci M. Early but not lasting improvement of recalcitrant subcorneal pustular dermatosis (Sneddon-Wilkinson disease) after infliximab therapy: relationships with variations in cytokine levels in suction blister fluids. *Clin Exp Dermatol*. 2005;30(6):662-665.
- Voigtländer C, Lüftl M, Schuler G, Hertl M. Infliximab (anti-tumor necrosis factor alpha antibody): a novel, highly effective treatment of recalcitrant subcorneal pustular dermatosis (Sneddon-Wilkinson disease). *Arch Dermatol*. 2001;137(12):1571-1574.
- Versini M, Mantoux F, Angeli K, Passeron T, Lacour JP. Maladie de Sneddon-Wilkinson échappant à l'infliximab et à l'éta nercept: efficacité de l'adalimumab [Sneddon-Wilkinson disease: efficacy of

- intermittent adalimumab therapy after lost response to infliximab and etanercept]. *Ann Dermatol Venereol*. 2013;140:797-800.
8. Guerin C Jr, Beylot-Barry M, Frouin E, Hainaut E, Masson RM. Treatment of subcorneal pustular dermatosis (Sneddon-Wilkinson disease) with anti-tumor necrosis factor alpha. *Cureus*. 2021;13(8): e17147.
 9. Naretto C, Baldovino S, Rossi E, Spriano M, Roccatello D. The case of SLE associated Sneddon-Wilkinson pustular disease successfully and safely treated with infliximab. *Lupus*. 2009;18:856-857.
 10. Kretschmer L, Maul JT, Hofer T, Navarini AA. Interruption of Sneddon-Wilkinson subcorneal Pustulation with infliximab. *Case Rep Dermatol*. 2017;9(1):140-144.