

Research article

Open Access

Current limitations of SNP data from the public domain for studies of complex disorders: a test for ten candidate genes for obesity and osteoporosis

Volodymyr Dvornyk^{†1}, Ji-Rong Long^{†1}, Dong-Hai Xiong^{†1}, Peng-Yuan Liu¹, Lan-Juan Zhao¹, Hui Shen¹, Yuan-Yuan Zhang¹, Yong-Jun Liu¹, Sonia Rocha-Sanchez¹, Peng Xiao¹, Robert R Recker¹ and Hong-Wen Deng^{*1,2}

Address: ¹Osteoporosis Research Center and Department of Biomedical Sciences, Creighton University, 601 N. 30th St., Suite 6730, Omaha, NE 68131, USA and ²Laboratory of Molecular and Statistical Genetics, College of Life Sciences, Hunan Normal University, Changsha, Hunan 410081, P. R. China

Email: Volodymyr Dvornyk - dvornyk@creighton.edu; Ji-Rong Long - jrlongyou@creighton.edu; Dong-Hai Xiong - donghai@creighton.edu; Peng-Yuan Liu - [pyliu@creighton.edu](mailto:pylui@creighton.edu); Lan-Juan Zhao - zhaolanjuan@creighton.edu; Hui Shen - huishen@creighton.edu; Yuan-Yuan Zhang - zhangyuan2@creighton.edu; Yong-Jun Liu - jun@creighton.edu; Sonia Rocha-Sanchez - saraujo@creighton.edu; Peng Xiao - peng@creighton.edu; Robert R Recker - rrecker@creighton.edu; Hong-Wen Deng* - deng@creighton.edu

* Corresponding author †Equal contributors

Published: 25 February 2004

Received: 24 November 2003

BMC Genetics 2004, 5:4

Accepted: 25 February 2004

This article is available from: <http://www.biomedcentral.com/1471-2156/5/4>

© 2004 Dvornyk et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Public SNP databases are frequently used to choose SNPs for candidate genes in the association and linkage studies of complex disorders. However, their utility for such studies of diseases with ethnic-dependent background has never been evaluated.

Results: To estimate the accuracy and completeness of SNP public databases, we analyzed the allele frequencies of 41 SNPs in 10 candidate genes for obesity and/or osteoporosis in a large American-Caucasian sample (1,873 individuals from 405 nuclear families) by PCR-invader assay. We compared our results with those from the databases and other published studies. Of the 41 SNPs, 8 were monomorphic in our sample. Twelve were reported for the first time for Caucasians and the other 29 SNPs in our sample essentially confirmed the respective allele frequencies for Caucasians in the databases and previous studies. The comparison of our data with other ethnic groups showed significant differentiation between the three major world ethnic groups at some SNPs (Caucasians and Africans differed at 3 of the 18 shared SNPs, and Caucasians and Asians differed at 13 of the 22 shared SNPs). This genetic differentiation may have an important implication for studying the well-known ethnic differences in the prevalence of obesity and osteoporosis, and complex disorders in general.

Conclusion: A comparative analysis of the SNP data of the candidate genes obtained in the present study, as well as those retrieved from the public domain, suggests that the databases may currently have serious limitations for studying complex disorders with an ethnic-dependent background due to the incomplete and uneven representation of the candidate SNPs in the databases for the major ethnic groups. This conclusion attests to the imperative necessity of large-scale and accurate characterization of these SNPs in different ethnic groups.

Background

A single nucleotide polymorphism (SNP) is generally defined as a stable substitution of a single base with a frequency of more than 0.01 in at least one population [1]. In human genetic studies, SNPs are simply referred to as bi-allelic markers since the other types (tri-allelic and tetra-allelic SNPs) are very rare in the human genome [2]. SNPs have been recognized as an important tool in human genetics and medicine [3,4]. Because SNPs are abundant and scattered throughout the whole human genome, they have been widely used in the genetic association studies of various complex diseases such as obesity, osteoporosis, asthma, hypertension (see, e.g., [5-7]). Various analyses of SNPs across the human genome have been also conducted to determine haplotype patterns in human populations [8-12]. These data are very useful to study the genetic basis of common complex diseases [13].

At present there are several SNP public databases. The largest are dbSNP and HGVbase containing together several million SNPs [14]. There also exist other relatively small or specific SNP databases, e.g., TSC [15], JSNP [16], HOWDY [17], GeneSNPs [18]. While the databases have been continuously expanded, the quality and completeness of the deposited SNP data remain to be of particular importance and to be assessed. Recent evaluation of quality and comprehensiveness of about four million candidate SNPs from some public and Celera databases showed that about 6–12% SNPs could not be validated [19]. They represent rare variants, population-specific SNPs and sequencing errors. The other studies reported that only about 50% SNPs are common in any given population. Therefore, measurement of allele frequency and linkage disequilibrium for SNPs in databases should be done to efficiently select a minimal subset of SNPs for population association studies of complex diseases [20,21].

Obesity and osteoporosis are common complex disorders with continuously growing burden and cost for their prevention and treatment. For example, the most recent data for the United States, derived from the third National Health and Nutrition Examination Survey (1988–94), showed ~20% of US men and ~25% of US women are obese and these proportions are increasing [22]. The same tendency has been observed in other human populations [23]. Osteoporosis is another major health problem, particularly in the elderly. More than 40% of postmenopausal women, on average, will suffer at least one osteoporotic fracture [24]. This disorder incurred an estimated direct cost of ~14 billion dollars in the USA alone in 1995 [25].

SNPs of candidate genes for obesity and osteoporosis may be of particular importance in genetic studies of these disorders. The allele frequencies are important in the selec-

tion of SNPs for studying complex diseases [26]. For example, association studies of osteoporosis usually generate inconsistent results in different ethnic groups [27]. One important reason is that polymorphisms associated significantly with certain osteoporotic phenotypes in a given ethnic group may be absent or rare in another ethnic group [28-31]. Moreover, racial differences in the prevalence of certain allele could account for certain proportion of disease trait variation between different ethnicities [32]. Finally, a comparison of SNP allele frequencies among different ethnicities can provide valuable information for mapping by admixture linkage disequilibrium (MALD) [33].

Despite the importance of the above data, no attempts have been made to check the quality and completeness of SNPs of candidate genes for obesity and osteoporosis in the public databases. In fact, most allele frequencies in the databases were obtained from studies of relative small samples (dozens or so), which could yield large sampling errors. In addition to the databases, there are abundant SNP data from many association/linkage studies of these two common complex diseases. These individually published data have not yet been assessed in reference to each other, or to those in the databases. In most association studies, the regular sample sizes used to detect allele frequencies have only been of the order of a hundred or so. However, to obtain reliable estimates of allele frequency distributions, relatively larger sample sizes are needed. For example, for an SNP, to let allele frequency estimate's $|error| = 0.05$, the required sample sizes for frequency distribution (0.5, 0.5), (0.7, 0.3), and (0.9, 0.1) should be at least 1150, 1000, and 400, respectively [34]. Otherwise, the estimates of allele frequencies may be biased to a large extent.

In the present study, we determined allele frequencies of 41 SNPs in 10 candidate genes for obesity and/or osteoporosis in a large American-Caucasian sample (1,873 subjects from 405 nuclear families). We then compared them with the corresponding data for the major ethnic groups obtained from the public databases and various individual studies to check their accuracy and completeness, and to compare the SNP allele frequencies in different ethnic groups.

Results

SNP polymorphism in the studied Caucasian sample

Of the 41 candidate SNPs for obesity and/or osteoporosis identified in the studied Caucasian sample (Table 1), 29 (70.7%) had minor allele frequency ≥ 0.1 . Eight SNPs (19.5%) were monomorphic. Allele frequencies of twelve SNPs in Caucasians were reported for the first time (Table 2). Five of these SNPs were monomorphic and the other seven were polymorphic with minor allele frequencies ranged 0.019–0.398.

Table 1: Summary information about the studied SNPs

SNP ^a	Gene ^b	Name ^c	dbSNP Accession	Mutation ^d	Domain	Minor allele frequency	Forward primer	Reverse primer
1	APOE		ss12568587	G-C	5' UTR	0.357	TCCCCAGGAGCCGGTGA	CCCCAAGCCCAGCCCC
2	APOE		ss12568609	G-A	Intron 2	0.399	CCTCAGGTGATCTGCCCGTTTC	ACTCCTGGGCTCAAGTGATCCTC
3	APOE		ss12568607	T-C	Exon 4	0.149	CGGGCACGGCTGTCCAA	CGAGCATGGCCTGCACCTC
4	APOE		ss12568612	C-T	Exon 4	0.087	GCTGCGTAAGCGGCTCC	GCGGCCCTGTTCCACC
5	COL1 α 1		ss12568606	G-T	5' UTR	0.154	GCACCCTGCCCTAGACCAC	CCTAGTGCCAGCGACTGCA
6	COL1 α 1	SpI	ss12568597	G-T	Intron 1	0.188	CCAATCAGCCGCTCCCATTC	CATCGGGAGGGCAGGCTC
7	COL1 α 1		ss12568598	G-T	Exon 8	0.000	GGAAGACTGGGATGAGGGCA	GGCTCGCCAGGCTCACC
8	COL1 α 1		ss12568584	G-A	Exon 45	0.019	CTCAGCCTTCCCTGGCCAA	AGGCGGAAGTTCATTGGCATC
9	ER- α		ss12568579	A-G	Exon 1	0.484	TTGAGCTGCGGACGGTTCA	CGCCGGTTTCTGAGCCTTC
10	ER- α	PvuII	ss12568596	T-C	Intron 1	0.449	TGGGATTCCAGGCATGAACCAC	TGGCGTCGATTATCTGAATTTGGCC
11	ER- α		ss12568619	G-A	Intron 3	0.257	CCCAGAAACAAGTCATCTGCTATTGACA	TGTAACAAAAGGTTAAACAATGGTTAGCCC
12	ER- α		ss12568618	G-C	Exon 4	0.218	ACAGCCTGGCCTTGTCCTC	CAGGTTGGTCAGTAAGCCCATCA
13	ER- α		ss12568585	G-A	Intron 4	0.098	GATCAATGAAGTGGTCTTGAAAAACCAA	GGTGACAAGCTGAAAATCTAAGCTTCA
14	ER- α		ss12568605	G-A	Intron 6	0.119	GGAAACGGCCCTTGAAATTGTAAA	CTGCCTACAGAATACAGTCAGCCA
15	ER- α		ss12568617	G-A	Exon 8	0.203	TCGCATTCTTGCAAAAGTATTACATCAC	CAAGCAATGAATGGCCACTCATCTAGAAA
16	IL-6		ss12568616	C-G	5' UTR	0.434	GGGCAGAATGAGCCTCAGACATC	GACATGCCAAAGTGCTGAGTCACTAATA
17	IL-6		ss12568586	A-T	Exon 4	0.005	CCTCCACTGCAAAGGATTTATTCAACA	CATGTCTGCGCCACTGGTTC
18	LEPR		ss12568615	A-G	Exon 4	0.290	AGATTTAAGTTGTCTTGCATGCCACC	TTAAGCCCAGCATCCATTAGCTATTCTTTTC
19	LEPR		ss12568604	G-C	Exon 14	0.182	GAGTAATTGGAGCAATCCAGCCTACA	GCTTACGCCACTGTACATCTTAGCTC
20	LEPR		ss12568614	G-A	Exon 20	0.353	GCCACGCTGATCAGCAACTC	CCCTTGACTTGTCTAGTCAAAAAGCAC
21	PTHRI		ss12568589	G-A	Intron 1	0.001	GACTTACATTAGGATTCAGGTTACTGCCA	GGGACGCAAGCCTGAGTCC
22	PTHRI		ss12568592	A-G	Intron 2	0.398	GCAGAACCCTAAGGGCTTGTC	GGCGGGACCCAGGATACA
23	PTHRI		ss12568591	C-T	Intron 8	0.374	CGAGCCTCAATTCAGGTGAATCTAACC	CCCGCCCCAAGTGAACA
24	PTHRI		ss12568588	G-A	Intron 10	0.397	CCTTGAGCCCTTGTTTTCTTTTC	GCTCCGGGAACAAAAGTGGATCA
25	PTHRI		ss12568590	G-A	Exon 13	0.380	CTACAAGGCTCAAATGCCCCAAA	TTGGCGTCCACTACATTGTCTTCA
26	TGF- β 1		ss12568613	C-T	5' UTR	0.311	GGGCCAGTTTCCCTATCTGTAAA	CTGGGCCACCGTCTCATC
27	TGF- β 1		ss12568603	+C/-C	Intron 4	0.021	CCAGCCCCACTTATCTATCCC	GGAAAGGCCGGTTCATGCCA
28	TGF- β 1		ss12568593	C-T	Exon 5	0.008	CAGGCTACAAGGCTCACCTGAA	GGTTCACTACCGGCCGC
29	TGF- β 1		ss12568602	C-T	Intron 5	0.274	GGCTTGTCTTAAGCATTGCGTGAAATTAA	GTACAGCTGCCGACGC
30	TNFR2		ss12568594	T-C	5' UTR	0.000	TGCACTCGGCCTGTTTAGACTC	CTGTTTCTGCCCCCTGCC
31	TNFR2		ss12568611	T-G	Exon 6	0.198	AGCCACCCAGCCACTC	GCTTGAGCAGTGCTGGGTTTC
32	UCP3		ss12568601	C-T	5' UTR	0.265	CACTGCCCTCACCAGCCA	GTGAGTCTGCCACGGCA
33	UCP3		ss12568600	G-A	Exon 2	0.000	GCCCTAAAGGGACTGGGCA	GAAAGGTAACGAGGTGAGCAAAAACA
34	UCP3		ss12568595	T-C	Exon 3	0.258	TGATTCCTGTAACATCTGGACTTTTCATC	CTGCCTAAATCCCCTTAGCAGAAAAA
35	UCP3		ss12568580	T-C	Exon 5	0.447	CCTAACAGGAACCTTGCCCAACATCA	TCCACGGAGTCTGGGTTCC
36	UCP3		ss12568599	C-T	Exon 7	0.000	CCTAACAGGAACCTTGCCCAACATCA	TCCACGGAGTCTGGGTTCC
37	VDR		ss12568583	G-A	5' UTR	0.281	CAGCATGCCTGTCTCAGC	CCAGTACTGCCAGCTCCCA
38	VDR	FokI	ss12568581	C-T	Exon 2	0.373	TGGCCCTGGCACTGACTC	GGCACGTTCCGGTCAAAGTC
39	VDR		ss12568582	C-T	Exon 4	0.000	GGACAGTCTGCGGCCCA	CCCTACTCCCTGGGCC
40	VDR	BsmI	ss12568610	G-A	Intron 8	0.419	GTGCCCTCACTGCCCTTA	CCTCAAATAACAGGAATGTTGAGCCCA
41	VDR	TaqI	ss12568608	T-C	Exon 9	0.408	GGCCAGGCAGTGGTATCAC	AGGTCGGCTAGCTTCTGGATCA

^aDesignation in the present study. ^bAbbreviations: APOE, apolipoprotein E; COL1 α 1, collagen type I α 1; ER- α , estrogen receptor- α ; IL-6, interleukin-6; LEPR, leptin receptor; PTHRI, parathyroid hormone (PTH)/PTH-related peptide receptor type I; TGF- α 1, transforming growth factor- β 1; TNFR2, tumor necrosis factor receptor 2; UCP3, uncoupling protein 3; VDR, vitamin D (1,25-dihydroxyvitamin D₃) receptor. ^cCommon name. ^dMinor alleles are given in bold.

Table 2: Minor allele frequencies of SNPs firstly reported for Caucasians by this study

Gene	SNP	Database or reference ^a	Sample size	Minor allele frequency
COL1 α 1	7	[52]	292	0.000
COL1 α 1	8	dbSNP	1,861	0.019
ER- α	14	JSNP	1,861	0.119
IL-6	17	dbSNP	190	0.005
PTHR1	21	JSNP	380	0.001
PTHR1	22	JSNP	1,861	0.398
PTHR1	23	JSNP	1,863	0.374
PTHR1	24	JSNP	1,855	0.397
PTHR1	25	JSNP	1,851	0.380
TNFR2	30	JSNP	281	0.000
VDR	37	JSNP	1,861	0.281
VDR	39	dbSNP	190	0.000

For gene abbreviations and SNP designations see Table 1. ^aThe source of information about the given SNP.

Data on the candidate SNPs for obesity and/or osteoporosis from major ethnic groups: an analysis of literature and databases

Caucasians

In available literature and public SNP databases, we found data about allele frequencies of 29 SNPs (representing 9 genes) in Caucasians, which were shared with the present study. The corresponding data are given in Table 3. The allele frequencies of only 14 SNPs had been previously reported for Caucasians to the public databases (dbSNP and/or HGvbase). No significant differences were found between the allele frequencies in our study and in the public databases, except for SNP31 in Australians ($P = 0.05$, Table 3).

In contrast to the databases, available literature contained data on allele frequencies of the 23 SNPs. We determined significant differences between our and literature data for allele frequencies of four SNPs (SNP16, SNP32, SNP35, and SNP36). We have not found any allele frequency data on the SNPs of the PTHR1 gene (SNP21-SNP25) for Caucasians. Some SNP indicated large discrepancies in allele frequencies between the data from the different databases. For example, the minor allele frequencies of SNP26 for two Caucasian samples reported to dbSNP and HGvbase are 0.43 and 0.26, respectively (Table 3). The differences between the values are not statistically significant due to

the small sample sizes ($n = 31$ and $n = 42$, respectively), so it is impossible to conclude what actual allele frequencies are in these populations and whether they are indeed significantly different.

African/African-Americans

Table 4 presents our SNP data in comparison with the corresponding data on African and/or African-American samples obtained from literature and the databases. In the public domain, we found in total only 18 SNPs from seven genes, which are shared with our study. The candidate genes for obesity and/or osteoporosis appear to be underrepresented for Africans/African-Americans in the public SNP databases and literature: only 11 shared SNPs were found in the databases and seven in the literature. Among those, one (SNP36) was monomorphic and four had minor allele frequency <0.1 . Significant differences in allele frequencies between Caucasians and Africans/African-Americans were found at three SNPs (SNP16, SNP34 and SNP35). Three more SNPs (SNP32, SNP33, and SNP41) manifested nearly significant differences in allele frequencies between these ethnic groups (Table 4). We found no data on the SNPs of three genes (COL1 α 1, PTHR1, and TNFR2) for Africans/African-Americans in the public domain, which would be shared with our study.

Similar to the case with Caucasians, there are large inconsistencies between some SNP data for Africans/African-Americans from the different databases. For example, the reported minor allele frequencies of SNP26 from the TSC and dbSNP databases are 0.27 and 0.15, respectively (Table 4). The former is not significantly different ($P = 0.61$) from the corresponding value in Caucasians, while the latter is nearly significantly different ($P = 0.07$). However, it is hard to determine, which value is true because of the small sample sizes ($n = 42$ and $n = 24$, respectively) in the databases.

Asians

As compared to our SNP data, this ethnic group had corresponding data on 22 SNPs (8 genes) in the public domain (Table 5). The data on 13 SNPs of the 6 genes were available from the public databases, and on 10 SNPs of the 5 genes were reported in the literature. Among those SNPs, two (SNP16 and SNP33) were monomorphic and one (SNP36) had minor allele frequency <0.1 . Two more SNPs of the VDR gene (SNP40 and SNP41) have a minor allele frequency about this value; however, this frequency varies in different Asian populations. We observed a significant inconsistency between the values of a minor allele frequency for SNP40 in Asians from the database and literature, respectively. In the database, this value (0.41) is about 3–10 times higher than that reported in the literature (0.043–0.13).

Table 3: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Caucasians from databases and literature

Gene	SNP	Reference population	Database or reference	Data from a database or reference		Caucasians ^a		
				Frequency ^b	n	Frequency	n	
APOE	1	American Caucasians	[53]	0.33	220	0.357	1,862	0.48
		American Caucasians	dbSNP	0.33	24	0.399	1,839	0.64
	3	Europeans from North Karelia, Finland	dbSNP	0.4	24	0.399	1,839	0.84
		European-Americans	dbSNP	0.12	24	0.149	1,858	0.91
	4	Caucasians ^c	HGVbase	0.16	152	0.149	1,858	0.81
		Europeans from North Karelia, Finland	dbSNP	0.23	24	0.149	1,858	0.37
		European-Americans	dbSNP	0.19	24	0.087	1,783	0.16
		Caucasians ^c	HGVbase	0.08	152	0.087	1,783	0.87
COL1 α 1	5	White women with Spanish ancestors	[54]	0.13	256	0.154	1,856	0.36
		American Caucasians	[31]	0.187	637	0.188	1,855	0.98
	6							
ER- α	9	Australian females	[55]	0.44	125	0.484	1,861	0.39
		Greek males	[56]	0.49	50	0.484	1,861	0.95
	10	American females	[57]	0.45	253	0.449	1,861	0.96
		Finnish females	[58]	0.42	322	0.449	1,861	0.38
	11	SANGER 12 DNAs of Caucasian origin	dbSNP	0.08	12	0.257	1,857	0.32
	12	Australian females	[55]	0.23	120	0.218	1,859	0.84
		Slovenian females	[59]	0.3	85	0.218	1,859	0.10
	13	Caucasians ^c	dbSNP	0.27	31	0.218	1,859	0.59
Caucasians ^c		HGVbase	0.08	42	0.098	1,863	1.00	
15	Caucasians ^c	dbSNP	0.18	31	0.203	1,863	0.90	
IL-6	16	Caucasians	dbSNP	0.5	31	0.434	1,860	0.58
		English	[60]	0.57	2,560	0.434	1,860	<0.01
		Italians	[61]	0.45	183	0.434	1,860	0.74
		Spaniards	[62]	0.56	118	0.434	1,860	0.01
		Finns	HGVbase	0.46	400	0.434	1,860	0.37
		Finns	HGVbase	0.48	50	0.434	1,860	0.62
		Finns	HGVbase	0.39	42	0.434	1,860	0.68
		Finns	[63]	0.45	400	0.434	1,860	0.60
LEPR	18	English males	HGVbase	0.27	322	0.290	1,839	0.41
		Caucasians ^c	[64]	0.28	56	0.290	1,839	0.98
		Swedish males	[65]	0.24	284	0.290	1,839	0.09
	19	Swedish males	[65]	0.15	284	0.182	1,776	0.21
		Caucasians ^c	[64]	0.38	56	0.353	1,860	0.79
	20	Caucasians ^c	dbSNP	0.4	42	0.353	1,860	0.64
		English males	HGVbase	0.36	322	0.353	1,860	0.77
TGF- β 1	26	Caucasians ^c	dbSNP	0.43	31	0.311	1,840	0.21
		Caucasians ^c	HGVbase	0.26	42	0.311	1,840	0.59
	27	Caucasians ^c	[66]	0.02	304	0.021	1,863	0.98
		Danes	[66]	0.027	302	0.008	190	0.33
	29	Caucasian ^c female DZ twins	[67]	0.25	1,802	0.274	1,852	0.08
TNFR2	31	UK Caucasians	[68]	0.23	192	0.198	1,842	0.39
		Germans	[69]	0.25	94	0.198	1,842	0.33
	Australians	HGVbase	0.26	197	0.198	1,842	0.05	
UCP3	32	American Caucasians	[70]	0.21	56	0.265	1,767	0.49
		Danes	[71]	0.27	857	0.265	1,767	0.85
		French	[72]	0.22	894	0.265	1,767	0.01
	33	Caucasians ^c	[73]	0.01	165	0.000	295	0.44
		American Caucasians	[70]	0.2	58	0.258	1,862	0.38
	34	American Caucasians	[70]	0.2	58	0.258	1,862	0.38
		Caucasians ^c	[74]	0.5	501	0.447	1,863	0.04

Table 3: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Caucasians from databases and literature (Continued)

	36	Caucasians ^c	[73]	0.02	165	0.000	295	0.05
VDR	38	American Caucasians	[75]	0.42	49	0.373	1,861	0.61
		English	[76]	0.4	241	0.373	1,861	0.47
		Australians	[77]	0.35	43	0.373	1,861	0.87
	40	Caucasians ^c	dbSNP	0.42	107	0.419	1,860	0.94
		English females	[76]	0.44	241	0.419	1,860	0.61
		Australian males	[77]	0.47	39	0.419	1,860	0.66
		Swiss	[78]	0.39	197	0.419	1,860	0.49
		French infants	[79]	0.38	589	0.419	1,860	0.07
	41	Caucasians ^c	dbSNP	0.48	31	0.408	1,859	0.51
		French	[80]	0.35	99	0.408	1,859	0.33
		Australians	[81]	0.41	68	0.408	1,859	0.94

For gene abbreviations and SNP designations see Table 1. ^a Present study. ^b Frequency of the allele corresponding to a minor allele in the present study. ^c Caucasians with unidentified ethnic background.

Table 4: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Africans/African- Americans from databases and literature

Gene	SNP	Reference population	Database or references	Data from a database or reference		Caucasians ^a		P
				Frequency ^b	n	Frequency	n	
APOE	2	African-Americans	dbSNP	0.48	24	0.399	1,839	0.52
	3	African-Americans	dbSNP	0.10	24	0.149	1,858	0.56
	4	African-Americans	dbSNP	0.04	24	0.087	1,783	0.25
ER-α	11	SANGER 12 DNAs of African-American origin	dbSNP	0.08	12	0.257	1,857	0.32
	12	African/African-Americans	dbSNP	0.27	24	0.218	1,859	0.61
	13	TSC panel of African-Americans	dbSNP	0.16	40	0.098	1,863	0.27
	15	African/African-Americans	dbSNP	0.28	23	0.203	1,863	0.42
IL-6	16	African/African-Americans	dbSNP	0.96	24	0.434	1,860	<0.01
LEPR	20	TSC panel of African-Americans	dbSNP	0.45	42	0.353	1,860	0.25
TGF-β1	26	TSC panel of African-Americans	TSC	0.27	42	0.311	1,840	0.61
		African/African-Americans	dbSNP	0.15	24	0.311	1,840	0.07
UCP3	32	African-American females	[70]	0.15	57	0.265	1,767	0.07
	33	African-Americans	[73]	0.08	18	0.000	295	0.06
	34	African-American females	[70]	0.45	63	0.258	1,862	<0.01
	35	African-American females	[70]	0.247	73	0.447	1,863	<0.01
		African/African-Americans	[74]	0.20	276	0.447	1,863	<0.01
	36	African-Americans	[70]	0.00	18	0.000	295	10.00
VDR	38	African-American males	[77]	0.244	37	0.373	1,861	0.15
	40	African-American males	[77]	0.312	32	0.419	1,860	0.30
		Nigerians	[82]	0.37	93	0.419	1,860	0.41
	41	African/African-Americans	dbSNP	0.35	24	0.408	1,859	0.71
	African-American males	[77]	0.244	39	0.408	1,859	0.06	

For gene abbreviations and SNP designations see Table 1. ^a Present study. ^b Frequency of the allele corresponding to a minor allele in the present study.

Table 5: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Asian populations from databases and literature

Gene	SNP	Reference population	Database or reference	Data from a database or reference		Caucasians ^a		P
				Frequency ^b	n	Frequency	n	
ER- α	9	Japanese	[83]	0.412	200	0.484	1,861	0.06
		Thai females	[84]	0.337	129	0.484	1,861	<0.01
	10	Korean males	[85]	0.39	219	0.449	1,861	0.11
		Korean females	[86]	0.415	248	0.449	1,861	0.35
		Japanese children	[87]	0.431	102	0.449	1,861	0.80
		Taiwanese	[88]	0.39	246	0.449	1,861	0.09
		Thai females	[89]	0.414	134	0.449	1,861	0.49
	11	SANGER 12 DNAs of Asian origin	dbSNP	0.36	12	0.257	1,857	0.49
	12	Japanese	[83]	0.483	200	0.218	1,859	<0.01
		Japanese females	[90]	0.481	306	0.218	1,859	<0.01
	13	Japanese	JSNP	0.34	744	0.098	1,863	<0.01
		TSC panel of unrelated Asians	TSC	0.27	32	0.098	1,863	0.01
	14	Japanese	JSNP	0.296	752	0.119	1,861	<0.01
	15	Japanese	JSNP	0.18	741	0.203	1,863	0.21
		Japanese	[83]	0.21	200	0.203	1,863	0.88
IL-6	16	Chinese	[31]	0.997	147	0.434	1,860	<0.01
		Chinese males	[31]	0.998	259	0.434	1,860	<0.01
		Japanese	[31]	1.000	388	0.434	1,860	<0.01
LEPR	20	TSC panel of unrelated Asians	dbSNP	0.13	42	0.353	1,860	<0.01
		Japanese	JSNP	0.15	748	0.353	1,860	<0.01
PTHR1	22	Japanese	JSNP	0.43	747	0.398	1,861	0.14
	23	Japanese	JSNP	0.43	748	0.374	1,863	0.01
	24	Japanese	JSNP	0.44	750	0.397	1,855	0.05
	25	Japanese	JSNP	0.43	746	0.380	1,851	0.02
TGF- β 1	26	TSC panel of unrelated Asians	dbSNP	0.47	36	0.311	1,840	0.06
TNFR2	31	Japanese	[91]	0.113	265	0.198	1,842	<0.01
		Thai	[92]	0.127	201	0.198	1,842	0.02
UCP3	33	Asians ^c	[73]	0.00	11	0.000	295	10.00
	35	Japanese	JSNP	0.45	750	0.447	1,863	0.92
	36	Asians ^c	[73]	0.04	11	0.000	295	0.54
VDR	37	Japanese	JSNP	0.23	750	0.281	1,861	0.01
	38	Japanese	[93]	0.37	195	0.373	1,861	0.99
	40	SANGER 12 DNAs of Asian origin	dbSNP	0.41	12	0.419	1,860	0.77
		Chinese females (Han nationality)	[94]	0.043	162	0.419	1,860	<0.01
		Taiwanese	[95]	0.083	90	0.419	1,860	<0.01
		Japanese	[93]	0.13	195	0.419	1,860	0.02
		Koreans	[96]	0.055	211	0.419	1,860	<0.01
		Thai females	[97]	0.107	84	0.419	1,860	0.01
	41	Chinese (Han nationality)	[98]	0.05	223	0.408	1,859	<0.01
		Japanese females	[99]	0.135	119	0.408	1,859	<0.01
		Koreans	[96]	0.055	120	0.408	1,859	<0.01

For gene abbreviations and SNP designations see Table 1. ^a Present study. ^b Frequency of the allele corresponding to a minor allele in the present study. ^c Asians with unidentified ethnic background.

A comparison of our Caucasian sample and the Asian populations revealed significant differences in allele frequencies of 13 SNPs located in the 6 genes (Table 5). Another SNP representing the TGF-β1 gene (SNP26) indicated the nearly significant difference ($P = 0.06$). No such differences were observed only in the UCP3 gene.

Hispanic and Pacific Rim populations

These populations had the poorest data on the SNPs from the databases and literature as compared to the corre-

sponding data from the present study. We found the data on only seven SNPs of the five candidate genes for Hispanics (Table 6) and five SNPs of the four genes for Pacific Rim populations (Table 7). SNP33 and SNP36 for the UCP3 gene were virtually monomorphic in Hispanics, similar to that in our Caucasian sample, while only one significant difference in allele frequencies was observed at SNP16 between these ethnic populations among the seven compared SNPs (Table 6).

Table 6: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Hispanics from databases and literature

Gene	SNP	Database or references	Data from a database or reference		Caucasians ^a		P
			Frequency ^b	n	Frequency	n	
ER-α	12	dbSNP	0.239	23	0.218	1,859	1.00
	15	dbSNP	0.196	23	0.203	1,863	1.00
IL-6	16	dbSNP	0.795	22	0.434	1,860	<0.01
TGF-β1	26	dbSNP	0.413	23	0.311	1,840	0.36
UCP3	33	[73]	0.00	27	0.000	295	1.00
	36	[73]	0.02	27	0.000	295	1.00
VDR	41	dbSNP	0.348	23	0.408	1,859	0.83

For gene abbreviations and SNP designations see Table 1. ^aPresent study. ^bFrequency of the allele corresponding to a minor allele in the present study.

Table 7: Comparison of SNP allele frequencies of the studied Caucasian sample with the data on Pacific Rim populations from databases

Gene	SNP	Database	Data from a database or reference		Caucasians ^a		P
			Frequency ^b	n	Frequency	n	
ER-α	12	dbSNP	0.478	23	0.218	1,859	0.02
	15	dbSNP	0.326	23	0.203	1,863	0.19
IL-6	16	dbSNP	0.957	23	0.434	1,860	<0.01
TGF-β1	26	dbSNP	0.458	24	0.311	1,840	0.19
VDR	41	dbSNP	0.13	23	0.408	1,859	<0.01

For gene abbreviations and SNP designations see Table 1. ^aPresent study. ^bFrequency of the allele corresponding to a minor allele in the present study.

No data on the SNPs of the 10 candidate genes for obesity and/or osteoporosis were found for Pacific Rim populations in the available literature. All the data presented in Table 7 were obtained from the dbSNP database. Three SNPs out of the five differed significantly in the allele frequencies between Pacific Rim populations and the Caucasian sample in our study.

Some new candidate SNPs for genetic studies of obesity and osteoporosis

In this study, we identified a number of SNPs, which may be suitable for genetic studies of obesity and osteoporosis. In particular, the six polymorphic SNPs with minor allele frequency ≥ 0.1 , which were firstly determined in Caucasians (Table 2), may be used in population association studies. A number of SNPs with significant variation in allele frequencies in populations of different ethnicity may be appropriate for studying a genetic basis of between-ethnic differences in the rates of obesity and/or osteoporosis. Examples are SNP34, SNP35, and SNP32 for Caucasians and African-Americans (Table 4) and several SNPs located in the ER- α , LEPR, PTHR1, TNFR2, and VDR genes for Caucasians and Asians (Table 5). Some of the SNPs may also be candidates for such studies, if their allele frequencies reported to the databases are validated by using sufficiently large sample sizes (e.g., SNP26 in Asians).

Discussion

Ethnic variation of SNP allele frequencies of the candidate genes for obesity and/or osteoporosis

We observed significant differences in allele frequencies of some studied SNPs for Caucasians (SNP16, SNP35, SNP32, and SNP36, Table 3) between our results and respective data from the databases and literature. This may be due to various factors, such as population admixture and relatively smaller sample size in the other studies [34]. For example, the sample in the present study consisted of the subjects of various ethnic backgrounds (German, French, Dutch, Swedish, some Portuguese and Italian backgrounds) and the relative proportions of these ethnic groups are not ascertained. In the mixed samples from other studies, these proportions may be different that thus influence the allele frequencies to a larger or lesser extent (e.g., SNP35 and SNP36, Table 3). Likewise, monoethnic samples may have allele frequencies significantly different from those in the mixed samples (e.g., SNP16 in English and Spaniards, SNP32 in French, Table 3). Our results showed that major ethnic groups have significantly different allele distribution at some SNP markers of candidate genes for complex disorders. The largest differences were observed between Asian and Caucasian populations, namely at 13 SNPs out of 22 compared (Table 5). Some genes indicated the significant differences at most SNPs compared (e.g., PTHR1 between Japanese

and Caucasians, Table 5). These results are consistent with the previously reported data about high differentiation between Asians and Caucasians at candidate genes for bone mass [35,36].

Another important finding is the observed significant differences between Caucasian and Pacific Rim populations in SNP allele frequencies of three important candidate genes for osteoporosis, ER- α , IL-6, and VDR (Table 7). Interestingly, the respective SNP allele frequencies in Asian and Pacific Rim populations have very similar values (Tables 5 and 7). This may suggest that the Pacific Rim sample from the database has mainly Asian ancestry [37] or, perhaps there are some implications for an evolutionary history of these populations [38]. In any event, much more comprehensive data should be obtained to clarify these issues.

There are abundant data about significant between-ethnic differentiation at loci, which may underlie complex diseases, including obesity and osteoporosis (see, e.g., [36,39-41]). Given that there is well-known different incidence of these disorders in various ethnic groups (e.g., [42,43]), such genetic differentiation may have an important implication for studying a genetic basis of ethnicity-specific definitions of obesity and osteoporosis.

Another important application of the SNP markers with high large between-ethnic and small within-ethnic differences is MALD studies of complex diseases in the admixed populations with known parental ethnicities [33,44]. Our results showed that such SNPs exist in the candidate genes for obesity and osteoporosis. However, the large-scale screening of the candidate genes is necessary to identify such markers with sufficient density.

The results of our study suggest that ethnic heterogeneity of large samples may notably affect the observed allele frequencies. It is supported by the fact that several SNP allele frequencies of some sufficiently large monoethnic samples (e.g., SNP16 in English, SNP32 in French) significantly differ from those determined in our sample of mixed Caucasians (Table 3).

SNP databases: current limitations for studying complex diseases

Our analyses showed that the SNP databases in their current status might have some limitations for studies of complex disorders, especially in different ethnic groups, due to incomplete and/or uneven representation of SNPs and/or candidate genes in these groups. As indicated above, of the ten candidate genes examined here, only four have corresponding but incomplete SNP data in the databases for all the major ethnic groups and may be used in the comparative studies of obesity and/or osteoporosis

in the populations of different ethnicity. The SNP data in the databases for the other six genes (60% examined) need substantial updating. To do this, large-scale studies should be performed for the other major ethnic populations. Given that many complex diseases have different rates in different ethnic groups, the extensive volume of the SNP data needs to be updated and validated. This conclusion essentially corresponds to the recently reported results of the SNP databases evaluation regarding their use for whole-genome association studies in humans [21]. Our study provides an example showing the incompleteness of the SNP data in the current databases for studying complex disorders with an ethnic-dependent background.

Another important problem is inconsistency of the data for some SNP markers between either the different databases or databases and literature. It makes selection of a right SNP for a study rather difficult. For example, a SNP is usually considered to be appropriate for association or linkage studies, if its minor allele frequency is ≥ 0.1 [45,46]. Based on the data of SNP11 from the dbSNP database (Table 3), this SNP is hardly appropriate for the studies in Caucasians, because its minor allele frequency is 0.08. In fact, as determined in the present study, its actual frequency is 0.257 that makes this SNP suitable for the population association and linkage research.

In terms of well-known ethnic differences in incidence of some complex diseases, the discrepancies in the SNP database data may yield wrong conclusion about suitability of particular SNPs for studying genetic basis of these differences. For example, as was mentioned above, SNP26 from the TGF- $\beta 1$ gene has the reported minor allele frequency 0.43 (dbSNP) and 0.26 (HGvbase) for Caucasians, 0.27 (TSC) and 0.15 (dbSNP) for African-Americans, and 0.47 (dbSNP) for Asians (Tables 3,4,5). This gene is a candidate for bone mineral density [47] and, thus, may contribute to the ethnic differences in bone mass and osteoporosis. However, from the above data on the SNP allele frequencies, it is impossible to infer whether this particular SNP may be related to these ethnic differences in bone mass, because none of the values significantly differ from any other due to small sample size and, accordingly, limited statistical power.

Large discrepancies in some allele frequency data between the different databases (e.g., SNP4 for Caucasians) are likely due to small sample sizes and respective large sampling errors of the estimates. The sample sizes in the SNP databases are usually smaller than 50. With such a sample size, a relative sampling error of the allele frequency estimates ranges from 0.12 to 0.67 (fig. 1). In this term, the estimates of the SNP allele frequencies in the present study with the sample size of 1,873 are far more reliable than the respective data from the most databases. Estima-




Figure 1
Relative sampling error of different minor allele frequencies under various sample sizes. The relative sampling error is obtained as $\sqrt{\text{Var}(\hat{p})} / \hat{p}$, where $\text{Var}(\hat{p})$ is the variance of estimated allele frequencies and is computed as $\text{Var}(\hat{p}) = \frac{1}{2n} p(1-p)$ [51]. We plotted relative sampling errors of estimated allele frequencies vs. sample sizes.

tion of allele frequencies in small samples is notably affected by heterogeneity of the samples. Given that information about an ethnic background of the samples in the SNP databases is usually scarce, the allele frequency estimates in the databases may be significantly biased due to the ethnic admixture.

The small size of the samples in the databases makes it difficult to powerfully test differences in SNP allele frequencies (Δf) between the data from various sources, especially if absolute values of these differences are not large. Fig. 2 illustrates a statistical power of such estimation. Given two samples of size 50 each, the probability to correctly determine $\Delta f = 0.1$ is only 27% (fig. 2A). Having a sample size of 1,800 (similar to ours) increases this probability only up to 40% (fig. 2B). Even a 3-fold difference in the absolute values (e.g., SNP11, Table 4) is not statistically significant due to the insufficient sample size in the databases. These examples suggest that some differences in SNP allele frequency data in the public databases may exist due to improper sampling or too small sample sizes [34]. These inconsistencies may be somewhat reduced by pooling the samples. However, this should be done with caution, because pooling data from small samples with a particular genetic background may introduce a large bias to the new values.




Figure 2
Power to detect differences in allele frequencies between two samples at $P = 0.05$. Since the power depends on the sample sizes of the two populations, we fixed one population at a sample size of n , while allowing another population to vary in sample sizes. Δf is a frequency difference between the sample of given size n and the samples of various sizes. A – $n = 50$; B – $n = 1,800$.

Furthermore, in many cases, even after pooling, the sample size may remain too small to gain sufficient reliability of the newly obtained allele frequencies.

The SNP data in literature may somewhat supplement those from the databases. They are sometimes obtained with larger samples and, therefore, have smaller sampling error. However, they are not systematized, which makes their collection and use difficult.

In conclusion, our study demonstrated that, although a large volume of SNP data is available in public databases and literature, a great portion of these data needs comprehensive updating and validating in order to be appropriate for genetic studies of complex disorders, such as obesity and/or osteoporosis. Such large-scale studies of the disease-associated SNPs in various ethnicities may provide important insights into the evolutionary history of human populations as well as in etiology of these diseases.

Methods

Subjects

All the study subjects came from ongoing genetic studies of complex traits that have been approved by the Creighton University Institutional Review Board. All the subjects were Caucasians of western or northern European origin (German, French, Dutch, Swedish), but some had Portuguese and Italian backgrounds. We have recruited

405 nuclear families, each composed of both parents and at least one child. The total sample size was 1,873 subjects, including 840 parents and 1,133 children. All individuals volunteered to participate in the research and signed informed-consent documents before entering the project.

The study SNPs

The candidate SNPs and genes for the present study were selected from publicly available sources based on some of the following criteria: 1) functional relevance and importance for obesity and/or osteoporosis; 2) degree of heterozygosity, i.e., allele frequencies, as reported in literature or databases; 3) position in or around the genes; and 4) their use in previous genetic epidemiology studies. After searching public SNP databases (dbSNP, JSNP, HGVbase, and TSC) and literature, we chose 41 SNPs in 10 genes shown to be associated with obesity and/or osteoporosis. The information about the SNPs is given in Table 1. Among them, SNP27 is an insertion/deletion polymorphism of a cytosine (+/- C) and the others are nucleotide substitutions.

SNP genotyping

DNA was extracted from whole blood using a commercial isolation kit (Gentra Systems, Minneapolis, MN, USA) following the procedure recommended by the manufacturer. The genotyping involved a polymerase chain reaction (PCR) and invader assay (Third Wave Technology,

Madison, WI, USA) and was essentially the same for all the SNPs. The PCR mix consisted of 35 ng genomic DNA, 0.2 mM each dNTP, 1 × PCR buffer and 1.5 mM MgCl₂, 0.4 μM each primer, and 0.35 units of Taq polymerase (ABI Applied Biosystems, Foster City, CA, USA) in a total 10 μl reaction volume. The PCR primers for all SNPs are given in Table 1. The PCR was performed on PE9700 Thermal Cycler (Perkin Elmer Cetus, Norwalk, CT) using the following profile: 95 °C for 5 min, 30 cycles of 94 °C for 1 min, 50 °C for 1 min, 72 °C for 1 min, and then 72 °C for 5 min. After the amplification, the product was diluted 1:20 in nuclease-free water. The invader reaction was carried out in 7.5 μl reaction volume containing 3.75 μl diluted PCR product, 1.5 μl probe mix, 1.75 μl Cleavase FRET mix, and 0.5 μl Cleavase enzyme/MgCl₂ solution (Third Wave Technology). The reaction mix was overlaid by 15 μl mineral oil and denatured at 95 °C for 5 min, and then incubated at 63 °C for 20 min on PE9700 Thermal Cycler. After the incubation, the fluorescence intensity for both colors (FAM dye and Red dye) was measured by Cytofluor 4000 (ABI). The data were then loaded to the Invader Analyzer software (Third Wave Technology), and the genotype for every sample was called according to the ratio of the fluorescence intensity of the two dyes.

We initially genotyped all the 41 SNPs in a random sample of 190 to 380 subjects to find SNPs with minor allele frequencies <0.01. Those were then excluded from the further genotyping. Finally, 33 SNPs were genotyped for the whole sample.

Data analysis

PedCheck software [48] was used to verify the accuracy of SNP genotyping in reference to Mendelian inheritance of the alleles within each family. For the 29 polymorphic SNPs the allele frequencies in the nuclear families were estimated by maximum-likelihood method [49] implemented in SOLAR <http://www.sfbr.org/sfbr/public/software/solar>. This method uses all available marker information by accounting the dependence between relatives. Differences in allele frequencies of each SNP among various populations were tested using the χ^2 test or Fisher's exact two-tailed test as implemented in Advisor software [50]. SNPs with minor allele frequency less than 0.01 were considered to be monomorphic.

Authors' contributions

VD combined results of the analysis and drafted the manuscript. JRL and DHX performed the literature search, SNP genotyping, and statistical analysis. PYL carried out the statistical analysis. LJZ, HS, YYZ, YJL, SA, and PX performed the SNP genotyping. RRR and HWD conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Links

Database of Single Nucleotide Polymorphism, dbSNP <http://www.ncbi.nlm.nih.gov/SNP>

Japanese Single Nucleotide Polymorphisms, JSNP <http://snp.ims.u-tokyo.ac.jp>

Human Genome Variation Database, HGVbase <http://hgibase.cgb.ki.se/>

The SNP Consortium, TSC http://snp.cshl.org/allele_frequency_project/

Acknowledgments

The investigators were partially supported by grants from Health Future Foundation of the USA, the National Institutes of Health (K01 AR02170-01, R01 GM60402-01A1), the State of Nebraska Cancer and Smoking Related Disease Research Program, and US Department of Energy (DE-FG03-00ER63000/A00). The study also benefited from support (to HWD) of Hunan Province Special Professor Start-up Fund (25000612), Chinese National Science Foundation (CNSF) Outstanding Young Scientist Award (30025025), CNSF Grant (30170504), Seed Fund from the Ministry of Education of P. R. China (25000106), and a key project grant from the Ministry of Education of P. R. China. We thank all the study subjects for volunteering to participate in the study.

References

1. Taylor JG, Choi EH, Foster CB, Chanock SJ: **Using genetic variation to study human disease.** *Trends Mol Med* 2001, **7**:507-512.
2. Brookes AJ: **The essence of SNPs.** *Gene* 1999, **234**:177-186.
3. Miller RD, Kwok PY: **The birth and death of human single-nucleotide polymorphisms: new experimental evidence and implications for human history and medicine.** *Hum Mol Genet* 2001, **10**:2195-2198.
4. Gray IC, Campbell DA, Spurr NK: **Single nucleotide polymorphisms as tools in human genetics.** *Hum Mol Genet* 2000, **9**:2403-2408.
5. Lin RCY, Wang XL, Dalziel B, Caterson ID, Morris BJ: **Association of obesity, but not diabetes or hypertension, with glucocorticoid receptor N363S variant.** *Obes Res* 2003, **11**:802-808.
6. Tamura K, Suzuki M, Arakawa H, Tokuyama K, Morikawa A: **Linkage and association studies of STAT6 gene polymorphisms and allergic diseases.** *Int Arch Allergy Immunol* 2003, **131**:33-38.
7. Yamada Y, Ando F, Niino N, Shimokata H: **Association of polymorphisms of interleukin-6, osteocalcin, and vitamin D receptor genes, alone or in combination, with bone mineral density in community-dwelling Japanese women and men.** *J Clin Endocrinol Metab* 2003, **88**:3372-3378.
8. Daly MJ, Rioux JD, Schaffner SF, Hudson TJ, Lander ES: **High-resolution haplotype structure in the human genome.** *Nat Genet* 2001, **29**:229-232.
9. Jeffreys AJ, Kauppi L, Neumann R: **Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex.** *Nat Genet* 2001, **29**:217-222.
10. Reich DE, Cargill M, Bolk S, Ireland J, Sabeti PC, Richter DJ, Lavery T, Kouyoumjian R, Farhadian SF, Ward R, Lander ES: **Linkage disequilibrium in the human genome.** *Nature* 2001, **411**:199-204.
11. Patil N, Berno AJ, Hinds DA, Barrett WA, Doshi JM, Hacker CR, Kautzer CR, Lee DH, Marjoribanks C, McDonough DP, Nguyen BT, Norris MC, Sheehan JB, Shen N, Stern D, Stokowski RP, Thomas DJ, Trulsson MO, Vyas KR, Frazer KA, Fodor SP, Cox DR: **Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21.** *Science* 2001, **294**:1719-1723.
12. Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D: **The structure of haplotype blocks in the human genome.** *Science* 2002, **296**:2225-2229.

13. Phillips MS, Lawrence R, Sachidanandam R, Morris AP, Balding DJ, Donaldson MA, Studebaker JF, Ankeney WM, Alfisi SV, Kuo FS, Camisa AL, Pazorov V, Scott KE, Carey BJ, Faith J, Katari G, Bhatti HA, Cyr JM, Derohannessian V, Elosua C, Forman AM, Grecco NM, Hock CR, Kuebler JM, Lathrop JA, Mockler MA, Nachtmann EP, Restine SL, Varde SA, Hozza MJ, Gelfand CA, Broxholme J, Abecasis GR, Boyce-Jacino MT, Cardon LR: **Chromosome-wide distribution of haplotype blocks and the role of recombination hot spots.** *Nat Genet* 2003, **33**:382-387.
14. Aerts J, Wetzels Y, Cohen N, Aerssens J: **Data mining of public SNP databases for the selection of intragenic SNPs.** *Hum Mutat* 2002, **20**:162-173.
15. Thorisson GA, Stein LD: **The SNP Consortium website: past, present and future.** *Nucleic Acids Res* 2003, **31**:124-127.
16. Hirakawa M, Tanaka T, Hashimoto Y, Kuroda M, Takagi T, Nakamura Y: **JSNP: a database of common gene variations in the Japanese population.** *Nucleic Acids Res* 2002, **30**:158-162.
17. Hirakawa M: **HOWDY: an integrated database system for human genome research.** *Nucleic Acids Res* 2002, **30**:152-157.
18. Marsh S, Kwok P, McLeod HL: **SNP databases and pharmacogenetics: great start, but a long way to go.** *Hum Mutat* 2002, **20**:174-179.
19. Reich DE, Gabriel SB, Altshuler D: **Quality and completeness of SNP databases.** *Nat Genet* 2003, **33**:457-458.
20. Marth G, Yeh R, Minton M, Donaldson R, Li Q, Duan S, Davenport R, Miller RD, Kwok PY: **Single-nucleotide polymorphisms in the public domain: how useful are they?** *Nat Genet* 2001, **27**:371-372.
21. Carlson CS, Eberle MA, Rieder MJ, Smith JD, Kruglyak L, Nickerson DA: **Additional SNPs and linkage-disequilibrium analyses are necessary for whole-genome association studies in humans.** *Nat Genet* 2003, **33**:518-521.
22. Kuczmarski RJ, Flegal KM, Campbell SM, Johnson CL: **Increasing prevalence of overweight among US adults. The National Health and Nutrition Examination Surveys, 1960 to 1991.** *JAMA* 1994, **272**:205-211.
23. Kopelman PG: **Obesity as a medical problem.** *Nature* 2000, **404**:635-643.
24. Spencer H, Kramer L: **NIH consensus conference: Osteoporosis, Factors contributing to osteoporosis.** *J Nutr* 1986, **116**:319-322.
25. Ray NF, Chan JK, Thamer M, Melton LJ III: **Medical expenditures for the treatment of osteoporotic fractures in the United States in 1995: report from the National Osteoporosis Foundation.** *J Bone Miner Res* 1997, **12**:24-35.
26. Goddard KA, Hopkins PJ, Hall JM, Witte JS: **Linkage disequilibrium and allele-frequency distributions for 114 single-nucleotide polymorphisms in five populations.** *Am J Hum Genet* 2000, **66**:216-234.
27. Liu YZ, Liu YJ, Recker RR, Deng HW: **Molecular studies of identification of genes for osteoporosis: the 2002 update.** *J Endocrinol* 2003, **177**:147-196.
28. Lim SK, Park YS, Park JM, Song YD, Lee EJ, Kim KR, Lee HC, Huh KB: **Lack of association between vitamin D receptor genotypes and osteoporosis in Koreans.** *J Clin Endocrinol Metab* 1995, **80**:3677-3681.
29. Nakajima T, Ota N, Shirai Y, Hata A, Yoshida H, Suzuki T, Hosoi T, Orimo H, Emi M: **Ethnic difference in contribution of Sp1 site variation of COL1A1 gene in genetic predisposition to osteoporosis.** *Calcif Tissue Int* 1999, **65**:352-353.
30. Morrison NA, Qi JC, Tokita A, Kelly PJ, Crofts L, Nguyen TV, Sambrook PN, Eisman JA: **Prediction of bone density from vitamin D receptor alleles.** *Nature* 1994, **367**:284-287.
31. Lei SF, Deng FY, Liu XH, Huang QR, Qin Y, Zhou Q, Jiang DK, Li YM, Mo XY, Liu MY, Chen XD, Wu XS, Shen H, Dvornyk V, Zhao L, Recker RR, Deng HW: **Polymorphisms of four bone mineral density candidate genes in Chinese populations and the comparison with the other populations of different ethnicity.** *J Bone Miner Metab* 2003, **21**:34-42.
32. Beavan S, Prentice A, Dibba B, Yan L, Coper C, Ralston SH: **Polymorphism of the collagen type I $\alpha 1$ gene and ethnic differences in hip-fracture rates.** *N Engl J Med* 1998, **339**:351-352.
33. Collins-Schramm HE, Phillips CM, Operario DJ, Lee JS, Weber JL, Hanson RL, Knowler WC, Cooper R, Li H, Seldin MF: **Ethnic-difference markers for use in mapping by admixture linkage disequilibrium.** *Am J Hum Genet* 2002, **70**:737-750.
34. B-Rao C: **Sample size considerations in genetic polymorphism studies.** *Hum Hered* 2001, **52**:191-200.
35. Han KO, Moon IG, Hwang CS, Choi JT, Yoon HK, Min HK, Han IK: **Lack of an intronic Sp1 binding-site polymorphism at the collagen type I $\alpha 1$ gene in healthy Korean women.** *Bone* 1999, **24**:135-137.
36. Dvornyk V, Liu H, Shen H, Lei SF, Zhao L, Huang QR, Qin Y, Jiang DK, Long J, Zhang Y, Gong G, Recker RR, Deng HW: **Differentiation of Caucasians and Chinese at bone mass candidate genes: implication for ethnic difference of bone mass.** *Ann Hum Genet* 2003, **67**:216-227.
37. Yao YG, Watkins WS, Zhang YP: **Evolutionary history of the mtDNA 9-bp deletion in Chinese populations and its relevance to the peopling of east and southeast Asia.** *Hum Genet* 2000, **107**:504-512.
38. Lum JK, Cann RL: **mtDNA lineage analyses: origins and migrations of Micronesians and Polynesians.** *Am J Phys Anthropol* 2000, **113**:151-168.
39. Padyukov L, Hahn-Zoric M, Lau YL, Hanson LA: **Different allelic frequencies of several cytokine genes in Hong Kong Chinese and Swedish Caucasians.** *Genes Immun* 2001, **2**:280-283.
40. Bridges SL Jr, Jenq G, Moran M, Kuffner T, Whitworth WC, McNicholl J: **Single-nucleotide polymorphisms in tumor necrosis factor receptor genes: definition of novel haplotypes and racial/ethnic differences.** *Arthritis Rheum* 2002, **46**:2045-2050.
41. Lamsis F, Flannery GR, White NG, Muratore R, Kaelan C, Mitchell RJ: **Alleles and haplotypes of tumor necrosis factor (TNF) α and β genes in three ethnic populations of Sulawesi Indonesia.** *Hum Biol* 2002, **74**:381-396.
42. Davis JW, Novotny R, Wasnich RD, Ross PD: **Ethnic, anthropometric, and lifestyle associations with regional variations in peak bone mass.** *Calcif Tissue Int* 1999, **65**:100-105.
43. Colin-Bell A, Adair LS, Popkin BM: **Ethnic differences in the association between body mass index and hypertension.** *Am J Epidemiol* 2002, **155**:346-353.
44. Collins-Schramm HE, Kittles RA, Operario DJ, Weber JL, Criswell LA, Cooper RS, Seldin MF: **Markers that discriminate between European and African ancestry show limited variation within Africa.** *Hum Genet* 2002, **111**:566-569.
45. Kaessmann H, Zollner S, Gustafsson AC, Wiebe V, Laan M, Lundberg J, Uhlen M, Paabo S: **Extensive linkage disequilibrium in small human populations in Eurasia.** *Am J Hum Genet* 2002, **70**:673-685.
46. Sasaki T, Tahira T, Suzuki A, Higasa K, Kukita Y, Baba S, Hayashi K: **Precise estimation of allele frequencies of single-nucleotide polymorphisms by a quantitative SSCP analysis of pooled DNA.** *Am J Hum Genet* 2001, **68**:214-218.
47. Audi L, Garcia-Ramirez M, Carrasosa A: **Genetic determinants of bone mass.** *Horm Res* 1999, **51**:105-123.
48. O'Connell JR, Weeks DE: **PedCheck: a program for identification of genotype incompatibilities in linkage analysis.** *Am J Hum Genet* 1998, **63**:259-266.
49. Boehnke M: **Allele frequency estimation from data on relatives.** *Am J Hum Genet* 1991, **48**:22-25.
50. Elashoff JD, Query N: **Advisor version 3.0 user's guide.** *Cork, Ireland, Statistical Solutions Ltd.* 1999.
51. Weir BS: *Genetic Data Analysis 2: Methods for Discrete Population Genetic Data* 2nd edition. *Sunderland, MA: Sinauer Associates, Inc.*; 1996.
52. Lund AM, Jensen BL, Nielsen LA, Skovby F: **Dental manifestations of osteogenesis imperfecta and abnormalities of collagen I metabolism.** *J Craniofac Genet Dev Biol* 1998, **18**:30-37.
53. Martin ER, Lai EH, Gilbert JR, Rogala AR, Afshari AJ, Riley J, Finch KL, Stevens JF, Livak KJ, Slotterbeck BD, Slifer SH, Warren LL, Conneally PM, Schmechel DE, Purvis I, Pericak-Vance MA, Roses AD, Vance JM: **SNPing away at complex diseases: analysis of single-nucleotide polymorphisms around APOE in Alzheimer disease.** *Am J Hum Genet* 2000, **67**:383-394.
54. Garcia-Giralt N, Nogues X, Enjuanes A, Puig J, Mellibovsky L, Bay-Jensen A, Carreras R, Balcells S, Diez-Perez A, Grinberg D: **Two new single-nucleotide polymorphisms in the COL1A1 upstream regulatory region and their relationship to bone mineral density.** *J Bone Miner Res* 2002, **17**:384-393.
55. Curran JE, Lea RA, Rutherford S, Weinstein SR, Griffiths LR: **Association of estrogen receptor and glucocorticoid receptor gene polymorphisms with sporadic breast cancer.** *Int J Cancer* 2001, **95**:271-275.

56. Evangelopoulos D, Alevizaki M, Lekakis J, Cimponeriu A, Papamichael C, Kominakis A, Kalofoutis A, Moutsatsou P: **Molecular analysis of the estrogen receptor α gene in men with coronary artery disease: association with disease status.** *Clin Chim Acta* 2003, **331**:37-44.
57. Willing M, Sowers MR, Aron D, Clark MK, Burns TL, Bunten C, Crutchfield M, D'Agostino D, Jannausch M: **Bone mineral density and its change in white women: estrogen and vitamin D receptor genotypes and their interaction.** *J Bone Miner Res* 1998, **13**:695-705.
58. Salmen T, Heikkinen AM, Mahonen A, Kroger H, Komulainen M, Saarikoski S, Honkanen R, Maenpää PH: **Early postmenopausal bone loss is associated with PvuII estrogen receptor gene polymorphism in Finnish women: effect of hormone replacement therapy.** *J Bone Miner Res* 2000, **15**:315-321.
59. Jurada S, Marc J, Prezelj J, Kocijancic A, Komel R: **Codon 325 sequence polymorphism of the estrogen receptor α gene and bone mineral density in postmenopausal women.** *J Steroid Biochem Mol Biol* 2001, **78**:15-20.
60. Humphries SE, Luong LA, Ogg MS, Hawe E, Miller GJ: **The interleukin-6 -174 G/C promoter polymorphism is associated with risk of coronary heart disease and systolic blood pressure in healthy men.** *Eur Heart J* 2001, **22**:2243-2252.
61. Flex A, Gaetani E, Pola R, Santoliquido A, Aloï F, Papaleo P, Dal Lago A, Pola E, Serricchio M, Tondi P, Pola P: **The -174 G/C polymorphism of the interleukin-6 gene promoter is associated with peripheral artery occlusive disease.** *Eur J Vasc Endovasc Surg* 2002, **24**:264-268.
62. Vozarova B, Fernandez-Real JM, Knowler WC, Gallart L, Hanson RL, Gruber JD, Ricart W, Vendrell J, Richart C, Tataranni PA, Wolford JK: **The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians.** *Hum Genet* 2003, **112**:409-413.
63. Hulkkonen J, Pertovaara M, Anttonen J, Pasternack A, Hurme M: **Elevated interleukin-6 plasma levels are regulated by the promoter region polymorphism of the IL6 gene in primary Sjogren's syndrome and correlate with the clinical manifestations of the disease.** *Rheumatology (Oxford)* 2001, **40**:656-661.
64. Thompson DB, Ravussin E, Bennett PH, Bogardus C: **Structure and sequence variation at the human leptin receptor gene in lean and obese Pima Indians.** *Hum Mol Genet* 1997, **6**:675-679.
65. Rosmond R, Chagnon YC, Holm G, Chagnon M, Perusse L, Lindell K, Carlsson B, Bouchard C, Bjorntorp P: **Hypertension in obesity and the leptin receptor gene locus.** *J Clin Endocrinol Metab* 2000, **85**:3126-3131.
66. Langdahl BL, Carstens M, Stenkjær L, Eriksen EF: **Polymorphisms in the transforming growth factor β 1 gene and osteoporosis.** *Bone* 2003, **32**:297-310.
67. Keen RW, Snieder H, Molloy H, Daniels J, Chiano M, Gibson F, Fairbairn L, Smith P, MacGregor AJ, Gewert D, Spector TD: **Evidence of association and linkage disequilibrium between a novel polymorphism in the transforming growth factor β 1 gene and hip bone mineral density: a study of female twins.** *Rheumatology* 2001, **40**:48-54.
68. Tsuchiya N, Kawasaki A, Tsao BP, Komata T, Grossman JM, Tokunaga K: **Analysis of the association of HLA-DRB1, TNF α promoter and TNFR2 (TNFRSF1B) polymorphisms with SLE using transmission disequilibrium test.** *Genes Immun* 2001, **2**:317-322.
69. Wiczorek S, Dahmen N, Jagiello P, Epplen JT, Gencik M: **Polymorphisms of the tumor necrosis factor receptors: no association with narcolepsy in German patients.** *J Mol Med* 2003, **81**:87-90.
70. Kimm SY, Glynn NW, Aston CE, Damcott CM, Poehlman ET, Daniels SR, Ferrell RE: **Racial differences in the relation between uncoupling protein genes and resting energy expenditure.** *Am J Clin Nutr* 2002, **75**:714-719.
71. Dalgaard LT, Sørensen TI, Drivsholm T, Borch-Johnsen K, Andersen T, Hansen T, Pedersen O: **A prevalent polymorphism in the promoter of the UCP3 gene and its relationship to body mass index and long term body weight change in the Danish population.** *J Clin Endocrinol Metab* 2001, **86**:1398-1402.
72. Meirhaeghe A, Amouyel P, Helbecque N, Cotel D, Otabe S, Froguel P, Vasseur F: **An uncoupling protein 3 gene polymorphism associated with a lower risk of developing Type II diabetes and with atherogenic lipid profile in a French cohort.** *Diabetologia* 2000, **43**:1424-1428.
73. Chung WK, Luke A, Cooper RS, Rotini C, Vidal-Puig A, Rosenbaum M, Chua M, Solanes G, Zheng M, Zhao L, LeDuc C, Eisberg A, Chu F, Murphy E, Schreier M, Aronne L, Caprio S, Kahle B, Gordon D, Leal SM, Goldsmith R, Andreu AL, Bruno C, DiMauro S, Leibel RL: **Genetic and physiologic analysis of the role of uncoupling protein 3 in human energy homeostasis.** *Diabetes* 1999, **48**:1890-1895.
74. Lanouette CM, Chagnon YC, Rice T, Perusse L, Muzzin P, Giacobino JP, Gagnon J, Wilmore JH, Leon AS, Skinner JS, Rao DC, Bouchard C: **Uncoupling protein 3 gene is associated with body composition changes with training in HERITAGE study.** *J Appl Physiol* 2002, **92**:1111-1118.
75. Chiu KC, Chuang LM, Yoon C: **The vitamin D receptor polymorphism in the translation initiation codon is a risk factor for insulin resistance in glucose tolerant Caucasians.** *BMC Med Genet* 2001, **2**:2.
76. Bretherton-Watt D, Given-Wilson R, Mansi JL, Thomas V, Carter N, Colston KW: **Vitamin D receptor gene polymorphisms are associated with breast cancer risk in a UK Caucasian population.** *Br J Cancer* 2001, **85**:171-175.
77. Bell NH, Morrison NA, Nguyen TV, Eisman J, Hollis BW: **Apal polymorphisms of the vitamin D receptor predict bone density of the lumbar spine and not racial difference in bone density in young men.** *J Lab Clin Med* 2001, **137**:133-140.
78. Ferrari SL, Rizzoli R, Slosman DO, Bonjour JP: **Do dietary calcium and age explain the controversy surrounding the relationship between bone mineral density and vitamin D receptor gene polymorphisms?** *J Bone Miner Res* 1998, **13**:363-370.
79. Suarez F, Zeghoud F, Rossignol C, Walrant O, Garabedian M: **Association between vitamin D receptor gene polymorphism and sex-dependent growth during the first two years of life.** *J Clin Endocrinol Metab* 1997, **82**:2966-2970.
80. Taverna MJ, Sola A, Guyot-Argenton C, Pacher N, Bruzzo F, Slama G, Reach G, Selam JL: **TaqI polymorphism of the vitamin D receptor and risk of severe diabetic retinopathy.** *Diabetologia* 2002, **45**:436-442.
81. Tao C, Yu T, Garnett S, Briody J, Knight J, Woodhead H, Cowell CT: **Vitamin D receptor alleles predict growth and bone density in girls.** *Arch Dis Child* 1998, **79**:488-493.
82. Fischer PR, Thacher TD, Pettifor JM, Jorde LB, Eccleshall TR, Feldman D: **Vitamin D receptor polymorphisms and nutritional rickets in Nigerian children.** *J Bone Miner Res* 2000, **15**:2206-2210.
83. Sasaki M, Tanaka Y, Kaneuchi M, Sakuragi N, Dahiya R: **Polymorphisms of estrogen receptor α gene in endometrial cancer.** *Biochem Biophys Res Commun* 2002, **297**:558-564.
84. Ongphiphadhanakul B, Chanprasertyothin S, Payattikul P, Saetung S, Piaseu N, Chailurkit L, Chansirikarn S, Puavilai G, Rajatanavin R: **Association of a T262C transition in exon I of estrogen receptor- α gene with skeletal responsiveness to estrogen in post-menopausal women.** *J Endocrinol Invest* 2001, **24**:749-755.
85. Koh JM, Kim DJ, Hong JS, Park JY, Lee KU, Kim SY, Kim GS: **Estrogen receptor α gene polymorphisms PvuII and XbaI influence association between leptin receptor gene polymorphism (Gln223Arg) and bone mineral density in young men.** *Eur J Endocrinol* 2002, **147**:777-783.
86. Han KO, Moon IG, Kang YS, Chung HY, Min HK, Han IK: **Nonassociation of estrogen receptor genotypes with bone mineral density and estrogen responsiveness to hormone replacement therapy in Korean postmenopausal women.** *J Clin Endocrinol Metab* 1997, **82**:991-995.
87. Kikuchi T, Hashimoto N, Kawasaki T, Uchiyama M: **Association of serum low-density lipoprotein metabolism with oestrogen receptor gene polymorphisms in healthy children.** *Acta Paediatr* 2000, **89**:42-45.
88. Lai IC, Liao DL, Bai YM, Lin CC, Yu SC, Chen JY, Wang YC: **Association study of the estrogen receptor polymorphisms with tardive dyskinesia in schizophrenia.** *Neuropsychobiology* 2002, **46**:173-175.
89. Ongphiphadhanakul B, Rajatanavin R, Chanprasertyothin S, Piaseu N, Chailurkit L, Sirisriro R, Komindr S: **Estrogen receptor gene polymorphism is associated with bone mineral density in premenopausal women but not in postmenopausal women.** *J Endocrinol Invest* 1998, **21**:487-493.
90. Hoshino S, Hosoi T, Miyao M, Shiraki M, Orimo H, Ouchi Y, Inoue S: **Identification of a novel polymorphism of estrogen receptor-**

- α gene that is associated with calcium excretion in urine. *J Bone Miner Metab* 2000, **18**:153-157.
91. Shibue T, Tsuchiya N, Komata T, Matsushita M, Shiota M, Ohashi J, Wakui M, Matsuta K, Tokunaga K: **Tumor necrosis factor α 5'-flanking region, tumor necrosis factor receptor II, and HLA-DRB1 polymorphisms in Japanese patients with rheumatoid arthritis.** *Arthritis Rheum* 2000, **43**:753-757.
 92. Hananantachai H, Patarapotikul J, Looareesuwan S, Ohashi J, Naka I, Tokunaga K: **Lack of association of -308A/G TNFA promoter and 196R/M TNFR2 polymorphisms with disease severity in Thai adult malaria patients.** *Am J Med Genet* 2001, **102**:391-392.
 93. Ban Y, Taniyama M, Ban Y: **Vitamin D receptor gene polymorphism is associated with Graves' disease in the Japanese population.** *J Clin Endocrinol Metab* 2000, **85**:4639-4643.
 94. Zhang H, Tao G, Wu Q, Liu J, Gao Y, Chen R, Leng X: **Vitamin D receptor gene polymorphism in postmenopausal women of the Han and Uygur nationalities in China.** *Chin Med J (Engl)* 2000, **113**:787-789.
 95. Huang CM, Wu MC, Wu JY, Tsai FJ: **Association of vitamin D receptor gene BsmI polymorphisms in Chinese patients with systemic lupus erythematosus.** *Lupus* 2002, **11**:31-34.
 96. Lee CK, Hong JS, Cho YS, Yoo B, Kim GS, Moon HB: **Lack of relationship between vitamin D receptor polymorphism and bone erosion in rheumatoid arthritis.** *J Korean Med Sci* 2001, **16**:188-192.
 97. Ongphiphadhanakul B, Rajatanavin R, Chanprasertyothin S, Chailurkit L, Piaseu N, Teerarungsikul K, Sirisriro R, Komindr S, Puavilai G: **Vitamin D receptor gene polymorphism is associated with urinary calcium excretion but not with bone mineral density in postmenopausal women.** *J Endocrinol Invest* 1997, **20**:592-596.
 98. Zhao J, Zhou X, Meng X, Liu G, Xing X, Liu H, Xu L: **Polymorphisms of vitamin D receptor gene and its association with bone mineral density and osteocalcin in Chinese.** *Chin Med J (Engl)* 1997, **110**:366-371.
 99. Iki M, Saito Y, Dohi Y, Kajita E, Nishino H, Yonemasu K, Kusaka Y: **Greater trunk muscle torque reduces postmenopausal bone loss at the spine independently of age, body size, and vitamin D receptor genotype in Japanese women.** *Calcif Tissue Int* 2002, **71**:300-307.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

